

A quantitative analysis to objectively appraise drought indicators and model drought impacts

S. Bachmair¹, C. Svensson², J. Hannaford², L.J. Barker², and K. Stahl¹

[1]{Hydrology, Faculty of Environment and Natural Resources, University of Freiburg, Freiburg, Germany}

[2]{Centre for Ecology and Hydrology, Wallingford, United Kingdom}

Correspondence to: S. Bachmair (sophie.bachmair@hydrology.uni-freiburg.de)

Abstract

Drought monitoring and early warning is an important measure to enhance resilience towards drought. While there are numerous operational systems using different drought indicators, there is no consensus on which indicator best represents drought impact occurrence for any given sector. Furthermore, thresholds are widely applied in these indicators but, to date, little empirical evidence exists as to which indicator thresholds trigger impacts on society, the economy, and ecosystems. The main obstacle for evaluating commonly used drought indicators is a lack of information on drought impacts. Our aim was therefore to exploit text-based data from the European Drought Impact report Inventory (EDII) to identify indicators which are meaningful for region-, sector-, and season-specific impact occurrence, and to empirically determine indicator thresholds. In addition, we tested the predictability of impact occurrence based on the best performing indicators. To achieve these aims we applied a correlation analysis and an ensemble regression tree approach, using Germany and the UK (the most data-rich countries in the EDII) as a testbed. As candidate indicators we chose two meteorological indicators (Standardized Precipitation Index (SPI) and Standardized Precipitation Evaporation Index (SPEI)) and two hydrological indicators (streamflow and groundwater level percentiles). The analysis revealed that accumulation periods of SPI and SPEI best linked to impact occurrence are longer for the UK compared with Germany, but there is variability within each country, among impact categories and, to some degree, seasons. The median of regression tree splitting values, which we regard as estimates of

1 thresholds of impact occurrence, was around -1 for SPI and SPEI in the UK; distinct
2 differences between northern/northeastern versus southern/central regions were found for
3 Germany. Predictions with the ensemble regression tree approach yielded reasonable results
4 for regions with good impact data coverage. The predictions also provided insights into the
5 EDII, in particular highlighting drought events where missing impact reports may reflect a
6 lack of recording rather than true absence of impacts. Overall, the presented quantitative
7 framework proved to be a useful tool for evaluating drought indicators, and to model impact
8 occurrence. In summary, this study demonstrates the information gain for drought monitoring
9 and early warning through impact data collection and analysis. It highlights the important role
10 that quantitative analysis with impacts data can have in providing “ground truth” for drought
11 indicators, alongside more traditional stakeholder-led approaches.

12

13 **1 Introduction**

14 Drought is less tangible than other natural hazards, such as earthquakes or floods, due to its
15 slow onset, insidious nature, and complex, often non-structural impacts (Gillette, 1950;
16 Wilhite et al., 2007). Nonetheless, drought is known to affect more people than any other
17 hazard, and to cause high economic loss (Loayza et al., 2012; Wilhite et al., 2007). While
18 droughts cannot be prevented, societal vulnerability can be reduced, with monitoring and
19 early warning (hereafter, M&EW) being one important measure to enhance drought
20 resilience. The aim of M&EW is to provide adequate and timely information on drought
21 conditions to enable people and organizations to be better prepared and react accordingly
22 (Svoboda et al., 2002; Wilhite and Svoboda, 2000). Such systems are usually based on several
23 drought indicators representing different domains of the hydrological cycle, i.e. indicators for
24 meteorological drought, soil moisture drought and vegetation stress, hydrological drought,
25 and groundwater drought.

26 A recent trend has been the design of “combined” or “multivariate” indicators consisting of a
27 blend of individual ones. The rationale behind the construction of blended indicators is that a
28 single indicator is not sufficient to adequately capture different types of drought, and the
29 corresponding multiplicity of drought impacts that differ markedly in response time (Hao and
30 Singh, 2015). There have been several studies assessing the link between indicators of
31 different types of droughts, e.g. between meteorological drought and streamflow, soil
32 moisture, or remotely sensed vegetation stress indicators (Haslinger et al., 2014; Ji and Peters,

1 2003; Martínez-Fernández et al., 2015; Vicente-Serrano and López-Moreno, 2005; Vicente-
2 Serrano et al., 2012). These are useful when there is an assumption that the lag between, say,
3 meteorological and hydrological drought represents the response time for impact occurrence
4 in, say, riverine ecosystems. Drought indicator choices can be substantiated by stakeholder
5 consultation or expert judgement, as has been implemented for the operational US Drought
6 Monitor (Svoboda et al., 2002). Similar initiatives have been developed in research project
7 settings in southwest Germany (Stölzle and Stahl, 2011) and Switzerland (Kruse et al., 2010).

8 However, while indicators representing different types of drought are commonly used as
9 proxies for impact occurrence, there is, to date, little empirical evidence as to which indicator
10 best represents drought impact occurrence for any given sector. Lackstrom et al. (2013)
11 identified an impact-driven perspective as the “missing piece” of drought monitoring; what is
12 of ultimate interest is knowledge of when and where a precipitation shortfall or low
13 streamflow or groundwater level will translate into impacts on society, the economy, and
14 ecosystems. A direct, empirical evaluation of drought indicators with impact information
15 would obviate the need for assumptions based on intercomparing different drought indicators.

16 Aside from identifying indicators important for drought impacts, there is a need for a better
17 understanding of the meaning of indicator thresholds used for drought declaration and as
18 triggers for management actions in drought plans. Such thresholds are mostly based on hazard
19 intensity classes corresponding to a certain frequency of occurrence, e.g. following the widely
20 accepted Standardized Precipitation Index scheme, with classes ranging from 0 to -0.99 (mild
21 drought), -1 to -1.49 (moderate drought), -1.5 to -2 (severe drought), and < -2 (extreme
22 drought) (McKee et al., 1993). The US Drought Monitor (USDMD) differentiates between five
23 drought severity classes based on several indicators and corresponding thresholds (Svoboda et
24 al., 2002). Different thresholds again are used for delineating alert classes of the Combined
25 Drought indicator of the European Drought Observatory (European Drought Observatory,
26 2013).

27 Common to all thresholds is that they are arbitrary cut-off points (e.g. McKee et al., 1993;
28 Svoboda et al., 2002). A survey among drought managers in the US on drought plans and
29 respective indicators and triggers revealed that there is large uncertainty in the selection of
30 thresholds, with one survey reply uncovering that most states selected their indicators “out of
31 a hat” without knowing whether they “worked” (Steinemann, 2014). There is currently no

1 consensus on appropriate drought indicators and thresholds meaningful for practitioners of
2 different sectors.

3 Regarding drought prediction, a substantial body of research has been dedicated to forecasting
4 drought indicators with sufficient lead time (e.g. Dutra et al., 2014; Mehta et al., 2014;
5 Trambauer et al., 2014; Wetterhall et al., 2015). However, while the models used for
6 forecasting may propagate the climate signal into soils and hydrology, they do not include a
7 further link to the tangible negative environmental and socio-economic impacts of a particular
8 drought. Models bridging the gap between drought indicators and impacts are rare. While
9 predictions of crop yield are more common (e.g. Hlavinka et al., 2009; Mavromatis, 2007;
10 Quiring and Papakryiakou, 2003), very few studies have tested approaches for modeling other
11 types of drought impacts such as wildfires, or impacts on public water supply or the energy
12 and industry sector (e.g. Blauhut et al. (2015), Stagge et al. (2014), Gudmundsson et al.
13 (2014), and Vicente-Serrano et al. (2012)). The complexity of processes and the
14 interconnectedness of the multitude of drought impacts, which may occur with much delay
15 and even outside of the hazard affected area (Logar and van den Bergh, 2013; Wilhite et al.,
16 2007), may be one reason why few drought impact models have been presented.

17 The most important obstacle, however, is a paucity of information on drought impacts.
18 Initiatives to rectify this include the US Drought Impact Reporter (DIR) (Wilhite et al., 2007),
19 and the more recently developed European Drought Impact report Inventory (EDII) (Stahl et
20 al., 2015a). Both provide text-based, categorized information on reported drought impacts.
21 The majority of impacts of the US DIR stem from online media clipping (Wilhite et al.,
22 2007), meaning that it can be used as a real-time monitoring tool. In contrast, the EDII is
23 designed as a research database with a focus on past drought events. Other potential sources
24 of drought impact data are reported crop yields, or losses assembled in the Emergency Events
25 Database EM-DAT (www.emdat.be) or by re-insurance companies. Nevertheless, crop yield
26 reductions may not necessarily be due to drought and loss data mostly provides aggregated
27 information on large events without details on the temporal and spatial evolution of impacts,
28 which is essential for empirically validating indicators and developing drought impact
29 models.

30 Only very few studies to date have exploited text-based impact datasets. Dieker et al. (2010)
31 qualitatively and quantitatively compared the USDM to impact data from the US DIR. Stagge
32 et al. (2014) and Blauhut et al. (2015) both worked with EDII data at the country- or macro-

1 region-scale across Europe, with impacts coded as a binary response variable (impact versus
2 no impact) to determine the likelihood of impact occurrence for different impact types.
3 Bachmair et al. (2015) also used EDII data to test the feasibility of evaluating drought
4 indicators with impacts at smaller spatial scales in Germany. As an extension to Stagge et al.
5 (2014) and Blauhut et al. (2015), they replaced the binary data with the number of impact
6 occurrences, thus providing a measure of impact severity. A correlation analysis and
7 extraction of indicator values concurrent with past impact onset showed variability in
8 indicator performance and onset thresholds at the sub-country scale and between drought
9 events. The effect of different impact categories or types was not assessed (Bachmair et al.,
10 2015).

11 Building on these previous efforts, the aim of this study is to exploit the EDII to link drought
12 indicators to impacts using quantitative methodologies. Germany (DE) and the UK were
13 selected as a test-bed, since they represent the countries with most impact data in the EDII
14 database, but the aim is to develop methods that can be extended to other geographical areas
15 in future applications. Specifically, the aims are to

- 16 • evaluate different drought indicators using text-based impact information to identify
17 indicators that are meaningful for region-, sector-, and season-specific impact occurrence,
- 18 • to empirically determine indicator thresholds representative for impact occurrence, as an
19 alternative to using the default, arbitrarily selected hazard class thresholds intrinsic to
20 indicators such as the SPI,
- 21 • to model impact occurrence via machine learning to assess the potential for predictive
22 purposes (i.e. predicting impacts based on indicators alone), and exploit the relationships
23 between indicators and text-based impact data to “backwards learn” about the nature of
24 the impact data itself.

25

26 **2 Data**

27 **2.1 Spatial and temporal resolution**

28 As temporal and spatial resolution of the drought indicator and impact data we selected
29 monthly time series for the period 1970-2012, aggregated at the NUTS1 level (level 1 of the
30 Nomenclature of Units for Territorial Statistics, a spatial unit used in the European Union).

1 NUTS1 regions represent major socio-economic regions. This level of spatial aggregation was
2 chosen because of a lack of sufficient data for analysis with finer-scale resolution. However,
3 studies have shown that drought signals typically cover areas larger than NUTS1 regions (e.g.
4 Hannaford et al. (2011)). In Germany NUTS1 regions correspond to the federal states. In the
5 UK there are 12 NUTS1 regions, in Germany 16 (see Table 1 for a list of NUTS1 regions
6 considered for analysis and abbreviations used in this study, and Figure 1 for the size of
7 NUTS1 regions). Note that two NUTS1 regions in the UK and three in Germany were
8 excluded from the analysis due to having insufficient impact data (see section 2.3 for details).

9 **2.2 Drought indicators**

10 As drought indicators we selected the Standardized Precipitation Index (SPI) (McKee et al.,
11 1993), the Standardized Precipitation Evaporation Index (SPEI) (Vicente-Serrano et al.,
12 2010), and streamflow percentiles (Q). In addition, groundwater level percentiles (G) were
13 included for Germany. For the SPI and SPEI, accumulation periods of 1-8, 12, and 24 months
14 were chosen. Gridded SPI and SPEI data were calculated based on E-OBS gridded data
15 (version 9.0; 0.25° regular spatial grid (Haylock et al., 2008)) using the R Package ‘SCI’
16 (Stagge et al., 2014). For the UK and Germany the underlying station density of the gridded
17 data is relatively high within Europe, and the dataset is based on more European observing
18 stations than in other European or global datasets (Haylock et al., 2008). The gamma
19 distribution was used for the computation of the SPIs and the generalized logistic distribution
20 for the SPEIs (reference period: 1971-2010). Potential evapotranspiration for the SPEI was
21 estimated using the Hargreaves method (Hargreaves, 1994). For each NUTS1 region, regional
22 averages of mean monthly SPI-*n* or SPEI-*n* were calculated. Here, *n* denotes the accumulation
23 period. The mean was chosen since Bachmair et al. (2015) found little differences between
24 the performance of different indicator metrics per spatial unit (e.g. mean vs. minimum, or 10th
25 percentile vs. percent area with SPI or SPEI below a threshold). The reference period for
26 calculation of streamflow percentiles is 1960-2012 in the UK, and 1970-2011 in Germany
27 (also for groundwater).

28 The monthly streamflow percentiles are based on monthly mean streamflows. In Germany
29 these are calculated from daily streamflow records for several gauging stations per federal
30 state; monthly groundwater percentiles come from weekly to monthly readings of
31 groundwater levels or spring discharge for several monitoring stations per state (data

1 provision by different agencies of the German federal states, see Kohn et al. (2014)). Many of
2 these stations are used for the federal states' hydrological forecasting systems and thus
3 represent stations with good data quality. Monthly streamflow records for the UK were taken
4 from daily river flow records held on the UK National River Flow Archive (NRFA)
5 (<http://nrfa.ceh.ac.uk/>). The UK Benchmark Network (Bradford and Marsh, 2003) of near-
6 natural catchments was used, alongside the network of sites used in the National Hydrological
7 Monitoring Programme (NHMP: <http://nrfa.ceh.ac.uk/nhmp>). No groundwater measurements
8 were used from the UK due to the limited number of NHMP borehole records available in
9 many NUTS regions, reflecting the concentration of productive aquifers in the south and east
10 of the country.

11 The streamflow gauging stations in the UK and Germany encompass both near-natural and
12 anthropogenically influenced streamflow records. Since different drought impacts may occur
13 in near-natural and regulated rivers it is necessary to have an indicator reflecting the drought
14 signal in both natural and managed systems. Figure 1 displays the spatial location of Q and G
15 measurement stations and the boundaries of the NUTS1 regions in the UK and Germany. The
16 number of stations per NUTS1 region is displayed in Table 1. Regional average mean
17 monthly Q and G values were calculated for each NUTS1 region, provided there was at least
18 one station with non-missing observations in the region. As further predictors that may
19 modify the drought indicators' power to explain drought impact occurrence we also selected
20 the month of impact occurrence (M) and the year of impact occurrence (Y). For this purpose
21 the series of months (1-12) was transformed into a sinusoidal curve shifted by four months
22 (peak in July and lowest value in January).

23 **2.3 Drought impacts**

24 Drought impact data come from the European Drought Impact report Inventory (EDII) (Stahl
25 et al., 2015a), which can be viewed online at <http://www.geo.uio.no/edc/droughtdb/> (data
26 extraction for this study: October 2014). The EDII defines a “drought impact” as a negative
27 environmental, economic or social effect experienced under drought conditions. Examples of
28 drought impacts are crop losses, water supply shortages and hosepipe bans, increased
29 mortality of aquatic species, reduced production at thermal or nuclear power plants due to a
30 lack of cooling water, or impaired navigability of streams, to name a few. Drought conditions
31 themselves (anomaly in precipitation, soil moisture, streamflow, groundwater levels etc.),

1 without a negative consequence or at least evoking serious concerns, are not considered an
2 impact. The source of EDII entries is text-based reports on drought impacts, e.g.
3 governmental or NGO reports, books, newspapers, digital media or scientific papers. Each
4 impact report in the EDII contains the following information: 1) a spatial reference (different
5 levels of geographical regionalization, including the European Union NUTS regions
6 standard), 2) a temporal reference (at least the year of occurrence), and 3) an assigned impact
7 category. The 15 categories, e.g. agriculture, water supply, etc., are shown in Figure 2. Each
8 category subsumes several impact type subcategories (see Stahl et al. (2015a) for details).

9 For the analysis the qualitative information on drought impacts was transformed into monthly
10 time series of number of drought impact occurrences per NUTS1 region. The same
11 methodology as in Bachmair et al. (2015) was applied during the conversion of a “drought
12 impact report” (EDII entry) into “drought impact occurrence” (hereafter termed I). In short,
13 this entails the following (see Bachmair et al. (2015) for details):

- 14 • Each impact report was assigned to a NUTS1 region. Impact reports with country-level
15 information only were omitted from the analysis. An impact report was converted into
16 several I if 1) the impact report stated impact occurrence in several NUTS1 regions or 2)
17 an impact fell into several impact subtypes.
- 18 • Each I is temporally referenced by specifying a start and end month. Impact reports only
19 stating the year of occurrence were omitted from the analysis. In case only the season was
20 provided in the report, we assumed the drought impact occurred during each month of this
21 season (winter= DJF, spring= MAM, summer= JJA, fall= SON).

22 For each NUTS1 region and month the total number of I was determined, hereafter termed N_I .
23 Table 1 shows the N_I per NUTS1 region included in the analysis, which sum up to 4551 N_I
24 (UK) and 1534 N_I (DE) in total for each country. Some analyses were undertaken for impacts
25 separated into the 15 impact categories. However, a different kind of split of the data was also
26 made, into two larger groups:

- 27 • hydrological drought impacts (I_h), i.e. impacts resulting from drought conditions of
28 surface waters or groundwater,
- 29 • impacts due to other types of drought (I_o), i.e. impacts associated with meteorological and
30 soil moisture drought and concurrent extremes (e.g. heat waves).

31 The differentiation between I_h and I_o is based on a keyword search of the impact description
32 field in the database and therefore does not strictly follow any impact category or impact

1 subtype. Examples of I_h include impaired navigability of streams, increased temperature in
2 surface waters negatively affecting aquatic species, drying up of reservoirs, or reduced fishery
3 production. I_o comprises most agricultural and forestry impacts, impacts on recreation or
4 human health, soil subsidence, or wildfire. Figure 2 shows the total number of I , I_h and I_o per
5 NUTS1 region and season, as well as their categorical distributions.

6 **2.4 Selection of periods for analysis**

7 All the analyses were carried out using time series with monthly resolution. However,
8 different subsets of periods for analysis were used. For each NUTS1 region separately, a
9 subset of years within 1970-2012 was selected for analysis based on drought impact
10 occurrence. Years with at least one impact occurrence in the region were selected. All months
11 of the selected years were included in this censored time series. The censoring was undertaken
12 to exclude years with drought conditions yet no impact reports in the EDII, similar to
13 Bachmair et al. (2015). The search for impact reports in both countries focused on known
14 drought events; the absence of impact reports in the EDII for years with drought conditions
15 may therefore be attributable to either a lack of impact occurrence or simply a lack of drought
16 impact reports, whether through not being discovered or not being published in the first place.
17 Table 1 shows the length of time series per region and the percentage of months with impact
18 occurrence in this censored time series. Despite the above-described censoring approach a
19 considerable percentage of months with zero impact occurrence remained. The data analysis
20 was only applied to regions with at least 10 months with impact occurrence, which led to the
21 exclusion of Northern Ireland and Scotland (UK), and the Hanseatic City of Bremen,
22 Hanseatic City of Hamburg, and Thuringia (DE). Additionally, a seasonal analysis was
23 carried out based on the censored time series per NUTS1 region. For this purpose, only data
24 for months occurring in the selected season were retained, in correct temporal sequence from
25 one year to the next (see 2.3 for the definition of seasons).

26

27 **3 Methods linking indicators and impacts**

28 **3.1 Correlation analysis**

29 First, we carried out a cross-correlation analysis between different drought indicators and the
30 number of impacts, accounting for temporal autocorrelation in the indicator and/or impact

1 time series. Spearman rank correlation coefficients (ρ) were calculated between time series of
2 drought indicators and number of impact occurrences, for each NUTS1 region separately.
3 Rank correlation was chosen over Pearson correlation since the counts of the impact data are
4 not normally distributed. Correlations were undertaken between time series of different
5 indicators on the one hand (mean SPI and SPEI for 1-8, 12, and 24 months; Q; G (DE only);
6 month (M) and year (Y) of impact occurrence), and time series of number of impact
7 occurrences for different impact subsets on the other:

- 8 • total impacts (N_I)
- 9 • hydrological drought impacts (N_{Ih})
- 10 • impacts due to other types of drought (N_{Io})
- 11 • impacts per impact category, and
- 12 • impacts per season (DJF, MAM, JJA, SON).

13 A subset of impact data was only included in the analysis if there were at least 10 months with
14 impact occurrence. Since there was temporal autocorrelation present in the time series of SPI
15 and SPEI of longer accumulation periods, in time series of Q and G, and in the impact time
16 series for most UK and some German NUTS1 regions, significance levels of the cross-
17 correlation analysis had to be corrected. Temporal autocorrelation of time series used in cross-
18 correlation analysis violates the assumption of serial independence and increases the
19 likelihood of type I error (Hurlbert, 1984; Jenkins, 2005). We applied the “Modified Chelton
20 method” by Pyper and Peterman (1998), which adjusts the “effective” number of degrees of
21 freedom used for determining significance levels. While we use Spearman’s ρ for the cross-
22 correlation analysis, autocorrelation coefficients represent Pearson’s r (based on square root
23 transformed data for the counts of impact occurrence). We define strength of correlation as
24 follows: 0-0.1 (no correlation), >0.1-0.3 (weak), >0.3-0.6 (moderate), >0.6-0.9 (strong), and
25 >0.9 (very strong).

26 **3.2 Random forest modeling**

27 Second, we employed a machine learning approach utilizing an ensemble regression tree
28 approach called “random forest” (Breiman, 2001). Similar to the cross-correlation analysis,
29 the random forest approach also identifies drought indicators best linked to impact
30 occurrence. In addition to extracting predictor importance, the random forest approach is used

1 for obtaining splitting values as estimates of thresholds of impact occurrence, and to model
2 drought impact occurrence.

3 A “random forest” (Breiman, 2001) is a machine learning algorithm, which constructs a large
4 number of classification or regression trees (CARTs) on bootstrapped subsamples of the data.
5 For our analysis we applied the R package ‘randomForest’ developed by Liaw and Wiener
6 (2002). Details about the random forest (RF) methodology and model parameterization are
7 given in the appendix. The RF predictors for each NUTS1 region included the same indicators
8 as used in the correlation analysis. The response variable is the square root transformed
9 monthly counts of impact data per NUTS1 region. We then ran models for the same subsets of
10 impacts as in the correlation analysis if there were at least 10 months with impact occurrence:
11 total impacts (N_I), hydrological drought impacts (N_{Ih}), non-hydrological drought impacts
12 (N_{Io}), and impacts per impact category.

13 To identify the drought indicators best linked to impact occurrence we used the “variable
14 importance” feature of the RF algorithm described in Liaw and Wiener (2002), which enabled
15 us to use the ranks of percent decrease in accuracy as variable importance measure (e.g. Strobl
16 et al., 2009). Another output from the RF analysis are the splitting values for each predictor.
17 The construction of each regression tree is based on recursively splitting the data into more
18 homogenous groups (nodes). At each node, the best splitting variable and splitting value are
19 determined, with multiple splits possible for the same variable (Strobl et al., 2009). For our
20 analysis we extracted the splitting values corresponding to each predictor, considering all
21 trees and nodes, and visualized their distribution as boxplot. We regard these splitting values
22 as estimates of thresholds of impact occurrence. All RF models are based on multiple
23 indicators. Therefore, indicator thresholds of individual indicators are conditional on predictor
24 interactions.

25 The predictive potential of the random forest models was assessed in two ways. First, the
26 overall model performance was evaluated based on a 10-fold cross-validation. The goal of
27 this assessment (hereafter “RF Predictions”) is to test the performance of RF models as a
28 potential tool for predictive purposes, and to learn about the indicator-impact relationship.
29 The data for cross-validation is the censored time series for each NUTS1 region, i.e. the time
30 series based on the sub-selection of years with drought impact occurrence within 1970-2012.
31 For each of the ten model runs the censored time series was split into 90% for training and
32 10% for prediction; impact occurrence of the left-out 10% is predicted with a random forest

1 model constructed on the training data. The cross-validation procedure allows evaluation of
2 the predictive performance for “unseen” data excluded from model fitting. As model
3 performance metrics we computed mean absolute error (MAE), root mean squared error
4 (RMSE), and error components according to the Kling-Gupta-Efficiency (Gupta et al., 2009)
5 modified by Gudmundsson (2012): relative difference in mean ($\Delta\mu$), relative difference in
6 standard deviation ($\Delta\sigma$), and strength of correlation between observed versus modeled number
7 of impacts (r). Zero is the optimal value of $\Delta\mu$ and $\Delta\sigma$; negative and positive values indicate
8 under- and over-prediction, respectively (Gudmundsson et al., 2012).

9 The second assessment (hereafter “RF Backwards Learning”) is the application of the RF
10 models that were fitted to the censored time series to predict N_I per NUTS1 region to those
11 years that had been excluded, i.e. the years within 1970-2012 that have zero impact
12 occurrence. The purpose of this second assessment is to scrutinize the impact data in the EDII
13 database to backwards learn where a year without impacts may either be due to no impacts or
14 due to the lack of reporting or finding reports. As the observations themselves are examined
15 no model performance metrics are presented.

16

17 **4 Results**

18 **4.1 Correlation of indicators with impacts**

19 In the UK the strength of correlation between times series of N_I and different indicators ranges
20 between -0.65 and 0.51 (Figure 3). Lower indicator values coincide with higher N_I (negative
21 correlation) for all drought indicators except for M, where positive values in summer concur
22 with a higher N_I (positive correlation). Overall, SPI and SPEI are very similar in terms of
23 strength of correlation. For southern and central UK, accumulation periods of SPI and SPEI
24 exceeding about 6 months show the strongest correlation with N_I , whereas the more northern
25 regions show the strongest correlation for short to intermediate accumulation periods. SPI-24
26 and SPEI-24 are the indicators with the strongest correlation for half of the NUTS1 regions
27 (WAL, CWE, EE, SWE, and SEE), with ρ ranging between -0.38 and -0.65. Streamflow
28 percentiles display a moderate and significant ρ in parts of eastern England, but for the other
29 regions the correlation is weak to moderate and not significant at the 5% level (two-sided
30 test). There is mainly no or a weak (non-significant) correlation with Y, which varies in sign.

1 A split into I_h versus I_o , and a split by impact category reveal distinct differences in
2 correlation patterns for some impact subsets (Figure 3). The difference between I and I_h is
3 rather minor. As can be seen in Figure 2, I_h is the dominant impact type in the UK. Other
4 drought impacts (I_o) show a distinctly different pattern. With weak to moderate ρ for all
5 indicators, no best SPI and SPEI time scale can be singled out. For agriculture, which mostly
6 represents I_o , only CEE and CWE show strong relationships, but for all accumulation periods.
7 While the correlation patterns for water supply and freshwater ecosystem impacts are similar
8 to I_h , shorter to intermediate accumulation periods of SPI and SPEI (4 to 8 months, for a few
9 cases also 12 months) show highest correlation with water quality impacts. For other impact
10 categories correlation could only be determined for very few regions (wildfire, tourism,
11 waterborne transportation), or not at all due to too few months with impact occurrence. A split
12 by season (Figure 4) also shows distinct differences, yet could not be determined for all
13 regions given limited impact data if partitioned seasonally.

14 In Germany, the overall strength of correlation between times series of N_I and different
15 indicators is in a similar range as in the UK (-0.62 to 0.74). Contrary to the UK, shorter to
16 intermediate accumulation periods of SPI and SPEI best correlate with impact occurrence
17 (Figure 5). Eleven of the 13 analyzed regions show the highest ρ for SPEI-2 to SPEI-4; for
18 SPI-24 and SPEI-24 a non-significant correlation in inverse direction is found. The difference
19 between SPI and SPEI is slightly more pronounced in Germany, with SPEI performing
20 somewhat better (absolute difference in ρ up to 0.13). Q performs similar to SPI in many
21 cases. Groundwater level percentiles show no or non-significant weak correlation with N_I . In
22 contrast, the sine expression of the month shows a higher and often significant ρ , especially in
23 the northern NUTS1 regions. Similar to the UK, there is no or only a weak correlation with Y.
24 As in the UK, there are regional differences, yet mostly regarding the strength of correlation.
25 Most regions in the north and northeast of Germany display a noticeably lower strength of
26 correlation (mostly weak ρ) than the central and southern regions.

27 Similar to the UK, a split into I_h and I_o reveals differences in correlation patterns compared
28 with I , yet the picture for I_h and I_o is the opposite: while the correlation pattern for N_{I_o} is rather
29 similar to N_I , there is a noticeable shift towards higher correlation with longer SPI and SPEI
30 time scales for I_h . N_{I_o} dominates over N_{I_h} in some German regions, in contrast to the situation
31 in the UK (Figure 2). A further split by impact category uncovered the following: agricultural
32 impacts show highest ρ for SPI and SPEI time scales of 1-4 months, yet most correlations are

1 weak and not significant; there is a shift towards higher correlation with longer SPI and SPEI
2 timescales for impacts on waterborne transportation in some NUTS1 regions; for all other
3 impact categories correlations could only be determined for one or two regions (BV or BW or
4 RP) due to too little impact data. A seasonal split was also not possible to assess due to too
5 few months with I in spring, fall, and winter; the majority of impacts in Germany occurred in
6 summer (Figure 2).

7 **4.2 Indicator importance in random forest models**

8 For the UK, the general picture from the random forest approach is very similar to the
9 findings from the correlation analysis, both regarding I and different impact subsets (I_h , I_o , and
10 I per impact category) (Figure 6). Long accumulation periods of SPI and SPEI (12 and 24
11 months) appear as the highest ranking predictors for most regions, except the more northern
12 regions NEE, NEW and YHU. Q does not show up as important predictor. Distinct
13 differences compared with the correlation analysis include the following: 1) Y plays an
14 important role for I and most impact subsets; 2) for I_o , the RF predictor importance shows a
15 shift to intermediate accumulation periods of SPI and SPEI (7/8 months). This shift is not as
16 clearly discernible in the correlation patterns. The same holds true for the agricultural impacts.
17 In contrast to the UK, where the RF predictor importance plots look very similar to the
18 correlation analysis plots, there is more variation for Germany (Figure 7). The RF predictor
19 importance patterns are spottier than the correlation analysis patterns with less smooth
20 transitions between adjacent indicators. Nevertheless, the general tendency of best predictors
21 is confirmed.

22 **4.3 Indicator thresholds in random forest models**

23 While splitting values of all indicators for all impact subsets (I , I_h , I_o , different impact
24 categories) were extracted, we only show the threshold distribution, i.e. splitting value
25 distribution, for selected SPI and SPEI time scales (best performance for different regions
26 and/or impact subsets) and streamflow and groundwater level percentiles (Figures 8 and 9).
27 While we display the threshold distribution of individual indicators, it is important to
28 remember they are conditional on multi-predictor interactions in the RF model.

29 For the UK, the threshold distribution for both meteorological indicators generally shows a
30 considerable range, which decreases with increasing accumulation period. For the same

1 accumulation periods of SPEI the range extends to less negative values. Apart from this, the
2 differences between SPI and SPEI are negligible with interquartile ranges (IQR)
3 predominantly between 0 and -2. When only focusing on the median of the distribution, SPI-8
4 and SPEI-8 values scatter around -1 for most NUTS1 regions. For SPI and SPEI of 12 and
5 especially 24 months duration the scatter around -1 is slightly more variable. Regarding
6 streamflow percentiles the splitting values cover almost the entire range, the IQR is distinctly
7 larger than for SPI and SPEI, and the median ranges between 0.2 and 0.37. The split by
8 impact category results in slightly narrower ranges of threshold distributions for many impact
9 categories, and often a more negative median (not shown). All indicators show regional
10 differences, however without systematic patterns.

11 For Germany, the splitting values in the different federal states range from roughly +1.5 to -
12 2/-3 for both SPI and SPEI (Figure 9). Absolute values of the IQR of German regions are
13 similar to the UK. Contrary to the UK, a regional pattern exists regarding the median of the
14 SPI and SPEI threshold distributions. The southern and most central federal states display a
15 more negative median (mainly between -1 and -1.5) than the northern/northeastern states
16 (with a median between 0 and -1). A small but noticeable gradient from SH to BV can be seen
17 in Figure 9. Streamflow percentiles show a similarly large spread of splitting values to the
18 UK, yet the IQR is mostly smaller and the median is slightly lower (0.14-0.29). For
19 groundwater level percentiles, the median per region ranges between 0.2 and 0.68; no regional
20 pattern is found. The low amount of impact data for the RF analysis for several impact
21 categories prevented a systematic intercomparison among impact categories.

22 **4.4 Impact predictions with random forest models**

23 RF Predictions for the UK show that observed and modeled impacts agree well for the
24 NUTS1 regions SWE, SEE, and EE (Figure 10). In most central regions and LND there is
25 more spread. The northern regions NEE and NEW show least agreement. The R^2 ranges
26 between 0.16 (NWE) and 0.73 (WAL) (Table 2). Due to the random component in the RF
27 algorithm, model performance varies marginally for replications. Regional differences more
28 or less reflect the length of each time series and the percentage of months with impact
29 occurrence. That is, regions with $R^2 > 0.6$ generally have longer time series and a higher
30 percentage of months with I than regions with lower R^2 (Tables 1 and 2). For Germany,
31 observed and modeled impacts agree less well than for many UK regions (Table 2). However,

1 much fewer data points for Germany than for the UK make a comparison difficult (Figures 10
2 and 11). Among the federal states of Germany, BV and BB show better agreement than other
3 regions. The majority of federal states yielded an R^2 between 0.33 and 0.54 (Table 2). Only
4 four states show an $R^2 > 0.6$. Overall, the lower agreement between observations and
5 predictions than in the UK concurs with the shorter time series of indicator and impact time
6 series in Germany, and a smaller percentage of months with $N_I > 0$ (Table 1).

7 The generally small difference in the mean ($\Delta\mu$) of observed versus modeled impacts for both
8 the UK and Germany (Table 2) suggests that the central tendency is well modeled. However,
9 a closer look at the time series of observed and modeled number of impact occurrences
10 (Figure 12, time series with gray background (RF Predictions)) reveals that small values are
11 generally over-predicted and large values often under-predicted. The under-prediction of N_I
12 causes lower standard deviations for the modelled values than for the observed ($\Delta\sigma$ between -
13 0.22 and -0.52, see Table 2).

14 Furthermore, Figure 12 shows that predictions and observations in the UK and Germany
15 generally agree well both regarding initiation of impact occurrence and its subsequent
16 temporal evolution. This is also reflected by a moderate to strong correlation between
17 predictions and observations (Table 2). The blue line in Figure 12 represents an impact
18 threshold of one, as guidance for interpretation: modeled impacts smaller than one may be
19 regarded as an absence of impacts. Taking this into account the temporal dynamics agree even
20 better, especially regarding impact onset. An obvious disagreement between dynamics of
21 observations and predictions is found in many regions in the UK in 1991/1992, where
22 modeled N_I is more dynamic than the observed static “block” of N_I following an impact peak.
23 The block-shaped data represent impacts due to a contraction of the stream network in large
24 parts of the south and east of the UK during these years. In Germany, states with larger
25 amplitude of N_I (BV, BW, RP, and NW) tend to have a better agreement of temporal
26 dynamics, especially when only focusing on values above the one-impact-threshold line. For
27 states with low amplitude of N_I , which often concurs with less negative splitting values (see
28 section 4.3), the temporal dynamics are less well modeled (not shown).

29 The RF Backwards Learning predictions for all years with zero impact occurrence according
30 to the EDII database are shown on white background in Figure 12. They expose instances of
31 potentially “false-positive impacts”, i.e. a positive number of impacts is modeled while there
32 are no observed impacts. A clear example for the UK is the period 1972-74, when drought

1 conditions occurred, which would have caused impacts in many UK regions according to the
2 RF model trained on the censored time series. Another example of false-positive impacts in
3 the UK is found for many southern and central regions in the second half of the 1990s after a
4 peak of N_I in 1995. While for the UK two major, spatially coherent cases of false-positive
5 impacts exist, the pattern for Germany is more diverse and region-specific.

6

1 **5 Discussion**

2 **5.1 Performance of drought indicators**

3 The correlation analysis and the random forest approach revealed the following insights about
4 the performance of drought indicators, which will be discussed in the context of expectations
5 and literature: 1) the best-performing drought indicators are region and impact category
6 specific, and in the UK season specific to some degree. While in the UK generally long
7 accumulation periods of SPI and SPEI (12-24 months) performed best, short to intermediate
8 accumulation periods (2-4 months) were best linked with drought impacts in Germany.
9 However, there is spatial variability within each country, and differences among impact
10 categories. 2) A comparison among indicators (SPI vs. SPEI vs. Q (vs. G in Germany))
11 uncovered that in the UK SPI and SPEI perform similarly to each other, and Q performs less
12 well. In Germany SPEI often performed slightly better than SPI, the linkage with Q is better
13 than in the UK, and there is low agreement between G and impact occurrence. 3) The largely
14 congruent findings from the two different statistical approaches independently validate the
15 results.

16 While much can be speculated about the drivers of region-, impact type-, and season-specific
17 variability, it is nonetheless necessary to explore the underlying mechanisms for the observed
18 differences to rule out purely data-driven, yet physically meaningless, indicator-impact-
19 relationships. Regional differences can result from both 1) “real” physical, spatial differences
20 in geographic properties (e.g. climate, geology, soil, land use), vulnerability towards drought,
21 and hazard characteristics, triggering impacts differing in type and response time, and 2)
22 differences due to inherent spatial and temporal biases in the impact data (see Bachmair et al.
23 (2015) on potential EDII error sources).

24 In the UK we found differences in best SPI and SPEI accumulation periods between most
25 southern/central regions (long periods) versus more northern regions (shorter periods). This
26 corresponds well to known differences in the nature of the drought hazard, and impacts.
27 Strong regional contrasts in drought occurrence across the UK have been noted previously,
28 with a particular contrast between the upland northern and western UK, which is susceptible
29 to short-term (6 month) summer half-year droughts, and the lowlands of the south-eastern UK
30 that are susceptible to longer-term (18 month or greater) multi-annual droughts (Jones and
31 Lister, 1998; Marsh et al., 2007; Parry et al., 2011). These findings reflect both the

1 climatological rainfall gradient across the UK and the predominance of groundwater
2 dominated catchments in the south-east (Folland et al., 2015). While we also found regional
3 differences in indicator-impact-linkage patterns in Germany, they mostly relate to differences
4 in strength of correlation (weaker correlation in northern/northeastern states). The smaller
5 amplitude of impact time series in these states may explain weaker correlation in contrast to
6 southern/central states with predominantly larger amplitude, i.e. pronounced impact peaks, as
7 hypothesized by Bachmair et al. (2015).

8 The differences in indicator-impact-relationships between the UK and Germany, and some of
9 the within-country variability, are also very likely a result of the regional composition of
10 drought impact types. It is common knowledge that impacts caused by different types of
11 drought have different response times due to propagation through the hydrological cycle (e.g.
12 Mishra and Singh, 2011; National Drought Mitigation Center, 2015; Wilhite and Glantz,
13 1985). In the UK impacts associated with drought conditions of surface waters and
14 groundwater (I_h) clearly dominate (see Figure 2). This agrees well with longer SPI and SPEI
15 accumulation periods as best predictors in the UK compared with Germany. There, the
16 fraction of non-hydrological drought impacts (I_o) is distinctly larger than in the UK.
17 Agricultural and forestry impacts in Germany account for roughly 20-70 percent of impacts
18 depending on the region, and this may explain why short to intermediate SPI and SPEI
19 accumulation periods are the best predictors.

20 The identification of best-performing indicators for specific impact types is a key outcome of
21 this study. For instance, agricultural and hydrological drought impacts were generally best
22 linked to shorter and longer SPI and SPEI time scales, respectively. Here, “shorter” and
23 “longer” refers to different absolute values: 1-4 (DE) and 7-8 months (UK) for agriculture,
24 and 7/8 (DE) and 12/24 months (UK) for I_h . Perhaps unsurprisingly, a universal
25 recommendation about best indicators hence cannot be inferred. However, the similar relative
26 shift in best SPI and SPEI time scales suggests that there are likely to be typical patterns of
27 response for given impact types, but that these are mediated by regional cause-effect-
28 mechanisms. This is in line with results of the studies by Blauhut et al. (2015) and Stagge et
29 al. (2014). Seasonal variation in linkage patterns as observed in our study for the UK further
30 complicates recommendations regarding a single best drought indicator. Part of the variation
31 across the seasons is likely to reflect differences in impact type distribution between the
32 seasons (see Figure 2). For example, the long SPI and SPEI time scales for winter and spring

1 in permeable catchments in the southeastern lowlands (Figure 4) reflect long groundwater
2 droughts, which in turn affects groundwater-fed rivers. The winter half-year is the main
3 recharge season and failure to recharge will trigger water use restrictions, while shrinking
4 headwaters will result in freshwater ecosystem impacts. However, less permeable catchments
5 are likely to respond more readily to winter rainfall as the evapotranspiration is low in this
6 season. For the bulk of rivers, the SPI and SPEI time scales are therefore shorter, with impacts
7 related to low absolute water levels mainly in summer and fall, although effects can be long-
8 lasting.

9 A surprising result is that streamflow did not appear as an important drought indicator in the
10 UK, even after a separation of hydrological drought impacts. In Germany, groundwater level
11 percentiles played only a minor role. There are several possible reasons for these
12 discrepancies. For groundwater level percentiles the mismatch is likely attributable to a
13 lagged groundwater response (Bachmair et al., 2015). One probable reason for the lack of
14 relationship between I and Q may be the nature of the spatially aggregated streamflow data,
15 which represents different catchments varying in size and characteristics (including degree of
16 human influence), lumped over a large administrative area, which does not coincide with
17 catchment boundaries. A further reason may be the nature of the EDII data, especially
18 regarding the subdivisions of I_h . While in Germany the fraction of instream impacts of I_h is
19 larger (e.g. impaired navigability of streams, water quality, and reduced power production due
20 to a lack of cooling water), water supply impacts dominate I_h in the UK. For groundwater or
21 reservoir-fed water supply systems these impacts are, to a certain extent, disconnected from
22 river flows (the purpose of reservoirs being to smooth out variations in instream water
23 availability).

24 Overall, despite a rather complex picture in terms of best drought indicator for impact
25 occurrence, the empirical evaluation of drought indicators with text-based impact information
26 proved to be a feasible approach. Given the minor differences in the outcomes of the
27 correlation and the random forest analysis for the UK, both methods appear recommendable.
28 Generally, the strength of the random forest algorithm is that it can handle interactions and
29 nonlinearities among variables, and thus identify non-intuitive relationships (Evans et al.,
30 2011; Hastie et al., 2009). Furthermore, random forests are robust to noise (Breiman, 2001;
31 Hastie et al., 2009), and the bootstrap sampling provides a way to account for the uncertainty
32 of the impact data. Nevertheless, the “black-box” nature of the RF model (Breiman, 2001)

1 may not be as useful when an intuitive method for the choice of best drought indicator is
2 needed (e.g. when working with a wide range of stakeholders from different backgrounds).
3 For Germany, systematic differences in indicator-impact-linkage patterns were easier to
4 perceive in the correlation plots than in the RF predictor importance plots. For large data sets
5 the RF algorithm has the potential to detect relatively complex structures; for small data sets,
6 however, this is unlikely to be the case (Maindonald and Braun, 2006).

7 **5.2 Indicator thresholds for impact occurrence**

8 The analysis of splitting values used in the random forest construction highlighted a large
9 spread. Yet, when focusing on the median there are differences between the countries and
10 among the regions (medians around -1 for SPI and SPEI of different accumulation periods in
11 the UK, and in DE between ca. 0 and -1 (north/northeast) and -1 and -1.5 (southern/central
12 states)), and, to some extent, impact categories.

13 We regard splitting values during recursive partitioning as estimates of thresholds of impact
14 occurrence because they provide guidance on critical predictor values triggering a
15 consequence. Nevertheless, the uncertainty of the text-based impact data clearly must be
16 taken into account in the search for meaningful thresholds. One cause of the large spread of
17 the threshold distributions is the uncertain timing of actual impact occurrence, especially
18 regarding its termination. Second, in the UK there are impacts appearing as “blocks”
19 following an impact peak in 1990. They arise from EDII reports citing long-lasting impacts
20 without an exact known end-point or temporal evolution of the severity of the impact (i.e. low
21 flow anomalies in eastern and southern Britain causing contraction of the stream network and
22 thus impacts on aquatic species reported for the years 1990 to 1992). Third, hosepipe bans and
23 drought orders do not represent direct impacts of drought, but are triggered (and canceled) by
24 an administrative/political decision as an intermediate step. The onset and termination of the
25 impacts they are meant to reflect may therefore be more uncertain than those for other, more
26 direct impacts. These issues highlight the necessity to separately consider phases of drought
27 development and recovery for drought M&EW (Parry et al., in review; Steinemann and
28 Cavalcanti, 2006). Fourth, differences in impact reporting between Germany and the UK also
29 need to be considered. In the UK, a significant proportion of impacts for later droughts (2004-
30 2006 and 2010-2012) were sourced from weekly Drought Management Briefs by the

1 Environment Agency (EA). In Germany there is no continuous information on drought
2 impacts, and no unifying impact reporting scheme exists within the federal state structure.

3 A reason why we consider tree splitting values as meaningful thresholds of impact occurrence
4 is because Bachmair et al. (2015) found similar threshold patterns for Germany using the
5 same impact data but a different methodological approach based on extracting indicator
6 values concurrent with past impact onset. Both approaches revealed differences in indicator
7 thresholds between northern/northeastern versus southern/central German federal states.
8 These differences were speculated to result from differences in geographic properties and thus
9 different vulnerability to drought (Bachmair et al., 2015). The northern/north-eastern states
10 tend to have more sandy soils with lower water holding capacity than in the south, and lower
11 natural water availability (Bundesamt für Gewässerkunde, 2003; Bundesanstalt für
12 Geowissenschaften und Rohstoffe, 2007). This could explain impact occurrence for less
13 negative SPI and SPEI thresholds.

14 Despite possible shortcomings of EDII data and the method to derive indicator thresholds, we
15 recommend further efforts to empirically validate indicator thresholds with impact data.
16 Drought indicator thresholds informed by impact data may complement and allow comparison
17 with local-scale decision making on drought triggers, which is usually based on past
18 hydrological data, stakeholder knowledge and the experience of individuals (e.g. Steinemann,
19 2014). In our study the median of the SPI and SPEI threshold distribution ranged around -1 in
20 the UK, which correspond to the transition between mild and moderate drought according to
21 the SPI classification by McKee (1993). At the same time, the differences in median of the
22 SPI and SPEI threshold distributions for Germany (lower values for SPEI) demonstrate that,
23 despite the standardized nature of such indices, the same thresholds (and corresponding
24 statistical return periods) are not necessarily equally meaningful for drought impact
25 occurrence. To improve that knowledge base, more studies should systematically evaluate and
26 make public the delineation rules of different drought severity classes by using drought
27 impacts (as e.g. Sepulcre-Canto et al., 2012) or by stakeholders' experience (as e.g.
28 Steinemann and Cavalcanti, 2006).

29 **5.3 Lessons learned from random forest predictions**

30 The two parts of the random forest modeling exercise exposed that: 1) there are differences
31 among regions in terms of predictive power, with RF models for regions with better impact

1 data (longer censored time series, a higher percentage of non-zero data, and larger amplitude
2 of the impact time series) showing good agreement between observations and predictions. 2)
3 While the temporal dynamics of impact occurrence were reasonably reproduced, over- and
4 under-prediction of small and large values, respectively, are an issue. 3) Backwards leaning
5 about impact occurrence for years with no observations (through RF models trained on
6 drought years) provided valuable insights into time periods which potentially lack impact data
7 in the EDII.

8 Overall, the analysis revealed that RF models generally represent a suitable tool for drought
9 M&EW, yet further model tuning is possible (e.g. reduction of predictors, grouping several
10 regions for increasing the number of observations, and impact category specific models). The
11 finding that there is good agreement between observed and predicted number of drought
12 impact occurrences for regions with good data availability is promising. It underlines the
13 benefit of spending time and resources on impact data collection. The necessity of expanding
14 impact data collection and its benefit for drought M&EW has also been reported by others
15 (Lackstrom et al., 2013; Stahl et al., 2015b; Wilhite et al., 2007).

16 Despite the promising predictive capability of RF models for some regions, the under-
17 prediction of peaks is an issue. One reason for this may be an inherent bias of the random
18 forest algorithm with high values being under-predicted and low values being over-predicted,
19 as observed by others (Ordoyne and Friedl, 2008). This is because the RF algorithm computes
20 averages over a large number of model predictions and hence reduces the range and variance
21 of predictions compared with observed values (Liaw and Wiener, 2002; Ordoyne and Friedl,
22 2008). Another reason may be an impact-reporting bias caused by impact-reporting increasing
23 during peaks of events. We hypothesize that drought impacts may go unreported during the
24 early stages of a drought, but once a certain threshold of public attention and media coverage
25 is exceeded there is a tendency for more complete reporting. Also, the chances of finding
26 information on drought impacts are higher for recent events due to better online availability of
27 reports and new media channels compared with decades ago.

28 The RF Backwards Learning assessment provided a way to scrutinize whether an absence of
29 data in the EDII for certain time periods reflects a true absence of drought impacts, or simply
30 missing data. For the UK, we discovered two prominent examples of droughts that are more
31 severe in modeled impacts than observed EDII impacts: the early 1970s and late 1990s. Both
32 are well documented droughts, but previous studies suggest the former genuinely had less

1 impacts (Cole and Marsh, 2006), in part due to a wet summer in 1973. In contrast, the late
2 1990s is likely to represent missing impact data. For the 1995-1997 drought, impacts from the
3 hot, dry summer of 1995 are captured in the EDII; the summer drought had very severe water
4 supply impacts, triggering public enquiries, and was thus very extensively reported. However,
5 a protracted groundwater drought, with water restrictions in some areas, extended into 1997
6 (Cole and Marsh, 2006). However no “formal” drought report was issued on the latter phases
7 of the drought so these impacts have not been captured by the EDII. Altogether, false-positive
8 impacts identified with the RF Backwards Learning assessment provide guidance on which
9 time periods to focus on when searching for additional impact information.

10

11 **6 Conclusions**

12 The broad goal of our analysis was two-fold: to learn about the relationship between drought
13 indicators and text-based impact information, to advance drought monitoring and early
14 warning practices; and to test methodologies that can be extended to other locations in future
15 applications. We found that drought indicators best linked to impact occurrence are generally
16 SPI and SPEI with long accumulation periods (12-24 months) for the UK, and with short to
17 intermediate accumulation periods (2-4 months) for Germany. Additionally, the indicator-
18 impact-response varies within the countries. This calls for evaluating continental drought
19 M&EW systems at smaller spatial scales. Also, our analysis provided additional empirical
20 evidence that impacts associated with different types of drought (e.g. agricultural versus
21 hydrological drought) have different response times, as reflected by distinct differences in
22 indicator-impact-linkage patterns for each impact category. For regions with sufficient data, a
23 random forest machine learning approach proved to be a suitable tool for objectively
24 identifying indicator thresholds of impact occurrence, and for predicting the number of
25 drought impact occurrences. The regression tree splitting values, which we regard as
26 estimates of thresholds of impact occurrence, showed a considerable spread, yet the median
27 revealed differences among regions and, to a lesser extent, impact categories. In the UK the
28 median of threshold values was around -1 for SPI and SPEI. For Germany, distinct
29 differences in threshold values were found between northern/northeastern versus
30 southern/central regions. Such insight into indicator thresholds could provide guidance when
31 designing and validating drought triggers, and complements existing approaches like
32 stakeholder consultation. While there are certainly caveats given the uncertainty in exact

1 timing, number, and severity of impacts, the text-based reports served as a reasonable basis
2 for quantifying impacts. A comparison of time series of observed versus modeled impacts
3 additionally yielded valuable insights into the contents of the European Drought Impact report
4 Inventory and allowed us to identify potential gaps in the temporal coverage of the impact
5 database. Overall, the information gain from evaluating drought indicators with impacts
6 underlines the strong benefits of impact data collection, and is an important step towards
7 closing the gap between knowledge about hazard intensity and on-the-ground drought
8 conditions.

9

10 **Acknowledgements**

11 This study is an outcome of the Belmont Forum project DrIVER (Drought Impacts:
12 Vulnerability thresholds in monitoring and Early warning Research). Funding to the project
13 DrIVER by the German Research Foundation DFG under the international Belmont
14 Forum/G8HORC's Freshwater Security programme (project no. STA-632/2-1) and to the EU-
15 FP7 DROUGHT R&SPI project (contract no. 282769) is gratefully acknowledged. Financial
16 support for DrIVER for the UK-based authors was also under the Belmont Forum and was
17 provided by the UK Natural Environment Research Council (Grant NE/L010038/1). The
18 article processing charge was funded by the German Research Foundation (DFG) and the
19 Albert Ludwigs University Freiburg in the funding programme Open Access Publishing.

20 We thank Lukas Gudmundsson for the provision of SPI and SPEI gridded data developed
21 within the DROUGHT R&SPI project, and the UK National River Flow Archive for
22 provision of river flow data. We further thank the following agencies of the German federal
23 states for supplying streamflow and groundwater level data through the Bundesanstalt für
24 Gewässerkunde-funded project "Extremjahr 2011": Bayerisches Landesamt für Umwelt
25 (LfU), Hessisches Landesamt für Umwelt und Geologie (HLUG), Landesamt für Natur,
26 Umwelt und Verbraucherschutz Nordrhein-Westfalen (LANUV), Landesamt für Umwelt und
27 Arbeitsschutz Saarland (LUA), Landesamt für Umwelt, Naturschutz und Geologie
28 Mecklenburg-Vorpommern (LUNG), Landesamt für Umwelt, Wasserwirtschaft und
29 Gewerbeaufsicht Rheinland-Pfalz (LUWG), Landesanstalt für Umwelt, Gesundheit und
30 Verbraucherschutz Brandenburg (LUGV, Regionalabteilungen Ost, Süd, West), Landesanstalt
31 für Umwelt, Messungen und Naturschutz Baden-Württemberg (LUBW), Landesbetrieb für
32 Hochwasserschutz und Wasserwirtschaft Sachsen-Anhalt (LHW), Landesbetrieb für

1 Küstenschutz, Nationalpark und Meeresschutz Schleswig-Holstein (LKNM),
2 Niedersächsischer Landesbetrieb für Wasserwirtschaft, Küsten- und Naturschutz (NLWKN),
3 Ruhrverband, Sächsisches Landesamt für Umwelt, Landwirtschaft und Geologie (LfULG),
4 Staatliches Amt für Landwirtschaft und Umwelt Vorpommern (StALU-VP), Thüringer
5 Landesamt für Umwelt und Geologie (TLUG), Landesamt für Landwirtschaft, Umwelt und
6 ländliche Räume (LLUR), Wasser- und Schifffahrtsverwaltung des Bundes (WSV).

7 **Appendix**

8 Details about the applied random forest methodology: Non-parametric regression using
9 random forest (RF) consists of the following steps (see Liaw and Wiener (2002) for details):
10 1) n_{tree} bootstrap samples are used. The individual cases making up the sample are drawn
11 randomly with replacement from the original data, preserving each month's pairing of
12 predictand and predictors. The size of each sample is about two-thirds of the size of the total
13 dataset; 2) for each bootstrap sample, an unpruned tree is grown. That is, for each node in
14 turn, a split-in-two of the data is performed for each of m_{try} randomly chosen predictor
15 variables, and the predictor whose split results in the two most homogeneous groups
16 (minimizing the residual sum of squares) of the predictand is chosen as the splitting variable
17 for that node; 3) new data is predicted by averaging predictions over n_{tree} regression trees
18 (Liaw and Wiener, 2002). The user-defined variable n_{tree} was set to 1000. The model
19 parameter m_{try} (number of predictors randomly sampled as candidates at each split) was left as
20 default: one third of the total number of predictors (Liaw and Wiener, 2002). For all other
21 parameters the default was kept as well. The model error is determined by predicting the
22 excluded data ("out-of-bag" data according to Breiman (2001)) at each bootstrap iteration
23 using the tree grown with the bootstrap sample and averaging all errors (Liaw and Wiener,
24 2002).

25 In this study, the response variable is the square root transformed monthly counts of impact
26 data per NUTS1 region. This transformation yielded a near normal distribution of the non-
27 zero data in many regions. Some British NUTS1 regions, however, showed a bi-modal
28 distribution of NI (NEE, NEW, YHU, and SEE with varying extent), and in some German
29 states the distribution of NI remained positively skewed after the square root transformation.
30 Results for a log-transform were similar.

31

1 **References**

- 2 Bachmair, S., Kohn, I. and Stahl, K.: Exploring the link between drought indicators and
3 impacts, *Nat. Hazards Earth Syst. Sci.*, 15(6), 1381–1397, doi:10.5194/nhess-15-1381-2015,
4 2015.
- 5 Blauhut, V., Gudmundsson, L. and Stahl, K.: Towards pan-European drought risk maps:
6 quantifying the link between drought indices and reported drought impacts, *Environ. Res.*
7 *Lett.*, 10(1), 014008, doi:10.1088/1748-9326/10/1/014008, 2015.
- 8 Bradford, R. B., Marsh, T. J.: Defining a network of benchmark catchments for the UK. *P. I.*
9 *Civil Eng. – Water & Mar. En.*, 156(WM2), 109-116.
- 10 Breiman, L.: Random Forests, *Mach. Learn.*, 45(1), 5–32, doi:10.1023/A:1010933404324,
11 2001.
- 12 Bundesamt für Gewässerkunde: Hydrologischer Atlas von Deutschland., 2003.
- 13 Bundesanstalt für Geowissenschaften und Rohstoffe: Bodenarten in Oberböden Deutschlands
14 1:1000000, Hannover., 2007.
- 15 Cole, G. A. and Marsh, T. J.: Major droughts in England and Wales from 1800 and evidence
16 of impact, Environment Agency, Bristol., 2006.
- 17 Dieker, E., van Lanen, H. A. J., and Svoboda, M.: Comparison of three drought monitoring
18 tools in the USA, WATCH Technical Report No. 25, available at: [http://www.eu-](http://www.eu-watch.org/publications/technical-reports/3)
19 [watch.org/publications/technical-reports/3](http://www.eu-watch.org/publications/technical-reports/3), last access: 11 September 2015, 2010.
- 20 Dutra, E., Pozzi, W., Wetterhall, F., Di Giuseppe, F., Magnusson, L., Naumann, G., Barbosa,
21 P., Vogt, J. and Pappenberger, F.: Global meteorological drought – Part 2: Seasonal forecasts,
22 *Hydrol. Earth Syst. Sci.*, 18(7), 2669–2678, doi:10.5194/hess-18-2669-2014, 2014.
- 23 European Drought Observatory: PRODUCT FACT SHEET: Combined Drought Indicator –
24 EUROPE, available at: <http://edo.jrc.ec.europa.eu/edov2/php/index.php?id=1101> (last access:
25 20 August 2015), 2013.

- 1 Evans, J. S., Murphy, M. A., Holden, Z. A. and and Cushman, S. A.: Modeling Species
2 Distribution and Change Using Random Forest, in Predictive Species and Habitat Modeling
3 in Landscape Ecology, edited by C. A. Drew, Y. F. Wiersma, and F. Huettmann, p. 313,
4 Springer New York, New York, NY., 2011.
- 5 Folland, C. K., Hannaford, J., Bloomfield, J. P., Kendon, M., Svensson, C., Marchant, B. P.,
6 Prior, J. and Wallace, E.: Multi-annual droughts in the English Lowlands: a review of their
7 characteristics and climate drivers in the winter half-year, *Hydrol. Earth Syst. Sci.*, 19(5),
8 2353–2375, doi:10.5194/hess-19-2353-2015, 2015.
- 9 Gillette, H.: A creeping drought under way, *Water Sew. Work.*, 97, 104–105, 1950.
- 10 Gudmundsson, L., Rego, F. C., Rocha, M. and Seneviratne, S. I.: Predicting above normal
11 wildfire activity in southern Europe as a function of meteorological drought, *Environ. Res.*
12 *Lett.*, 9(8), 084008, doi:10.1088/1748-9326/9/8/084008, 2014.
- 13 Gudmundsson, L., Wagener, T., Tallaksen, L. M. and Engeland, K.: Evaluation of nine large-
14 scale hydrological models with respect to the seasonal runoff climatology in Europe, *Water*
15 *Resour. Res.*, 48(11), n/a–n/a, doi:10.1029/2011WR010911, 2012.
- 16 Gupta, H. V., Kling, H., Yilmaz, K. K. and Martinez, G. F.: Decomposition of the mean
17 squared error and NSE performance criteria: Implications for improving hydrological
18 modelling, *J. Hydrol.*, 377(1-2), 80–91, doi:10.1016/j.jhydrol.2009.08.003, 2009.
- 19 Hannaford, J., Lloyd-Hughes, B., Keef, C., Parry, S. and Prudhomme, C.: Examining the
20 large-scale spatial coherence of European drought using regional indicators of precipitation
21 and streamflow deficit, *Hydrol. Process.*, 25(7), 1146–1162, doi:10.1002/hyp.7725, 2011.
- 22 Hao, Z. and Singh, V. P.: Drought characterization from a multivariate perspective: A review,
23 *J. Hydrol.*, 527, 668–678, doi:10.1016/j.jhydrol.2015.05.031, 2015.
- 24 Hargreaves, G. H.: Defining and Using Reference Evapotranspiration, *J. Irrig. Drain. Eng.*,
25 120(6), 1132–1139, doi:10.1061/(ASCE)0733-9437(1994)120:6(1132), 1994.

- 1 Haslinger, K., Koffler, D., Schöner, W. and Laaha, G.: Exploring the link between
2 meteorological drought and streamflow: Effects of climate-catchment interaction, *Water*
3 *Resour. Res.*, 50(3), 2468–2487, doi:10.1002/2013WR015051, 2014.
- 4 Hastie, T., Tibshirani, R. and Friedman, J.: *The Elements of Statistical Learning*, Springer
5 New York, New York, NY., 2009.
- 6 Haylock, M. R., Hofstra, N., Klein Tank, A. M. G., Klok, E. J., Jones, P. D. and New, M.: A
7 European daily high-resolution gridded data set of surface temperature and precipitation for
8 1950–2006, *J. Geophys. Res.*, 113(D20), D20119, doi:10.1029/2008JD010201, 2008.
- 9 Hlavinka, P., Trnka, M., Semerádová, D., Dubrovský, M., Žalud, Z. and Možný, M.: Effect of
10 drought on yield variability of key crops in Czech Republic, *Agric. For. Meteorol.*, 149(3-4),
11 431–442, doi:10.1016/j.agrformet.2008.09.004, 2009.
- 12 Hurlbert, S. H.: Pseudoreplication and the Design of Ecological Field Experiments, *Ecol.*
13 *Monogr.*, 54(2), 187, doi:10.2307/1942661, 1984.
- 14 Jenkins, G.: The influence of climate on the fishery recruitment of a temperate, seagrass-
15 associated fish, the King George whiting *Sillaginodes punctata*, *Mar. Ecol. Prog. Ser.*, 288,
16 263–271, 2005.
- 17 Ji, L. and Peters, A. J.: Assessing vegetation response to drought in the northern Great Plains
18 using vegetation and drought indices, *Remote Sens. Environ.*, 87(1), 85–98, 2003.
- 19 Jones, P. D. and Lister, D. H.: Riverflow reconstructions for 15 catchments over England and
20 Wales and an assessment of hydrologic drought since 1865, *Int. J. Climatol.*, 18(9), 999–
21 1013, doi:10.1002/(SICI)1097-0088(199807)18:9<999::AID-JOC300>3.0.CO;2-8, 1998.
- 22 Kohn, I., Rosin, K., Freudiger, D., Belz, J. U., Stahl, K. and Weiler, M.: Niedrigwasser in
23 Deutschland 2011, *Hydrol. und Wasserbewirtschaftung*, 58(1), 4–17,
24 doi:10.5675/HyWa_2014, 2014.
- 25 Kruse, S., Seidl, I. and Staehli, M.: Informationsbedarf zur Früherkennung von Trockenheit in
26 der Schweiz, *Wasser Energ. Luft*, 102(4), 4, 2010.

- 1 Lackstrom, K., Brennan, A., Ferguson, D., Crimmins, M., Darby, L., Dow, K., Ingram, K.,
2 Meadow, A., Reges, H., Shafer, M. and Smith, K.: The Missing Piece: Drought Impacts
3 Monitoring. Workshop report produced by the Carolinas Integrated Sciences & Assessments
4 program and the Climate Assessment for the Southwest., 2013.
- 5 Liaw, A. and Wiener, M.: Classification and Regression by randomForest, R news, available
6 at: [ftp://131.252.97.79/Transfer/Treg/WFRE_Articles/Liaw_02_Classification and regression](ftp://131.252.97.79/Transfer/Treg/WFRE_Articles/Liaw_02_Classification%20and%20regression%20by%20randomForest.pdf)
7 [by randomForest.pdf](ftp://131.252.97.79/Transfer/Treg/WFRE_Articles/Liaw_02_Classification%20and%20regression%20by%20randomForest.pdf) (last access: 20 August 2015), 2002.
- 8 Loayza, N. V., Olaberria, E., Rigolini, J. and Christiaensen, L.: Natural Disasters and Growth:
9 Going Beyond the Averages, *World Dev.*, 40(7), 1317–1336,
10 doi:10.1016/j.worlddev.2012.03.002, 2012.
- 11 Logar, I. and van den Bergh, J. C. J. M.: Methods to Assess Costs of Drought Damages and
12 Policies for Drought Mitigation and Adaptation: Review and Recommendations, *Water*
13 *Resour. Manag.*, 27(6), 1707–1720, doi:10.1007/s11269-012-0119-9, 2013.
- 14 Maindonald, J. and Braun, J.: Data analysis and graphics using R: an example-based
15 approach, Third Edit., Cambridge University Press., 2006.
- 16 Marsh, T., Cole, G. and Wilby, R.: Major droughts in England and Wales, 1800–2006,
17 *Weather*, 62(4), 87–93, doi:10.1002/wea.67, 2007.
- 18 Martínez-Fernández, J., González-Zamora, A., Sánchez, N. and Gumuzzio, A.: A soil water
19 based index as a suitable agricultural drought indicator, *J. Hydrol.*, 522, 265–273,
20 doi:10.1016/j.jhydrol.2014.12.051, 2015.
- 21 Mavromatis, T.: Drought index evaluation for assessing future wheat production in Greece,
22 *Int. J. Climatol.*, 27(7), 911–924, 2007.
- 23 McKee, T. B., Doesken, N. J. and Kleist, J.: The relationship of drought frequency and
24 duration to time scales, in Preprints, 8th Conference on Applied Climatology, pp. 179–184,
25 Anaheim, California., 1993.

- 1 Mehta, V. M., Wang, H., Mendoza, K. and Rosenberg, N. J.: Predictability and prediction of
2 decadal hydrologic cycles: A case study in Southern Africa, *Weather Clim. Extrem.*, 3, 47–53,
3 doi:10.1016/j.wace.2014.04.002, 2014.
- 4 Mishra, A. K. and Singh, V. P.: Drought modeling—A review, *J. Hydrol.*, 403(1), 157–175,
5 2011.
- 6 National Drought Mitigation Center: Types of Drought, available at:
7 <http://drought.unl.edu/DroughtBasics/TypesofDrought.aspx> (last access: 20 August 2015),
8 2015.
- 9 Ordoyne, C. and Friedl, M.: Using MODIS data to characterize seasonal inundation patterns
10 in the Florida Everglades, *Remote Sens. Environ.*, 112, 4107–4119,
11 doi:10.1016/j.rse.2007.08.027, 2008.
- 12 Parry, S., Hannaford, J., Prudhomme, C., Lloyd-Hughes, B. and Williamson, J.: Objective
13 drought and high flow catalogues for Europe. [online] Available from: [http://www.eu-](http://www.eu-watch.org/publications/technical-reports)
14 [watch.org/publications/technical-reports](http://www.eu-watch.org/publications/technical-reports), 2011.
- 15 Parry, S., Prudhomme, C., Wilby, R. L. and Wood, P. J.: Drought termination: concept and
16 characterisation, *Prog. Phys. Geogr.*, in review.
- 17 Pyper, B. J. and Peterman, R. M.: Comparison of methods to account for autocorrelation in
18 correlation analyses of fish data, *Can. J. Fish. Aquat. Sci.*, 55, 2127–2140, 1998.
- 19 Quiring, S. M. and Papakryiakou, T. N.: An evaluation of agricultural drought indices for the
20 Canadian prairies, *Agric. For. Meteorol.*, 118(1), 49–62, 2003.
- 21 Sepulcre-Canto, G., Horion, S., Singleton, A., Carrao, H. and Vogt, J.: Development of a
22 Combined Drought Indicator to detect agricultural drought in Europe, *Nat. Hazards Earth*
23 *Syst. Sci.*, 12(11), 3519–3531, doi:10.5194/nhess-12-3519-2012, 2012.
- 24 Stagge, J. H., Kohn, I., Tallaksen, L. M. and Stahl, K.: Modeling drought impact occurrence
25 based on climatological drought indices for Europe., *J. Hydrol.*, 2014.

- 1 Stagge, J. H., Tallaksen, L. M., Gudmundsson, L., Van Loon, A. F. and Stahl, K.: Candidate
2 Distributions for Climatological Drought Indices (SPI and SPEI), *Int. J. Climatol.*, n/a–n/a,
3 doi:10.1002/joc.4267, 2015.
- 4 Stahl, K., Kohn, I., Blauhut, V., Urquijo, J., De Stefano, L., Acacio, V., Dias, S., Stagge, J.
5 H., Tallaksen, L. M., Kampragou, E., Van Loon, A. F., Barker, L. J., Melsen, L. A., Bifulco,
6 C., Musolino, D., de Carli, A., Massarutto, A., Assimacopoulos, D., and Van Lanen, H. A. J.:
7 Impacts of European drought events: insights from an international database of text-based
8 reports, *Nat. Hazards Earth Syst. Sci. Discuss.*, 3, 5453–5492, doi:10.5194/nhessd-3-5453
9 2015, 2015a.
- 10 Stahl, K., Kohn, I., De Stefano, L., Tallaksen, L. . . , Rego, F. C., Seneviratne, S. I., Andreu, J.
11 and Van Lanen, H. A. .: An impact perspective on pan-European drought sensitivity, in
12 *Drought: Research and Science-Policy Interfacing*, edited by Joaquin Andreu, A. Solera, J.
13 Paredes-Arquiola, D. Haro-Monteagudo, and H. van Lanen, pp. 329–334, CRC Press,
14 London, 2015b.
- 15 Steinemann, A.: Drought Information for Improving Preparedness in the Western States, *Bull.*
16 *Am. Meteorol. Soc.*, 95(6), 843–847, doi:10.1175/bams-d-13-00067.1, 2014.
- 17 Steinemann, A. C. and Cavalcanti, L. F. N.: Developing multiple indicators and triggers for
18 drought plans, *J. Water Resour. Plan. Manag.*, 132(3), 164–174, 2006.
- 19 Stölzle, M. and Stahl, K.: Wassernutzung und Trockenheitsindikatoren in Baden-
20 Württemberg, *Standort-Zeitschrift für Angew. Geogr.*, 35(3), 94–101, 2011.
- 21 Strobl, C., Malley, J. and Tutz, G.: An introduction to recursive partitioning: rationale,
22 application, and characteristics of classification and regression trees, bagging, and random
23 forests., *Psychol. Methods*, 14(4), 323–348, 2009.
- 24 Svoboda, M., LeCompte, D., Hayes, M., Heim, R., Gleason, K., Angel, J., Rippey, B., Tinker,
25 R., Palecki, M. and Stooksbury, D.: The drought monitor, *Bull. Am. Meteorol. Soc.*, 83(8),
26 1181–1190, 2002.

- 1 Trambauer, P., Werner, M., Winsemius, H. C., Maskey, S., Dutra, E. and Uhlenbrook, S.:
2 Hydrological drought forecasting and skill assessment for the Limpopo river basin, Southern
3 Africa, *Hydrol. Earth Syst. Sci. Discuss.*, 11(8), 9961–10000, doi:10.5194/hessd-11-9961-
4 2014, 2014.
- 5 Vicente-Serrano, S. M., Beguería, S. and López-Moreno, J. I.: A multiscale drought index
6 sensitive to global warming: the standardized precipitation evapotranspiration index, *J. Clim.*,
7 23(7), 1696–1718, 2010.
- 8 Vicente-Serrano, S. M., Beguería, S., Lorenzo-Lacruz, J., Camarero, J. J., López-Moreno, J.
9 I., Azorin-Molina, C., Revuelto, J., Morán-Tejeda, E. and Sanchez-Lorenzo, A.: Performance
10 of drought indices for ecological, agricultural, and hydrological applications, *Earth Interact.*,
11 16(10), 1–27, 2012.
- 12 Vicente-Serrano, S. M. and López-Moreno, J. I.: Hydrological response to different time
13 scales of climatological drought: an evaluation of the Standardized Precipitation Index in a
14 mountainous Mediterranean basin, *Hydrol. Earth Syst. Sci.*, 9(5), 523–533, 2005.
- 15 Wetterhall, F., Winsemius, H. C., Dutra, E., Werner, M. and Pappenberger, E.: Seasonal
16 predictions of agro-meteorological drought indicators for the Limpopo basin, *Hydrol. Earth
17 Syst. Sci.*, 19(6), 2577–2586, doi:10.5194/hess-19-2577-2015, 2015.
- 18 Wilhite, D. A., Svoboda, M. D. and Hayes, M. J.: Understanding the complex impacts of
19 drought: A key to enhancing drought mitigation and preparedness, *Water Resour. Manag.*, 21,
20 763–774, doi:10.1007/s11269-006-9076-5, 2007.
- 21 Wilhite, D. and Glantz, M.: Understanding: the drought phenomenon: the role of definitions,
22 *Water Int.*, 10(3), 111–120, doi:10.1080/02508068508686328, 1985.
- 23 Wilhite, D. and Svoboda, M.: Drought early warning systems in the context of drought
24 preparedness and mitigation, in *Early Warning Systems for Drought Preparedness and
25 Drought Management*, edited by D. A. Wilhite, M. V. K. Sivakumar, and D. A. Wood, pp. 1–
26 21, World Meteorological Organization, Geneva., 2000.

27

1 Table 1. Information on NUTS1 regions in the UK and Germany (DE) considered for analysis

Country	NUTS1 region name	NUTS1 region abbr.	N_T	Length of censored timeseries (months)	Percentage of months with $N_T > 0$	No. streamflow stations	No. groundwater stations
UK	North East	NEE	28	48	22.9	9	-
UK	North West	NWE	400	120	35.8	16	-
UK	Yorkshire and the Humber	YHU	213	108	32.4	11	-
UK	East Midlands	CEE	345	120	37.5	13	-
UK	Wales	WAL	884	156	35.9	20	-
UK	West Midlands	CWE	310	96	42.7	12	-
UK	East of England	EE	545	156	50.0	12	-
UK	South West	SWE	456	156	57.1	23	-
UK	South East	SEE	1079	168	57.1	23	-
UK	London	LND	291	144	45.1	1	-
DE	Schleswig-Holstein	SH	34	60	25.0	9	9
DE	Mecklenburg-Western Pomerania	MP	54	96	28.1	7	4
DE	Lower Saxony	LS	107	132	28.0	38	42
DE	Saxony-Anhalt	ST	46	96	22.9	16	14
DE	Brandenburg	BB	114	96	30.2	21	18
DE	Berlin	BE	57	72	23.6	-	-
DE	North Rhine-Westphalia	NW	143	84	34.5	23	18
DE	Hesse	HE	95	60	43.3	19	18
DE	Saxony	SX	50	96	31.3	23	10
DE	Rhineland-Palatinate	RP	182	84	35.7	20	18
DE	Saarland	SL	42	36	30.6	3	-
DE	Baden-Wuerttemberg	BW	228	84	39.3	28	15
DE	Bavaria	BV	382	72	33.3	69	26


2

1 Table 2. Model performance metrics of cross-validated random forest models per NUTS1
 2 region.

Country	NUTS1	MAE	RMSE	$\Delta\mu$	$\Delta\sigma$	r	R ²
UK	NEE	0.44	0.58	0.03	-0.49	0.51	0.26
UK	NWE	1.01	1.48	0.06	-0.51	0.40	0.16
UK	YHU	0.57	0.77	0.00	-0.32	0.76	0.58
UK	CEE	0.72	0.96	-0.01	-0.31	0.74	0.54
UK	WAL	0.82	1.25	-0.01	-0.42	0.85	0.73
UK	CWE	0.59	0.88	0.00	-0.22	0.79	0.62
UK	EE	0.71	0.92	-0.02	-0.40	0.79	0.62
UK	SWE	0.55	0.70	0.01	-0.25	0.84	0.70
UK	SEE	0.92	1.23	0.01	-0.38	0.79	0.62
UK	LND	0.67	0.84	0.02	-0.42	0.67	0.45
DE	SH	0.19	0.31	0.08	-0.25	0.90	0.81
DE	MP	0.35	0.48	0.05	-0.46	0.68	0.46
DE	LS	0.38	0.56	0.04	-0.45	0.73	0.53
DE	ST	0.30	0.45	0.10	-0.40	0.68	0.46
DE	BB	0.43	0.62	-0.02	-0.40	0.78	0.61
DE	BE	0.26	0.50	0.08	-0.30	0.79	0.62
DE	NW	0.57	0.87	0.00	-0.52	0.69	0.48
DE	HE	0.61	0.82	0.08	-0.51	0.61	0.37
DE	SN	0.31	0.43	0.00	-0.41	0.71	0.50
DE	RP	0.68	1.03	0.06	-0.44	0.58	0.34
DE	SL	0.56	0.72	0.13	-0.48	0.65	0.42
DE	BW	0.74	1.16	0.02	-0.32	0.58	0.34
DE	BV	0.68	1.21	0.04	-0.27	0.82	0.67

3


4



1

2 Figure 1: Maps displaying NUTS1 regions in the UK (left) and Germany (right), and the
 3 location of streamflow gauging and groundwater monitoring stations. See Table 1 for NUTS1
 4 region abbreviations.


5



1


2 Figure 2: Number of impact occurrences and distribution of impacts per impact category per
 3 NUTS1 region and season for the UK (top four plots) and Germany (bottom four plots).

4




1
 2 Figure 3: UK: rank correlation coefficients (ρ) between drought indicators and number of
 3 impact occurrences for total impacts, hydrological drought impacts (I_h), impacts due to other
 4 types of drought (I_o), and selected impact categories per NUTS1 region.

5




1
 2 Figure 4: UK: rank correlation coefficients (ρ) between drought indicators and number of
 3 impact occurrences per NUTS1 region and season.
 4

1




2

3 Figure 5: Germany: rank correlation coefficients (ρ) between drought indicators and number
 4 of impact occurrences for total impacts, hydrological drought impacts (I_h), impacts due to
 5 other types of drought (I_o), and selected impact categories per NUTS1 region.
 6




1
2
3
4
5

Figure 6: UK: ranks of predictor importance during random forest construction for total impacts, hydrological drought impacts (I_h), impacts due to other types of drought (I_o), and selected impact categories per NUTS1 region.




1

2 Figure 7: Germany: ranks of predictor importance during random forest construction for total
 3 impacts, hydrological drought impacts (I_h), impacts due to other types of drought (I_o), and
 4 selected impact categories per NUTS1 region.



1
 2 Figure 8: UK: distribution of splitting values during random forest construction (i.e.
 3 thresholds of impact occurrence) for selected drought indicator variables for each NUTS1
 4 region. The boxplot whiskers extend to the minimum and the maximum of the distribution,
 5 the box encompasses the interquartile range, and the line inside the box displays the median.



1

2 Figure 9: Germany: distribution of splitting values during random forest construction (i.e.
 3 thresholds of selected drought indicator variables for each NUTS1 region. Boxplots as Figure
 4 8.

5





Figure 10: RF Predictions for different regions in the UK (transformed variables).



1
2
3
4

Figure 11: RF Predictions for different regions in Germany (transformed variables).




Figure 12: Time series of observed and modeled number of impact occurrences for a selection of NUTS1 regions in the UK and Germany (transformed variables). Grey background: RF Predictions, white background: RF Backwards Learning. The blue line indicates an impact threshold of one: modeled impacts smaller than one should be regarded as absent impact.