

Assessing changes ~~in~~ urban flood vulnerability through mapping land use from historical information

M. Boudou¹, B. Danière¹, M. Lang¹

¹ Irstea, UR HHLY, Hydrology-Hydraulics, 5 rue de la Doua, Villeurbanne F-69626, France

Correspondance to: M. Boudou (martin.boudou@irstea.fr)

Abstract

This paper presents an ~~diachronic~~ appraisal of the temporal evolution of flood vulnerability of two French cities, Besançon and Moissac, which ~~werehave been~~ largely impacted by ~~two-past~~ floods in January 1910 and March 1930, respectively. Both flood events figured among the most significant events recorded in France during the ~~XXth~~-20th century, ~~as a function in~~ terms of certain parameters such as the intensity and severity of the flood and spatial extension of the damage. An analysis of historical sources allows the mapping of land use and occupation within the ~~flood-areas affected by~~~~extent of~~ the two ~~historical~~-floods, both in past and present contexts, ~~providing-It gives~~ an insight of the complexity of flood risk evolution, at a local scale.

1 Introduction

Directive 2007/60/EC on the assessment and management of flood risks draws up a new framework for the promotion of historical information. It aims to reduce and ~~to~~-manage the risks that floods pose to human health, the environment, cultural heritage and economic activity. The Directive requires Member States to ~~first first~~ carry out a preliminary assessment by 2011 to identify ~~the the~~-river basins and then the associated coastal areas which are at risk of flooding. For such zones, ~~the following subsequent~~ steps would consist in~~involve~~ drawing up flood risk maps by 2013 and establishing flood risk management plans focused on prevention, protection and preparedness by 2015. The Directive applies to inland waters as well as all coastal waters across the whole territory of the EU. In France, a national Historical Database ~~on floods~~ (<http://bdhi.fr/>), ~~has been opened to the public in 2015,~~ based on the inventory of major floods, ~~in France produced~~was produced in 2011 within the framework of

the EU Flood Directive (Lang and Coeur, 2014; Lang et al., 2012) and was made available to the public in 2015. It contains a description of 176 “remarkable” flood events from 1770 to 2011.

A key issue of the Flood Directive is ~~the to~~ accurately assessment of the flood risk. A commonly accepted definition of flood risk is the combination ~~between of~~ a flood hazard and the vulnerability of ~~the~~ assets that are exposed (~~de~~ Bruijn, K.M., 2005; Schanze, 2006; Cardona et al., 2012). ~~In suit with~~ Following this definition, the French Government distinguished two main steps for flood risk assessment. A first step consists ~~ed of in~~ mapping the potential flood extent ~~to in order to~~ evaluate the number of infrastructure assets exposed. Starting from this data, a second step consists ~~s of determinined in censng~~ the ~~asset~~ asset-exposure and vulnerability of the asset. For this purpose, some indicators ~~haved~~ been adopted, according to the potential impacts on human health, economic activity, the environment and cultural heritage within the potential flood extent. To ~~mention just name~~ a few, ~~they are for instance~~ these indicators include the number of ~~inhabitants affected population exposed~~, the number of ~~single-one~~ storey buildings, the number of employed ~~personsments~~, the number of nuclear power stations, the area of remarkable built heritage, etc. Following this approach, ~~the~~ flood risk assessment ~~drew up leads to~~ a contrasted overview of the actual flood risk. The results indicate a strong and unequal ~~assets~~ asset-exposure of assets over the French territory, and raise some concerns in a context of increasing flood damages (SwissRe, 2015) and global change.

The term “vulnerability” has long been a subject of debate in the scientific literature, being covered by several definitions (Birkmann, 2006; Wisner et al., 1994). A commonly used definition of vulnerability is the likelihood of the elements at risk to produce damage. Based on that definition, assessing the vulnerability and its evolution can be broken down into two main steps: firstly, assessing the exposure by listing the elements at risk and secondly, assessing the susceptibility of the elements at risk (Merz et al., 2007). To carry out these two steps, we identify a series of indicators adapted for a retrospective analysis.

On the one hand, the exposure analysis is supported by quantifying the number of buildings and inhabitants at risk. On the other hand, the susceptibility analysis is based on identifying the building use type, providing some keys for understanding the kind of damage to be expected during floods (Barroca et al., 2006). For example, some building types are especially likely to trigger major damage (industrial or commercial activities) or cause disturbances for

society (e.g. public infrastructures such as hospitals or schools), thus requiring special attention from risk managers (Merz et al., 2007).

Many authors have already highlighted the importance of historical data as a tool for risk assessment (Glade *et al.*, 2001; Brazdil *et al.*, 2006; Coeur and Lang, 2008; Kjeldsen *et al.*, 2014). A general survey of flood mapping techniques in Europe by de Moel et al. (2009) provides evidence that flood maps are available in almost all countries, based on historical floods or design-basis floods. As an example, Barnikel (2004), Tropeano and Turconi (2004) or Luino *et al.* (2012) reported past flood extents in relation to present-day land use, which allows the development of prospective analyses of flood risk.

Assessing flood impacts and understanding the past vulnerability of a territory is an essential step towards a long-term mitigation strategy (Changnon *et al.*, 2000). Firstly, it allows a better understanding of the circumstances that lead to a disaster. Secondly, it helps to shed the light on the actual state of the vulnerability within a territory. This vulnerability (especially visible through the exposure of the assets) has to be seen as the result of a complex historical evolution, partly related to the occurrence of past-damaging flood events in the past (Barrera *et al.*, 2006).

To take account of a potential increase in flood risk, the Flood Directive assessment has to be considered in terms of a long time scale. The indicators developed during the preliminary phase are in fact closely correlated with the present-day situation and raise some questions about the past situation of vulnerability. How do we assess the vulnerability and exposure situations during past flood events based on uncertain and sparse historical sources? Can we confirm an increase in the exposure and vulnerability of stakeholders' exposure and vulnerability based on a temporal analysis of past disasters? Are these disasters still relevant and easily integrated into risk management policies as indicated in the Flood Directive text?

To address these issues, the present study sets out to highlight the interest of historical information through applying a transdisciplinary multidisciplinary and mapping approach (Danière, 2014). The study is based on the set of 176 major French floods in France, which offers an opportunity to explore the vulnerability associated with past flood events. We applied this methodology to two case studies selected for their "remarkability": the January 1910 flood event (generalized over all the North-East of France) and the March 1930 flood event

(~~concentrated~~focused on the Tarn River valley). We ~~focused~~ ~~the-our~~ analysis on two cities, Besançon and Moissac, ~~which were each one~~ largely affected by the floods of 1910 and 1930, respectively~~one of these two events~~. After a brief presentation of the two flood events (section 2), we present the methodological framework used for mapping ~~the~~ ~~the~~-vulnerability (section 3). ~~It-This approach is~~~~has been~~ applied ~~to~~~~on~~ the two case studies (section 4), illustrating the past and present vulnerability situations in the two cities. Finally, some key ~~pointss~~ are given (section 5) ~~about-concerning~~ the ~~interest-importance~~ of historical information for assessing vulnerability changes during the ~~XX~~th 20th century.

2 Case studies

2.1 Selection of two remarkable flood events

During the inventory work carried out for the Flood Directive in 2011, we selected a total of 176 major floods in France since 1770 (see Lang and Coeur, 2014) based on the following considerations: diversity of flood types, strong flood hazard or spatial extent, important socio-economic impacts, in addition to reference events used in planning documents (flood mapping area) or last significant flood in living memory. Using a ~~ransdisciplinary~~ multidisciplinary methodology, ~~we established~~ an evaluation grid based on three main features ~~was established~~ (Boudou *et al.*, 2015): 1/ flood intensity (score between 3.5 to 14) according to several criteria (return period of ~~the~~-maximum peak discharge; duration of submersion; ~~dyke~~ breaches or log jams); 2/ flood severity (score between 3 to 12), with two main indicators: ~~;~~ flood damages (number of fatalities, economic loss) and social, media or political impacts of the event (establishing a new risk policy, calling for international solidarity to face the crisis, ~~etc....~~); 3/ spatial extent~~sion~~ of damages (score between 2 to 8). ~~This grid allowed us~~~~It allowed~~ to rank the 176 major floods (Boudou, 2015). Then, a second level of ~~selection-led~~selection led us to focus on ~~the nine~~~~9~~-events show~~ned~~ in Fig. 1 (Jan. 1910, March 1930, Oct. 1940, Dec. 1947 / Jan. 1948, Dec. 1959, Jan. 1980, Nov. 1999 ~~and~~; Dec. 2000 / April 2001). These flood events cover all flood typologies (oceanic/snowmelt/Mediterranean floods, ~~marine-storm submersions~~surges, cyclones, dam ~~breaching~~eaking) and are considered as some of the most remarkable in accordance with the evaluation grid. Lang et al. (2012) presented the main characteristics of these nine events (except for the 1947-48 flood).

In this ~~study paper~~ we ~~will~~ investigate the two oldest selected events, ~~respectively which took place~~ in January 1910 and March 1930, focusing on the urban situation in ~~Besaneon~~ Besançon and Moissac (Fig. 2). The aim is to focus on two cities ~~that~~ which have been significantly flooded in the past and to understand how their vulnerability to flooding ing has changed ~~until now~~ up to the present day. A detailed inventory of documentary sources on these two events can be found in the online material.¹

2.2 The January 1910 flood event in ~~Besaneon~~ Besançon (Doubs River catchment)

The flood of January 1910 ranks ~~fifth among~~ fifth among the 9 floods selected as remarkable according to the evaluation grid (Fig. 1). This flood event is mostly known for being the most significant flood ~~that~~ affecting the city of Paris, with a return period of about one hundred years for several rivers of the Seine basin. After a very wet end to the year 1909 (450 mm of rainfall in 3 months), the Seine basin received a large amount of rain and snow in January 1910 (about 300 mm in the upper part, 110 mm in the central part and 280 mm in the downstream part). The water level at Paris-Austerlitz was 8.66 m, the second highest historical level after the flood of February 1658 (8.80 m) (Champion, 1858-1864; Goubet, 1997). There were a relatively small number of direct fatalities (7 deaths) plus 9 indirect deaths (several cavity collapses), but the impact within the Paris region was extremely high, with 150 000 persons affected ~~people~~ and economic losses of about 400 million gold francs ~~or~~ (1.5 billion euros, 2015) (Picard, 1910). Despite the fact that a large part of ~~the Northern France~~ the territory was also affected, most of the attention of society and recollection ~~the memory of this event~~ have been focused on Paris. ~~In order to~~ To demonstrate the remarkability of this event, not only for the Seine catchment area but also for more rural regions, we ~~then decided to focus~~ concentrate our study on the Doubs basin where the flood of January 1910 remains one of the most significant historical floods, with ~~and~~ the highest water level being recorded in the city of Besançon (see fig. 3, e.g. Z = 245.55 m at “Poterne, Place la Revolution”). While ~~As~~ the flood event ~~across~~ on the Seine basin ~~was~~ is characterized by a clustering of several oceanic rainfall events, the flood event ~~in~~ on the Doubs basin was triggered by an episode of a heavy rainfall ~~event~~ from ~~the~~ 18 to 21 ~~st of~~ January (between 150 and 250 mm), plus the presence of ~~a~~ extensive ~~large~~ snow cover after a wet winter which led

¹ Auxiliary material is available in the html. doi:XXX

1 ~~to~~ ~~to a~~ significant snow melting. A large part of the old city of ~~Besaneon~~Besançon was
2 flooded, with huge damages~~s~~. Many shops, houses and their basements were inundated,
3 causing important losses of furniture. The streets ~~of the town were~~ also particularly badly
4 ~~affected~~suffered due to the high flow velocity. In total, the cost of the flooding at Besançon is
5 estimated at around 2 million francs (DREAL Franche-Comté *et al.*, 2010), ~~actually~~
6 representing 7.7 million euros ~~2015~~in present-day money.

7 According to several documentary sources (Allard, 1910; Ministère de l'Ecologie, 2011), it
8 appears that the hydro-meteorological conditions of the event (peak discharge at Besançon of
9 about 1750 m³/s, with a return period of about 100 years; catchment area of 4379 km²) cannot
10 explain why the flood level was so high throughout the old city. Such exceptional water levels~~s~~
11 in the city centre ~~were~~is the consequence of energy losses ~~at~~along the bridges of the town.
12 These energy losses were larger than usual (cf. Fig. 3, in comparison with the 1882 and 1896
13 flood events) due to a jam–log jam (about 35 000 m³), resulting from the ~~submersion~~
14 inundation of a paper factory a few kilometres upstream ~~of~~to ~~Besaneon~~Besançon, contributing
15 significantly to a the–raisinge of the water level.

16 Archive sources (especially administrative reports produced by the Chief Engineer of the
17 Ponts-et-Chaussées, Serial S, Doubs departmental archives) also revealed some major failures
18 of ~~the~~–flood warning during the event. Surprised both by the ~~flood~~–arrival and ~~the~~its intensity
19 of the flood, the local authorities did not succeed ~~into~~ ~~establish~~–setting up temporary
20 protectiveng structures at the different opened city gates (“postern gates”), ~~which~~–~~and~~ directly
21 contributed to the ~~inundations~~submersion of the city (Fig. 4)

22 2.3 The March 1930 flood in Moissac (Tarn River catchment)

23 At the end of February 1930, ~~an intense~~–large Mediterranean rainfall event occurred in the
24 South-West of France, with hot and moist air from the Mediterranean Sea penetrating deep
25 into the Massif Central highlands. From 25 February to 4 March, a large area was affected by
26 heavy rainfall (e.g. more than 200 mm over 6000 km² during 4 days), with a maximum of 694
27 mm in 7 days at Saint-Gervais-sur-Mare (spring of the Orb river). The very serious adverse
28 consequences of this rainfall event can be explained by at least two factors. From October
29 1929 to February 1930, high rainfall totals were observed (e.g. 1 177 mm at Lodève, 840 mm
30 at Florac), thus favouring a strong reaction of the basins which were already saturated.

Moreover, a warming in temperature associated with intense rainfall was causing a large amount of snow melting (20 to 100 cm) above 600 m.

Due to its intensity and ~~its~~ unusual date of occurrence-~~date~~ (at the end of a wet winter) the rainfall event triggered ~~to~~ an exceptional flood event (Pardé, 1930). The following flood hazard intensity can be judged exceptional for the downstream part of the Tarn catchment (8000 m³/s at Moissac, 15 400 km²; mean annual discharge 230 m³/s), with a return period of about 250-300 years (Dreal Midi-Pyrénées, 2014). Between 210 and 230 fatalities were recorded during ~~this~~ Tarn River flood ~~event~~ (resp. Bichambis, 1930 and Boudou, 2015), ~~leading to which represents~~ one of the most ~~destructive~~ damaging flood events ever recorded in France and surely the most significant ~~during~~ for the ~~XXth~~ 20th century. The economic loss for the ~~entire surrounding all~~ region ~~around was estimated at~~ is estimating around 1 billion francs, which ~~represents~~ corresponds to 570 million euros 2015 (Journal Officiel de la République Française, 1930).

One of the striking ~~features~~ issues of the disaster can be found in the concentration of ~~the~~ damages in the town ~~city~~ of Moissac (120 deaths ~~out of~~ for a total of 210). Reconstructing and mapping the flood chronology using historical sources ~~provides us with~~ enhances a better understanding of the circumstances of the disaster (Fig. 5). ~~On The~~ 3rd of March 1930, the flood arrived in the town. Before 18:30 the Tarn River was already overflowing the main channel, ~~both on both the south left and north right~~ bank ~~sides~~. Fortunately, the ~~town~~ city centre was protected by three main dikes and the ~~embankment of the~~ railway line embankment. From 18:30 to 23:00, the water level ~~rose~~ raised and the flood extent covered the area between the main dikes at the eastern part of the town ~~city~~. Around 23:00, at the time of maximum discharge-~~value~~ (estimated at around 8000 m³/s), three breaches suddenly appeared along the ~~embankment~~ railway embankment. These ~~breaches~~ led ~~breaches led~~ to a sudden outburst of the dikes and ~~to the~~ final submersion ~~inundation~~ of the town ~~city~~.

According to the ~~death~~ locations of fatalities and the ~~disaster~~ feedback of information on the disasters, the explanation of the high ~~number of fatalities~~ death toll is twofold. Firstly, the rapid influx of water ~~into~~ within the city due to the flash flood and dike failures induced a surprise effect ~~on~~ for the inhabitants of Moissac. Secondly, the collapse of more than 600 houses ~~was~~ is related to the typical kind of housing ~~in~~ of this region, ~~being made built of with~~ raw bricks especially vulnerable to flooding and ~~sustained~~ durable contact with water.

3 Methodology for monitoring changes in flood vulnerability

3.1 Relevance of historical events in the present context?

One of the main requirements of the Flood Directive is to identify areas with a potential high level of flood risk, based on historical floods that would have significant adverse consequences if they occurred again. As the consequences are ~~both dependent on~~ the flood hazard ~~as well as~~ the personal, social and economic assets located in the flood risk zones, one of the main concerns is to assess the ~~changes in evolution over time of~~ local vulnerability of city centres ~~as a function of time~~. In ~~For~~ both case studies, the main casualties and/or economic losses within the catchment were located in ~~one a single municipal area~~. But some aggravating factors ~~are were~~ time depend~~ants~~, such as woody debris upstream ~~of~~ bridges at Besançon or ~~dyke~~ failures ~~to at~~ the east of Moissac. Other aggravating factors ~~are were~~ related to social vulnerability, such as failures ~~of~~ flood warning at Besançon or vulnerable building materials at Moissac.

~~In order~~ ~~To obtain a~~ ~~to~~ better ~~understand~~ understanding of the local disaster process, our study ~~we aims to~~ ~~will~~ monitor changes in flood vulnerability, comparing the past and ~~the~~ present situations. Several questions have to be addressed. Is it possible to ~~assesse~~ ~~correctly depict~~ correctly the changes in the vulnerability over time according to the available sources? Does ~~the~~ mapping of land use provide enough information to identify indicators of vulnerability? Can we establish ~~some~~ scenarios concerning ~~about~~ the impact of a future flood based on a historical flood?

After a preliminary analysis ~~that involves~~ by geo-referencing historical information in the present ~~day~~ context, we ~~then~~ ~~will~~ consider the mapping of land use and ~~the~~ ~~counting~~ ~~estimating of~~ the numbers of the population at risk, while comparing ~~from the~~ past ~~and the~~ ~~to~~ present situations.

3.2 ~~A~~ dynamic mapping to locate historical information

A preliminary step of this ~~studywork~~ consists ~~s of~~ ~~ed in the~~ ~~implement~~ carrying out ~~of a~~ dynamic mapping with a spatial display of the ~~historical information formerly previously~~ collected historical information. The historical corpus made up of various document formats and sources ~~is was~~ included in a GIS by locating the information available. However, ~~S~~ some

place names have ~~however~~ changed since the date of the flood event ~~date, which required thus~~
requiring supplementary ~~work~~ treatment of the data.

~~The~~ ~~Such~~ dynamic consultation of historical information is not only ~~offer~~ interest ~~for~~ ~~to~~
correctly locatinge the various sources of information on flood vulnerability, ~~but~~. ~~It~~ can also
be used to develop risk awareness and risk culture on an exposed territory. As an example, the
high-water mark inventory developed ~~for~~in the Seine river catchment ([www.reperesdecrues-
seine.fr/carte.php](http://www.reperesdecrues-seine.fr/carte.php)) provides ~~a~~ dynamic mapping which is easily understandable and interactive
for the general public, in contrast~~contrary~~ to the maps resulting from hydraulic or
hydromorphogenic modelling (de Moel *et al.*, 2009).

3.3 Evolution of land use

~~We will~~In this section, we address the exposure ~~and susceptibility~~and susceptibility to flood
risk (Fig. 6) using simplified descriptors which remain consistent with the level of data
availability and accuracy of historical information (Barnikel and Becht, 2003, Barnikel,
2004).

Firstly, the exposure analysis is based on the ~~evolution of the~~changes in the population living
per building and provides information about the evolution of ~~for~~ built-up areas ~~evolution~~.
Secondly, susceptibility analysis based on land-use classification provides relevant
information to evaluate the nature of buildings affected during ~~the~~ flooding. ~~Use of~~
~~h~~Historical information is required which at least describesing the land cover on different
dates ~~is required~~. For example, historical maps and aerial photos often depict the built-up
territory for a specific year.

~~In order to~~To perform a spatial analysis of historical maps, it is necessary to integrate them~~their~~
~~integration~~ into a GIS ~~was required~~. Three steps ~~are~~were executed: scanning, georeferencing
and digitalizationing supported by a spatial reference ~~system~~d ~~geometry~~ (Fig. 6a) (Rumsey
and Williams, 2002, Levin *et al.*, 2010). A set of historical maps and aerial photographs
produced by the French National Institute of Geographic and Forest Information (IGN) are
~~was~~ used to depict the extent of built-up areas~~urban extension~~ at the scale of a block of
houses ~~scale~~. A total of 7 topographic maps (from 1911 to 1988) are used for Besançon and
26 aerial photographs for Moissac (from 1947 to 1983). Aerial photographs are favoured in
the case of Moissac because of the inconvenient representation of the town on topographic
maps, which is split between four map plates. These raster data are then~~were~~ imported and

georeferenced. A spatial database (BD TOPO) ~~produced by coming from the~~ IGN, describing the present French territory and its infrastructures, ~~is was~~ used to select control points and ~~to~~ evaluate distortions during the digitizing step. During this last step, information from topographic maps ~~is was~~ vectorized into a unique “historical layer”. In this way, each object ~~is given gets~~ a spatial reality (via the GIS representation) and a temporal reality (by associating a temporal field to indicate its existence for a specific year). Consequently, the “historical layer” ~~makes it possible allows us~~ to ~~depict some obtain~~ “temporal snapshots” (Langran and Chrisman, 1988, Gregory and Healey, 2007) of the urban fabric: ~~the~~ space is discretized based on ~~available information at available at the time of event the event period~~.

Subsequently, the description of “historical layer” objects provides information on the ~~nature kind~~ of building exposure. A land-use classification ~~is was drawn up achieved~~ based on a nomenclature adapted from ~~the an~~ Urban Atlas of ~~the~~ European Environment Agency (<http://www.eea.europa.eu/data-and-maps/data/urban-atlas>), according to historical information constraints (Fig. 6b). A first geomatic processing ~~step is performed was run~~ to discretize the residential buildings on a 0.25 hectare grid. ~~In each mesh, a A~~ density criterion ~~is was~~ applied ~~in each grid cell~~, based on the ~~percentage contribution to the art of buildings~~ footprint, leading to a ~~partition distinction~~ between dense and sparse areas. ~~In order to~~ To enhance the classification, a second processing ~~step is carried out was then run~~, using a proximity criterion for each building, ~~based on by~~ the number of buildings within a 200-~~meters~~ radius (continuous and discontinuous buildings). Local information ~~is then added~~ related to the location and ~~the natures~~ of non-residential constructions ~~were added~~. BD TOPO data ~~are were~~ used to describe ~~the current situation time~~, and a ~~point-in-time unetual~~ layer ~~is was~~ built with our “historical corpus” information for ~~ancient time~~ earlier historical periods.

3.4 Census of the exposed population within the flood extent

General information is provided by the evolution of population at the scale of the municipality. Figure 7 presents the data derived from several population censuses during the 20th century. It shows than the number of inhabitants has grown by about + 100 % at Besançon (from 57 978 to 116 914, between 1911 and 2010) and + 60 % at Moissac (from 7 814 to 12 354, between 1911 and 2006). As only part of the built-up area was affected by floods, especially in the case of Besançon, it is necessary to cross two layers of information: the number of inhabitants per small block and the spatial extent of the historical flood (1910 or 1930 floods at Besançon and Moissac, respectively).

Human exposure ~~is taken into account ed for~~, by census or an estimation of ~~the~~ resident population. The aim ~~here is was to disperse distribute thea~~ raw demographic data throughout the blocks of houses by following its evolution at different scales (Wu *et al.*, 2008). ~~The mMaps so produced canould shed the~~ light on the evolution of human exposure within the ~~area affected by the~~ flood ~~extent~~.

To assess the current ~~population~~ living ~~population~~ within the flood extent, we ~~applied a formula to redistribute at block of house scale make use of~~ two ~~demographic data sets produced by the~~ French National Institute for Statistics and Economic Studies (INSEE) ~~demographic data sets, applying formula (1) to redistribute the population data at the scale of blocks of houses.~~ The first ~~datasetone~~ is defined at infra-municipal scale ~~withwith~~ IRIS data ~~use~~ (Infra-urban statistical area). The second ~~datasetone~~ is based on ~~an estimation of the~~ fiscal population ~~withestimation~~ in a 200 x 200 m grid. These datasets ~~arewere~~ distributed ~~through at the scale of~~ residential blocks of houses, based on a volumetric ~~trictie~~ method (Lwin and Murayama, 2009), in proportion ~~to theof~~ building footprint ~~area multiplied bytimes~~ the vertical density, ~~according tousing~~ the building height [provided by BD TOPO](#):

$$\text{Developped area} = \frac{\text{building height} \times \text{building floor area}}{\text{average storey height}} \quad (1)$$

Historical information, ~~in the form ofas an old a~~ census or ~~a~~ raw demographic data, ~~iswas~~ required to ~~census or to~~ estimate (Ekamper, 2010) the ~~numbers of the population~~ exposed ~~population~~ at the time of ~~the~~ disaster. General census reports are available for every French municipality (sometimes online), generally ~~compiled~~ every 5 years ~~up~~ until 1946, with some exceptions. These documents contain ~~nominative information about the~~ municipal population ~~in nominative list, gathered grouped~~ by building and street, ~~aton~~ different dates. The comparison between past and present exposed population within the flood extent should take account ~~of the~~ possible ~~changesevolutions~~ of census methodology over time.

4 Change of vulnerability based on two case studies

We ~~will~~ now consider the changes of vulnerability ~~inon~~ the two case studies, from past to present, using historical sources and current information.

4.1 Changes in vulnerability of Besançon vulnerability with respect to the January 1910 flood

Figure 8 displays the land use within the area affected by the 1910 flood ~~extent~~ in Besançon, based on the situations in 1911 and 2013 ~~contexts~~ (resp. dates of two censuses). No significant change can be seen in terms of ~~on~~ vulnerability, according to the spatial extent of the ~~-~~built-up area. Since the centre of ~~As Besançon~~ Besançon downtown is located within a meander of the Doubs River, with no opportunity ~~for~~ spatial expansion or urban densification, there has been ~~was~~ no increase of exposure, apart ~~from~~ the hospital area. ~~Although~~ Despite the city has experienced a spatial expansion towards the ~~n~~North, on the right bank, this area ~~is~~ is located outside our zoning at a larger scale.

According to the land use classification, we can not ~~ice~~ see significant changes ~~with~~ in the various activities. There has been ~~was~~ a fall in military ~~function~~ employment, in favour of an increase ~~in of the~~ administrative and public ~~facilities~~ function. While ~~the~~ military areas have decreased ~~by of~~ 74% between 1911 and 2013, ~~the~~ administrative areas have ~~were multiplied~~ grown by a factor of 12. A reduction of human exposure is noticeable between ~~1910~~ 1911 (the census year closest to the 1910 flood) and 2013, with a 24% decrease in the city-centre ~~of the~~ downtown population.

The demographic evolution is represented on Fig. 9 at the scale of a block of houses ~~scale~~, reflecting the ~~household~~ decrease in household size (~~reduction~~ decline in the number of inhabitants per building) and some decline in ~~removal of~~ residential function (reduction of inhabited buildings within the city centre ~~downtown~~).

4.2 Changes in vulnerability of Current Moissac with respect vulnerability to the March 1930 flood

The flood risk mapping of Moissac ~~cartography gives~~ yields an opposite diagnosis ~~istie~~, with an ~~important-major~~ increase of vulnerability within the area affected by the 1930 flood ~~extent~~ (Fig. 10). Built-up ~~surface~~ areas have expanded by 122% between 1930 and 2013. Such spatial extension is explained by new residential development (mainly housing estates) and economic buildings ~~on the East of the downtown city centre~~ and by a progressive densification of ~~the~~ the low-density area on the south ~~left~~ bank flood plain.

Despite a new distribution of the population (Table 1), the human exposure ~~did~~ has not ~~not~~ change significantly changed. The reduction ~~of of the downtown~~ population density in the city

centre is compensated by a spatial expansion (Fig. 11). The human exposure has mainly increased on the ~~downtown~~ east side of the city centreside, especially in the area located between the two levees. It should be noted that no general census report ~~is~~was available for Moissac in the 1930s. The~~refore, the -1930-exposed~~ population exposed to flood risk in 1930 was ~~therefore~~ estimated ~~through-from~~ a raw-rough demographic data set, ~~obtained~~provided from an internet database ~~holding-containing a~~ historical population census at ~~the~~a municipality scale (<http://cassini.ehess.fr/>), ~~which was~~and then ~~distributed according to~~persed ~~based-on~~ the volume-based~~trie~~ method.

4.3 ~~An~~ appraisal of the temporal evolution of flood risk

These two case studies shed ~~the~~ light on the complexity of flood-risk evolution. At ~~the a large scale of a country~~nation-wide scale, it is clearly ~~acknowledged~~admitted that the increase of flood damages ~~during-over~~ the last few decades is induced by a general increase ~~in~~ing-of flood vulnerability (Kron, 2002, Luino *et al.* 2012, Kundzewicz *et al.*, 2014, Smith *et al.*, 2014). At a local scale, where topographic, social and economic contextss are crucial, it is necessary to have a more detailed~~n in-depth~~ analysis.

In Besançon, there has been no extension of the urban area within the old city since 1910, but significant land-use changes have led to a decrease of flood vulnerability as some previously residential areas are now used as administrative buildings buildings~~flood-risk-vulnerability decreased since 1910, but with significant land use changes.. Submersion-~~The frequency of flooding has changed in the historical centre, due to the establishment of safety measuress establishment, especially with the construction of mitigation structures such as cofferdams to close the postern-gates. Some uncertainties remain ~~forte determining~~represent the flooded area in the case of an 1910-event flood~~comparable to the 1910 reference flood, since~~as opposite effects come into play. The log jams at the bridges are not expected to be repeated, but additional hydraulic losses have been introduced by new hydraulic structures since 1910. Nowadays, the reference flood selected in the regulatory documents is a simulated flood larger than the January 1910 flood.

In Moissac, the ~~trajectory-of-the~~changes in vulnerability ~~show~~follows a more contrasted ~~pattern~~evolution. As in various other French regions, the built-up areas ~~city-experienced-have grown-a-growth in~~ spatial extentsion since 1930, characterized by an important development of housing estatess-development. One critical point is the development of one-storey buildings,

leading to a higher human ~~and structural~~ vulnerability due to the lack of a refuge floor. ~~At the~~ ~~On the other hand~~ ~~opposite~~, building quality has improved. During the 1930 flood, the house collapses in Moissac and the ~~consequent~~ ~~related~~ fatalities were closely related to the construction materials ~~used for its construction~~. ~~In order to~~ ~~increase~~ ~~weakness~~ ~~the resistance of~~ ~~in~~ the structures, new materials and ~~architecture~~ ~~building~~ techniques were ~~then~~ used during the reconstruction ~~stage~~ age. Another positive ~~evolution~~ ~~change~~ is related to the improvement of safety measures, due to progress in ~~both~~ ~~flood~~ ~~warning~~ ~~decision~~ ~~making~~ as well as ~~and~~ regards emergency population evacuation schemes implemented by the civil protection services. ~~Today,~~ ~~the~~ 1930 flood in Moissac, ~~with~~ ~~which~~ return period ~~is~~ estimated at around 250 years, is nowadays considered as the reference flood hazard ~~both~~ for the local flood risk management strategy as well as ~~and~~ for planning and development documents. This territory ~~would~~ appears to remain vulnerable, especially to risks of ~~dyke failure~~ ~~risks~~.

5 Conclusion and perspectives

This ~~studypaper~~ ~~presented~~ a case study on the urban vulnerability of two French cities ~~which have~~ ~~that were~~ ~~been~~ largely impacted by ~~two~~ ~~past~~ floods occurring in January 1910 and March 1930. ~~This approach~~ ~~It~~ gives an insight ~~into~~ ~~of~~ the complexity of flood risk evolution, ~~while also taking~~ ~~with~~ local characteristics into account. Mapping historical sources can provide reliable information on the ~~past~~ flood vulnerability in the past, but this requires given some preliminary work. A first step is necessary to ~~correctly~~ locate and geo-reference the historical information within the present geographical reference system. Qualitative information (~~images~~ ~~pictures~~, technical reports, national and local newspaper articles, paintings, marble plaques, etc. historical accounts...) can be interpreted ~~as~~ ~~at~~ complement some to historical maps on land use. ~~An~~ ~~The~~ assessment of the population at risk within spatial units can be ~~deducted~~ inferred from technical documents with nominative lists of persons as well from old censuses. Historical information on past floods can therefore be useful when building scenarios on ~~the~~ future possible floods, providing a reliable reference of what might be possible in terms of water depth, flow velocity and flood extent. Additional work is needed to account for possible changes both in vulnerability and flood hazard over the past several decades (from historical floods to the present day) and for future decades (prospective studies). –It is also important to ~~bear~~ keep in mind the uncertainties associated ~~uncertainties~~ ~~with~~ ~~on~~ historical data and to use relevant scales when mapping vulnerability indicators.

As usual, ~~thea diachronic appraisal~~ temporal analysis of flood risk evolution at a local scale implies a good knowledge of the general context of the socio-economic development of territories, as well as ~~evolutions-changes in the of risk-memoryrecollection~~ and perception ~~of risk~~. According to ~~the~~ data availability, this ~~studypaper~~ focuses ~~ed onon only~~ a small ~~part component~~ of vulnerability. ~~However, In order to complete carry out a comprehensive total~~ flood vulnerability analysis, ~~some~~ other indicators should ~~however~~ be taken into account. After ~~the~~ Xynthia storm surges in 2010 (41 fatalities due to floods in France), Vinet *et al.* (2012) showed ~~for instance~~ that the age of the population ~~age~~ is a key component of local vulnerability. It is clear that the insurance system may benefit from similar analyses on urban flood vulnerability over the last few decades.

This ~~paper study addresses the issue offocused on~~ flood vulnerability, ~~which-that~~ is an important ~~componentpart~~ of the flood risk. ~~In Pparallel, research-work is-however-also necessary-on flood hazard is also necessary,-in-order~~ to simulate past floods in a present-day context, taking into account modifications of the river (morphological changes and river engineering) and new settlements on the flood plain.

6 Author contribution

M. Boudou established the evaluation grid used for the selection of “remarkable” flood events. He collected data on the two historical floods and produced thematic maps on flood hazard. B. Danière carried out dynamic mapping to locate historical information and thematic maps on flood vulnerability. M. Lang ~~did the-supervised~~ edion of the drafting-writing of the paper.

7 Acknowledgements

The authors ~~would like to~~ especially thank the DREAL of Besançon, the DDT of Moissac, and the IGN ~~Institute~~ for providing data. We are also ~~grateful tothank~~ Freddy Vinet and Denis Cœur for their advices. Maria-Carmen Llasat and two anonymous referees are acknowledged for their useful comments. Finally, the authors would like to thank the French Minister of Ecology, Sustainable ~~D~~development and Energy (MEDDE) for the financial support of Martin Boudou’s PhD. Michael Carpenter post-edited the English style and grammar.

8 References

Allard, M.: Les récentes inondations à Besançon, Bibliothèque et archives municipales de la ville de Besançon, 1910.

Barnikel, F., and Becht, M.: A historical analysis of hazardous events in the Alps? the case of Hindelang (Bavaria, Germany), *Natural Hazards and Earth System Science*, 3(6), 625-635, 2003. Barnikel, F.: The value of historical documents for hazard zone mapping, *Natural Hazards and Earth System Science*, 4(4), 599-613, 2004.

Barrera, A., Llasat, M. C., and Barriendos, M.: Estimation of extreme flash flood evolution in Barcelona County from 1351 to 2005, *Natural Hazards and Earth System Sciences*, 6, 505-518, 2006.

[Barroca, B., Bernardara, P., Mouchel, J.-M., and Hubert, G.: Indicators for identification of urban flooding vulnerability, *Natural Hazards and Earth System Science*, 6, 553-561, 2006.](#)

[Bichambis, P.: Inondations du midi en mars 1930 : les paisibles rivières devenues torrents de ruine et de mort. Les deuils, les ruines, les héros, 128p., 1930](#)

[Birkmann, J.: Measuring vulnerability to promote disaster-resilient societies: Conceptual frameworks and definitions, *Measuring vulnerability to natural hazards: Towards disaster resilient societies*, 9-54, 2006.](#)

Boudou, M.: Approche multidisciplinaire pour la caractérisation d'inondations remarquables : enseignements tirés de de neufs évènements en France (1910-2010). PhD, Univ. Montpellier 3, 2015.

[Boudou, M., Coeur, D., Lang, M., Vinet, F.: Grille de lecture pour la caractérisation d'événements remarquables d'inondation en France : exemple d'application pour la crue de mars 1930. In *Environnement, politiques publiques et pratiques locales*, Toulouse, 2015.](#)

[Brazdil, R., Kundzewicz, Z.W., Benito, G.: Historical hydrology for studying flood risk in Europe. *Hydrol. Sci. J.* 51 \(5\), 739–764, 2006.](#)

[Champion, M.: Les inondations en France depuis le VI^e siècle jusqu'à nos jours. Re-édition Cemagref Editions, Paris, 2000, 6 volumes, 1858-1864.](#)

[Cardona, O. D., Van Alast, M. K., Birkmann, M., Fordham, M., McGregor, G., Perez, R., Pulwarty, R. S., Schipper, E. L. F., and Sinh, B. T.: Determinants of risk: exposure and vulnerability, in: *Managing the Risks of Extreme Events and Disasters to Advance Climate*](#)

[Change Adaptation, edited by: Field, C. B., V. Barros, T.F. Stocker, D. Qin, D.J. Dokken, K.L. Ebi, M.D. Mastrandrea, K.J. Mach., and G.-K. Plattner, S. K. A., M. Tignor, and P.M. Midgley, A Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change \(IPCC\), Cambridge University Press, Cambridge, UK, and New York, NY, USA, 65-108, 2012.](#)

Changnon, S. A., Pielke, R. A., Changnon, D., Sylves, R. T., and Pulwarty, R.: Human Factors Explain the Increased Losses from Weather and Climate Extremes, Bulletin of the American Meteorological Society, 81, 437-442, 2000.

[Coeur, D., Lang M.: Use of documentary sources on past flood events for flood risk management and land planning. C.R. Geoscience, Thematic issue « Ecosystems and extreme climatic events », Académie des Sciences, Paris. 340, 644-650, 2008.](#)

Danière, B.: Analyse cartographique de l'évolution de la vulnérabilité en zone urbaine face aux inondations dites remarquables. Master 2 Univ. J. Monet Saint-Etienne, Irstea Lyon, 111 pp., 2014.

[de Bruijn, K.M.: Resilience and flood risk management: a systems approach applied to lowland rivers. PhD dissertation, Delft Univ., 210p., 2005.](#)

de Moel, H. D., Alphen, J. V., and Aerts, J. C. J. H.: Flood maps in Europe—methods, availability and use, Natural Hazards and Earth System Science, 9(2), 289-301, 2009.

[DREAL Franche-Comté, EPTB Saône-et-Doubs, Ville de Besançon: 1910: la Crue du siècle à Besançon - Dossier de Presse, www.franche-comte.developpement-durable.gouv.fr, 9 p., 2010.](#)

[Dreal Midi-Pyrénées: Mise en œuvre de la Directive Inondation. Rapport d'accompagnement des cartographies du TRI Montauban Moissac. 29p + annexes, 2014.](#)

Ekamper, P.: Using cadastral maps in historical demographic research: Some examples from the Netherlands, The History of the Family, 15(1), 1-12, 2010.

[Glade, T., Albini., P., Frances, F.: The use of historical data in natural hazard assessments Advances in Natural and Technological Hazards Research, Kluwer Academic Publishers, 220p, 2001.](#)

[Goubet, A.: Les crues historiques de la Seine à Paris, La Houille Blanche, 8, 23-27, 1997.](#)

- Gregory, I. N., and Healey, R. G.: Historical GIS: structuring, mapping and analysing geographies of the past, *Progress in Human Geography*, 31(5), 638-653, 2007.
- [Journal Officiel de la République Française: Loi portant création d'un fonds provisionnel d'un milliard de francs, en vue de la réparation des dommages de caractère exceptionnel causés par les orages et les crues du 1er au 30 mars 1930, 88, 3970, 11 avril 1930.](#)
- [Kjeldsen, T. R., Macdonald, N., Lang, M., Mediero, L., Albuquerque, T., Bogdanowicz, E., Brazdil, R., Castellarin, A., David, V., Fleig, A., Gül, G.O., Kriauciuniene, J., Kohnova, S., Merz, B., Nicholson, O., Roald, L.A., Salinas, J.L., Sarauskienel, D., Sraj, M., Strupczewski, W., Szolgay, J., Toumazis, A., Vanneuville, W., Veijalainen, N., Wilson, D.: Documentary evidence of past floods in Europe and their utility in flood frequency estimation, J. Hydrology, 517, 963-973, doi: 10.1016/j.jhydrol.2014.06.038, 2014.](#)
- Kron, W.: Keynote lecture: Flood risk= hazard× exposure× vulnerability, *Proceedings of the Flood Defence*, 82-97, 2002.
- Kundzewicz, Z. W., Kanae, S., Seneviratne, S. I., Handmer, J., Nicholls, N., Peduzzi, P., Mechler, R., Bouwer, L.M, Arnell, N., Mach, K., Muir-Wood, R., Brakenridge, G.R., Kron, W., Benito, G., Honda, Y., Takahashi, K. and Sherstyukov, B.: Flood risk and climate change: global and regional perspectives, *Hydrological Sciences Journal*, 59 (1), 1-28, 2014.
- Lang, M, Coeur, D., 2014. Les inondations remarquables en France. Inventaire 2011 pour la directive Inondation, Ed. Quae, 640p.
- Lang, M., Coeur, C., Bacq, B., Bard, A., Becker, T., Bignon, E., Blanchard, R., Bruckmann, L., Delserieys, M., Edelblutte, C. and Merle, C.: Preliminary Flood Risk Assessment for the European Directive: inventory of French past floods. In “Comprehensive Flood Risk Management”, Kjlin & Schweckendiek Ed., ISBN 978-0-415-62144-1, 1211-1217, 2012.
- Langran, G., and Chrisman, N. R.: A framework for temporal geographic information. *Cartographica, The International Journal for Geographic Information and Geovisualization*, 25(3), 1-14, 1988.
- Levin, N., Kark, R., and Galilee, E.: Maps and the settlement of southern Palestine, 1799–1948: an historical/GIS analysis, *Journal of Historical Geography*, 36(1), 1-18, 2010.

- 1 [Luino, F., Turconi, L., Petrea, C., Nigrelli, G.: Uncorrected land-use planning highlighted by](#)
- 2 [flooding: the Alba case study \(Piedmont, Italy\). Nat. Hazards Earth Syst. Sci., 12, 2329-2346,](#)
- 3 [2012.](#)
- 4 Lwin, K., and Murayama, Y.: A GIS Approach to Estimation of Building Population for
- 5 Micro spatial Analysis, Transactions in GIS, 13(4), 401-414, 2009.
- 6 [Merz, B., Thieken, A., and Gocht, M.: Flood risk mapping at the local scale: concepts and](#)
- 7 [challenges, in: Flood risk management in Europe, Springer, 231-251, 2007.](#)
- 8 [Ministère de l'Ecologie: L'évaluation préliminaire des risques d'inondation 2001. Bassin](#)
- 9 [Rhône Méditerranée - Partie III Unité de présentation du Doubs, 159-177, 2011](#)
- 10 Pardé, M.: La crue de mars 1930 dans le sud et le sud-ouest de la France: Genèse de la
- 11 catastrophe, Revue Géographique des Pyrénées et du sud-ouest, 1 (IV), 3-99, 1930.
- 12 [Picard, A.: Rapport de la commission chargée d'analyser les inondations sur le bassin de la](#)
- 13 [Seine de janvier 1910. Rapport au président du Conseil et au ministère de l'Intérieur, Paris,](#)
- 14 [IN, 1910.](#)
- 15 Rumsey, D. and Williams, M.: Historical maps in GIS, Knowles, A.K, editor Past time, past
- 16 place: GIS for history, Redlands, CA : ESRI Press, 1-18, 2002.
- 17 [Schanze, J.: Flood risk management – A basic framework, In Flood Risk Management:](#)
- 18 [Hazards, Vulnerability and Mitigation Measures, Chap I, Springer, 1-20, 2006.](#)
- 19 Smith, A., Martin, D., and Cockings, S.: Spatio-Temporal Population Modelling for Enhanced
- 20 Assessment of Urban Exposure to Flood Risk, Applied Spatial Analysis and Policy, 1-19,
- 21 2014.
- 22 SwissRe: Natural catastrophes and man-made disaster in 2014: convective and winter storms
- 23 generate most losses, Sigma, 2, 52, 2015.
- 24 [Tropeano, D., Turconi, L.: Using Historical Documents for Landslide, Debris Flow and](#)
- 25 [Stream Flood Prevention. Applications in Northern Italy. Nat. Hazards, 31, 663–679, 2004.](#)
- 26 Vinet, F., Lumbroso, D., Defossez, S., and Boissier, L.: A comparative analysis of the loss of
- 27 life during two recent floods in France: the sea surge caused by the storm Xynthia and the
- 28 flash flood in Var, Natural hazards, 61, 1179-1201, 2012.

1 [Wisner, B., Blaikie, P., Cannon, T., and Davis, I: At risk: natural hazards, people's](#)
2 [vulnerability and disasters, London, Routledge, 284p, 1994.](#)

3 Wu, S. S., Wang, L., & Qiu, X.: Incorporating GIS building data and census housing statistics
4 for sub-block-level population estimation, The Professional Geographer, 60(1), 121-135,
5 2008.

6
7

1 **Tables**

2 Table 1. Exposed population in 1930 and 2013 for each ~~submersion~~flooded area (cf. Fig.
3 11) in Moissac

4

Flooded area (Fig. 11)	1930	2013
(1)	4089	1160
(2)	1044	2880
(3)	2267	2000
Total	7400	6040

5

6

7

Figure captions

Figure 1. Location map of the ~~nine~~⁹ most remarkable French flooding events selected in this study and table ~~showing~~^{of} their related remarkability scores ~~related~~ (Boudou, 2015)

Figure 2: Location of the case studies: (left) Doubs basin and Besançon~~-city~~; (right) Tarn basin and Moissac~~-city~~

Figure 3. Longitudinal profile of the Doubs River within the old city of Besançon and ~~flood~~ inter-comparison of floods (sources: [Ville de Besançon – Service de la voirie et des eaux : Profil en long des crues du Doubs du 21 janvier 1910, 28 décembre 1882 et 10 mars 1896, 10 mars 1910, Bibliothèque et archives municipales de Besançon, série 0\).](#) [Locations of République and Battant bridges are shown on Fig. 4](#)

Figure 4: Old Besançon city centre with characteristic water inlets during the flood event on 17 to 21 February 1910~~flood event~~

Figure 5. Flood chronology and location of fatalities during the ~~3-March-1930~~ flood event in ~~the city of~~ Moissac on 3 March 1930

Figure 6. Evolution of vulnerability: (a) exposure; (b) susceptibility (building use type)


[Figure 7. Evolution of the number of inhabitants during the 20th century at Besançon and Moissac. Source: EHESS-Cassini before 1962, INSEE from 1968](#)

Figure 8. Land use types and soil occupation within the area affected by the 1910 flood ~~extent~~ in ~~Besaneon~~Besançon: a/ in 1911; b/ in 2013

Figure 9. Estimated number of inhabitants per building within the area affected by the 1910 flood ~~extent-area~~ in ~~Besaneon~~Besançon: (a) in ~~1910~~1911; (b) in 2013. Some blocks of houses are depicted only on one of the maps~~in only one map~~, because of land-use changes. Non-residential blocks of houses are not taken into account here

Figure 10. Land use types and soil occupation within the area affected by the 1930 flood ~~extent~~ in Moissac: (a) in 1930; (b) in 2013

Figure 11. Estimated number of inhabitants per building within the area affected by the 1930 flood ~~extent-area~~ in Moissac: (a) in 1930; (b) in 2013



3


4

5

6

Figure 1. Location map of the nine most remarkable French flood events selected in this study and table showing their related remarkability scores (Boudou, 2015)

1




2

3 **Figure 2: Location of the case studies: (left) Doubs basin and Besançon; (right) Tarn basin**
4 **and Moissac**

5 ~~(left) Doubs basin and Besançon city; (right) Tarn basin and Moissac city~~

6

1
2



3
4
5
6
7
8
9
10

Figure 3. Longitudinal profile of the Doubs River within the old city of Besançon and inter-comparison of floods (sources: Ville de Besançon – Service de la voirie et des eaux : *Profil en long des crues du Doubs du 21 janvier 1910, 28 décembre 1882 et 10 mars 1896, 10 mars 1910*, Bibliothèque et archives municipales de Besançon, série 0). Locations of Republique and Battant bridges are shown on Fig. 4





Figure 4: Old Besançon city centre with characteristic water inlets during the flood event on 17 to 21 February 1910

1

2



3

4 Figure 5. Flood chronology and location of fatalities during the flood event in Moissac on 3
5 March 1930

6




Figure 6. Evolution of vulnerability: (a) exposure; (b) susceptibility (building use type)




Figure 7. Evolution of the number of inhabitants during the 20th century in Besançon and Moissac. Source: EHESS-Cassini before 1962, INSEE from 1968




Figure 8. Land use types and soil occupation within the 1910 flood extent in Besançon:
a/ in 1911; b/ in 2013




Figure 9. Estimated number of inhabitants per building within the area affected by the 1910 flood in Besançon: (a) in 1910; (b) in 2013. Some blocks of houses are depicted on only one of the maps, because of land-use changes. Non-residential blocks of houses are not taken into account here





Figure 10. Land use types and soil occupation within the area affected by the 1930 flood in Moissac: (a) in 1930; (b) in 2013

1



Source : BDTopo IGN, IRIS Data (INSEE) (C) Danière, Boudou (2015)

Figure 11. Estimated number of inhabitants per building within the area affected by the 1930 flood ~~extent area~~ in Moissac: (a) in 1930; (b) in 2013