Hydrol. Earth Syst. Sci. Discuss., 12, C4926–C4927, 2015 www.hydrol-earth-syst-sci-discuss.net/12/C4926/2015/ © Author(s) 2015. This work is distributed under the Creative Commons Attribute 3.0 License.



HESSD

12, C4926–C4927, 2015

Interactive Comment

Interactive comment on "Impact of two different types of El Niño events on runoff over the conterminous United States" by T. Tang et al.

Anonymous Referee #2

Received and published: 18 November 2015

The author explored the impact of two different type of El Nino (canonical-type and modoki-type) on runoff over the U.S. The topic treated here is quite interesting from both scientific and societal viewpoints. The impact of the present study must be huge because people in the US are very keen about such information. The analysis seem to be good for understanding their impacts in the present-day. However, it does not seem to be enough for the future response in a warmer climate as discussed in the following. Therefore, additional analysis may be necessary prior to acceptance.

1)The composite analysis by Figs. 1-7 seems to be enough to explore the impact of two different type of El Nino in the present-day. However, only Table 2 doesn't seem to be enough for exploring the future response in a warmer climate. It is better to add some figures related to the response to global warming, for example, same as Fig. 4,





but for the period of 2050-2099. Also, the author should touch on the topic of how the two-types of El Nino will change in a warmer climate, for example, frequency, intensity, and its atmospheric response. Cai et al (2015) may be helpful for that discussion.

Cai et al., Nature Climate Change, 5, 849–859 (2015)

2)P3L17: Weng et al. (2007; 2009) explored the teleconnection by El Nino Modoki relative to canonical type.

Weng, H., K. Ashok, S. K. Behera, S. A. Rao, and T. Yamagata, 2007: Impacts of recent El Nino Modoki on dry/wet conditions in the Pacific rim during boreal summer. Climate Dyn., 29, 113–129. Weng, H., S. K. Behera, and T. Yamagata, 2009: Anomalous winter climate conditions in the Pacific rim during recent El Nino Modoki and El Nino events. Climate Dyn., 32, 663–674.

3)P5L11: The period of the analysis is 1999-2009? In the Fig. 2 caption, the period of 1990-2009 is written.

Interactive comment on Hydrol. Earth Syst. Sci. Discuss., 12, 8977, 2015.

HESSD

12, C4926-C4927, 2015

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

