Hydrol. Earth Syst. Sci. Discuss., 2, S564–S565, 2005 www.copernicus.org/EGU/hess/hessd/2/S564/ European Geosciences Union © 2005 Author(s). This work is licensed under a Creative Commons License.



HESSD

2, S564-S565, 2005

Interactive Comment

Interactive comment on "Satellite data interpretation of causes and controls on groundwater-seawater flow directions, Merseyside, UK: implications for assessing saline intrusions" by S. Mukherjee et al.

S. Mukherjee et al.

Received and published: 19 August 2005

I am thankful to the Anonymous Referee #1 in all his points to improve the paper "Satellite data interpretation of causes and controls on groundwater-seawater flow directions, Merseyside, UK: implications for assessing saline intrusions". Following minor changes may kindly be done:

- 1. Abstract, line 3: "by", not "my".
- 2. Abstract, line 4: "numerous faults with vertical displacements of 300m. (as much word removed)
- 3.Results. line 10: "than" not "that".

Full Screen / Esc

Print Version

Interactive Discussion

Discussion Paper

FGU

- 4.Discussion, line 22: This model coincide with the already established fact that in the areas where the water table falls below sea level, the flow is from the sea to the aguifer.
- 5.Chapter 7.3, line 8: The role of faults in controlling groundwater discharge (to the river) adjacent to areas with relatively elevated water tables has been confirmed using satellite data.
- 6. Chapter 7.3, line 16: Please read the entire sentence as following: Wherever sand is present in the opening of the fault, it allows the water to flow both sides, whereas, wherever clay is present, water is not allowed to move both sides. Thus presence of sand (in the fault opening) makes mixing of groundwater and estuary water possible.

Interactive comment on Hydrology and Earth System Sciences Discussions, 2, 887, 2005.

HESSD

2, S564-S565, 2005

Interactive Comment

Full Screen / Esc

Print Version

Interactive Discussion

Discussion Paper

EGU

S565