



Papers published in *Hydrology and Earth System Sciences Discussions* are under open-access review for the journal *Hydrology and Earth System Sciences*

Groundwater ecohydrology: GIScience tools to forecast change and sustainability of global ecosystems, studies in Africa, Europe and North America

D. R. Steward¹, W. J. de Lange², X. Yang¹, S. L. Vasak³, and T. N. Olschoorn⁴

¹Department of Civil Engineering, Kansas State University, 2118 Fiedler Hall, Manhattan, Kansas 66506-5000, USA

²Deltares, Princetonlaan 6, 3508AL Utrecht, The Netherlands

³International Groundwater Resources Assessment Centre, Princetonlaan 6, 3508TA Utrecht, The Netherlands

⁴Water Resources Section, Faculty of Civil Engineering and Geosciences, Delft University of Technology, Stevinweg 1, P.O. Box 5048, 2628 CN Delft, The Netherlands

Received: 19 February 2009 – Accepted: 28 February 2009 – Published: 31 March 2009

Correspondence to: D. R. Steward (steward@ksu.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Groundwater
ecohydrology studies
in Africa, Europe and
North America

D. R. Steward et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

This study examines the interface between groundwater hydrology and ecology, and addresses a scientific grand challenge to develop a comprehensive, systematic understanding of continental water dynamics by linking the hydrosphere and biosphere.

5 There exists a current lack of data interoperability between groundwater modeling tools due to differences in numerical techniques – Analytic Element Method (AEM), Finite Difference Method (FDM), and Finite Element Method (FEM) – which lend themselves well to either vector or raster data, and legacy input/output file formats that are not well suited across models. Nonetheless, investigative computational tools are all founded in

10 the same conceptualization of hydrologic properties associated with mass, flux, pathways and residence time. A consistent framework is developed using modern Geographic Information Science (GIScience) methods to organize and archive important information from international datasets and previous groundwater ecohydrology studies organized around aquifer and water point, line, polygon and raster features. Case

15 studies illustrate the efficacy of this platform to address existing data interoperability issues for representative groundwater ecohydrology problems of global significance including the impact of human-induced forcings, change in species, and forcings by natural processes on groundwater ecohydrology. In North America, we study the relationships between groundwater pumping in the Ogallala Aquifer and changes in riparian habitat and phreatophyte species composition. In Europe, we study the impacts of

20 changes in forest species composition on groundwater recharge and baseflow to biologically diverse fens and wetlands in the Veluwe sand hill region of The Netherlands. In Africa, we study the wetlands of the Okavango Delta in Botswana that forms an oasis in the midst of the Kalahari Desert and the role of groundwater in flushing salts from this freshwater ecosystem. In each study, we document the current state of knowledge, identify pertinent datasets and previous studies, develop new conceptual and computer models, and summarize findings. This computational platform provides a framework to study sustainability, to forecast the impacts of changes in forcings, and to provide a

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

1 Introduction

Ecohydrology is an interdisciplinary field of study crossing the interface between hydrologic and ecologic sciences (Hannah et al., 2004). Hydrologic surficial processes (precipitation, evapotranspiration, runoff and recharge) impact the integrity of freshwater ecosystems that depend upon adequate quantity, quality, timing, and temporal variability of water flow (Baron et al., 2002). Likewise, ecosystem processes impact groundwater hydrology through terrestrial recharge of rain that seeps past the root zone (Jobbág and Jackson, 2007) and through phreatophyte root uptake of groundwater (Dawson, 1993). These interrelated processes are influenced by the physiological characteristics of vegetation, the pedology of the soil, and the climate (Rodríguez-Iturbe, 2000).

The relationship between groundwater recharge and discharge is one of the most important aspects in the protection of ecologically valuable areas (Batelaan et al., 2003).

The rate of recharge to groundwater is strongly impacted by the land cover (Batelaan and De Smedt, 2007), and changes in recharge impact groundwater storage, depth to water, and the rate of base flow discharge to surface water (Brutsaert, 2008). This discharge to aquatic ecosystems moderates river/stream flow and alters surface water chemistry and temperature (Hayashi and Rosenberry, 2002). The severity of vegetation response to altered groundwater elevation is influenced by a plant's drought and flooding tolerance and its root system size and water uptake capacity (Naumburg et al., 2005).

Meeting human water needs while sustaining the services that aquatic ecosystems provide is a central research challenge for the 21st century (Palmer and Bernhardt, 2006). Freshwater ecosystems, which are functionally intact and biologically complex, provide economically valuable ecosystem goods and services (potable water, food supply, industrial use, habitat, instream uses as recreation, transportation and waste dis-

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



posal, and adaptivity to climate change) with intergenerational benefits to society (Jackson et al., 2001; Baron et al., 2002; Richter et al., 2003). Riparian and aquatic ecosystems have been altered or lost in many parts of the world through human-induced groundwater depletion and stream dewatering with particularly strong effects in arid and semiarid regions (Pringle, 2001).

Establishing methodology for scientific ecohydrologic investigation and its application to case studies is an emerging topic of international interest. The Consortium of Universities for the Advancement of Hydrologic Science, Inc. (CUAHSI, 2007) recently released a science plan that identifies “linking the hydrosphere and the biosphere” as the number one challenge in developing a more comprehensive, systematic understanding of continental water dynamics. The EU Water Framework Directive requires that every country introduce measures to improve and sustainably maintain good water quality and ecological status based on the best available scientific information and using an unbiased, independent and logical methodology (Bruen, 2008). It was found by the Internationally Shared Aquifer Resource Management (ISARM) Programme of UNESCO to be considerably more complicated to develop mutually beneficial and sustainable programs for transboundary aquifer development than for surface water, and that multidisciplinary approaches are required (Puri and Aureli, 2005).

Geographic Information Science (GIScience)-based tools and methodology provide a framework for quantitative analysis of the coupling between human and natural systems (Adriaens et al., 2003; Nellis, 2005). Fundamental process understanding of ecohydrology considers interactions operating at a range of spatial and temporal scales (Hannah et al., 2004). This need may be addressed through the rapid improvements in the availability of high-resolution geospatial data, which improves understanding of geomorphic drivers and the contexts of regional vulnerability to land-use change (Poff et al., 2006).

This paper puts forth a GIScience methodology to conceptualize groundwater and its interactions with ecology. The groundwater concepts data model addresses current lack of data interoperability across numerical modeling approaches, data input file for-

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶


[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



mats and shapefile field mapping techniques. Case studies are presented for ecologically important regions in Africa, Europe and North America that identify and utilize GIS data repositories from a variety of sources. These studies develop new understanding to address scientific challenges posed by Newman et al. (2006) for ecohydrology:

- 5 1. landscape interactions in the paleodominated and human-dominated ages,
2. vegetation and groundwater recharge, and
3. hydrological change and vegetation.

We examine the impact of human-induced forcings, change in species, and forcings by natural processes on groundwater ecohydrology.

10 2 Methods

A conceptual view of groundwater ecohydrology processes and important variables are shown in Fig. 1. Recharge is that fraction of precipitation/irrigation that seeps downward past the root zone of plants to the saturated groundwater zone, where voids between soil particles are fully filled with water. Recharged groundwater flows through 15 an aquifer from regions with high groundwater elevation to low groundwater elevation. It re-emerges to the land surface through human induced pumping, phreatophyte groundwater uptake by plant roots or baseflow to surface water. The depth to water (land surface minus groundwater elevation) and the height of the capillary fringe above groundwater in the unsaturated zone controls the proximity of groundwater to phreatophyte root zones.

20 A mathematical description of groundwater flow is provided by the following three equations. Darcy's Law is a constitutive relationship between the specific discharge vector q , which is a measure of the rate of flow, and the head, ϕ , which is a measure

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

of groundwater elevation (the level to which water would rise in a pipe located at the position (x,y,z) ,

$$\mathbf{q} = -k \nabla \phi \quad (1)$$

where the hydraulic conductivity, k , is a property of the aquifer medium and the fluid.

5 The Dupuit assumption that head is uniform in the vertical z -direction gives a relationship where the discharge per width, $\mathbf{Q} = (Q_x, Q_y)$, is equal to the saturated groundwater thickness times the specific discharge, which may be written as follows for unconfined groundwater:

$$Q_x = (\phi - B)q_x, \quad Q_y = (\phi - B)q_y \quad (2)$$

10 where B is the elevation of the base of the aquifer medium. Continuity of flow (conservation of mass) relates the divergence of \mathbf{Q} to the change of groundwater elevation over time

$$\frac{\partial Q_x}{\partial x} + \frac{\partial Q_y}{\partial y} = -S_y \frac{\partial \phi}{\partial t} + R \quad (3)$$

15 where S_y is the specific yield of the porous medium, R is the recharge rate, and water is assumed incompressible.

Predictive models of groundwater flow utilize these equations, with boundary and initial conditions related to known values of groundwater elevation and flow rates, to forecast existing conditions and future changes to groundwater stores. Three numerical techniques are commonly employed to implement these groundwater equations 20 and boundary/initial conditions in computer models. The Analytic Element Method (AEM) discretizes hydrogeologic features into points, lines and polygons and develops mathematical functions that exactly satisfy the governing equations and reproduces the flow generated by the feature (Strack, 1989). For example, a well has the geometry of a point in two-dimensions and the head is obtained by evaluating the Thiem 25 (1906) or Theis (1935) solution; similar solutions exist for line elements (Steward et al.,

Groundwater
ecohydrology studies
in Africa, Europe and
North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

2008) and polygon elements (Strack, 1989). The Finite Difference Method (FDM) discretizes the flow domain into a set of cells (normally rectangular) with grid points used to delineate cells, and utilizes finite differences to develop a set of equations that gives the groundwater elevation at each grid point (Bear and Verruijt, 1987). The head at locations between these points is obtained through interpolation. The Finite Element Method (FEM) discretizes the flow domain into a set of polygons with prescribed geometry and spatial variation of head across each element and develops a set of equations to obtain the head at collocation points on each element (Bear and Verruijt, 1987). The head at points between these collocation points are obtained by evaluating the prescribed shape functions.

Case studies will illustrate results obtained using these techniques with the computer programs:

- AEM: SPLIT (Janković and Barnes, 1999), available at <http://www.groundwater.buffalo.edu>
- FDM: MODFLOW (McDonald and Harbaugh, 1988), available at <http://water.usgs.gov/software>
- FEM: FEFLOW (Diersch, 2005), available at <http://www.wasy.de>

There is a current lack of data interoperability between such groundwater models. The individual numerical techniques lend themselves well to either vector (AEM/FEM) or raster (FDM) data types, which are both important for hydrological investigations (Whiteaker et al., 2007). These differences along with model assumptions lead to organization of input/output file formats that are not well suited for use across all three numerical methods. Individually, these computer models have begun to address this challenge by using vector shapefiles with fields associated with variables pertinent to the particular software needs. Yet still, a need exists in establishing a spatial groundwater data model that supports all models and allows greater flexibility in specifying groundwater properties; a need that is addressed here.

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

While computer models have difficulty using each other's data files, they are still founded in the same conceptualization of hydrologic properties associated with mass, flux, pathways and residence time (Reckhow et al., 2004). Furthermore, each model uses data related to groundwater ecohydrology that are common across approaches (borehole lithology and hydrogeology, DEM (Digital Elevation Model), surface water quantity and quality, soils, climate, etc.) An existing challenge is to organize data from a variety of sources and resolutions (Zheng et al., 2006) in a format that is conducive to groundwater investigations across a broad range of computational tools.

To address this challenge, a GIScience based framework has been developed to organize conceptual models of groundwater, which contain the hydrogeologic features important for a particular study (Yang et al., 2009). The data model in Fig. 2 is independent from the choice of computational tool. The geological layers are prescribed in AquiferLayer as aquifers and aquitards following Strassberg et al. (2007). Hydrogeology is then subdivided into features associated with the Aquifer (that through which water flows) and the Water (that which drives the flow of water), where features may be raster grids or have the geometry of points, lines, or polygons. This partitions groundwater hydrogeology into objects related to either the AquiferProperty associated with the geologic medium or the WaterProperty associated with the fluid. Each feature is related to the AquiferLayer (aquifer or aquitard) in which it resides. The Units of properties are specified in these tables, and the time at which properties are specified may be set to support transient studies. For example, a WaterLine may be used to represent a river that lies in the AquiferLayer at the land surface and the WaterProperty may be used to specify the head/elevation along the river with Units of meters above mean sea level (m.s.l.). The relationship classes enable a default property value to be associated with each feature and/or additional properties to be specified at points, lines, polygons or rasters.

This data model provides a relational database of data pertinent to groundwater ecohydrology, and a consistent mechanism to gather data and to store a conceptual view of groundwater ecohydrology that is common across and independent from ground-

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

water computer models. This builds upon existing endeavors to establish a consistent data model for vector based data (Steward et al., 2005; Bernard et al., 2005; Steward and Bernard, 2006, 2007; Yang et al., 2009) and extends these studies to incorporate raster data. It is complimentary with existing endeavors to store measurements for 5 observatory level hydrologic data based upon the ArcHydro platform (Maidment, 2002; Strassberg et al., 2007; Horsburgh et al., 2008) and provides a conceptual model of these data. This also provides a platform that may be integrated with the GetValue protocol of OpenMI (Moore and Tindall, 2005; Gregersen et al., 2007) to provide a means 10 of exchanging data across a variety of investigative tools associated with human and natural components of water resources (Steward et al., 2009a).

It should be noted that this data model is not intended to compete with existing Graphical User Interfaces (GUIs) for existing tools, many of which are quite excellent. The data model provides a central storage mechanism to structure and organize datasets for a groundwater conceptual model, making them syntactically similar and 15 ready to integrate for various purposes. With built-in database management capabilities, well-defined domain, relationships, and topological rules, spatial data models improve data storage efficiency and enforce data integrity. For example, data duplication is avoided by storing well locations in a spatial object and defining a relationship to join the wells with their pumping rates and schedule. Rules may be defined to prevent illegal values such as negative well pumping rates and to limit units of hydrological 20 parameters to those with valid dimensions (e.g., m^3/day for pumping rate). Data integrity ensures that, for example, if a well is deleted its associated pumping rates will be deleted also. Topological rules define how points, lines and polygons share geometry, for example, WaterBoundary points that specify values of head along a river must 25 fall on the WaterLine. Once published, the structure of a spatial data model is open and allows any third-party to develop customized tools and interfaces for its specific needs. It also provides a mechanism to archive data from a study for future use.

In summary, the data model in Fig. 2 puts forth a consistent representation for conceptual groundwater models that addresses current lack of data interoperability across

**Groundwater
ecohydrology studies
in Africa, Europe and
North America**

D. R. Steward et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

models, provides flexibility in management of data and naming conventions, provides enhanced data integrity and a consistent framework to archive data for future use. We next demonstrate the utility of these methods to capture a conceptual groundwater model for studies representative of ecologically and hydrologically important regions of the world. A general procedure for its implementation is followed:

5

1. Gather pertinent data and make assessable in GIS (scan and georeference maps, load into ArcHydro groundwater model, as appropriate).
2. Add aquifer/aquitards as per conceptualization of geologic media to the Aquifer-Layer table.
- 10 3. Identify important features in conceptualization (geology, surface water, recharge/discharge zones, wells, etc.), and construct those associated with geologic media (AquiferPoint, AquiferLine, AquiferPolygon, AquiferRaster) and the fluid (WaterPoint, WaterLine, WaterPolygon, WaterRaster), and associate with the AquiferLayer in which they reside.
- 15 4. Associate appropriate AquiferProperty or WaterProperty with features from information in step 1, and/or model calibration. This may be accomplished by setting a single value of properties important for a feature (e.g., an AquiferLayer may have a single value of Hydraulic conductivity), setting the values at a set of points (e.g., a WaterPoint feature for a well may specify the Discharge for pumping rates) or lines (e.g., An AquiferLine may be used to specify a contour of uniform Base elevation), or setting the values in a raster (e.g., an AquiferRaster may contain the Thickness of a layer obtained from interpolation between borehole data).
- 20

25

The data model for each case study is used within computational models (AEM, FDM, and/or FEM), the conceptualization of important groundwater ecohydrology drivers is refined based upon interpretation of results, and findings are presented. We invite readers interested in more technical details to explore the data model at <http://www.growe.ksu.edu>.

Groundwater
ecohydrology studies
in Africa, Europe and
North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

3 Case studies

The study regions and their soils are identified in Fig. 3. The Ogallala Aquifer lies in the High Plains region of central North America with wind-blown loess soils interspersed with fluvial sands and clays overlying thick sedimentary deposits (Gutentag et al., 1984). This semi-arid grassland ecosystem has been transformed by human-induced groundwater pumping over the past 50 years (Opie, 2000). The Veluwe in the delta region of western Europe contains ice-pushed ridges with soils consisting of glaciofluvial sand, gravel and clay sediments that overlie marine and fluvial sands and clays (Vasak et al., 1981). This region has been transformed from an agriculturally managed grassland to coniferous forest and is planned to be converted to deciduous forest (Gehrels, 1999). The Okavango Delta in the southwestern extension of the East African Rift System of sub-Saharan Africa forms a sedimentary alluvial fan with wind-blown aeolian deposits of Kalahari sands (Wilson and Dincer, 1976). This wetland forms one of the most biologically diverse freshwater ecosystems in the world (Warne, 2004).

In the following sections, we study the groundwater ecohydrology of these regions using the GIScience methods and computational approaches previously described. First, the groundwater ecohydrologic processes and current state of knowledge are reviewed and the ecosystem forcings from human activities, changes in species, and natural processes are identified. A conceptual model is then constructed using GIS data organization supplemented with domain specific understanding from previous groundwater ecohydrology studies to guide this development. A variety of data sources are identified and documented for each case study region; for example, different sources were used to construct the pictures of soils for the three areas in Fig. 3. Numerical models are applied to understand each ecosystem and findings are summarized.

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

3.1 North America: interrelationships between regional groundwater pumping and phreatophyte distribution over the Ogallala Aquifer

HESSD

6, 2795–2844, 2009

The Ogallala Aquifer is located in the High Plains region of the central United States of America. This region has been transformed with the advent of modern well hydraulics over the past 50 years from the “Great American Desert” in the “Dust Bowl” region into the “Breadbasket of the World”. Groundwater is a common pool resource that has been over appropriated in many regions with withdrawals exceeding the natural rate of recharge. This study examines the relationship between a declining groundwater elevation and the distribution of phreatophytes in riparian corridors.

10 3.1.1 Groundwater ecohydrologic processes

The Ogallala Aquifer spans 450 000 km² and underlies 27% of the irrigated land in the United States (Dennehy, 2000). This semi-arid grassland ecosystem was classified by Powell (1879) as an arid land lying west of the 100th meridian with a mean annual precipitation less than 500 mm (20 in) that would require irrigation for successful agriculture. This irrigation need is being met through groundwater extraction that provides society with a water supply that is reliable through drought but unsustainable through natural recharge (Custodio, 2002), a pertinent problem occurring in semi-arid regions of the world: e.g. in Africa, Asia, the Middle East and South America (Montaigne, 2002; UNESCO, 2003; Foster and Chilton, 2003). Within the study region, the average water-use over the past 25 years is 175 mm/yr while natural recharge averages 24 mm/yr (Hansen, 1991; Steward et al., 2009b). Attempts to stabilize this imbalance between water input and output include reducing water use, increasing irrigation efficiency, and adopting water-saving land-use practices (Sophocleous, 2005; Buletewicz et al., 2009).

25 Groundwater and surface water interact and groundwater depletion leads to reduced baseflow, induced groundwater recharge, and subsequent streamflow depletion (Sophocleous, 2002). Stream depletion is dependent upon the level of pumping induced drawdowns and the properties of the streambed (Butler Jr. et al., 2001) as well

Groundwater
ecohydrology studies
in Africa, Europe and
North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



as the properties and extent of underlying aquitards that may slow the rate of induced recharge and spread it over a large surface area (Butler Jr. et al., 2007b). Sustaining streamflow in freshwater ecosystems may result in a sustainable yield of an aquifer that is considerably less than recharge if adequate amounts of water are to be available to 5 sustain both the quantity and quality of streams, springs and wetlands (Sophocleous, 2000).

Evapotranspiration by phreatophytes provides a pathway for most of the 20–50% of total long-term water budget depletions in large river systems that are ascribed to 10 natural vegetation (Cleverly et al., 2006). Within the study region, cottonwoods and salt cedars have both been shown to withdraw groundwater from the capillary fringe that forms in the vadose zone above saturated groundwater (Butler Jr. et al., 2007a). The cumulative effect of diurnal pumping by these phreatophytes results in the capture 15 of groundwater from the upper regions of the aquifer (Steward and Ahring, 2009). This phreatophyte-induced redistribution of groundwater to drier surface layers significantly increases photosynthesis and evapotranspiration, and establishes a direct link 20 between plant root functioning and global climate (Lee et al., 2005). The two key environmental variables that influence survivorship of wetland and riparian vegetation are depth to groundwater and inundation frequency (Stromberg et al., 1996). Groundwater management practices coupled with drought have led to die-off of phreatophytes and subsequent exotic infestation when drought ended in California (Elmore et al., 2003) 25 and the Ogallala Aquifer region (Butler Jr. et al., 2007a).

3.1.2 Conceptual model and results

The conceptual model of the hydrogeologic features important for this study in Fig. 4 illustrates a drainage network of rivers, pumped wells used primarily for irrigation, and an 25 aquitard situated beneath alluvial deposits of the Arkansas River and above the underlying Ogallala Aquifer. The background 30 m raster National Elevation Dataset shows the gently sloping land surface from west to east incised by river valleys with ephemeral (west) to perennial (east) streams. In predevelopment conditions before wells were

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

introduced, the groundwater elevation was close to the land surface in the riparian Arkansas River corridor and phreatophytes (cottonwoods) became established. Human forcings through groundwater extraction has contributed to groundwater declines. Currently, this river rarely flows in this region and native cottonwood populations are becoming replaced by exotic salt cedars.

In this study, we demonstrate the capacity of the GIScience methods to address large-scale transient models and datasets. The conceptual model of the Ogallala Aquifer is captured in the GIScience data model presented in Fig. 2 using the data sources and previous studies listed in Table 1. The AquiferLayer table contains the three layers associated with the alluvium of the Arkansas River, the underlying aquitard, and the Ogallala Aquifer. Studies by the United States Geological Survey (USGS) (Gutentag et al., 1984) were augmented by lithology records from boreholes collected by the Kansas Geological Survey (KGS, 2008) to develop AquiferLine features containing contours of bedrock elevation (Macfarlane and Wilson, 2006). These studies also led to AquiferPolygon features used to depict the hydraulic conductivity and specific yield for the three AquiferLayer objects: alluvium in the Arkansas River corridor, underlying aquitard, and the Ogallala Aquifer. The Division of Water Resources in the Kansas Department of Agriculture developed recharge maps (Hansen, 1991) used to develop WaterPolygon features. The DWR also collects annual water-use reports for each point of diversion as mandated by Kansas law (K.S.A. 82a-732) that were used to develop WaterPoint features with WaterProperty of pumping rates. The USGS National Hydrography Dataset was used to construct WaterLine features for rivers and their WaterProperty was assigned water elevation obtained from DEMs and values for the resistance of river beds were obtained from previous studies and model calibration.

The vector features in this data model were used in the FEM based computer model FEFLOW to study the impacts of groundwater pumping. Note that this conceptual model also supports other computer models, e.g., MODFLOW (Yang and Steward, 2009). Thus, a major advantage of utilizing this conceptual data model is that it provides a mechanism to integrate data from existing sources while archiving data in a for-

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

mat that promotes interoperability for future use (e.g., researchers may use the data in their own different model for their specific analysis) and benefit to water-dependent communities.

Results are presented in Fig. 5 that show contours of groundwater elevation and the depth to groundwater obtained by subtracting the DEM raster from the head obtained in the groundwater model. The initial water table elevation was established for predevelopment conditions in 1959 before wells became broadly established, and a transient model was run using these initial conditions to provide groundwater elevations at the end of 2006, the last year of available data. Model results were calibrated against a network of observation wells maintained by the Kansas Geological Survey (Hausberger et al., 1998) and gave a root mean square error of 5.4 m for predevelopment conditions (over 24 observation wells) and 6.1 m for the last year of simulation (over 176 observation wells). Boundary conditions of specified head were established from a surface obtained by kriging yearly values of groundwater elevation across the network of observation wells; this kriging was performed yearly from 1959 to 2006 to provide the transient boundary condition and the WRIS dataset provides annual water use at each well. While most of the wells in the observatory network are used for irrigation with yearly fluctuations to 10 s of meters, measurements were only used for observations taken between 1 December and 31 January when the water level in wells had mostly recovered. The Kansas Department of Agriculture conducted a helicopter survey of salt cedar populations along the Upper Arkansas River in 2005 and classified riparian habitat based upon percentage salt cedar, which is also shown in Fig. 5.

These new model results illustrate that irrigation water used to support crops in the terrestrial ecosystem have led to declines in the groundwater elevation throughout the region. These results also illustrate that the aquitard in the Arkansas River corridor has partially mitigated the groundwater declines and is effectively holding groundwater in the alluvium near land surface. Within this riparian corridor, cottonwood die-offs have been observed as groundwater levels dropped and the helicopter survey illustrates subsequent establishment of relatively deep-rooted salt cedars that continue to

Groundwater
ecohydrology studies
in Africa, Europe and
North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

tap groundwater. While irrigated agriculture has sustained economic development in the region for the past half century, change for many western Kansas communities is inevitable (Leatherman et al., 2004). This groundwater ecohydrology model enables forecasts and prediction of the impact of proposed change in water-use practices, feedback to the groundwater due to ecological thresholds like die-off of the riparian vegetation, and provides a scientific basis to inform public policy planning and debate. The data model also facilitates integration of groundwater results with models of human drivers (Steward et al., 2009a).

10 3.2 Europe: interrelationships between plant species composition and ground-water recharge/discharge zones in the Veluwe

The Veluwe region of The Netherlands is located in the delta region of the international Rhine River system. With formation of the Hoge Veluwe National Park, this region has undergone transition over the first half of the last century from heather vegetation, managed agricultural grassland, bare land and moving sand dunes to a coniferous (spruce, fir) forest that both stabilized sand and provided support wood for the coal mining industry. This change of species has led to decreases in the rate of groundwater recharge with subsequent loss of groundwater fed and biologically diverse fens and wetlands. The EU Water Framework Directive obliges European nations to research and learn how to provide sustainable ecological approaches. This study examines 15 The Netherlands plans to reestablish fens and wetlands by replacing coniferous with deciduous (oak, beech, birch) forests.

20 3.2.1 Groundwater ecohydrologic processes

The Veluwe is a nutrient poor, sandy area covered by forests, heath lands, drift sand areas and some agricultural fields and constitutes one of the most important nature areas of the Netherlands in terms of both ecological quality and recreation (Turnhout et al., 2004). This temperate maritime climate has an average annual rainfall of 850 mm

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



and evapotranspiration around 550 mm (Gehrels et al., 1994). The hills in this region are the remains of ice-pushed ridges with glacial basins located on their concave eastern and northern sides (van der Meulen et al., 2005). These hills form a groundwater recharge zone with a sandy unsaturated zone of about 40 m in the center that provides 5 baseflow to surface water surrounding the Veluwe where the unsaturated zone is near zero (Vasak et al., 1981; Gehrels et al., 1994). The ice-scoured basins along the edge of the Veluwe form a stacked system of troughs filled with banded material consisting of an ill-sorted mixture of gravels, sands, silts and pebbles with dispersed outsize clasts (Postma et al., 1983). The spring waters that flow into these basins along the 10 eastern and southern Veluwe areas contain high concentrations of NO_3 attributed to enrichment during passage of groundwater through the nitrogen-rich soils (Higler and Verdonschot, 1993).

Fens and fen meadows are rich in endangered species and occur in complex landscape settings where groundwater discharges to low-lying areas, making them sensitive 15 to hydrological changes in the valley and surrounding area (Grootjans et al., 1996). Within the Veluwe area, naturally wet areas have become dessicated (Vermulst and de Lange, 1999) due to landscape changes involving centuries of human use from cutting forests for agriculture to degradation of forest into heath land to grazing and sod cutting leading many heath lands to degrade further into drift sand areas (Turnhout 20 et al., 2004). Dessicated fen mires have become severely eutrophied due to degradation (oxidation) of peat, and watering from other sources usually causes deterioration of the original flora and fauna due to changes in water quality (Best et al., 1993). It is important to study the hydrologic controls in vegetation composition through water and solute cycling and the impacts of water resources management on these regions 25 of species-rich gradients in series of soil and vegetation types that mirror the relief of the landform (Grootjans et al., 1996).

Presently, reforestation of coniferous forests has stabilized most of the drift sand through a succession of vegetation (lichens, mosses, trees) that is increasing biodiversity (Turnhout et al., 2004). The coniferous forests planted as even-aged stands in the

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

first half of the last century are undergoing conversion into mixed stands with indigenous deciduous tree species in the Veluwe (Kuiters and Slim, 2002). The recharge below a deciduous forest is greater than between a coniferous forest, with rates of interception averaging 10% (winter) to 26% (summer) for deciduous forests and 22–39% for coniferous forests in the Veluwe (Gehrels, 1999). Ecosystem prediction of the implications of such change in landcover and subsequent increase in rates of groundwater recharge and fen/stream baseflow may be facilitated by coupling hydrological and ecological process knowledge using GIS and ecohydrologic models using field data (Grootjans et al., 1996; Boumans et al., 2008). Within the Veluwe region, the dominant causes of changes in groundwater fluctuation are changes in groundwater recharge, large-scale dewatering in polders formed to the north in the former IJsselmeer and groundwater withdrawals by wells (Gehrels et al., 1994).

3.2.2 Conceptual model and results

The conceptual groundwater ecohydrologic model of the Veluwe is illustrated in Fig. 6 by depicting stream networks, inland waterways and wells. The background 25 m Actual Height of the Netherlands dataset shows the ice-pushed hills of the Veluwe and the surrounding valleys along the eastern edge where biologically diverse fens and wetlands are located. The recharge occurring in the Veluwe hills drives groundwater flows through aquifer layers with properties identified from boreholes and knowledge from previous investigations. The baseflow discharge to the aquatic ecosystems is impacted by the rate of recharge in the hilly regions, which is controlled by forest species composition and their relative efficiencies in intercepting/evaporating and transpiring precipitation.

In this study, we demonstrate the capacity of the GIScience methods to provide a data model for the NHI (Netherlands Hydrologic Instrument), an integrated system of models for national water management in The Netherlands. The NHI captures the ecological inputs of vegetation in unsaturated vadose zone models, providing input to this conceptual model and promising better integration with ecohydrologic drivers in

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



the future. The datasets and previous studies are identified in Table 2. The Aquifer-Layer table contains the four aquifers and three aquitards that exist in the Veluwe region (de Lange, 2006). The TNO-Geological Survey of The Netherlands maintains the REGIS/DINO datasets containing borehole lithography. These AquiferPoint features, together with AquiferPolygon features that depict geological units, have been used to develop AquiferRaster datasets related to hydraulic conductivity and aquifer elevation. TNO datasets also contain water-use information for wells, which is stored in the WaterPoint features with WaterProperty values associated with the pumping rate. WaterLine and WaterPolygon features associated with surface water have been used to develop WaterRaster datasets containing information about surface water properties and recharge. The discharge boundaries providing upward leakage through the less-pervious cover layer in the deep Flevo-polders to the north were specified using cells in the WaterRaster grid along the boundary.

The 250 m × 250 m raster datasets in this data model were used in the Finite Difference Method computer model MODFLOW to study the impact of change in tree species in the Veluwe. Note that the groundwater data model also has the capacity to store the complimentary Dutch national groundwater model that uses the AEM (de Lange, 2006), effectively bridging datasets and understanding between these models. New results from this model are presented in Fig. 7 that show the contours of groundwater elevation and the depth to water. This model was exercised by imposing recharge rates throughout the terrestrial ecosystem corresponding to coniferous and to deciduous forests. The results show a groundwater mound beneath the center of the Veluwe that raises from an elevation of 32 m beneath a coniferous forest to 40 m beneath a deciduous forest; this change began in the century before the foresting was completed (1850–1950) and is observed through changes in forest species. Within the lowlands to the east, there is an increase in the upstream portions of the river network that contains shallow depth to water, where fens and wetlands may become reestablished. The depth to water and length of upstream portions of stream bottoms that intersect the water table varies, not just under change in species as examined here,

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

but also due to non-linear feedbacks associated with climate change and water use.

The NAtional GROundwater Model (NAGROM) was developed to improve understanding of groundwater movement in multiple aquifer systems (de Lange, 2006), and forms a foundation of hydrogeologic understanding for the Netherlands Hydrologic Instrument. This NHI gives national impacts for policy making and to assess possible measures for sustainable ecology. This modeling approach, coupled with the GIScience data model presented here, provides an avenue to research sustainable ecological approaches as required by all European countries in the Water Framework Directive, and to understand feedbacks between the terrestrial ecology of forests, the groundwater hydrology of recharge and baseflow, and the aquatic ecology of fens and wetlands. Thus, the groundwater datamodel promotes future analysis with different modeling concepts to make use and hence show/stress different aspects of the future water system and its interaction/feedback with ecosystems. It also provides a mechanism to promote analysis and comparison of model results across EU countries with different modeling techniques, and more balanced overall conclusions.

3.3 Africa: interrelationships between plant groundwater uptake and salt accumulation and groundwater flow in the Okavango Delta

The Okavango Delta is located in the extreme southwestern extension of the East African Rift Valley. Its transboundary surface water inflows from precipitation occurring during the rainy season in the mountains of Angola and Namibia to arrive months later in northern Botswana during the dry season, providing a biological diverse oasis amidst the sands of the Kalahari Desert. The high evapotranspiration rates of this wetland results in salt accumulation within the delta. This study examines how the regional groundwater flow flushes salts to help sustain this freshwater ecosystem.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

3.3.1 Groundwater ecohydrologic processes

The Okavango Delta is an alluvial fan formed by the Okavango River discharging into a graben at the southern extremity of the East African Rift System (McCarthy and Ellery, 1994). This river enters through the panhandle, a tectonic graben 100 km long and 10 km wide, and then spreads atop 300 m of sediment (McCarthy et al., 2005; McCarthy, 2006) to form a wetland with surface area that varies seasonally between 6000–12 000 km² (Dincer et al., 1987). The waterborne sediments within the rift valley are bounded by a fault to the south and a parallel fault 15 km from the end that partially redistributes groundwater (Wilson and Dincer, 1976).

The wetlands of the Okavango Delta lie within the Kalahari Desert. While this region receives a mean annual precipitation of 500 mm/yr (Dincer et al., 1987), recharge is small due to grass/tree competition and sands that are of small enough grain size to hold precipitation (de Vries and von Hoyer, 1988). Tree roots have been measured to depths of 68 m (de Vries and von Hoyer, 1988) and it is likely that Acacia trees tap to tens of meters and perched water tables, limiting present recharge of the Kalahari to 1–5 mm/yr (de Vries et al., 2000). Within the wetlands of the Okavango Delta, evapotranspiration exceeds precipitation by a factor of 3–4 (McCarthy and Ellery, 1994; McCarthy, 2006). Inflow in the Okavango River is 10×10^9 m³/yr with a net salinity of 30 ppm (major dissolved species consisting of sodium, bicarbonates and silica) and precipitation adds 5×10^9 m³/yr (Dincer et al., 1987; Gieske, 1997). Evapotransport results in a loss of nearly 98% of this influx (Linn et al., 2003), resulting in annual salt accumulation in the delta of 300 000 tons/yr (Dincer et al., 1987; Bauer-Gottwein et al., 2007).

The surface flood inundates the Okavango Delta traveling through an alluvial fan of rivers that traverse the delta and the perennial swamps (Gieske, 1997). Large scale channel avulsions occur in this braided stream network where new channel systems form over tens to hundreds of years as main river channels become blocked by deposition of silts (Ellery et al., 1993b), sediments consisting mostly of fine sand (400 000 m³

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

annually) brought to the delta by the Okavango River (Dincer et al., 1987), and blockage formed by papyrus (Gieske, 1997). Within the wetlands, 150 000 vegetated islands have formed on termite mounds (Warne, 2004) and subsurface accumulation of silica and calcite (McCarthy et al., 1993). The islands constitute between 10–20% of the surface area of the perennial swamp with sizes ranging from a few square meters to several hectares (Ellery et al., 1993a; McCarthy et al., 1993). 75% of the total water loss from the wetland is caused by transpiration-driven lateral groundwater flow into the islands, with the remaining 25% lost to evaporation from the open water surface (Wolski and Savenije, 2006).

5 A zonation in plant species occurs across islands from evergreen trees to deciduous trees and palms to grasses to bare soil in salt pans at island centers (McCarthy and Ellery, 1994). This zonation is not related to depth of groundwater, with seasonal variation of 1.5 m in the panhandle, 20 cm in the central regions of perennial swamps, and 2 m in the lower reaches of the fan (Ellery et al., 1993a). Instead, zonation changes 10 across an island as a carbonate encrusted area free of vegetation forms in the island centers (Ellery et al., 1993a). Salt precipitates preferentially with the most soluble salts found in centers (McCarthy, 2006), where lignin stained (brown) salt water occur with enriched alkali carbonates indicate terminal evaporation of water in the capillary zone around the pans (McCarthy et al., 1991). It is supposed that the vegetation enhanced 15 salt accumulation process takes on the order of 100–200 years to form an area void of vegetation; this process may be interrupted by swamp abandonment leading to accumulation of silica and calcite in the subsurface and leaching of sodium salts to groundwater (McCarthy et al., 1993). Numerical simulation suggests that salts accumulate 20 to densities that, over hundreds to thousands of years, generate density driven fingers that transport high salinity groundwater downward towards the base of the aquifer (Zimmermann et al., 2006). This is supported by electrical soundings carried out on the fringes of the Delta that indicate a marked increase in salinity of the groundwater with depth, with a layer of lower salinity groundwater (50–70 m thick increasing towards 25 permanent swamp) floating on denser saline water (McCarthy, 2006).

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



3.3.2 Conceptual model and results

The conceptual groundwater ecohydrologic model of the Okavango Delta in Fig. 8 illustrates the regional groundwater system. At the surface, water inflows from the Okavango River in the northwest to an outlet leading towards salt pans in the Kalahari Desert to the southeast. The background 90 m DEM shows ridges surrounding the graben located in the southwestern extension of the East African Rift Valley, as well as the gently sloping land surface within the wetland ecosystem. The region of salt accumulation by phreatophyte driven evapotranspiration of groundwater in the perennial swamp and vegetated islands are averaged over polygons, where tens of thousands of small islands lie within an area of shallow water table. The regional groundwater gradient that flushes salts from beneath the Okavango Delta are driven by surficial flow and recharge depicted by polygons. The connected faults that transport water and the dikes at the southwestern boundary of the Okavango Delta that impede the movement of water are depicted by lines. Groundwater observations have high salt concentrations in areas south of the Okavango Delta, and surficial salts are found in salt pans to the south.

In this study, we illustrate the capacity of the GIScience methods to develop conceptual groundwater ecohydrology models for problems broad in area with relatively sparse data. In such cases, alternative conceptual models may be constructed and the GIS system may be used to help and thus reduce uncertainty and increase robustness. The conceptual model is captured in the GIScience data model using the datasets and previous investigations identified in Table 3. A geological map of Botswana was developed by the Department of Geological Survey and Mines (Hepworth et al., 1973), and a land system map was developed by the Ministry of Agriculture (De Wit et al., 1990). These, together with remote sensed imagery, were used to develop a single AquiferLayer model with AquiferLine features for connected fault systems and Aquifer-Polygon features for the graben and ridges. The Okavango Delta management plan of the Department of Environmental Affairs (DEA, 2008) was used to develop Aquifer-

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Line features for the dikes. WaterLine features for rivers were developed from maps and previous studies and the WaterProperty values for head were obtained from the DEM. The WaterPolygon features for the perennial wetland, areas of high evapotranspiration, and areas of recharge were developed from previous studies and remotely sensed data, and their respective rates of recharge/discharge were specified in the WaterProperty table.

The vector data in this data model was used in the Analytic Element Method based computer model SPLIT to study the steady regional groundwater flow system. The steady assumption is adopted because, while seasonal variation in groundwater elevation fluctuate about a mean elevation immediately beneath the wetlands, the present regional hydraulic gradient is more or less in steady-state (de Vries et al., 2000). The groundwater elevation and depth to water in the Okavango Delta region are shown in Fig. 9. These results compare well with observed values of groundwater elevation and depth to water from both regional (de Vries et al., 2000; Staudt, 2003) and local studies within the Okavango Delta (McCarthy et al., 1998). This model provides new understanding of a low hydraulic gradient in groundwater within the delta extending to the southeast. This groundwater flow provides a mechanism to flush the salts that accumulate within and beneath the wetlands to locations southeast of the Okavango Delta beneath the salt pans in the Kalahari Desert, and thus enable the freshwater zone to continue to exist.

This GIScience data model and groundwater model also enables prediction of the outcome of how future changes in natural forcings or human activities could change this equilibrium between salt accumulation within and groundwater movement away from the Okavango Delta. Driving agents that could change the system dynamics include: climatic changes, geological changes, hydrological changes, vegetational-geomorphological changes, zoological activities, and anthropogenic changes (Ringrose et al., 2005). Human activities such as increased fertilizer application or reservoir construction could result in changes to river flow and plant species composition (Warne, 2004; Murray-Hudson et al., 2006). This model helps establish

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

current conditions and provides a platform to evaluate impacts on regional transport of the salts that accumulate within the Okavango Delta that could result from possible future change of natural and human forcings.

4 Conclusions

5 Modern society faces challenges in meeting human water needs while maintaining functionally intact freshwater ecosystems that provide goods and services with inter-generational benefits to society. A GIScience methodology is presented here to conceptualize groundwater and its interactions with ecology. Approaches are developed to study the impact of human-induced forcings, change in species, and forcings by natural 10 processes on groundwater ecohydrology.

The GIScience methods for conceptual models and datasets promote a framework to integrate data from existing sources while archiving data in a format that promotes interoperability for future use and benefit to water-dependent communities. International datasets and results from previous studies are organized and archived within the 15 conceptual data model presented in Fig. 2. Hydrogeology is organized about aquifer features (that through which water flows) and water features (that which drives the flow of water). Relationships are developed between vector (point, line, polygon) and raster (grid) data types and their properties. This addresses current lack of interoperability of data across computational groundwater models and provides a platform that is shown 20 to support numerical approaches based upon the Analytic Element Method, the Finite Difference Method, and the Finite Element Method. The data model provides flexibility in management of data and naming conventions and provides rules and topological relationships that enhance data integrity over existing methodology using shapefiles. Further information may be found at <http://www.growe.ksu.edu>.

25 Case studies deal with globally important ecosystems in Africa, Europe, and North America. In each study, the groundwater ecology processes and current state of knowledge are reviewed and important ecosystem forcings are identified. We then develop

Groundwater
ecohydrology studies
in Africa, Europe and
North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

new conceptual and computational models of each ecosystem to study and understand ecohydrologic processes. In North America, we study interrelationships between groundwater pumping and phreatophyte distribution over the Ogallala Aquifer using the conceptual model in Fig. 4 with data sources and previous studies in Table 1. Results in Fig. 5 illustrate while groundwater pumping has led to regional groundwater declines, the aquitard beneath the alluvium in the Arkansas River corridor holds groundwater near the surface where phreatophytes continue to exist. In Europe, we study interrelationships between forest species composition and groundwater baseflow discharge to biologically diverse fens and wetlands using the conceptual model in Fig. 6 with data sources and previous studies in Table 2. Results in Fig. 7 illustrate the potential for reestablishment of fens as coniferous forest are replaced by deciduous forests with lower interception rates, and the NHI-based groundwater model supports scenario testing to quantify the impacts of these change on baseflow discharge to fens and wetlands. We also demonstrate the capacity of these methods to serve as a computational platform for the National Hydrological Instrument of The Netherlands, and discuss the capacity of this framework to bridge datasets between existing AEM-based national groundwater models and ongoing FDM-based MODFLOW models. In Africa, we study the regional role of groundwater in transport of salts that accumulate in the Okavango Delta region using the conceptual model in Fig. 8 with data sources and previous studies in Table 3. Results in Fig. 9 identify a low gradient in groundwater that contributes to flushing of salts deposited through vegetation enhanced evapotranspiration and enable this freshwater ecosystem to exist.

The grand challenge posed by CUAHSI of linking the hydrosphere and biosphere, and the societal obligation to meet human water needs while sustaining functionally intact and biologically diverse ecosystems, each require new developments in the interdisciplinary science of groundwater ecohydrology. In this paper, we reviewed and documented three important problems that cross this divide. We organized international datasets supplemented by domain specific knowledge gained from previous studies to develop a conceptualization of the groundwater ecohydrology. While a plethora of

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



mechanisms exist to organize this data, we chose to adopt a simple data model to support this process. This approach to conceptualizing groundwater ecohydrology supports a broad range of computational tools to investigate and develop understanding of these ecosystems. This provides a framework to study sustainability, to forecast the impacts of changes in forcings, and to provide a scientific underpinning that informs management and public policy debate.

Acknowledgements. The author gratefully acknowledges financial support provided by TNO-Geological Survey of The Netherlands, the National Science Foundation (grant EPS0553722), the United States Department of Agriculture/Agriculture Research Service (Cooperative Agreement 10 58-6209-3-018), and the Provost Office's Targeted Excellence Program at Kansas State University. Special thanks to J. Delsman, B. Goes, J. Verkaik and R. Vernes at Deltares for providing access to data for the Veluwe model.

References

Adriaens, P., Goovaerts, P., Skerlos, S., Edwards, E., and Egli, T.: Intelligent infrastructure for sustainable potable water: A roundtable for emerging transnational research and technology development needs, *Biotechnol. Adv.*, 22, 119–134, 2003. 2798

Baron, J. S., Poff, N. L., Angermeier, P. L., Dahm, C. N., Gleik, P. H., Hairston, Jr., N. G., Jackson, R. B., Johnston, C. A., Richter, B. D., and Steinman, A. D.: Meeting ecological and societal needs for freshwater, *Ecol. Appl.*, 12, 1247–1260, 2002. 2797, 2798

Batelaan, O. and De Smedt, F.: GIS-based recharge estimation by coupling surface-subsurface water balances, *J. Hydrol.*, 337, 337–355, 2007. 2797

Batelaan, O., De Smedt, F., and Triest, L.: Regional groundwater discharge: phreatophyte mapping, groundwater modeling and impact analysis of land-use change, *J. Hydrol.*, 275, 86–108, 2003. 2797

Batjes, N. H.: SOTER-based soil parameter estimates for Southern Africa, ISRIC – World Soil Information, Wageningen Report 2004/04, 2004. 2838

Bauer-Gottwein, P., Langer, T., Prommer, H., Wolski, P., and Kinzelbach, W.: Okavango Delta Islands: Interaction between density-driven flow and geochemical reactions under evapo-concentration, *J. Hydrol.*, 335, 389–405, 2007. 2815

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Bear, J. and Verruijt, A.: Modeling Groundwater Flow and Pollution: With Computer Programs for Sample Cases, Springer, New York, 1987. 2801

Bekker, R. P., De Wit, P. V., Fernando, K. S., Turnisang, D. M., Radcliffe, D. J., and Mphathi, M.: Vegetation map of the Republic of Botswana. Soil mapping and advisory services project, 5 Ministry of Agriculture, Gaborone, 1991. 2835

Bernard, E. A., Steward, D. R., and Le Grand, P.: A geodatabase for groundwater modeling in MLAEM and MODFLOW, in: ESRI International User Conference 2005, Paper 1633: 1–25, ESRI, San Diego, CA, 25–29 July 2005. 2803

Best, E. P. H., Verhoeven, J. T. A., and Wolff, W. J.: The ecology of the Netherlands wetlands: 10 characteristics, threats, prospects and perspectives for ecological research, *Hydrobiologia*, 265, 305–320, 1993. 2811

Boumans, L., Fraters, D., and van Drecht, G.: Mapping nitrate leaching to upper groundwater in the sandy regions of the Netherlands, using conceptual knowledge, *Environ. Monit. Assess.*, 137, 243–249, doi:10.1007/s10661-007-9756-5, 2008. 2812

Bruen, M.: Systems analysis – a new paradigm and decision support tools for the water framework directive, *Hydrol. Earth Syst. Sci.*, 12, 739–749, 2008, <http://www.hydrol-earth-syst-sci.net/12/739/2008/>. 2798

Brunner, P., Franssen, H. J. H., Kgatlhang, L., Bauer-Gottwien, P., and Kinzelbach, W.: How 15 can remote sensing contribute in groundwater modeling?, *Hydrogeol. J.*, 15, 5–18, doi:10.1007/s10040-0060-0127-z, 2007. 2835

Brutsaert, W.: Long-term groundwater storage trends estimated from streamflow records: Climatic perspective, *Water Resour. Res.*, 44, W02409, doi:10.1029/2007WR006518, 2008. 2797

Bulatowicz, T., Jin, W., Staggenborg, S., Lauwo, S. Y., Miller, M., Das, S., Andrensen, D., Peter- 20 son, J., Steward, D. R., and Welch, S. M.: Calibration of a crop model to irrigated water use using a genetic algorithm, *Hydrol. Earth Syst. Sci.*, accepted, 2009. 2806

Butler Jr., J. J., Zlotnik, V. A., and Tsou, M.-S.: Drawdown and stream depletion produced by pumping in the vicinity of a partially penetrating stream, *Ground Water*, 39, 651–659, 2001. 2806

Butler Jr., J. J., Kluitenberg, G. J., Whittemore, D. O., Loheide II, S. P., Jin, W., Billinger, M. A., 30 and Zhan, X.: A field investigation of phreatophyte-induced fluctuations in the water table, *Water Resour. Res.*, 43, W02404, doi:10.1029/2005WR004627, 2007a. 2807

Butler Jr., J. J., Zhan, X., and Zlotnik, V. A.: Pumping-induced drawdown and stream depletion

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

in a leaky aquifer system, *Ground Water*, 45, 178–186, 2007b. 2807

5 Cederstrand, J. R. and Becker, M. F.: Digital map of base of aquifer for the High Plains Aquifer in parts of Colorado, Kansas, Nebraska, New Mexico, Oklahoma, South Dakota, Texas, and Wyoming, Open File Report 98-393, US Geological Survey, available online at <http://water.usgs.gov/pubs/ofr98-393>, 1998a. 2833

10 Cederstrand, J. R. and Becker, M. F.: Digital map of specific yield for the High Plains Aquifer in parts of Colorado, Kansas, Nebraska, New Mexico, Oklahoma, South Dakota, Texas, and Wyoming, Open File Report 98-414, US Geological Survey, available online at <http://water.usgs.gov/pubs/ofr98-414>, 1998b. 2833

15 10 Cederstrand, J. R. and Becker, M. F.: Digital map of hydraulic conductivity for the High Plains Aquifer in parts of Colorado, Kansas, Nebraska, New Mexico, Oklahoma, South Dakota, Texas, and Wyoming, Open File Report 98-548, US Geological Survey, available online at <http://water.usgs.gov/pubs/ofr98-548>, 1998c. 2833

15 Cleverly, J. R., Dahm, C. N., Thibault, J. R., McDonnell, D. E., and Coonrod, J. E. A.: Riparian ecohydrology: Regulation of water flux from the ground to the atmosphere in the Middle Rio Grande, New Mexico, *Hydrol. Process.*, 20, 3207–3225, doi:10.1002/hyp.6328, 2006. 2807

20 CUAHSI: Hydrology of a Dynamic Earth A decadal research plan for hydrologic science, Consortium of Universities for the Advancement of Hydrologic Science, Inc., <http://www.cuahsi.org>, 2007. 2798

20 Custodio, E.: Aquifer overexploitation: what does it mean?, *Hydrogeol. J.*, 10, 254–277, doi: 10.1007/s10040-002-0188-6, 2002. 2806

25 Dawson, T. E.: Hydraulic lift and water use by plants: implications for water balance, performance and plant-plant interactions, *Oecologia*, 95, 575–574, 1993. 2797

25 de Lange, W. J.: Development of an Analytic Element ground water model of the Netherlands, *Ground Water*, 44, 111–115, 2006. 2813, 2814, 2834

30 de Lange, W. J., Vergroesen, A. J., Snepvangers, J., Pastoors, M. J. H., and van Bakel, P. J. T.: Part 5: Freatisch lekweerstanden (surface waters into lumped resistances), *Nationaal Hydrologisch Instrumentarium Phase 1 Report Series NHI-5*, Deltares, Utrecht, The Netherlands, 2008b. 2834

30 de Lange, W. J., Vergroesen, A. J., Snepvangers, J., Pastoors, M. J. H., van Bakel, P. J. T., and Massop, H.: Part 4: Regionale oppervlakte wateren (regional surface waters), *Nationaal Hydrologisch Instrumentarium Phase 1 Report Series NHI-4*, Deltares, Utrecht, The Netherlands, 2008a. 2834

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

de Vries, J. J. and von Hoyer, M.: Groundwater recharge studies in semi-arid Botswana – a review, in: *Estimation of Natural Groundwater Recharge*, edited by: Simmers, I., 339–347, D. Reidel Publishing Company, 1988. 2815

de Vries, J. J., Selaolo, E. T., and Beekman, H. E.: Groundwater recharge in the Kalahari, with reference to paleo-hydrologic conditions, *J. Hydrol.*, 238, 110–123, 2000. 2815, 2818, 2835

5 De Wit, P., Bekker, R., Fernando, K. S., Malsela, T. M., Tselakgsi, K. O., Radcliffe, D. J., and Mphathi, M.: Land systems map of the Republic of Botswana. Soil mapping and advisory services project, Ministry of Agriculture, Gaborone, 1990. 2817, 2835

10 DEA: Okavango Delta Management Plan, Department of Environmental Affairs, Gaborone, Botswana, 2008. 2817, 2835

Dennehy, K. F.: High Plains regional ground-water study, Fact Sheet FS-091-00, US Geological Survey, <http://co.water.usgs.gov/nawqa/hpgw/factsheets/DENNEHYFS1.html>, 2000. 2806

Diersch, H. J. G.: FEFLOW Finite Element subsurface flow and transport simulation system, reference manual, WASY GmbH, Berlin, 2005. 2801

15 Dincer, T., Child, S., and Khupe, B.: A simple mathematical model of a complex hydrologic system – Okavango Swamp, Botswana, *J. Hydrol.*, 93, 41–65, 1987. 2815, 2816

Dunlap, L. E., Lindgren, R. J., and Sauer, C.: Geohydrology and model analysis of stream-aquifer system along the Arkansas River in Kearny and Finney counties, Southwestern Kansas, USGS Water-Supply Paper 2253, 1985. 2833

20 Ellery, W. N., Ellery, K., and McCarthy, T. S.: Plant distribution in islands of the Okavango Delta, Botswana: determinants and feedback interactions, *African J. Ecol.*, 31, 118–134, 1993a. 2816

Ellery, W. N., Ellery, K., Rogers, K. H., McCarthy, T. S., and Walker, B. H.: Vegetation, hydrology and sedimentation processes as determinants of channel form and dynamics in the northeastern Okavango Delta, Botswana, *African J. Ecol.*, 31, 10–25, 1993b. 2815, 2835

25 Elmore, A. J., Mustard, J. F., and Manning, S. J.: Regional patterns of plant community response to changes in water: Owens Valley, California, *Ecol. Appl.*, 13, 443–460, 2003. 2807

Foster, S. S. D. and Chilton, P. J.: Groundwater: The processes and global significance of aquifer degradation, *Phil. Trans. R. Soc. Lond. B*, 358, 1957–1972, doi:10.1098/rstb.2003. 30 1380, 2003. 2806

Gehrels, H.: Groundwater level fluctuations: Separation of natural from anthropogenic influences and determination of groundwater recharge in the Veluwe area, The Netherlands, Ph.D. thesis, Vrije University, Amsterdam, 1999. 2805, 2812

Groundwater
ecohydrology studies
in Africa, Europe and
North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Gehrels, J. C., van Geer, F. C., and de Vries, J. J.: Decomposition of groundwater level fluctuations using transfer modelling in an area with shallow to deep unsaturated zones, *J. Hydrol.*, 157, 105–138, 1994. 2811, 2812, 2834

5 Gieske, A.: Modelling outflow from the Jao/Boro River system in the Okavango Delta, Botswana, *J. Hydrol.*, 193, 214–239, 1997. 2815, 2816, 2835

Goes, B. J. M., Snepvangers, J., de Lange, W., Goorden, N., Pastoors, M. J. H., and Masseop, H. T. L.: Part 3: Maaiveld (Land elevation), Nationaal Hydrologisch Instrumentarium Phase 1 Report Series NHI-3, Deltares, Utrecht, The Netherlands, 2008a. 2834

10 Goes, B. J. M., Snepvangers, J., de Lange, W. J., Goorden, N., Pastoors, M. J. H., and Masseop, H. T. L.: Part 1: Ondergrond (sub-soil), Nationaal Hydrologisch Instrumentarium Phase 1 Report Series NHI-1, Deltares, Utrecht, The Netherlands, 2008b. 2834

Goodall, J. L., Horsburgh, J. S., Whiteaker, T. L., Maidment, D. R., and Zaslavsky, I.: A first approach to web services for the National Water Information System, *Environ. Model. Softw.*, 23, 404–411, 2008. 2833

15 Gregersen, J. B., Gijsbers, P. J. A., and Westen, S. J. P.: OpenMI: Open Modeling Interface, *J. Hydroinform.*, 9, 175–191, 2007. 2803

Grootjans, A. B., van Wirdum, G., Kemmers, R., and van Diggelens, R.: Ecohydrology in The Netherlands: principles of an application-driven interdiscipline, *Acta Botanica Neerlandica*, 45, 491–516, 1996. 2811, 2812

20 Gumbrecht, T., Wolski, P., Frost, P., and McCarthy, T. S.: Forecasting the spatial extent of the annual flood in the Okavango Delta, Botswana, *J. Hydrol.*, 290, 178–191, 2004. 2835

Gutentag, E. D., Heimes, F. J., Krothe, N. C., Luckey, R. R., and Weeks, J. B.: Geohydrology of the High Plains Aquifer in parts of Colorado, Kansas, Nebraska, New Mexico, Oklahoma, South Dakota, Texas and Wyoming, Professional Paper 1400-B, US Geological Survey, 1984. 2805, 2808, 2833

Hannah, D. M., Wood, P. J., and Sadler, J. P.: Ecohydrology and hydroecology: A “new paradigm”? *Hydrol. Process.*, 18, 3439–3445, doi:10.1002/hyp.5761, 2004. 2797, 2798

Hansen, C. V.: Estimates of Freshwater Storage and Potential Natural Recharge for Principal Aquifers in Kansas, Water Resources Investigations 87-4230, US Geological Survey, 1991. 2806, 2808, 2833

30 Hausberger, G., Davis, J. C., Miller, R. D., Look, K., Adkins-Helgeson, D., Ross, J. A., Bennett, B., Scholss, J., and Bohling, G. C.: WIZARD; Water Information Storage and Retrieval Database, Open File Report 98-13, Kansas Geological Survey, available online at

Groundwater
ecohydrology studies
in Africa, Europe and
North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

http://www.kgs.ku.edu/Magellan/WaterLevels/index.html, 1998. 2809, 2833

Hayashi, M. and Rosenberry, D. O.: Effects of ground water exchange on the hydrology and ecology of surface water, *Ground Water*, 40, 309–316, 2002. 2797

Hepworth, J. V., Jones, C. R., and Jennings, K.: Geological map of Botswana, Department of Geological Survey and Mines, Botswana, 1973. 2817, 2835

Higler, L. W. G. and Verdonschot, P. F. M.: Stream valleys as wetlands, *Hydrobiologia*, 265, 265–279, 1993. 2811

Horsburgh, J. S., Tarboton, D. G., Maidment, D. R., and Zaslavsky, I.: A relational model for environmental and water resources data, *Water Resour. Res.*, 44, W05406, doi:10.1029/2007WR006392, 2008. 2803

Jackson, R. B., Carpenter, S. R., Dahm, C. N., McKnight, D. M., Naiman, R. J., Postel, S. L., and Running, S. W.: Water in a changing world, *Ecol. Appl.*, 11, 1027–1045, 2001. 2798

Janković, I. and Barnes, R. J.: High-order line elements in modeling two-dimensional ground-water flow, *J. Hydrol.*, 226, 211–223, 1999. 2801

Jarvis, A., Reuter, H., Nelson, A., and Guevara, E.: Hole-filled seamless SRTM data V3, International Centre for Tropical Agriculture, <http://srtm.csi.cgiar.org>, 2006. 2835

Jobbágy, E. G. and Jackson, R. B.: Groundwater and soil chemical changes under phreatophytic tree plantations, *J. Geophys. Res.*, 112, G02013, doi:10.1029/2006JG000246, 2007. 2797

KGS: Water Well Completion Records (WWC5), Kansas Geological Survey <http://www.kgs.ku.edu/Magellan/WaterWell>, 2008. 2808, 2833

Kuiters, A. T. and Slim, P. A.: Regeneration of mixed deciduous forest in a Dutch forest-heathland following a reduction of ungulate densities, *Biol. Conserv.*, 105, 65–74, 2002. 2812

Leatherman, J. C., Cader, H. A., and Bloomquist, L. E.: When the Well Runs Dry: The Value of Irrigation to the Western Kansas Economy, *Kansas Pol. Rev.*, 26, 7–21, 2004. 2810

Lee, J.-E., Oliveira, R. S., Dawson, T. E., and Fung, I.: Root functioning modifies seasonal climate, *Proc. Nat. Acad. Sci USA*, 102, 17576–17581, 2005. 2807

Linn, F., Masie, M., and Rana, A.: The impacts on groundwater development on shallow aquifers in the lower Okavango Delta, northwestern Botswana, *Environ. Geol.*, 44, 112–118, doi:10.1007/s002-54-002-0706-9, 2003. 2815

Macfarlane, P. A. and Wilson, B. B.: Enhancement of the bedrock-surface-elevation map beneath the Ogallala portion of the High Plains Aquifer, western Kansas, Technical Series 20,

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Kansas Geological Survey, 2006. 2808, 2833

Maidment, D. R.: Arc Hydro: GIS for water resources, ESRI Press, Redlands, CA, 1 edn., 2002. 2803

McCarthy, J., Gumbrecht, T., and McCarthy, T. S.: Ecoregion classification in the Okavango Delta, Botswana from multitemporal remote sensing, *Int. J. Remote Sens.*, 26, 4339–4357, 2005. 2815

McCarthy, T. S.: Groundwater in wetlands of the Okavango Delta, Botswana, and its contribution to the structure and function of the ecosystem, *J. Hydrol.*, 320, 264–282, 2006. 2815, 2816

McCarthy, T. S. and Ellery, W. N.: The effect of vegetation on soil and ground water chemistry and hydrology of islands in the seasonal swamps of the Okavango Fan, Botswana, *J. Hydrol.*, 154, 169–193, 1994. 2815, 2816

McCarthy, T. S., McIver, J. R., and Verhagen, B. T.: Groundwater elevation, chemical sedimentation and carbonate brine formation on an island in the Okavango Delta swamp, Botswana, *Appl. Geochem.*, 6, 577–595, 1991. 2816

McCarthy, T. S., Ellery, W. N., and Ellery, K.: Vegetation-induced, subsurface precipitation of carbonate as an aggradational process in the permanent swamps of the Okavango (Delta) fan, Botswana, *Chem. Geol.*, 107, 111–131, 1993. 2816

McCarthy, T. S., Bloem, A., and Larkin, P. A.: Observations on the hydrology and geohydrology of the Okavango Delta, *Verhandelinge van die Geologiese Vereniging van Suid-Afrika*, 101, 101–117, 1998. 2818, 2835

McDonald, M. G. and Harbaugh, A. W.: A modular three-dimensional ground-water flow model, *Techniques of Water-Resource Investigation 06-A1*, USGS, US Government Printing Office, Washington, 1988. 2801

Montaigne, F.: Water pressure, *National Geographic*, 2–33, 2002. 2806

Moore, R. V. and Tindall, C. I.: An overview of the open modelling interface and environment (the OpenMI), *Environ. Sci. Policy*, 8, 279–286, 2005. 2803

Murray-Hudson, M., Wolski, P., and Ringrose, S.: Scenarios of the impact of local and upstream changes in climate and water use on the hydro-ecology in the Okavango Delta, Botswana, *J. Hydrol.*, 331, 73–84, 2006. 2818

Naumburg, E., Mata-Gonzalez, R., Hunter, R. G., McLendon, T., and Martin, D. W.: Phreatophytic vegetation and groundwater fluctuations: a review of current research and application of ecosystem response modeling with an emphasis on Great Basin vegetation, *Environ.*

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Nellis, M. D.: Geospatial Information Technology, Rural Resource Development, and Future Geographies, *Ann. Assoc. Am. Geogr.*, 95, 1–10, 2005. 2798

5 Newman, B. D., Wilcox, B. P., Archer, S. R., Breshears, D. D., Dahm, C. N., Duffy, C. J., McDowell, N. G., Phillips, F. M., Scanlon, B. R., and Vivoni, E. R.: Ecohydrology of water-limited environments: A scientific vision, *Water Resour. Res.*, 42, W06302, doi: 10.1029/2005WR004141, 2006. 2799

Opie, J.: *Ogallala: Water for a dry land*, University of Nebraska Press, Lincoln, 2nd edn., 2000. 2805

10 Palmer, M. A. and Bernhardt, E. S.: Hydroecology and river restoration: Ripe for research and synthesis, *Water Resour. Res.*, 42, W03S07, doi:10.1029/2005WR004354, 2006. 2797

Pastoors, M. J. H.: Part 13: Onttrekkingen (water supply wells), *Nationaal Hydrologisch Instrumentarium Phase 1 Report Series NHI-13*, Deltas, Utrecht, The Netherlands, 2008. 2834

15 Poff, N. L., Bledsoe, B. P., and Cuhaciyan, C. O.: Hydrologic variation with land use across the contiguous United States: Geomorphic and ecological consequences for stream ecosystems, *Geomorphology*, 79, 264–285, 2006. 2798

Postma, G., Roep, T. B., and Ruegg, G. H. J.: Sandy-gravelly mass-flow deposits in an ice-marginal lake (Saalian, Leuvenumsche Beek Valley, Veluwe, The Netherlands), with emphasis on plug-flow deposits, *Sediment. Geol.*, 34, 59–92, 1983. 2811

20 Powell, J. W.: Report on the lands of the arid region of the United States, Government Printing Office, Washington, second edn., 1879. 2806

Pringle, C. M.: Hydrologic connectivity and the management of a biological reserves: A global perspective, *Ecol. Appl.*, 11, 981–998, 2001. 2798

25 Puri, S. and Aureli, A.: Transboundary Aquifers: A Global Program to Assess, Evaluate, and Develop Policy, *Ground Water*, 43, 661–668, 2005. 2798

Reckhow, K., Band, L., Duffy, C., Famiglietti, J., Genereux, D., Helly, J., Hooper, R., Krajewski, W., McKnight, D., Ogden, F., Scanlon, B., and Shabman, L.: Designing Hydrologic Observatories: A paper prototype of the Neuse Watershed, Technical Report 6, Consortium of Universities for the Advancement of Hydrologic Sciences, Inc., Washington, DC, 2004. 2802

30 Richter, B. D., Mathews, R., Harrison, D. L., and Wigington, R.: Ecologically sustainable water management: Managing river flows for ecological integrity, *Ecol. Appl.*, 13, 206–224, 2003. 2798

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Rijkswaterstaat: Actueel Hoogtebestand Nederland (Actual Height model of the Netherlands), <http://www.ahn.nl>, Ministry of Transport, Public Works and Water Management, 2008. 2834

Ringrose, S., Jellema, A., Huntsman-Mapila, P., Baker, L., and Brubaker, K.: Use of remotely sensed data in the analysis of soil-vegetation changes along a drying gradient peripheral to the Okavango Delta, Botswana, *Int. J. Remote Sens.*, 26, 4293–4319, 2005. 2818

5 Rodriguez-Iturbe, I.: Ecohydrology: A hydrologic perspective of climate-soil-vegetation dynamics, *Water Resour. Res.*, 36, 3–9, 2000. 2797

Selaolo, E. T., Beekman, H. E., Gieske, A. S., and de Vries, J. J.: Multiple tracer profiling in Botswana GRES findings, in: *Groundwater recharge estimation in Southern Africa*, edited by 10 Xu, Y. and Beekman, H. E., IHP Series 64, chap. 3, 33–49, UNESCO, Paris, France, 2003. 2835

Sophocleous, M.: From safe yield to sustainable development of water resources – the Kansas experience, *J. Hydrol.*, 235, 27–43, 2000. 2807

15 Sophocleous, M.: Interactions between groundwater and surface water: the state of the science, *Hydrogeol. J.*, 10, 52–67, doi:10.1007/s10040-001-0170-8, 2002. 2806

Sophocleous, M.: Groundwater recharge and sustainability in the High Plains aquifer in Kansas, USA, *Hydrogeol. J.*, 13, 351–365, doi:10.1007/s10040-004-0385-6, 2005. 2806

20 Staudt, M.: Environmental hydrogeology of Ramotswa South East District, Republic of Botswana, Tech. rep., Botswana Department of Geological Survey, Environmental Geology Division, 2003. 2818, 2835

Steward, D. R. and Ahring, T.: An analytic solution for groundwater uptake by phreatophytes spanning spatial scales from plant to field to regional, *J. Eng. Math.*, doi:10.1007/s10665-008-9255-x, in press, 2009. 2807

25 Steward, D. R. and Bernard, E. A.: The synergistic powers of AEM and GIS geodatabase models in water resources studies, *Ground Water*, 44, 56–61, 2006. 2803

Steward, D. R. and Bernard, E. A.: Integrated engineering and landscape architecture approaches to address groundwater declines in the High Plains Aquifer, in: *Case Studies in Environmental Engineering and Science*, edited by Bhandari, A. and Butkus, M. A., chap. 9, 121–134, Association of Environmental Engineering & Science Professors, 2007. 2803

30 Steward, D. R., Le Grand, P., and Bernard, E. A.: The Analytic Element Method and supporting GIS geodatabase model, in: *Proc. 27th World Conference on BEM/MRM*, pp. 187–196, WIT Press, Orlando, FL, 15–17 March 2005. 2803

Steward, D. R., Le Grand, P., Janković, I., and Strack, O. D. L.: Analytic formulation of Cauchy

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶


[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



5 van Bakel, P. J. T. and Veldhuizen, A. A.: Part 10: Beregening (recharge), Nationaal Hydrologisch Instrumentarium Phase 1 Report Series NHI-10, Deltares, Utrecht, The Netherlands, 2008. 2834

van der Meulen, M. J., van Gessel, S. F., and Veldkamp, J. G.: Aggregate resources in the Netherlands, Netherlands J. Geosci., 84, 379–387, 2005. 2811

10 Vasak, L., Krajenbrink, G. J. W., and Appelo, C. A. J.: The spatial distribution of polluted groundwater from rural centres in a recharge area in the Netherlands – the Veluwe, The Science of the Total Environment, 21, 105–112, 1981. 2805, 2811

Vermulst, J. A. P. H. and de Lange, W. J.: An analytic-based approach for coupling models for unsaturated and saturated groundwater flow at different scales, J. Hydrol., 226, 262–273, 1999. 2811, 2834

15 Vernes, R. W. and van Doorn, T. H. M.: REGIS II, A 3D hydrogeological model of The Netherlands, vol. 38, p. 107, Geological Society of America, Philadelphia, 2006. 2834

Warne, K.: Okavango: Africa's Miracle Delta, National Geographic, pp. 42–67, <http://ngm.nationalgeographic.com/ngm/0412/feature3>, 2004. 2805, 2816, 2818

20 Whiteaker, T. L., Maidment, D. R., Gopalan, H., Patino, C., and McKinney, D. C.: Raster-network regionalization for watershed data processing, Int. J. Geogr. Inform. Sci., 21, 341–353, 2007. 2801

Wilson, B., Bartley, J., Emmons, K., Bagley, J., Wason, J., and Stankiewicz, S.: Water Information Management and Analysis System, Version 5, for the Web. User Manual, Open File Report 2005-30, Kansas Geological Survey, Lawrence, Kansas, <http://hercules.kgs.ku.edu/geohydro/wimas>, 2005. 2833

25 Wilson, B. H. and Dincer, T.: An introduction to the hydrology and hydrography of the Okavango Delta, Symposium on the Okavango Delta, 33–48, 1976. 2805, 2815

Wolski, P. and Savenije, H. H. G.: Dynamics of floodplain-island groundwater flow in the Okavango Delta, Botswana, J. Hydrol., 320, 283–301, doi:10.1016/j.jhydrol.2005.07.027, 2006. 2816

30 Yang, X. and Steward, D. R.: The Interactions among Human-induced Groundwater Decline, Salt Transport, and Salt Cedar Invasion in the Upper Arkansas River Corridor, Kansas, Hydrogeol. J., in review, 2009. 2808

Yang, X., Steward, D. R., de Lange, W. J., Lauwo, S. Y., Chubb, R. M., and Bernard, E. A.: Data

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶


[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



model for system conceptualization: A case example in groundwater studies, *Int. J. GIS*, in review, 2009. 2802, 2803

Zheng, C., Lin, J., and Maidment, D. R.: Internet data sources for ground water modeling, *Ground Water*, 44, 136–138, 2006. 2802

5 Zimmermann, S., Bauer, P., Held, R., Kinzelbach, W., and Walther, J. H.: Salt transport on islands in the Okavango Delta: Numerical investigations, *Adv. Water Resour.*, 29, 11–29, 2006. 2816

Groundwater
ecohydrology studies
in Africa, Europe and
North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Groundwater
ecohydrology studies
in Africa, Europe and
North America

D. R. Steward et al.

Table 1. Data resources for the Ogallala Aquifer region in North America.

Object	Feature	Data Sources
AquiferPoint	Borehole	KGS (2008)
AquiferLine	Bedrock elevation	Gutentag et al. (1984); Cederstrand and Becker (1998a); Macfarlane and Wilson (2006)
AquiferPolygon	Hydraulic conductivity	Dunlap et al. (1985); Gutentag et al. (1984); Cederstrand and Becker (1998c)
AquiferPolygon	Specific yield	Dunlap et al. (1985); Gutentag et al. (1984); Cederstrand and Becker (1998b)
AquiferRaster	Land elevation	USGS (2008a)
WaterPoint	Well	Hausberger et al. (1998); Wilson et al. (2005)
WaterLine	River	Tsou et al. (2006); Goodall et al. (2008); USGS (2008b,a)
WaterPolygon	Recharge	Hansen (1991)

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.

Table 2. Data resources for the Veluwe region in Europe.

Object	Feature	Data Sources
AquiferPoint	Borehole	TNO (2008)
AquiferLine	Fault	Vernes and van Doorn (2006); Goes et al. (2008b)
AquiferPolygon	Geological units	de Lange (2006); Vernes and van Doorn (2006); Goes et al. (2008b)
AquiferRaster	Hydraulic conductivity	Vernes and van Doorn (2006); Goes et al. (2008b)
AquiferRaster	Land elevation	Goes et al. (2008a); Rijkswaterstaat (2008)
AquiferRaster	Layer elevation	Vernes and van Doorn (2006); Goes et al. (2008b)
WaterPoint	Well	Pastoors (2008)
WaterLine	River	de Lange et al. (2008a)
WaterPolygon	Lake, River	de Lange et al. (2008a)
WaterRaster	Recharge	Gehrels et al. (1994); van Bakel and Veldhuizen (2008)
WaterRaster	Surface water properties	Vermulst and de Lange (1999); de Lange et al. (2008b)

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Table 3. Data resources for the Okavango Delta region in Africa.

Object	Feature	Data Sources
AquiferLine	Dike	DEA (2008)
AquiferLine	Fault	Hepworth et al. (1973)
AquiferPolygon	Graben	Hepworth et al. (1973); De Wit et al. (1990); Brunner et al. (2007)
AquiferPolygon	Ridge	Hepworth et al. (1973); Staudt (2003)
AquiferRaster	Land elevation	Jarvis et al. (2006)
WaterLine	River	Ellery et al. (1993b); Gieske (1997); Jarvis et al. (2006)
WaterPolygon	Evapotranspiration	Gumbricht et al. (2004); Brunner et al. (2007)
WaterPolygon	Recharge	de Vries et al. (2000); Selaolo et al. (2003)
WaterPolygon	Wetland	De Wit et al. (1990); Bekker et al. (1991); McCarthy et al. (1998)

[Title Page](#) [Abstract](#) [Introduction](#)
[Conclusions](#) [References](#) [Tables](#) [Figures](#)
[◀](#) [▶](#) [◀](#) [▶](#)
[Back](#) [Close](#) [Full Screen / Esc](#)
[Printer-friendly Version](#) [Interactive Discussion](#)

Steward et al.: Groundwater ecohydrology GIScience tools and studies in Africa, Europe and North America




Fig. 1. Conceptual view of groundwater ecology: water that seeps downward past the root zone of the terrestrial ecosystem recharges the groundwater aquifer and eventually discharges to the surface through well pumping, phreatophyte root uptake, or baseflow to surface water in the aquatic ecosystem.

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)


[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.



Objects include points, , lines, , polygons, , raster catalogs, , and tables, ; relationships indicate redundancy (e.g.,  relates one-to-many).

Fig. 2. Groundwater concepts data model.






Fig. 3. The three study regions and their soil classifications obtained for the Ogallala Aquifer in the United States of America from the Digital General Soil Map (USDA, 2006), for the Veluwe in The Netherlands from CORINE (UNEP, 1985) and for the Okavango Delta in Botswana from ISRIC (Batjes, 2004). Note that soil classifications and color schemes are used from each of the respective data repositories.

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)




Fig. 4. Conceptual model of the Ogallala Aquifer, western Kansas, USA.

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.




Fig. 5. Groundwater elevation (m above m.s.l.) and depth to groundwater (m) in the Ogallala Aquifer, western Kansas, USA for predevelopment (top) and current (bottom) times.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.




Fig. 6. Conceptual model of the Veluwe, The Netherlands.

[Printer-friendly Version](#)

[Interactive Discussion](#)

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.




Fig. 7. Groundwater elevation (m above m.s.l.) and depth to groundwater (m) in the Veluwe, The Netherlands with coniferous (top) or deciduous (bottom) forests.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.




Fig. 8. Conceptual model of the Okavango Delta, Botswana.

Groundwater ecohydrology studies in Africa, Europe and North America

D. R. Steward et al.




Fig. 9. Groundwater elevation (m above m.s.l.) and depth to water (m) in the Okavango Delta, Botswana.