

Papers published in *Hydrology and Earth System Sciences Discussions* are under open-access review for the journal *Hydrology and Earth System Sciences*

Modelling the inorganic nitrogen behaviour in a small Mediterranean forested catchment, Fuirosos (Catalonia)

C. Medici¹, S. Bernal², A. Butturini³, F. Sabater³, A. J. Wade⁴, and F. Frances¹

¹Department of Hydraulic and Environmental Engineering, Polytechnic University of Valencia, Spain

²Centre d'Estudis Avancats de Blanes (CEAB-CSIC), Girona, Spain

³Department of Ecology, University of Barcelona, Spain

⁴Aquatic Environments Research Centre, University of Reading, Reading, UK

Received: 29 July 2009 – Accepted: 9 August 2009 – Published: 4 September 2009

Correspondence to: C. Medici (chme1@doctor.upv.es)

Published by Copernicus Publications on behalf of the European Geosciences Union.

HESSD

6, 5665–5703, 2009

Modelling inorganic
nitrogen behaviour in
Mediterranean
forested catchment

C. Medici et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

The aim of this work was to couple a nitrogen (N) sub-model to already existent hydrological lumped (LU4-N) and semi-distributed (LU4-R-N and SD4-R-N) conceptual models, to improve our understanding of the factors and processes controlling nitrogen cycling and losses in Mediterranean catchments. The N model adopted provides a simplified conceptualization of the soil nitrogen cycle considering mineralization, nitrification, immobilization, denitrification, plant uptake, and ammonium adsorption/desorption. It also includes nitrification and denitrification in the shallow perched aquifer. We included a soil moisture threshold for all the considered soil biological processes. The results suggested that all the nitrogen processes were highly influenced by the rain episodes and that soil microbial processes occurred in pulses stimulated by soil moisture increasing after rain. Our simulation highlighted the riparian zone as a possible source of nitrate, especially after the summer drought period, but it can also act as an important sink of nitrate due to denitrification, in particular during the wettest period of the year. The riparian zone was a key element to simulate the catchment nitrate behaviour. The lumped LU4-N model (which does not include the riparian zone) could not be validated, while both the semi-distributed LU4-R-N and SD4-R-N model (which include the riparian zone) gave satisfactory results for the calibration process and acceptable results for the temporal validation process.

1 Introduction

Nitrogen is present in both terrestrial and aquatic ecosystems and research is needed to understand its storage, transportation and transformations in river catchments worldwide because of its importance in controlling plant growth and freshwater trophic status (Vitousek et al., 2009; Chu et al., 2008; Schlesinger et al., 2006; Ocampo et al., 2006; Green et al., 2004; Arheimer et al., 1996). Numerous mathematical models have been developed to describe the nitrogen dynamics in cool temperate river-systems, but fur-

HESSD

6, 5665–5703, 2009

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

ther work is needed to understand and model the main processes controlling the nitrogen cycle in Mediterranean and semi-arid ecosystems since these are not well understood (Gelfand et al., 2008; Bernal et al., 2005; Avila et al., 1995; Wade et al., 2005). Mediterranean catchments are characterized by a complex hydrological behaviour that causes high inter and intra-annual variability in flow (Gallart et al., 2002). Consequently, models developed for temperate climates generally fail when applied to Mediterranean catchments (Bernal et al., 2004). Mediterranean ecosystems are subjected to severe drought periods followed by intense rainfall events, which produce alternate dry and humid conditions that influence the soil microbial activity (Austin et al., 2004; Reynolds et al., 2004; Schwinning et al., 2004b). Models based on a representation of temperate climates do not represent this rapid transition from dry to wet periods well. Birch (1959, 1960, 1964) was one of the first to characterize the impacts of soil drying and wetting cycles on mineralization and nitrification, demonstrating that rapid mineralization follows rewetting of dry soil and that in continuously moist conditions there is a release of nitrogen, much of it as nitrate. Many other authors stressed the influence that wet-dry cycles have on microbial biomass (Van Gestel et al., 1993), denitrification (Mummey et al., 1994; Peterjhon and Schlesinger, 1991) and ammonia volatilization (Heckathorn and Delucia, 1995). Schiwinning et al. (2004a, b) spoke about a “pulse dynamic” in arid and semi-arid ecosystems, considering the rainfall inputs to a dry soil as triggers of a cascade of biogeochemical and biological transformations. According to Schiwinning, precipitation applied to a dry soil surface creates a pulse of soil moisture that can be characterized by the depth to which soil water potentials are elevated to levels that promote biological activity and the length of time over which water potentials remain at biologically relevant levels. Intermittent streams and their associated riparian zone have been highlighted as “hot spots” for biogeochemical processes in arid and semi-arid regions (McIntyre et al., 2009). Bernal et al. (2007) suggested that Mediterranean riparian soils act as source or sink of dissolved nitrogen depending on the period of the year, mainly due to contrasting soil moisture condition between the dry and the wet period. Moreover, Butturini et al. (2003) suggested the unsaturated riparian soil of the

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Fuirosos catchment, a small intermittent Mediterranean stream in Catalonia (Spain), as a possible source of nitrate, especially after the summer drought, which can be rapidly mobilized due to the formation of a rising riparian groundwater table into the unsaturated upper soil layer adjacent to the stream channel. The nitrogen dynamics of the Fuirosos catchment were analysed previously with the process-based Integrated Catchment Model of Nitrogen (INCA-N) model (Whitehead et al., 1998; Wade et al., 2002; Bernal et al., 2004). INCA-N was developed for temperate regions and has been demonstrated to simulate properly the hydrology and nitrogen dynamics observed in these types of ecosystems (Wade et al., 2004). The model gave unsatisfactory result for the Fuirosos catchment suggesting that key processes were missing (Bernal et al., 2004). The present research aims to develop a new model to represent the inorganic nitrogen response in Mediterranean catchments using INCA-N as a basis, but including additional mechanisms to take into account the ideas and results pointed out before and obtained in previous studies in semi-arid and Mediterranean catchments. Namely, these new elements are: biological thresholds responses to soil moisture in order to reproduce the pulse dynamic observed in such environment; a specific function for the soil moisture correction factor for the mineralization process; nitrification and denitrification processes associated to the shallow perched water table and finally, the introduction of a riparian zone compartment. The nitrogen model scheme developed in this study was coupled to already existent hydrological conceptual models previously applied to the Fuirosos catchment (Medici et al., 2008).

2 Study site

The Fuirosos catchment (41°42' N, longitude 2°34' E) is located in the northern slopes of Catalan Littoral Range, near Barcelona (Spain) and it is a tributary of the Tordera River. The drainage area of Fuirosos is approximately 13 km² and the altitude ranges from 50 to 770 m a.s.l. (Fig. 1). The catchment is almost pristine, the predominant land cover being undisturbed forest; there is little agricultural activity and no urban areas.

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Within the catchment, there are four small reservoirs for human and cattle water supply (Fig. 1). This water consumption can be considered insignificant during the study period. The storage volume of these reservoirs ranges approximately from 5000 to 18 000 m³. The climate is typically Mediterranean, with temperature ranging from a monthly mean of 3°C in January to 24°C in August. Winter temperatures below 0°C are infrequent. During the observed period (from October 1999 to June 2003), the mean annual precipitation at Fuirosos was approximately 750 mm. The first hydrological year (1999/2000) was the driest of the four considered in this study (annual precipitation 454 mm) and the third (2001/2002) was the wettest (annual precipitation 850 mm). The mean annual potential evapotranspiration (PET) computed with the Penman equation considering the period from October 1999 to June 2003, was approximately 975 mm. The observed number of consecutive days during which the Fuirosos stream was completely dry reached values of 76 (summer 2000) and 98 (summer 2001). The predominant rock type in the Fuirosos catchment is leucogranite (50.9%). Other rock types include granodiorite (21.1%) and sericitic schists (23.5%) (IGME, 1983; Fig. 1). At the valley bottom there is an alluvial zone, where a well-developed riparian area flanks the Fuirosos stream channel. The forest covers the 90% of the total catchment area where perennial cork oak (*Quercus suber*) and pine tree (*Pinus halapensis* and *Pinus pinaster*) predominate. However, at the valley headwaters, mixed deciduous woodland of chestnut (*Castanea sativa*), hazel (*Corylus avellana*) and oak (*Quercus pubescens*) prevail. The discharge was measured from 13 October 1999 to 30 June 2003. Daily streamwater nitrate (NO₃) concentrations were also measured in water samples taken from the catchment outlet during the period from October 1999 to April 2003 and daily ammonium (NH₄) concentrations were also measured during the period from January 2001 to August 2002. For a complete description of the Fuirosos chemical water analyses see Bernal et al. (2004, 2005).

3 N-model description

The hydrological behaviour of the Fuirosos catchment has been successfully modelled previously (Medici et al., 2008). A key result of this previous study is that the perceptual model including four different catchment hydrological responses (direct flow, interflow, quick and slow base flow) is the most suitable to simulate the discharge at Fuirosos. The initial lumped conceptual model proposed (LU4) was developed into a semi-distributed form (SD4-R) in which the spatial variability of the evapotranspiration according to the vegetation cover and the local aspect was considered. In the final semi-distributed structure of the hydrological model (which gave a best fit of 0.78 in term of Nash & Sutcliffe index) an additional conceptual store representing the riparian zone was included, as well as the four reservoirs present in the catchment. In the current work, the previous cited models were extended to include processes representing the inorganic nitrogen cycle to simulate the nitrate and ammonium concentration observed in the Fuirosos stream. Therefore, the progressive perceptual approach adopted led from an initial lumped structure (LU4-N) to a very simple semidistributed one (LU4-R-N) that included the riparian tank along with the four small reservoirs and eventually to a more complex semidistributed one (SD4-R-N) that included the riparian zone, the four reservoirs as well as catchment spatial variability to some extent. The first approach to simulate the transport, storage and transformations of nitrogen in the terrestrial and aquatic components of the catchment was done using the lumped hydrological (LU4) model as a basis. The LU4-N model integrates hydrology, soil and shallow aquifer N processes, and simulates daily $\text{NO}_3\text{-N}$ and $\text{NH}_4\text{-N}$ concentrations in the stream. The soil nitrogen cycle conceptual model includes the mineralization process and non-biological nitrate fixation modelled using zero order kinematic equations. The processes of nitrification, ammonium bacterial immobilisation, ammonium and nitrate soil plant uptake, abiotic absorption and denitrification are included and represented using first order kinematic equations. The total number of parameter to be calibrated for the LU4-N model is 28 of which 9 are for the rainfall-runoff sub-model and 19 for the

HESSD

6, 5665–5703, 2009

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



N sub-model. A perceptual model which shows the key nitrogen stores and pathways is presented in Fig. 2. At present, the only source of N is atmospheric deposition as this is the main input of nitrogen in the catchment but other anthropogenic sources could be included in future versions if required. For the deposition, the estimated values obtained by Rodá et al. (2002, after Bernal et al., 2004) were used. Namely: the wet deposition of inorganic N was $5.7 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ (52% as ammonium and 48% as nitrate), while the dry deposition of inorganic N was $9.2 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ (45% as ammonium and 55% as nitrate). The model equations were written in terms of N mass and water volume and a daily time step was adopted. The equations were solved sequentially (i.e. for the soil ammonium cycle: first of all mineralization, secondly immobilization then plant uptake and finally nitrification) and it was verified, taking into account several different sequences, that the particular one adopted did not significantly affect the model results. In both shallow and deeper aquifer, N uptake associated with the transpiration flux is assumed to occur, which depends on the simulated NH_4 and NO_3 concentration in each aquifer, on the amount of water transpired by plants and finally on the annual maximum solute uptake. All the soil processes are adjusted by a soil moisture factor ($S_{1_Process}$) to represent the moisture control on bacterial processes and are temperature dependent (Whitehead et al., 1998; Wade et al., 2002). Moreover, a different soil moisture threshold (U) has been introduced for each soil process to determine activation. The concept of a threshold response is not new in arid land ecology (Reynolds et al., 2004; Schwinning et al., 2004a). Traditionally this concept has been related with the ecosystem primary production, though Schwinning and Sala (2004) generalized the threshold paradigm to a wide range of ecosystem processes. In fact, they suggested that the hierarchy of pulse events has a corresponding hierarchy of ecological responses that is determined by the ability of organism to utilize soil moisture pulses of different duration, infiltration depths and soil water potential. As a matter of example, the mineralization processes is described as:

$$M_{\text{NH}_4_Miner}(t) = K_{\text{Miner}} \cdot S_{1_Miner}(t) \cdot TF \quad (1)$$

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

where: $M_{\text{NH}_4\text{-Miner}}$ is the ammonium mineralized mass ($\text{kg ha}^{-1} \text{ day}^{-1}$) in a time step; K_{miner} is mineralization rate constant ($\text{kg ha}^{-1} \text{ day}^{-1}$) and TF is temperature factor, according to Wade et al. (2002) and $S_{1\text{-Miner}}$ is the soil moisture factor, which is calculated as follows:

$$S_{1\text{-Miner}}(t) = \frac{H_1(t) - IA}{U_{\text{Miner}}} \quad \text{if } 0 \leq H_1(t) - IA \leq U_{\text{Miner}} (\leq H_u^* - IA) \quad (2)$$

$$S_{1\text{-Miner}}(t) = 1 - \frac{(H_1(t) - IA - U_{\text{Miner}})}{(H_u^* - IA - U_{\text{Miner}})} \quad \text{if } U_{\text{Miner}} < H_1(t) - IA \leq H_u^* - IA$$

where: H_1 is the actual static storage water content (mm) and H_u^* is the maximum static storage water content (mm) (where the static tank represents water that can leave the catchment only by evapotranspiration); IA are the initial abstractions (interception and water detention in puddles) which were (approximately) estimated as 19 mm day^{-1} ; t is the time step (day) and U_{Miner} is the soil moisture threshold for mineralization (mm), which is expressed as a percentage of H_u^* .

According to Eq. (2), the $S_{1\text{-Miner}}$ factor has a triangular shape with a maximum value when the soil moisture content is equal to U_{Miner} . This is consistent with McIntyre et al. (2009), who found that mineralization is reduced under soil moisture content close to saturation, but increases under moderate soil moisture content. For the other soil nitrogen processes, the corresponding soil moisture factors are computed according the following general expression:

$$S_{1\text{-Process}}(t) = 0 \quad \text{if } 0 \leq H_1(t) - IA \leq U_{\text{Process}} (\leq H_u^* - IA)$$

$$S_{1\text{-Process}}(t) = \frac{(H_1(t) - IA - U_{\text{Process}})}{(H_u^* - IA - U_{\text{Process}})} \quad \text{if } U_{\text{Process}} < H_1(t) - IA \leq H_u^* - IA \quad (3)$$

where: U_{Process} is the generic soil moisture threshold for the soil process included in the model (except mineralisation); $S_{1\text{-Process}}$ is the soil moisture factor for any soil nitrogen process. Thus for any soil N process, except mineralization, a minimum soil moisture content is needed for the process to be activated.

The LU4-N model was then progressed to a simple semi-distributed structure splitting the catchment into two Hydrological Representative Units (HRUs): (1) the riparian

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



zone that represents only 0.5% of the total catchment area and (2) the rest of the catchment (hill-slope hereafter). In this way two different parameters sets were considered, one for each HRU. The LU4-R-N considers neither the spatial variability of the evapotranspiration nor that of the lithology. The LU4-R-N model requires 42 parameters to be calibrated, of which 11 for the rainfall-runoff model and 31 for the N sub-model (12 specific for each HRU and 7 common for the whole catchment) (Table 1). The aim with this model structure was to analyze the possible effect of the riparian zone on nitrate release to the stream. The hydrological conceptual scheme adopted for the semi-distributed model differs slightly from that published in Medici et al. (2008). In this case, part of the hill-slope discharge (corresponding to the area not drained by the four small reservoirs, which represents approximately 37% of the total catchment area) is routed through the riparian storage before reaching the stream channel (Fig. 3). This change does not affect the hydrology simulation considerably, but is thought to be relevant for simulating solute behaviour. We assumed that the main effect of the four reservoirs mainly was dilution on nitrate and ammonium concentration. In a next phase of development, the LU4-R-N was extended to include the spatial variation in evapotranspiration and lithology (SD4-R-N). As such, the catchment was divided into 4 HRUs: the three main catchment lithological units (leucogranite, granodiorite and sericitic schists, all together cited in this paper as hill-slope zone) and the riparian zone, as those used in the application of the SD4-R hydrological model (Medici et al., 2008). Thus, the PET spatial variability for the actual evapotranspiration computation was included taking into account the representative vegetation cover and the potential sunshine arriving to each lithological unit according to its representative aspect and surrounding relief. The parameterization of the 4-HRUs was done for the rainfall-runoff sub-model only; for the N sub-model, only the riparian and reminder of the catchment HRUs were considered for parameterization (Table 1). In this case, the total number of parameters to be calibrated for the hydrological model is 28, while for the N model is still 31 as for the LU4-R-N model.

4 Results

The calibration period covers approximately three hydrological years from October 1999 to August 2002, while the validation one considers the period from August 2002 to June 2003. Only nitrate concentrations were available for the temporal validation process. Parameters were optimized taking into account the Nash and Sutcliffe efficiency index (E), the balance error in terms of observed and simulated global loads (BE), the graphical fit between observed and simulated N time-series, the relative Root Mean Square Error (RMSE) index and the coefficient of determination (r^2). The calibration was done by an automatic process, namely Evolver 4.0 for Excel (32-bit) and then by final manual adjustment of the parameters to check the behaviour of the model. For the LU4-N model the same parameters determined in the study by Medici et al. (2008) were adopted for the hydrology simulation, so therefore only the 19 N-model parameters were calibrated in this study (Table 1). On the other hand, in the case of the semidistributed models (LU4-R-N and SD4-R-N), the rainfall-runoff model was calibrated firstly and afterwards the N sub-model. Because of the different hydrological scheme adopted for this study, the parameters set for the hydrology slightly differed to that proposed in Medici et al. (2008) without representing any relevant change worthy of attention. The parameter values determined in the calibration of each of the three nitrogen sub-model structures are shown in Table 1. The goodness-of-fit measures for the calibration and validation periods are summarized in Tables 2 and 3, respectively.

4.1 LU4-N calibration and validation results

Observed nitrate and ammonium daily stream concentrations at Fuirosos and the corresponding simulated ones, obtained with the LU4-N model structure, are shown in Fig. 4a. The LU4-N model reproduced quite satisfactorily the observed daily nitrate concentrations for the calibration period ($E=0.46$). According to this model conceptualization, the main pathway controlling nitrate flushing is the flow derived from the shallow aquifer. As a matter of example, to reproduce the highest nitrate peak observed

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



during March 2002 (Fig. 4a) the LU4-N model simulated, during the previous months, a huge accumulation of ammonium in soil that due to a significant rainfall event (almost 40 mm/day) percolated to the shallow aquifer where it was rapidly nitrified to nitrate. This nitrate rapidly reached the stream being transported by with the water flowing from the shallow aquifer to the stream. The LU4-N model rarely generates interflow, which in general is associated with rainfall largest events ($>40 \text{ mm day}^{-1}$) during the wet period, so it is the responsible for the nitrate flushing just in very few occasions. E.g.: the observed nitrate peak of the second year simulated (December 2000) it was a large simulated pulse of nitrification in the soil (almost $130 \text{ Kg Km}^{-2} \text{ day}^{-1}$) that caused a major flush of nitrate transported with interflow. In fact, the model simulated an earlier ammonium increase in soil that was rapidly nitrified when the soil moisture content exceeded the threshold for nitrification as a result of a large rainfall event (43 mm day^{-1}) (Fig. 5). This nitrification pulse dynamic reproduced in terms of average annual loads a Mineralisation:Nitrification (M:N) ratio of 10:1, which is consistent with the results of Serrasolses et al. (1999). On the other hand, it is worthy to notice that the daily simulated M:N ratio can achieve much higher values or it can also take values between zero and one (that means that nitrification overcomes mineralization), when a huge pick of nitrification takes place (Fig. 6). Concerning the simulation of streamwater ammonium concentrations, the LU4-N model could not reproduce the observations ($E < 0$) and the statistical relation between the simulated and observed data was not significant (Table 2). Despite the good results obtained for the calibration of the stream daily nitrate concentrations, the LU4-N model gave poor results for the validation period (Table 3). The model overestimated the nitrate concentration from August to October 2002, due to excessive nitrate amount carried by the base flow and the streamwater nitrate concentrations observed during late autumn and winter 2002–2003 were underestimated (Fig. 7a). A simple one-at-a-time perturbation sensitivity analysis highlighted that the mineralization related parameters (K_{\min} and U_{\min}), along with the maximum static storage water content (H_u^*) and the maximum annual ammonium plant uptake ($\text{MaxUP}_{\text{NH}_4}$) had the major impact on the nitrate related objective functions. Ammonium soil adsorp-

tion rate (K_{ads}) and the nitrification soil moisture threshold (U_{nitr}) were also highlighted as quite sensitive parameters considering the ammonium related objective functions.

4.2 LU4-R-N calibration and validation results

Observed nitrate and ammonium daily stream concentrations at Fuirosos and the corresponding simulated ones, obtained with the LU4-R-N model structure, are shown Fig. 4b. The obtained discharge efficiency and goodness indexes for the calibration period are similar to those obtained from the simulations done using the LU4-N model (Table 2). This occurs in part because the calibrated parameters for the hydrological components of the models are similar. Though, the nitrate simulation for the calibration period improved: the global E index for the daily nitrate concentration increased to 0.56, and the global BE error decreased to approximately -15% , despite the fact that the LU4-R-N model largely underestimated the highest nitrate concentration peak observed during March 2002 (Fig. 4b). The LU4-R-N model reproduced the nitrate concentration peak observed during April 2002 that was not simulated by the LU4-N model. During this occasion, because of a large rainfall event (almost 64 mm day^{-1}) the two models could generate nitrate that washed from the soil with interflow at approximately the same rate. However, in the case of the LU4-R-N model, part of the interflow passed through the riparian zone soil (Fig. 3) mobilizing nitrate previously accumulated in this pool. It has to be noticed that in the riparian soil, the simulated mineralization process occurred at a significantly higher rate than in the hill-slope soil and the nitrification process followed more closely the pattern of simulated mineralization being activated more easily than in the hill-slope area (Fig. 8). Therefore, the simulated annual M:N ratio in the riparian zone was almost 1:1 as well as the daily M:N ratio, while in the hill-slope zone the M:N ratio showed a higher variability as in the case of the lumped LU4-N model (Fig. 6). This dynamic allowed a significant amount of nitrate to be accumulated in the riparian soil, which was available to be rapidly flushed away by interflow derived from the hill-slope soil, as observed in April 2002. The temporal validation process gave better results for the LU4-R-N model than for the LU4-N

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



model (Table 3 and Fig. 7b). In particular, the introduction of the riparian zone allowed reproducing the nitrate concentration peak observed during November 2002 due to the same mechanism aforementioned (i.e.: previous nitrate accumulation in the riparian upper soil that is afterwards flushed away by interflow derived from the hill-slope soil).

5 Finally, the LU4-R-N model failed to reproduce the observed stream daily ammonium concentration. There was only a weak statistical relation between the observed and simulated streamwater ammonium concentrations ($r^2=0.02$; $p<0.1$). The positive E index for the first hydrological year (Table 2) represents a slight improvement from the result obtained for ammonium simulations with the LU4-N model. A simple one-at-a-time perturbation sensitivity analysis highlighted that hillslope and riparian mineraliza-
 10 tion related parameters (K_{\min} and U_{\min}), maximum static storage water contents (H_u^*), riparian denitrification related parameters (K_{denitr} and U_{denitr}) and the maximum annual ammonium plant uptake ($\text{MaxUP}_{\text{NH}_4}$) had the major impact on the nitrate related ob-
 15 jective functions. Ammonium soil adsorption rate (K_{ads}) and the hillslope nitrification soil moisture threshold (U_{nitr}) were also highlighted as quite sensitive parameters considering the ammonium related objective functions.

4.3 SD4-R-N calibration and validation results

Observed nitrate and ammonium daily stream concentrations at Fuirosos and the cor-
 20 responding simulated ones, obtained with the SD4-R-N model structure, are shown Fig. 4c. The global discharge E index for the calibration period was 0.78, while for the first, second and third years respectively the E -index was 0.5, 0.4 and 0.86 (Table 2). The BE error was less than 8%. Concerning the nitrate simulation, the E index for the whole period was approximately 0.68 and the BE error less than -9% (Table 2).
 25 Interestingly, this model structure could improve the simulation of the discharge peak flow observed on March 2002 (Fig. 9), which corresponded with the highest nitrate con-
 30 centration peak observed during the calibration period (Fig. 4). This discharge event can be classified as “intermediate flow” ($0.05 \text{ m}^3 \text{ s}^{-1} \leq Q < 1 \text{ m}^3 \text{ s}^{-1}$) according to Medici et al. (2008), which means that interflow was likely to have contributed along with the

quick base flow. This suggestion is also supported by the slope steepness of the hydrograph recession. Neither the lumped LU4-N model nor the semi-distributed LU4-R-N model could reproduce this discharge event because no interflow was generated in that instance and the only flow contributing to the discharge was the quick base flow. This improvement was reflected by the SD4-N-R model's ability to simulate satisfactorily the corresponding nitrate peak concentration which resulted in an E index for the third year of 0.66 for the streamwater nitrate concentration simulations (Fig. 4c and Table 2).

Also in this case, the riparian zone was highlighted as a quite active zone where both the annual and daily M:N ratio were most of the time quite close to 1:1, as in the case of the LU4-R-N model (Fig. 6). The M:N ratio behaviour for the leucogranite and granodiorite units was quite similar to the one obtained with the LU4-R-N model for the so called hill-slope area, while in the sericitic schists unit the nitrification process could take place more easily than in the rest of the hill-slope giving in general smaller values for the M:N ratio (Figs. 5 and 10). The sericitic unit is mainly facing North and it is largely covered by a deciduous woodland (chestnut (*Castanea sativa*), hazel (*Corylus avellana*) and oak (*Quercus pubescens*) with well-developed litter layers which could bring about higher nitrification rates than in the granitic units.

Finally, concerning the ammonium daily concentrations, the SD4-R-N model could not reproduce satisfactorily the daily NH_4 concentration for the calibration period (Fig. 4c, Table 2). The temporal validation results for this model structure are shown in Fig. 4c and Table 3. The E index improved to 0.3, in spite the larger discharge BE error. Also in this case, a simple one-at-a-time perturbation sensitivity analysis highlighted in general the mineralization related parameters as the most sensitive, as well as the maximum static storage water contents of each HRUs (H_{ij}^*) and the annual maximum ammonium plant uptake ($\text{MaxUP}_{\text{NH}_4}$). Moreover, also the ammonium soil adsorption rate (K_{ads}) and both hillslope and riparian zone nitrification soil moisture threshold (U_{nitr}) were highlighted as quite influent parameters especially considering the ammonium related objective functions.

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

5 Discussion

The LU4-N model performance for the calibration period could be considered satisfactory in terms of daily nitrate concentration. However, the temporal validation process call for caution when considering the result obtained, even if just one year for the validation may not be sufficient to accept or reject a model conceptualization. Inspection of the validation results pointed out that the LU4-N model simulated adequately the discharge event observed during November 2002 (Medici et al., 2008), but was unable to reproduce the associated nitrate peak. In fact, there was insufficient nitrate left in soil to be washed into the stream by the interflow to create a peak in the streamwater nitrate concentrations. Lowering the nitrate plant uptake from 50 (day^{-1}) to 0.3 (day^{-1}), which would be the maximum rate allowed to increase stream nitrate concentration during the validation period, increased the *BE* error for the calibration period to approximately 169% without significantly improving the model validation performance (*E* remained negative and *BE* increased to 75%). Alternatively, the problem may be related to the nitrification dynamic; a more continuous nitrification process instead of a pulsed response could help to improve nitrate simulation during the validation period. However, problems arose when a permanent nitrification dynamic for the whole catchment was invoked. Specifically, it became impossible to simulate a M:N ratio consistent with the one observed by Serrasolses (1999), unless the nitrification rate was kept extremely low but this resulted in a failure to represent the observed nitrate peaks. Also when considering a high mineralization rate that caused extreme high stream ammonium concentration, the annual immobilization rate became largely beyond the range expected from literature values (i.e.: approximately $0.1 \text{ kg ha}^{-1} \text{ yr}^{-1}$ according to Bonilla, 1990, after Bernal et al., 2004). The impossibility of obtaining acceptable results with the LU4-N model for the validation process forced us to explore different model structures. To this end, several authors (Butturini et al., 2003; Bernal et al., 2007; McIntyre et al., 2009) noted the importance of the riparian zone as a “hot spot” for nitrate removal/production in Mediterranean catchments. It was also highlighted

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



that the mechanism of mineralization-nitrification can be essentially different from the rest of the catchment due to the specific moisture condition and different organic matter that can be found there. Therefore, it was thought the role played by the riparian zone should have been taken into account, even if it is well known that adding model components and parameters to reproduce specific aspects of catchment behaviour does not necessarily lead to better results. Therefore, the lumped LU4-N model was progressed to a semidistributed model that was applied considering firstly 2 HRUs (LU4-R-N) and then taking into account 4 HRUs (SD4-R-N), as previously explained.

According to the LU4-R-N and SD4-R-N models conceptualization, microbial processes in the hill-slope occur in pulses stimulated by soil moisture increasing after rain (Figs. 8 and 9), as it was for the whole catchment with the LU4-N model (Fig. 5). Namely, simulated nitrification, immobilisation and denitrification were allowed to occur only after exceeding their respective soil moisture thresholds (Table 1). This threshold mechanism gives rise in the hill-slope to pulses that are particularly significant for nitrification. The LU4-R-N and SD4-R-N models, due to the threshold mechanism, reproduced in the hill-slope soil an annual average M:N ratio of approximately 8:1, which is consistent with the ratio (10:1) founded in other Mediterranean areas (e.g., Serrasolses et al., 1999), which was explained considering soil moisture limitation of nitrification. Interestingly, when considering the riparian zone alone the simulated M:N ratio decreased in both cases to almost 1:1 (Fig. 6). Supporting our simulations, Merrill (2006) found out that measured net mineralisation and net nitrification rates were similar in riparian zone ecosystem types. Moreover, it was found that in four of the five ecosystems considered in the study by Merrill (2006), net mineralization rates explained over 60% of the variation in net nitrification. This specific behaviour of the riparian soil allowed to easily accumulating nitrate that could be washed away by the interflow derived from the hill-slope causing significant increase in nitrate streamwater concentrations. Butturini et al. (2003) previously pointed out the unsaturated riparian soil layer at Fuirosos as a possible source of nitrate. In this study, it was observed that the rise of the local riparian groundwater table, after the summer drought, resulted

in the rapid flushing of nitrate stored in the soil during the long dry period. Our results suggested also a higher mineralization rate in the riparian area than in the rest of the catchment. A possible explanation for that, may be that the major tree species at the hill-slope of Fuirosos are perennial cork oak (*Quercus suber*) and pine (*Pinus halepensis* and *Pinus pinaster*), therefore the mineralization rates are expected to be low as a consequence of allelopathic compounds leached from plants and the quality of sclerophyllous leaf (Gallardo and Merino, 1992; Castaldi et al., 2002). The stream channel is flanked by a well-developed riparian area where alder (*Alnus glutinosa*) – a tree species with high quality litter, and exotic plane tree (*Platanus acerifolia*) predominates, allowing for higher decomposition and mineralization rates of litter accumulated on the stream bed and stream edge zone. Moreover, Acuña et al. (2007) observed that in the Fuirosos stream, leaf fall may extend from late summer to autumn (August to November) during dry years, due to hydrologic stress. Therefore, large inputs of organic matter accumulate on the streambed and riparian zone may fuel heterotrophic activity during the transition and wet periods (Von Schiller et al., 2008). Simulated mineralization was highest immediately after the summer drought period, when the soil moisture content was approximately 50% or less of the maximum soil static water content. This is consistent with the study of McIntyre et al. (2009) which noted that, for a semiarid intermittent stream, mineralization would be reduced under soil moisture conditions closed to saturation, while it would increase under moderate saturation. Other authors observed a high rate of humus decomposition and rapid mineralization following rewetting of dry soils and it was also observed that soils subject to wetting and drying cycles, release more nitrogen than continuously moist soil (Birch, 1964; Dick et al., 2005; Rey et al., 2005). Bernal et al. (2005) observed, at Fuirosos, that mineralization activity existed in the mineral soil and/or in the stream channel particularly during the transition period from dry to wet conditions and in a previous study performed in the soil of the riparian area of Fuirosos, Bernal et al. (2003) reported the highest mineralization rates in autumn. Interestingly, the SD4-R-N model reproduced a huge pulse of nitrification in the riparian soil just after the summer drought 2001 because of a sud-

Modelling inorganic nitrogen behaviour in Mediterranean forested catchmentC. Medici et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

den increase in soil moisture content due to the reverse flux, which is characteristic of arid and semiarid areas (Fig. 9). This is consistent with Butturini et al. (2003) that pointed out the reverse flux as a possible mechanism responsible for nitrate release in the riparian zone.

All the model structures considered included denitrification and nitrification in the shallow aquifer. This was necessary to represent the nitrate behaviour. These processes controlled the rate of reduction in the streamwater nitrate and ammonium concentrations during base flow conditions. This is consistent with previous studies of biogeochemical activities in the unsaturated zone of weathered granite (Legout et al., 2005) which demonstrated potential for bacterial activity and biogeochemical reaction in the lower soil horizons associated with lower carbon content. In particular, Legout et al. (2005) suggested that both nitrification and denitrification are likely to take place in the unsaturated weathered granite below the soil organic horizon. The denitrification process occurring in the riparian groundwater was especially relevant for the SD4-R-N model (Fig. 11), while for the LU4-R-N model denitrification occurred mainly in the riparian upper soil (Fig. 8). In our model the riparian interflow eventually percolates to the local riparian aquifer due to the extremely low slope in this catchment area and thus, it is nearly impossible to distinguish between soil and aquifer riparian denitrification (Fig. 3). Nonetheless, our results highlighted that the denitrification process in the riparian zone is a key mechanism to the reduction of groundwater nitrate in particular during the wettest period of the year. This is consistent with previous studies in Mediterranean areas (Peterjohn and Correl, 1984; Butturini et al., 2003; Rassam et al., 2006 and Bernal et al., 2007). Finally, none of the considered models could reproduce satisfactorily the daily stream ammonium concentration, which was low even during precipitation events. Nevertheless, they could represent at least the ammonium general trend and order of magnitude, which taking into account its erratic behaviour it can be considered an acceptable result. In particular, differently from the INCA-N model, they did not simulate ammonium leaching during storm flow because we included the adsorption/desorption mechanism in the soil compartment.

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

6 Conclusions

The aim of this study was to improve our understanding of the main processes that govern the inorganic nitrogen fate and losses in Mediterranean catchments by means of mathematical modelling. The results highlighted that in those ecosystems a pulse dynamic for most of the soil biological processes, related with the rainfall pattern occurs as previously suggested by Schiwinning (2004b). We reproduced this pulse dynamic by introducing a moisture threshold for each simulated soil-biological process. The concept of response thresholds is recurrent in the ecology of arid/semi-arid systems (Beatty 1974), and it has been used to explain the decoupling of nutrient gain and losses mechanisms (Schwinning et al., 2004). Our simulations suggested that nitrification shows a pulse dynamic in the hillslope soil, while it occurs more continuously in the riparian soil, which together with the interflow flushing effect can give rise to important stream nitrate concentration peaks during some periods of the year. These results point towards the riparian upper soil as a possible source of nitrate in this type of ecosystems, consistently with that observed in previous empirical studies (e.g., Butturini et al., 2003). Interestingly, the model reproduced by means of calibration the so-called “Birch effect”, which implies higher mineralization rate just after the summer drought. Finally, the results indicate the importance of the nitrification and denitrification processes in the unsaturated weathered granite below the soil organic horizon. The LU4-R-N and the SD4-R-N semi-distributed models could be calibrated to simulate flow and nitrate dynamic in Fuirosos and gave acceptable result for the temporal validation process. This suggests that the key processes controlling flow and nitrate behaviour are included within these models conceptual schemes and their mathematical representation seems reasonable. Thus, we believe that the conceptual models applied in the present study represent useful tools to investigate the inorganic nitrogen behaviour not just at Fuirosos. In fact, they can be easily generalized and the key mechanisms highlighted in this research could be by far extrapolated to other catchments.

Acknowledgements. This study was supported by a Formación de Personal Investigador (FPI)

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



grant and funds provided by the Generalitat Valenciana, Conselleria d'Educació and by the Ministerio de Educación Español (MEC) (projects: GCL2007-60144/HID; CGL2005-06219/HID CGL2008-03076-E/BTE).

References

- 5 Acuña, V., Giorgi, A., Muñoz, I., Sabater, F., and Sabater, S.: Meteorological and riparian influences on organic matter dynamics in a forested Mediterranean stream, *J. N. Am. Benthol. Soc.*, 26, 54–69, 2007.
- Arheimer, B., Andersson, L., and Lepistö, A.: Variation of nitrogen concentration in forest streams, Influences of flow, seasonality and catchment characteristics, *J. Hydrol.*, 179, 281–304, 1996.
- 10 Austin, A. T., Yahdjian, L., Stark, J. M., Belnap, J., Porporato, A., Norton, U., Ravetta, D. A., and Schaeffer, S. M.: Water pulses and biogeochemical cycles in arid and semiarid ecosystems, *Oecologia*, 141, 221–235, 2004.
- Ávila, A., Bonilla, D., Rodá, F., Piñol, J., and Neal, C.: Soil water chemistry in a holm oak (*Quercus ilex*) forest: interferences on biogeochemical processes for a montane Mediterranean area, *J. Hydrol.*, 166, 15–35, 1995.
- 15 Beatley, J. C.: Phenological events and their environmental triggers in Mojave-Desert, *Ecosyst. Ecol.*, 55, 856–863, 1974.
- Bernal, S., Butturini, A., Nin, E., Sabater, F., and Sabater, S.: Leaf litter dynamics and nitrous oxide emission in a Mediterranean riparian forest: implications for soil nitrogen dynamics. *J. Environ. Qual.*, 32, 191–197, 2003.
- Bernal, S., Butturini, A., Riera, J. L., Vázquez, E., and Sabater, F.: Calibration of the INCA model in a Mediterranean forested catchment: the effect of hydrological inter-annual variability in an intermittent stream, *Hydrol. Earth Syst. Sci.*, 8, 729–741, 2004,
- 20 <http://www.hydrol-earth-syst-sci.net/8/729/2004/>.
- Bernal, S., Butturini, A., and Sabater, F.: Seasonal variation of dissolved nitrogen and DOC:DON ratios in an intermittent Mediterranean stream, *Biogeochemistry*, 75, 351–372, 2005.
- 25 Bernal, S., Sabater, F., Butturini, A., Nin, E., and Sabater, S.: Factors limiting denitrification in a Mediterranean riparian forest, *Soil Biol. Biochem.*, 39, 2685–2688, 2007.
- 30

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Birch, H. F.: Further observations on humus decomposition and nitrification, *Plant Soil*, 11, 262–286, 1959.
- Birch, H. F.: Nitrification in soil after different period of dryness, *Plant Soil*, 12, 81–96, 1960.
- Birch, H. F.: Mineralization of plant nitrogen following alternate wet and dry conditions, *Plant Soil*, 12, 81–96, 1964.
- Butturini, A., Bernal, S., Nin, E., Hellin, C., Rivero, L., Sabater, S., and Sabater, F.: Influence of the stream groundwater hydrology on nitrate concentration in unsaturated riparian area bounded by an intermittent Mediterranean stream, *Water Resour. Res.*, 39(4), 1110, doi:10.1029/2001WR001260, 2003.
- Castaldi, S. and Aragosa, D.: Factors influencing nitrification and denitrification variability in a natural and fire-disturbed Mediterranean shrubland, *Biol. Fert. Soils*, 36, 418–425, 2002.
- Chapman, P. J., Edwards, A. C., and Cresser, M. S.: The nitrogen composition of streams in upland Scotland: some regional seasonal differences. *Sci. Total Environ.*, 265, 65–83, 2001.
- Chu, Y., Salles, C., Cernesson, F., Perrin, J. L., and Tournoud, M. G.: Nutrient load modelling during floods in intermittent rivers: An operational approach, *Environ. Modell. Softw.*, 23, 768–781, 2008.
- Dick, J., Skiba, U., Munro, R., and Deans, D.: Effect of N-fixing trees and crops on NO and N₂O emissions from Senegal soils, *J. Biogeogr.*, 33, 416–423, 2005.
- Gallardo, A. and Merino, J.: Nitrogen immobilization in leaf litter at two Mediterranean ecosystems of SW Spain, *Soil Biol. Biochem.*, 30, 1349–1358, 1992.
- Gallart, F., Llorens, P., Latron, J., and Regúes, D.: Hydrological processes and their seasonal controls in a small Mediterranean mountain catchment in the Pyrenees, *Hydrol. Earth Syst. Sci.*, 6, 527–537, 2002, <http://www.hydrol-earth-syst-sci.net/6/527/2002/>.
- Gelfand, I. and Yakir, D.: Influence of nitrite accumulation in association with seasonal patterns and mineralization of soil nitrogen in a semi-arid pine forest, *Soil Biol. Biochem.*, 40(2), 415–424, 2008.
- Green, P. A., Vörösmarty, C. J., Meybeck, M., Galloway, J. N., Peterson, B. J., and Boyer E. W.: Pre-industrial and contemporary fluxes through rivers: a global assessment based on typology, *Biogeochemistry*, 68, 71–105, 2004.
- Heckathorn, S. A. and Delucia, E. H.: Ammonia volatilization during drought in perennial C₄ grasses of tallgrass prairie, *Oecologia*, 101, 361–365, 1995.
- Hedin, L. O., Armesto, J. J., and Johnson, A. H.: Patterns of nutrients loss from unpolluted, old-growth temperature forests, *Evaluation of biogeochemical theory, Ecology*, 76, 493–509,

Modelling inorganic nitrogen behaviour in Mediterranean forested catchmentC. Medici et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

1995.

Jarvis, P., Rey, A., Petsikos, C., Wingte, L., Rayment, M., Pereira, J., Banza, J., David, J., Miglietta, F., Borghetti, M., Manca, G., and Valentini, R.: Drying and wetting of Mediterranean soils stimulates decomposition and carbon dioxide emission: the “Birch effect”, *Tree Physiol.*, 27, 929–940, 2007.

Legout, C., Molenat, J., Lefebvre, S., Marmonier, P., and Aquilina, L.: Investigation of biogeochemical activities in the soil and unsaturated zone of weathered granite, *Biogeochemistry*, 75, 329–350, 2005.

McIntyre, R., Adams, M., Ford, D., and Grierson, F.: Rewetting and litter addition influence mineralisation and microbial communities in soils from a semi-arid intermittent stream, *Soil Biol. Biochem.*, 41, 92–101, 2009.

Medici, C., Butturini, A., Bernal, S., Vázquez, E., Sabater, F., Vélez, J. I. and Francés, F.: Modelling the non-linear hydrological behaviour of a small Mediterranean forested catchment, *Hydrol. Process.*, 22, 3814–3828, 2008.

Merrill, A. G. and Benning, T. L.: Ecosystem type differences in nitrogen process rates and controls in the riparian zone of a montane landscape, *Forest Ecol. Manag.*, 222, 145–161, 2006.

Mummey, D. L., Smith, J. L., and Bolton, H. J.: Nitrous oxide flux from a shrub-steppe ecosystem: sources and regulation, *Soil Biol. Biochem.*, 26, 279–286, 1994.

Ocampo, C. J., Oldham, C. E., and Sivapalan, M.: Nitrate attenuation in agricultural catchments: shifting balances between transport and reaction, *Water Resour. Res.*, 42, W01408, doi:10.1029/2004WR003773, 2006.

Peterjohn, W. T. and Schlesinger, W. H.: Factors controlling denitrification in a Chihuahuan desert ecosystem, *Soil Sci. Soc. Am. J.*, 55, 1694–1701, 1991.

Peterjohn, W. T. and Correll, D. L.: Nutrient dynamics in an agricultural watershed: observations on the role of the riparian forest, *Ecology*, 65, 1466–1475, 1984.

Rassam, D. W., Fellows, C. S., De Hayr, R., Hunter, H., and Bloesch, P.: The hydrology of riparian buffer zones: two case studies in an ephemeral and perennial stream, *J. Hydrol.*, 325, 308–324, 2006.

Rey, A., Petsikos, C., Jarvis, P. G., and Grace, J.: Effect of temperature and moisture on rates of carbon mineralization in a Mediterranean oak forest soil under controlled and field conditions, *Eur. J. Soil Sci.*, 56, 589–599, 2005.

Reynolds, J. F., Kemp, P. R., Ogle, K., and Fernández, R. J.: Modifying the “pulse-reserve”

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



paradigm for deserts of North America: precipitation pulses, soil water, and plant responses, *Oecologia*, 141, 194–210, 2004.

Schlesinger, W. H., Reckhow, K. H., and Bernhardt, E. S.: Global Change: The nitrogen cycle and rivers, *Water Resour. Res.*, 42, W03S06, doi:10.1029/2005WR004300, 2006.

5 Schwinning, S. and Sala, E. O.: Hierarchy of responses to resource pulses in arid and semi-arid ecosystems, *Oecologia*, 141, 211–220, 2004a.

Schwinning, S., Sala, O. E., Loik, M. E., and Ehleringer, J. R.: Thresholds, memory, and seasonality: understanding pulse dynamics in arid/semi-arid ecosystems, *Oecologia*, 141, 191–193, 2004b.

10 Serrasolses, I., Diego V., and Bomilla, D.: Soil nitrogen dynamics, in: *Ecological of Mediterranean evergreen oak forests*, edited by: Roda, F., *Ecological Studies* 137, Springer, Berlin, Germany, 1999.

Stark, J. M., Smart, D. M., Hart, S. C., and Haubensak, K. A.: Regulation of nitrite oxide emissions from forest and rangeland soils of western North America, *Ecology*, 83, 2278–2292, 2002.

15 Stieglitz, M., Shaman, J., McNamara, J., Engel, V., Shanley, J., and Kling, G. W.: An approach to understanding hydrological connectivity on the hillslope and the implications for nutrient transport, *Global Biogeochem. Cy.*, 17, 1105, doi:10.1029/2003GB002041, 2003.

Van Gestel, M., Merckx, R., and Vlassak, K.: Microbial biomass responses to oil drying and rewetting: the fate of fast- and slow-growing microorganisms in soils from different climates, *Soil Biol. Biochem.*, 25, 109–123, 1993.

20 Vitousek, P. M., Naylor, R., Crews, T., David, M. B., Drinkwater, L. E., Holland, E., Johnes, P. J., Katzenberger, J., Martinelli, L. A., Matson, P. A., Nziyueheba, G., Ojima, D., Palm, C. A., Robertson, G. P., Sanchez, P. A., Townsend, A. R., and Zhang, F. S.: Nutrient imbalances in Agricultural Development, *Science*, 324, 5937, 1519–1520, 2009.

Von Schiller, D., Martí, E., Riera, J. L., Ribot, M., Argerich, A., Fonollá, P., and Sabater, F.: Inter-annual, annual and seasonal variation of P and N retention in a perennial and an intermittent stream, *Ecosystems*, 11, 670–687, doi:10.1007/s10021-008-9150-3, 2008.

25 Wade, A. J., Durand, P., Beaujouan, V., Wessel, W. W., Raat, K. J., Whitehead, P. G., Butterfield, D., Rankinen, K., and Lepisto, A.: A nitrogen model for European catchments: INCA, new model structure and equations, *Hydrol. Earth Syst. Sci.*, 6, 559–582, 2002, <http://www.hydrol-earth-syst-sci.net/6/559/2002/>.

Wade, A. J., Neal, C., Butterfield, D., and Futter, M. N.: Assessing nitrogen dynamics in Euro-

pean ecosystems, integrating measurement and modelling: conclusions, Hydrol. Earth Syst. Sci., 8, 846–857, 2004, <http://www.hydrol-earth-syst-sci.net/8/846/2004/>.

Whitehead, P. G., Wilson, P. G., and Butterfield, D.: A semi-distributed Nitrogen Model for Multiple Source Assessments in Catchments (INCA): Part 1 – Model structure and Process Equations, Sci. Total Environ., 201/211, 547–558, 1998.

HESSD

6, 5665–5703, 2009

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.

Table 1. Parameters considered in each of the three structures (LU4-N, LU4-R-N and SD4-R-N) and their effective values after calibration process.

Parameters	Description	LU4-N	LU4-R-N		SD4-R-N			
		Basin	Hill-slope	Rip. Z	Leucogr.	Granod.	Schist	Rip. Z
<i>Nitrogen model calibrated parameters</i>								
1	K_{\min}	Hillslope mineralization rate [$\text{Kg ha}^{-1} \text{day}^{-1}$]	0.51	0.51	3.5	0.5		3.3
2	K_{nitr}	Nitrification rate [day^{-1}]	1.0	0.6	1.0	1.0		2.2
3	K_{denitr}	Denitrification rate [day^{-1}]	0.1	0.08	1.8	0.04		1.17
4	K_{imm}	Immobilization rate [day^{-1}]	0.15	0.1	0.01	0.34		0.53
5	$K_{\text{up NO}_3}$	Nitrate plant uptake rate [day^{-1}]	50	2.04	2.04	44.15		63.04
6	$K_{\text{up NH}_4}$	Ammonium plant uptake rate [day^{-1}]	50	4.39	4.39	8.28		77.34
7	$K_{\text{denitr.aquif}}$	Shallow aquifer denitrification rate [day^{-1}]	0.06	0.06	0.03	0.22		0.11
8	$K_{\text{nitr.aquif}}$	Shallow aquifer nitrification rate [day^{-1}]	1.84	1.84	1.97	0.97		0.18
9	K_{ads}	Ammonium soil adsorption rate [day^{-1}]	0.88		0.88		0.82	
10	K_{des}	Ammonium soil desorption rate [day^{-1}]	0.05		0.05		0.5	
11	U_{\min}	Mineralization soil moisture threshold (% Hu^*)	48.2	48.2	36.8	56.0		22.9
12	U_{nitr}	Nitrification soil moisture threshold (% Hu^*)	57.2	57.2	34.0	63.0		34.0
13	U_{denitr}	Denitrification soil moisture threshold (% Hu^*)	89.7	78.6	67.0	85.0		93.7
14	U_{immob}	Immobilization soil moisture threshold (% Hu^*)	41.6	41.6	68.8	92.0		94.0
15	C_g	Maximum temperature difference ($^{\circ}\text{C}$)	6.15		6.15		6.15	
16	$\text{MaxAds}_{\text{NH}_4}$	Daily max. NH_4 adsorption [$\text{kg day}^{-1} \text{km}^{-2}$]	14.5		14.5		36.14	
17	$\text{MaxUP}_{\text{NH}_4}$	Annual max. NH_4 uptake [$\text{Kg ha}^{-1} \text{day}^{-1}$]	90.1		90.1		97.9	
18	$\text{MaxUP}_{\text{NO}_3}$ (1)	Annual max. NO_3 uptake [$\text{Kg ha}^{-1} \text{day}^{-1}$] (December, January and February)	21.6		21.6		18.5	
19	$\text{MaxUP}_{\text{NO}_3}$ (2)	Annual max. NO_3 uptake [$\text{Kg ha}^{-1} \text{day}^{-1}$] (Rest of the year)	118.0		118.0		54.44	

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.

Table 2. Calibration goodness of fit indexes (from 13 October 1999 to 22 August 2002): the global and annual Nash index (E ; where $E=1$ is the optimum); the global balance volume errors (BE); the coefficient of determination (r^2) (only shown when $p<0.01$) and the relative Root mean square error (Relative RMSE; where RMSE=0 is the optimum).

Index	LU4-N			LU4-R-N			SD4-R-N		
	Q	NO_3	NH_4	Q	NO_3	NH_4	Q	NO_3	NH_4
E_{TOT}	0.71	0.46	<0	0.70	0.56	<0	0.78	0.68	<0
$E_{1\text{yr}}$	0.62	<0	<0	0.50	0.45	0.2	0.52	0.49	0.35
$E_{2\text{yr}}$	0.62	0.47	<0	0.61	0.43	<0	0.43	0.57	<0
$E_{3\text{yr}}$	0.74	0.43	<0	0.72	0.48	<0	0.86	0.66	<0
$BE_{\text{TOT}}\%$	-1.24	-21.95	-30.2	-3.07	-0.48	-14.9	7.52	-8.22	21.03
r^2	0.77	0.5	n.s.	0.78	0.58	0.02	0.78	0.69	0.05
Relative RMSE	0.53	0.56	1.25	0.55	0.50	0.92	0.47	0.43	0.90

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.

Table 3. Validation goodness of fit indexes (from 1 August 2002 to 30 June 2003): the global and annual Nash index (E ; where $E=1$ is the optimum); the global balance volume errors (BE); the coefficient of determination (r^2) (only shown when $p<0.01$) and the relative Root mean square error (Relative RMSE; where RMSE=0 is the optimum).

Index	LU4-N		LU4R-N		SD4R-N	
	Q	NO_3	Q	NO_3	Q	NO_3
E_{TOT}	0.48	<0	0.61	0.13	0.5	0.32
BE_{TOT} %	28.37	-30.2	24.17	27.85	47	7.12
r^2	0.78	0.2	0.77	0.35	0.74	0.4
Relative RMSE	0.69	0.96	0.60	0.60	0.68	0.53

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.

Table 4. Nitrogen annual process rates.

N Processes	Measured values [Kg ha ⁻¹ day ⁻¹]*	Sim. values [Kg ha ⁻¹ day ⁻¹] LU4-N	Sim. values [Kg ha ⁻¹ day ⁻¹] LU4R-N	Sim. values [Kg ha ⁻¹ day ⁻¹] SD4R-N
Net mineralization	32.4–80.1	62.9	64.18	61.94
Net nitrification	4.4–7.5	6.19	7.83	8.84
Immobilization	0.08	4.8	4.52	0.08
Nitrate uptake by vegetation	10.3–58	13.42	13.51	14.94
Ammonium uptake by vegetation	53–80.5	59.17	58.79	60.67

* After Bernal et al. (2004).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.




Fig. 1. Geographical location of the Fuirosos catchment (Catalonia, NE Spain). Lithological units are shown in different shadings. The little triangles represent the four small reservoirs present in the catchment.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.





Fig. 2. Nitrogen cycle in the soil and aquifers systems for the LU4-N model (modified from Whitehead et al., 1998).

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

a. Hillslope area draining to the small reservoirs



b. Hillslope area draining through the riparian zone

c. Riparian zone




Fig. 3. LU4-R-N and SD4-R-N conceptual scheme, where **(a)** represents the part of the catchment that drains to the four small reservoirs located at the catchment; **(b)** represents the part of the catchment that drains through the riparian zone before reaching the stream channel and finally **(c)** represents the riparian zone which presents a bidirectional flow with the channel.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	




Fig. 4. Simulated and observed nitrate and ammonium (gr m^{-3}) for the calibration period (1999–2002) with **(a)** LU4-N; **(b)** LU4-R-N (with 2 HRUs) and **(c)** SD4-R-N (with 4 HRUs).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.




Fig. 5. Simulated soil moisture content (H_1) and nitrification soil moisture threshold (U_{nitr}) in mm, plus mineralization, nitrification and denitrification processes (kg) for the calibration period (1999–2002) with the LU4-N model.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.




Fig. 6. Simulated Mineralization:Nitrification ratio (M:N) variation according to the different models structures and to each HRU considered.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	




Fig. 7. Simulated and observed stream nitrate concentration (gr m^{-3}) for the validation period (1999–2002) with **(a)** LU4-N; **(b)** LU4R-N (with 2 HRUs) and **(c)** SD4R-N (with 4 HRUs).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.




Fig. 8. Simulated soil moisture content (H_1), nitrification soil moisture threshold (U_{nitr}) and denitrification soil moisture threshold (U_{denitr}) in mm, plus simulated mineralization, nitrification and denitrification processes (kg) for the riparian zone (calibration period 1999–2002) with the LU4R-N model.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.





Fig. 9. Simulated and observed discharges ($\text{m}^3 \text{s}^{-1}$) for the event of March 2002 obtained with: (a) the LU4-R-N model and (b) the SD4-R-N model.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.




Fig. 10. Simulated soil moisture content (H_1), nitrification soil moisture threshold (U_{nitr}) and denitrification soil moisture threshold (U_{denitr}) in mm, plus simulated mineralization, nitrification and denitrification processes (kg km^{-2}) for the riparian zone (calibration period 1999–2002) with the LU4R-N model.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Modelling inorganic nitrogen behaviour in Mediterranean forested catchment

C. Medici et al.




Fig. 11. Simulated denitrification process (kg) in the riparian local aquifer with the LU4-R-N and SD4-R-N models.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion