

Papers published in *Hydrology and Earth System Sciences Discussions* are under open-access review for the journal *Hydrology and Earth System Sciences*

Estimating surface fluxes over middle and upper streams of the Heihe River Basin with ASTER imagery

W. Ma¹, Y. Ma^{2,1}, Z. Hu¹, B. Su³, J. Wang¹, and H. Ishikawa⁴

¹Laboratory for Climate Environment and Disasters of Western China, Cold and Arid Regions Environmental and Engineering Research Institute, Chinese Academy of Sciences, Lanzhou, Gansu 730000, China

²Laboratory of Tibetan Environment Changes and Land Surface Processes, Institute of Tibetan Plateau Research, Chinese Academy of Sciences, Beijing, China

³International Institute for Geo-Information Science and Earth Observation, Enschede, The Netherlands

⁴Disaster Prevention Research Institute, Kyoto University, Kyoto, Japan

Received: 9 June 2009 – Accepted: 15 June 2009 – Published: 29 June 2009

Correspondence to: W. Ma (wqma@lzb.ac.cn)

Published by Copernicus Publications on behalf of the European Geosciences Union.

HESSD

6, 4619–4635, 2009

**Estimating surface
fluxes over the Heihe
River Basin with
ASTER imagery**

W. Ma et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

Surface fluxes are important boundary conditions for climatological modeling and the Asian monsoon system. Recent availability of high-resolution, multi-band imagery from the ASTER (Advanced Space-borne Thermal Emission and Reflection Radiometer) sensor has enabled us to estimate surface fluxes to bridge the gap between local scale flux measurements using micrometeorological instruments and regional scale land-atmosphere exchanges of water and heat fluxes that are fundamental for the understanding of the water cycle in the Asian monsoon system. A Surface Energy Balance System (SEBS) method based on ASTER data and field observations has been proposed and tested for deriving net radiation flux (R_n), soil heat flux (G_0), sensible heat flux (H) and latent heat flux (λE) over heterogeneous land surface in this paper. As a case study, the methodology was applied to the experimental area of the WATER (Watershed Allied Telemetry Experimental Research), located at the mid-to-upstream sections of the Heihe River, northwest China. The ASTER data of 3 May and 4 June in 2008 was used in this paper for the case of mid-to-upstream sections of the Heihe River Basin. To validate the proposed methodology, the ground-measured land surface heat fluxes (net radiation flux (R_n), soil heat flux (G_0), sensible heat flux (H) and latent heat flux (λE)) were compared to the ASTER derived values. The results show that the derived surface variables and land surface heat fluxes in different months over the study area are in good accordance with the land surface status. It is therefore concluded that the proposed methodology is successful for the retrieval of land surface heat fluxes using the ASTER data and field observation over the study area.

1 Introduction

Among many land surface experiments having been carried out so far, arid and cold regions were paid little attention. The land surface observations in arid and cold regions, both remotely sensed and in-situ, need to be strengthened for a better understand-

HESSD

6, 4619–4635, 2009

Estimating surface fluxes over the Heihe River Basin with ASTER imagery

W. Ma et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

ing of hydrological and ecological processes at different scales. The Watershed Allied Telemetry Experimental Research (WATER) is a simultaneous air-borne, satellite-borne, and ground-based remote sensing experiment conducted in the Heihe Basin, the second largest inland river basin in the northwest arid regions of China. The WATER is aiming at the research on water cycles, eco-hydrological and other land surface processes in catchment-scale. Data sets with high-resolution and spatiotemporal consistency will be generated based on this experiment. An integrated watershed model and a catchment-scale land/hydrological data assimilation system is proposed to be developed. The mission of WATER is to improve the observability, understanding, and predictability of hydrological and related ecological processes at catchmental scale, accumulate basic data for the development of watershed science and promote the applicability of quantitative remote sensing in watershed science studies (Li, 2008).

Remote sensing offers the possibility to derive regional distribution of land surface heat fluxes over heterogeneous land surface in combination with sparse field experimental stations. Remote sensing data provided by satellites are a means of obtaining consistent and frequent observations of spectral albedo and emittance of radiation at elements in a patch landscape and on a global scale (Sellers et al., 1990). The land surface variables and vegetation variables, such as surface temperature T_{sfc} , surface hemispherical albedo r_0 , NDVI, MSAVI, LAI and surface thermal emissivity ε can be derived directly from satellite observations (e.g., Susskind et al., 1984; Che'din et al., 1985; Tucker, 1986; Wan and Dozier, 1989; Menenti et al., 1989; Becker and Li, 1990, 1995; Watson et al., 1990; Baret and Guyot, 1997; Price, 1992; Kahle and Alley, 1992; Li and Becker, 1993; Qi et al., 1994; Norman et al., 1995; Schmugge et al., 1995; Kustas and Norman, 1997; Sobrino and Raissouni, 2000; Su, 2002; Ma et al., 2003a, b; Oku and Ishikawa, 2004; Kato, 2005; Ma, 2006b, 2007, 2009). The regional heat fluxes can be determined indirectly with the aid of these land surface variables and vegetation variables (Pinker, 1990).

Studies have explored several approaches to estimate the regional distribution of surface heat fluxes in recent years. These methods require specification of the vertical

Estimating surface fluxes over the Heihe River Basin with ASTER imagery

W. Ma et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

temperature difference between the surface temperature and the air temperature and an exchange resistance (e.g., Kustas et al., 1989; Kustas, 1990; Wang et al., 1995; Menenti et al., 1991; Menenti and Choudhury, 1993; Bastiaanssen, 1995; Kustas and Norman, 1997; Su, 2002). However, these remote sensing retrieval methods have been performed in homogeneous moist or semiarid regions, and investigations in heterogeneous landscape of arid and cold regions (e.g., the WATER area) are rare.

NOAA/AVHRR, GMS and Landsat-7 ETM data were used to determine regional land surface heat fluxes over heterogeneous landscape of the Tibetan Plateau (Ma et al., 2003a, b, 2005, 2006; Oku et al., 2007). However, the resolution of the NOAA/AVHRR and GMS data is about 1 km×1 km and sub-pixel heterogeneity has been omitted. So have Landsat-7 ETM data. The aim of this research is to upscale in-situ point observations of land surface variables and land surface heat fluxes to the regional scale using high-resolution (15 m×15 m) ASTER data.

2 Data and methodology

2.1 Data

The recent availability of high-resolution, multi-band imagery from the ASTER sensor has enabled us to estimate surface fluxes. ASTER covers a wide spectral region with 14 bands from the visible to the thermal infrared with high spatial, spectral and radiometric resolution. The spatial resolution varies with wavelength: 15 m in the visible and near-infrared (VNIR, 0.52–0.86 μm), 30 m in the short wave infrared (SWIR, 1.6–2.43 μm), and 90 m in the thermal infrared (TIR, 8.1–11.6 μm) (Yamaguchi, 1998).

The most relevant data, collected at the WATER (Fig. 1) surface stations (sites) to support the parameterization of land surface heat fluxes and analysis of ASTER images in this paper, consist of surface radiation budget components, surface radiation temperature, surface albedo, humidity, wind speed and direction measured at the Atmospheric Boundary Layer (ABL) towers, Automatic Weather Stations (AWSs), turbu-

Estimating surface fluxes over the Heihe River Basin with ASTER imagery

W. Ma et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

lent fluxes measured by eddy correlation technique, soil heat flux, soil temperature profiles, soil moisture profiles, and the vegetation state.

2.2 Theory and scheme

A Surface Energy Balance System (SEBS, Su, 2002) is proposed for the estimation of atmospheric turbulent fluxes and evaporative fraction using satellite earth observation data, in combination with meteorological information at proper scales. SEBS consists of: a set of tools for the determination of the land surface physical parameters, such as albedo, emissivity, temperature, vegetation coverage etc., from spectral reflectance and radiance measurements; a model for the determination of the roughness length for heat transfer (Su, 2002).

In this study, the Surface Energy Balance System (SEBS) retrieval algorithm is used for the ASTER data (Su, 2002). The general concept of the methodology is shown in a diagram (Fig. 2). The surface albedo for shortwave radiation (r_0) is retrieved from narrowband-broadband conversion by Liang (Liang, 2001). The land surface temperature (T_{sfc}) is derived using a method developed by Juan (2006) from multispectral thermal infrared data. Juan (2006) also evaluated a technique to extract emissivity information from multispectral thermal infrared data adding vegetation information. The radiative transfer model SMAC (Rahman and Dedieu, 1994) computes the downward shortwave and longwave radiation at the surface. With these results the surface net radiation flux (R_n) is determined. On the basis of the field observations, the soil heat flux (G_0) is estimated from the net radiation flux (R_n). The sensible heat flux (H) is estimated from T_{sfc} , and regional latent heat flux (λE) is derived as the residual of the energy budget theorem (Liou, 2004; Ma, 2006) for land surface.

The net radiation flux R_n is estimated as

$$\begin{aligned} R_n(x, y) &= K_{\downarrow}(x, y) - K_{\uparrow}(x, y) + L_{\downarrow}(x, y) - L_{\uparrow}(x, y) \\ &= (1 - r_0(x, y)) \cdot K_{\downarrow}(x, y) + L_{\downarrow}(x, y) - \varepsilon_0(x, y) \sigma T_{sfc}^4(x, y) \end{aligned} \quad (1)$$

Estimating surface fluxes over the Heihe River Basin with ASTER imagery

W. Ma et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Estimating surface fluxes over the Heihe River Basin with ASTER imagery

W. Ma et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where $\varepsilon_0(x, y)$ is surface emissivity, K_{\downarrow} (Wm^{-2}) represents the shortwave ($0.3\text{--}3\ \mu\text{m}$) and L_{\downarrow} (Wm^{-2}) is the longwave ($3\text{--}100\ \mu\text{m}$) radiation components, respectively. Surface albedo $r_0(x, y)$ is derived from narrowband-broadband conversion method by Liang (2001). Since ASTER has nine bands, it is expected that so many bands should enable us to convert narrowband to broadband albedos effectively. Liang (2001) found that the conversions are quite linear. The resultant linear equations are collated in the following.

$$r_0 = 0.484\alpha_1 + 0.335\alpha_3 - 0.324\alpha_5 + 0.551\alpha_6 + 0.305\alpha_8 - 0.367\alpha_9 - 0.0015 \quad (2)$$

Where i ($i=1\text{--}9$) are the correspondent ASTER band surface reflectance.

The equation to calculate soil heat flux is parameterized as (Su, 2002):

$$G_0 = R_n[\Gamma_c + (1 - f_c) \cdot (\Gamma_s - \Gamma_c)] \quad (3)$$

in which it is assumed that the ratio of soil heat flux to net radiation $\Gamma_c=0.05$ for full vegetation canopy (Monteith, 1973) and $\Gamma_s=0.315$ for bare soil (Kustas and Daughtry, 1989). An interpolation is then performed between these limiting cases using the fractional canopy coverage, f_c .

In order to derive the sensible and latent heat flux, the similarity theory will be made use of. In ASL (Atmospheric Surface Layer), the similarity relationships for the profiles of the mean wind speed, u , and the mean temperature, $\theta_0 - \theta_a$, are usually written in integral form as

$$u = \frac{u_*}{k} \left[\ln \left(\frac{z - d_0}{z_{0m}} \right) - \Psi_m \left(\frac{z - d_0}{L} \right) + \Psi_m \left(\frac{z_{0m}}{L} \right) \right] \quad (4)$$

$$\theta_0 - \theta_a = \frac{H}{ku_*\rho C_p} \left[\ln \left(\frac{z - d_0}{z_{0h}} \right) - \Psi_h \left(\frac{z - d_0}{L} \right) + \Psi_h \left(\frac{z_{0h}}{L} \right) \right] \quad (5)$$

where z is the height above the surface, $u_*=(\tau_0/\rho)^{1/2}$ is the friction velocity, τ_0 is the surface shear stress, ρ is the density of air, $k=0.4$ is von Karman's constant, d_0 is the

Estimating surface fluxes over the Heihe River Basin with ASTER imagery

W. Ma et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



zero plane displacement height, z_{0m} is the roughness height for momentum transfer, θ_0 is the potential temperature at the surface, θ_a is the potential air temperature at height z , z_{0h} is the scalar roughness height for heat transfer, ψ_m and ψ_h are the stability correction functions for momentum and sensible heat transfer respectively, and L is the

5 Obukhov length defined as:

$$L = \frac{\rho C_p u_*^3 \theta_v}{kgH} \quad (6)$$

where g is the acceleration due to gravity and θ_v is the potential virtual temperature near the surface.

The latent heat flux λE is the residual resulting from an application of the energy budget theorem to the land surface (Ma, 2006):

$$\lambda E = R_n - H - G_0 \quad (7)$$

3 Case studies and validation

As a case study, 4 scenes of ASTER data over the mid-to-upstream sections of the Heihe River Basin are used. Figure 3 shows the distribution maps of surface heat fluxes around the WATER area. Figure 4 shows the validation of the derived net radiation R_n , soil heat flux G_0 , sensible heat flux H and latent heat flux λE against ground measurements over the Watershed Airborne Telemetry Experimental Research (WATER) stations, Yingke, Huazhaizi, Guantan, Maliantan, A'rou, Binggou and Yakou with 1:1 line.

20 The results show the following:

1. The derived surface heat fluxes (net radiation flux R_n , soil heat flux G_0 , sensible heat flux H and latent heat flux λE) in different months over the study area are in good accordance with the land surface status. The experimental area includes a variety of land surfaces, such as a large area of grassy marshland, some

Estimating surface fluxes over the Heihe River Basin with ASTER imagery

W. Ma et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



desertification grassland areas, many small rivers; therefore these derived parameters show a wide range due to the strong contrast of surface features. Net radiation flux changed from 410 Wm^{-2} to 830 Wm^{-2} in May and from 410 Wm^{-2} to 870 Wm^{-2} in June. Soil heat flux varied from 30 Wm^{-2} to 260 Wm^{-2} in May and from 25 Wm^{-2} to 270 Wm^{-2} in June. Sensible heat flux is from 30 Wm^{-2} to 480 Wm^{-2} in May and from 10 Wm^{-2} to 470 Wm^{-2} in June, and latent heat flux varied from 60 Wm^{-2} to 430 Wm^{-2} in May and 10 Wm^{-2} to 560 Wm^{-2} in June (Fig. 3).

2. The derived net radiation flux over the study area is very close to the field measurement. It is the result of the improvement on surface albedo and surface temperature.
3. The regional soil heat flux derived from the relationship between soil heat flux and net radiation flux is suitable for heterogeneous land surface of the WATER area, because the relationship itself was derived from the same area.
4. The derived regional sensible heat flux and latent heat flux at the validation sites in the WATER area is in good agreement with field measurements (Fig. 4). This is due to the fact that atmospheric boundary layer processes have been considered in more detail in our methodology and the proposed parameterization for sensible heat flux and latent heat flux can be used over the upper streams of the Heihe River Basin area.

4 Concluding remarks

In this study, the regional distributions of land surface heat fluxes (net radiation flux, soil heat flux, sensible heat flux and latent heat flux) over middle and upper streams of the Heihe River Basin were derived with the aid of ASTER data and field observations. Reasonable results of land surface heat fluxes were gained in this study.

The retrieval of regional land surface heat fluxes over heterogeneous landscape is not an easy task.

1. Only three ASTER images are used in this study. To obtain more accurate regional land surface fluxes (daily to seasonal variations) over a larger area (the Heihe River Basin), more field observations (ABL tower and radiation measurement system, radiosonde system, turbulent fluxes measured by eddy correlation technique, soil moisture and soil temperature measurement system, etc.) and other satellite sensors such as MODIS (Moderate Resolution Imaging Spectroradiometer) and NOAA (National Oceanic and Atmospheric Administration)/AVHRR (Advanced Very High Resolution Radiometer) with more frequent temporal coverage have to be used.

2. This study implies the SEBS method is only applicable to clear-sky days. In order to extend its applicability to cloudy skies, we should consider using microwave remote sensing data to derive surface temperature and other land surface variables.

SEBS has been developed to estimate atmospheric turbulent fluxes using satellite earth observation data, in combination with meteorological data from a proper reference height given by either in-situ measurements for application to a point, and radiosonde or meteorological forecasts for application at larger scales. On the basis of these experimental validations, SEBS can be used to estimate turbulent heat fluxes at different scales with acceptable accuracy.

Acknowledgements. This work was under the CAS Action Plan for West Development Program (grant number: KZCX2-XB2-09-04) and Chinese State Key Basic Research Project (grant number: 2007CB714400). The auspices of the National Natural Science Foundation of China (40705004, 40825015 and 40810059006), Chinese National Key Programme for Developing Basic Sciences (2005CB422003), the Key Projects of International Cooperation, Chinese Academy of Sciences (GJHZ0735), the Opening Foundation of Institute of Plateau Meteorology, CMA(LPM2007014).

Estimating surface fluxes over the Heihe River Basin with ASTER imagery

W. Ma et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

References

- Baret, F. and Guyot, G.: Potentials and limits of vegetation indices for LAI and APAR assessment, *Remote Sens. Environ.*, 35, 161–173, 1997.
- Bastiaanssen, W. G. M.: Regionalization of surface fluxes and moisture indicators in composite terrain, Ph.D. thesis, Wageningen Agric. Univ., Wageningen, Netherlands, 143–161, p. 273, 1995.
- Becker, F. and Li, Z.-L.: Surface temperature and emissivity at various scales: Definition, measurement and related problems, *Remote Sensing Reviews*, 12, 225–253, 1995.
- Berk, A., Bernstein, L. S., and Robertson, D. C.: MODTRAN: A moderate resolution model for LOTRAN-7, GL-TR-89-0122, Geophys. Lab., Hanscom Air Force Base, Mass, 1989.
- Che'din, A., Scott, N. A., Wahiche, C., and Moulinier, P.: The improved initialisation inversion method: A high resolution physical method for temperature retrievals from Tiros-N series, *J. Clim. Appl. Meteorol.*, 24, 124–143, 1985.
- Juan, C., Sobrino, J., Gillespie, A., et al.: Improved land surface emissivities over agricultural areas using ASTER NDVI, *Remote Sens. Environ.*, 103(4), 474–487, 2006.
- Liou, K. N.: An introduction to atmospheric radiation, China Meteorol. Press, Beijing, China, 2nd edn., 2004.
- Kahle, A. B. and Alley, R. E.: Separation of temperature and emittance in remotely sensed radiance measurements, *Remote Sens. Environ.*, 42, 107–112, 1992.
- Yamaguchi, Y. and Kato, S.: Analysis of urban heat-island effect using ASTER and ETM+ Data: Separation of anthropogenic heat discharge and natural heat radiation from sensible heat flux, *Remote Sens. Environ.*, 99(1), 44–54, 2005.
- Kustas, W. P.: Estimates of evapotranspiration with a one- and twolayer model of heat transfer over partial canopy cover, *J. Appl. Meteorol.*, 29, 704–715, 1990.
- Kustas, W. P. and Daughtry, C. S. T.: Estimation of the soil heat flux/net radiation ratio from spectral data, *Agr. Forest Meteorol.*, 39, 205–223, 1990.
- Kustas, W. P. and Norman, J. M.: A two-source approach for estimating turbulent fluxes using multiple angle thermal infrared observations, *Water Resour. Res.*, 33, 1495–1508, 1997.
- Kustas, W. P., Choudhury, B. J., Moran, M. S., Reginato, R. J., Jackson, R. D., Gay, L. W., and Weaver, H. L.: Determination of sensible heat flux over sparse canopy using thermal infrared data, *Agr. Forest Meteorol.*, 44, 197–216, 1989.
- Li, X., Wang, J., Ma, M., et al.: Simultaneous remote sensing and ground-based experiment

HESSD

6, 4619–4635, 2009

Estimating surface fluxes over the Heihe River Basin with ASTER imagery

W. Ma et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Estimating surface fluxes over the Heihe River Basin with ASTER imagery

W. Ma et al.

- in the Heihe River Basin: Scientific objectives and experiment design, *Advance in Earth Sciences*, 23(9), 897–914, 2008 (in Chinese).
- Li, Z.-L. and Becker, F.: Feasibility of land surface temperature and emissivity determination from AVHRR data, *Remote Sens. Environ.*, 43, 67–85, 1993.
- 5 Liang, S.: Narrowband to broadband conversions of land surface albedo I Algorithms, *Remote Sens. Environ.*, 76, 213–238, 2000.
- Ma, W.: Estimating regional heat surface fluxes over heterogeneous landscapes of the Tibetan Plateau by using ASTER data, Ph.D. thesis, Chinese Academy of Sciences.
- Ma, W. and Ma, Y.: The annual variations on land surface energy in the northern Tibetan Plateau, *Environ. Geol.*, 50(5), 645–650, 2006b.
- 10 Ma, Weiqiang, Ma, Yaoming, Li, Maoshan, Hu, Zeyong, Zhong, Lei, Su, Zhongbo, Ishikawa, Hirohiko, and Wang, Jiemin: Estimating surface fluxes over the north Tibetan Plateau area with ASTER imagery, *Hydrol. Earth Syst. Sci.*, 13, 57–67, 2009, <http://www.hydrol-earth-syst-sci.net/13/57/2009/>.
- 15 Ma, Y.: Determination of regional surface heat fluxes over heterogeneous landscapes by integrating satellite remote sensing with boundary layer observations, Ph.D. thesis, Wageningen University, 2006.
- Ma, Y., Ishikawa, H., Tsukamoto, O., Menenti, M., Su, Z., Yao, T., Koike, T., and Yasunari, T.: Regionalization of surface fluxes over heterogeneous landscape of the Tibetan Plateau by using satellite remote sensing, *J. Meteorol. Soc. Jpn.*, 81, 277–293, 2003b.
- 20 Ma, Y., Zhong, L., Su, Z., Ishikawa, H., Menenti, M., and Koike, T.: Determination of regional distributions and seasonal variations of land surface heat fluxes from Landsat-7 Enhanced Thematic Mapper data over the central Tibetan Plateau area, *J. Geophys. Res.*, 111, D10305, doi:10.1029/2005JD006742, 2006a.
- 25 Ma, Y., Su, Z., Koike, T., Yao, T., Ishikawa, H., Ueno, K., and Menenti, M.: On measuring and remote sensing surface energy partitioning over the Tibetan Plateau – From GAME/Tibet to CAMP/Tibet, *Phys. Chem. Earth*, 28, 63–74, 2003a.
- Menenti, M. and Choudhury, B. J.: Parameterization of land surface evaporation by means of location dependent potential evaporation and surface temperature range, in: *Exchange Processes at the Land Surface for a Range of Space and Time Scales*, edited by: Bolle, H. J., Feddes, R. A., and Kalma, J. D., IAHS Publ., 212, 561–568, 1993.
- 30 Menenti, M., Bastiaanssen, W. G. M., and Van Eick, D.: Determination of hemispheric albedo with Thematic Mapper data, *Remote Sens. Environ.*, 28, 327–337, 1989.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Estimating surface fluxes over the Heihe River Basin with ASTER imagery

W. Ma et al.

- Menenti, M., Bastiaanssen, W. G. M., Hefny, K., and Abd El Karim, M. H.: Mapping of ground water losses by evaporation in the Western Desert of Egypt, DLO Winand Staring Centre, Wageningen, Netherlands, Rep. 43, 116 pp., 1991.
- Monteith, J. L.: Principles of environmental physics, Edward Arnold Press, 241 pp., 1973.
- 5 Norman, J. M., Kustas, W. P., and Humes, K. S.: A two-source approach for estimating soil and Vegetation energy fluxes from observations of directional radiometric surface temperature, *Agr. Forest Meteorol.*, 77, 263–293, 1995.
- Pinker, R. T.: Satellites and our understanding of the surface energy balance, *Palaeogeogr. Palaeoclimatol.*, 82, 321–342, 1990.
- 10 Price, J. C.: Estimating vegetation amount from visible and near infrared albedo, *Remote Sens. Environ.*, 41, 29–34, 1992.
- Qi, J., Chehbouni, A., Huete, A. R., Kerr, Y. H., and Sorooshian, S.: A modified soil adjusted vegetation index, *Remote Sens. Environ.*, 48, 119–126, 1994.
- Rahman, H. and Dedieu, G.: SMAC: a simplified method for atmospheric correction of satellite measurements in the solar spectrum, *Int. J. Remote Sens.*, 15, 123–143, 1994.
- 15 Schugge, T. J., Hook, S., and Kahle, A.: TIMS observation of surface emissivity in HAPEX-Sahel, paper presented at International Geoscience and Remote Sensing Symposium, Inst. of Electr. and Electron. Eng., Florence, Italy, July, 1995.
- Sellers, P. J., Rasool, S. I., and Bolle, H. J.: A review of satellite data algorithms for studies of the land surface, *B. Am. Meteorol. Soc.*, 71, 1429–1447, 1990.
- 20 Liang, S.: Narrowband to broadband conversions of land surface albedo, *Remote Sens. Environ.*, 76(2), 213–238, 2001.
- Sobrino, J. A. and Raissouni, N.: Toward remote sensing methods for land cover dynamic monitoring: application to Morocco, *Int. J. Remote Sens.*, 21(2), 353–366, 2000.
- 25 Su, Z.: The Surface Energy Balance System (SEBS) for estimation of turbulent heat fluxes, *Hydrol. Earth Syst. Sci.*, 6, 85–100, 2002, <http://www.hydrol-earth-syst-sci.net/6/85/2002/>.
- Susskind, J., Rosenfield, J., Renter, D., and Chahine, M. T.: Remote sensing of weather and climate parameters from HIRS2/MSU on TIROS-N, *J. Geophys. Res.*, 89, 4677–4697, 1984.
- 30 Tucker, C. J. (Ed.): Monitoring the grasslands of semi-arid Africa using NOAA AVHRR data, *Int. J. Remote Sens.*, 7–11, 1383–1622, 1986.
- Wan, Z. and Dozier, J.: Land surface temperature measurement from space: Physical principles and inverse modelling, *IEEE T. Geosci. Remote*, 27, 268–278, 1989.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Wang, J., Ma, Y., Menenti, M., Bastiaanssen, W. G. M., and Mistsuta, Y: The scaling-up of processes in the heterogeneous landscape of HEIFE with the aid of satellite remote sensing, J. Meteorol. Soc. Jpn., 73, 1235–1244, 1995.

Watson, K., Kruse, F., and Hummer-Miler, S.: Thermal infrared exploration in the Carlin trend, Geophysics, 55, 70–79, 1990.

Oku, Y. and Ishikawa, H.: Estimation of Land Surface Temperature over the Tibetan Plateau Using GMS Data, J. Appl. meteorology, 43, 548–561, 2004.

Oku, Y., Ishikawa, H., and Su, Z.: Estimation of Land Surface Energy Fluxes over the Tibetan Plateau using GMS Data, J. Appl. Meteorol. Clim., 46(2), 183–195, 2007.

Yamaguchi, Y., Kahle, A. B., Tsu, H., Kawakami, T., and Priel, M.: Overview of the Advanced Spaceborne Thermal Emission and Reflection Radiometer (ASTER), IEEE T. Geosci. Remote, 36(4), 1062–1071, 1998.

HESSD

6, 4619–4635, 2009

Estimating surface fluxes over the Heihe River Basin with ASTER imagery

W. Ma et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Estimating surface fluxes over the Heihe River Basin with ASTER imagery

W. Ma et al.




Fig. 1. Sketch map of studying area and the sites during the WATER. Network of the automatic meteorological stations and flux towers in WATER (For regional meteorological stations, two elements indicate air temperature and precipitation; three elements indicate wind direction plus two elements; four elements indicate wind speed plus three elements; six elements indicate air pressure, global radiation plus four elements. SMTMS: Soil Moisture and Temperature Measurement System).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Estimating surface fluxes over the Heihe River Basin with ASTER imagery

W. Ma et al.




Fig. 2. Diagram of parameterization procedure by combining ASTER data with field observations.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Estimating surface fluxes over the Heihe River Basin with ASTER imagery

W. Ma et al.




Fig. 3. Distribution maps of land surface heat fluxes over the WATER area.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Estimating surface fluxes over the Heihe River Basin with ASTER imagery

W. Ma et al.




Fig. 4. Validation of the derived net radiation R_n , soil heat flux G_0 , sensible heat flux H and latent heat flux λE against ground measurements over the Watershed Airborne Telemetry Experimental Research (WATER) stations, Yingke, Huazhaizi, Guantan, Maliantan, A'rou, Binggou and Yakou with 1:1 line.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion