Hydrol. Earth Syst. Sci. Discuss., 6, C3577-C3582, 2010

www.hydrol-earth-syst-sci-discuss.net/6/C3577/2010/ © Author(s) 2010. This work is distributed under the Creative Commons Attribute 3.0 License.



HESSD

6, C3577–C3582, 2010

Interactive Comment

Interactive comment on "Improving the rainfall rate estimation in the midstream of the Heihe River Basin using rain drop size distribution" by G. Zhao et al.

G. Zhao et al.

guozh@lzb.ac.cn

Received and published: 21 July 2010

The combination of polarimetric variables such as R (Z, ZDR) and R (KDP, Z, ZDR) are superior to R(Z) because of the less sensitivity of ZDR an KDP to DSD variations. The differential phase (KDP) can be used to correct the reflectivity factor for loss due to beam blockage by topography, attenuation, and anomalous propagations (Ryzhkov and Zrnic 1996; Ryzhkov et al. 2000). The lower sensitivity of R(KDP, Z, ZDR) and the higher sensitivity of R(Z) to variations in DSD can be explained by the fact that the difference between the forward-scattering amplitudes at horizontal (H) and vertical (V)





polarizations fH(D)-fV(D) in the definition of KDP is proportional to the 3rd power of the diameter of a raindrop for the mono-disperse DSD model, while the reflectivity factor Z is proportional to the 6th power of the diameter. There is an X-band polarimetric radar and several rain gauges available for testing these rainfall rate estimation. The attenuation has been corrected before we used the radar data. The follow table is given for testing. Fig.3 shows the radar reflectivity of precipitationaAC The OTT Parsivel has not be applied in other areas in China, but Zhang(1989) got the rain drop size distribution data in Piliang and got a conclusion that the relationship between the R and Z was $R=a \times zb$, but our conclusion was $R=a \times 10b \times Z$. It revealed that the Characteristic of rain drop size distribution in this area is different from the area of Qinghai-Tibet Plateau. Fig. 2 shows the DSD for a precipitation of stratiform rainfall cloud, the number of data is 236. Fig. 3 shows the DSD for a convective rainfall cloud, the number of data is 285. Fig. 3 shows that the size distribution for convective rain is broader than that for stratiform. Similar to Steinerde(1987)' conclusion, there is a multi-peak structure. Most peaks appear in the D < 3 mm region for convective rainfall cloud. In manuscript Fig.1 is not the average values for raindrop diameter. Table.1 shows the terminal velocity of the rain drop with different diameters. There the terminal velocity and rain drop diameters are got from PASIVEL directly. Because PASIVEL got the terminal velocity are rang of diameter, In Table 1 and Figure 1, one point contained lots of raindrop diameters. I agree with your conclusion that "4 types estimator" better instead of "4 relationships of rain-rate and radar parameters"

Please also note the supplement to this comment: http://www.hydrol-earth-syst-sci-discuss.net/6/C3577/2010/hessd-6-C3577-2010supplement.pdf

Interactive comment on Hydrol. Earth Syst. Sci. Discuss., 6, 6107, 2009.

HESSD

6, C3577–C3582, 2010

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



雷达站名:中科院寒旱所 雷达型号:714XDP 海拔高度:1669.0 = 日期: 2008/06/13 B† (ii]: 18:44:47 显示类型:无订正 重复频率: 1000/800 Hz 显示距离:75km 天线仰角:3.00* 35 40 45 50 55 60 70 极化方式: 双极化 极化方式: 双极化 雷达站名:中科院寒旱所 雷达型号:714XDP 海拔高度:1669.0m Full Screen / Esc 日期: 2008/06/13 时间: 19:09:04 Printer-friendly Version 显示类型:无订正 重复频率: 1000/800 Hz 显示距离:75km Interactive Discussion 天线仰角:2.00* Discussion Paper $(\mathbf{\hat{H}})$ (cc) 10 15 50 55 35 40 极化方式: 双极化 C3579 极化方式:双极化

Fig. 1 The radar reflectivity of precipitation

HESSD

6, C3577–C3582, 2010

Interactive

Comment

HESSD

6, C3577-C3582, 2010

Interactive Comment

Fig. 2 DSD for the stratiform rainfall cloud



Full Screen / Esc

Printer-friendly Version

Interactive Discussion



HESSD

6, C3577-C3582, 2010

Interactive Comment



2

4

D/mm

6

8

ő

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Fig. 3 DSD for the convective rainfall cloud

HESSD

6, C3577–C3582, 2010

Interactive Comment

Rain	$Z_H(dBZ)$	$Z_{DR}(dB)$	$K_{DP}(\text{degkm}^{-1})$	R(Z)/Err	$R(Z,Z_{DR})/\text{Err}$	$R(Z,Z_{DR},K_{DP})$ /Err	R _{gauge}	
gauge								
NO.03	29	0.8	1.2	1.9/17%	2.1/8%	2.0/13%	2.3	
NO.05	35	1.4	0.3	4.1/46%	4.9/75%	2.0/28%	2.8	
NO.11	35	1.9	1.4	4.1/2%	4.2/5%	4.5/12%	4.0	
NO.12	31	1.0	0.3	2.5/21%	2.8/12%	3.8/18%	3.2	
NO.14	38	2.0	2.2	6.1/39%	6.3/43%	4.7/7%	4.4	on / Eco
NO.19	35	1.3	2.9	4.1/29%	5.1/12%	5.9/2%	5.8	en / Esc
NO.26	23	1.3	0.6	0.7/75%	0.38/5%	0.31/22%	Printer-frier 0.4	dly Version
NO.29	35	4.6	1.6	4.5/26%	5.3/13%	6.0/1%	Interactive 6.1	Discussion

Table 1 The reliability of the rainfall estimators

