

Interactive comment on “Assessment of soil moisture fields from imperfect climate models with uncertain satellite observations” by G. Schumann et al.

Anonymous Referee #3

Received and published: 22 April 2009

General comments

The paper addresses the evaluation of seasonal soil moisture predictions of the LAM and GCM model against satellite observations, both averaged over a number of years. The fuzzy logic theory proposed in the paper is appealing as it explicitly accounts for uncertainty of satellite observations and as far as I know, it has not been used to evaluate Land Surface Schemes (LSS) before. However, I agree with the other reviewers that the paper lacks an elaboration of the sources of uncertainty and how these are encompassed in the fuzzy framework that is proposed. For example, from the paper it is unclear why the accuracy of the satellite observations has not been

C376

included and why the variance of the observation has been taken as a measure of uncertainty instead.

I also encourage including an elaboration and justification on the evaluation criterion that has been used, embedded in a discussion on the role of soil moisture in climate models, addressing issues such as: Why is the average soil moisture used and not e.g. the temporal variation of soil moisture? Are there other satellite products, that may be included in the evaluation? And how important is the soil moisture content in the upper 5-10 cm of the soil column for the the LSS performance, as the main source of evaporation is the root water uptake of vegetation, taken from the entire soil column?

Furthermore, as already stated by the other reviewers, some of the statements are not well justified or unclear. E.g. in Section 4.1 it is suggested that improved parameterization of runoff might improve the performance of LSS. In the model description, it is not explained which runoff parameterization is applied by MOSES and for the reader it is unclear, why the remark is made and how it fits in the overall aim of the paper.

Detailed comments:

I miss in the introduction a reference to other satellite products that may be used for LSS evaluation. See for example the SEBAL algorithm that derives turbulent heat fluxes from satellite images (e.g. Bastiaanssen et al., 2005).

Fig 2: The figure 2 shows the statistics of the absolute difference between RS1 and RS2. It would be informative to include some statistics of RS1 and RS2 as well

I agree with referee 2 that the improvement of the soil moisture simulation by LAM as compared to the GCM is not justified by the results presented in figure 5.

Technical corrections:

Eq.(1): The equation should be revised as c_1 and c_2 can be either $RS1 + \sigma$, $RS2 + \sigma$ or $RS1 - \sigma$ and $RS2 - \sigma$.

C377

I suggest reconsidering the title of the y-axis of figure 2.

References: Bastiaanssen, W.G.M., E.J.M. Noordman, H. Pelgrum, G. Davids, B.P. Thoreson and R.G. Allen, 2005: SEBAL model with remotely sensed data to improve water-resources management under actual field conditions. *ASCE Journal of Irrigation and Drainage Engineering*, 131, 85-93.

Interactive comment on Hydrol. Earth Syst. Sci. Discuss., 6, 2733, 2009.