

This discussion paper is/has been under review for the journal Hydrology and Earth System Sciences (HESS). Please refer to the corresponding final paper in HESS if available.

# Interannual variations of the terrestrial water storage in the Lower Ob' basin from a multisatellite approach

**F. Frappart<sup>1</sup>, F. Papa<sup>2</sup>, A. Güntner<sup>3</sup>, S. Werth<sup>3</sup>, G. Ramillien<sup>4</sup>, C. Prigent<sup>5</sup>,  
W. B. Rossow<sup>2</sup>, and M.-P. Bonnet<sup>1</sup>**

<sup>1</sup>Université de Toulouse, CNRS, IRD, OMP-LMTG, UMR 5563, Toulouse, France

<sup>2</sup>NOAA Cooperative Remote Sensing Science and Technology Center, City College of New York, New York, USA

<sup>3</sup>GFZ, Telegrafenberg, Potsdam, Germany

<sup>4</sup>Université de Toulouse, CNES, CNRS, IRD, OMP-DTP, Toulouse, France

<sup>5</sup>Laboratoire d'Etudes du Rayonnement et de la Matière en Astrophysique, Observatoire de Paris, CNRS, Paris, France

Received: 23 July 2010 – Accepted: 13 August 2010 – Published: 1 September 2010

Correspondence to: F. Frappart (frederic.frappart@lmtg.obs-mip.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

## Abstract

Temporal variations of surface water volume over inundated areas of the Lower Ob' basin in Siberia, one of the largest contributor of freshwater to the Arctic Ocean, are estimated using combined observations from a multisatellite inundation dataset and water levels over rivers and floodplains derived from the TOPEX/POSEIDON (T/P) altimetry satellite. We computed time-series of monthly maps of surface water volume over the period of common availability of T/P and the multisatellite data (1993–2004). The results exhibit similar interannual variabilities with precipitation estimates and river discharge observations. This study also presents monthly estimates of groundwater and permafrost mass anomalies during 2003–2004 based on a synergistic analysis using multisatellite observations and hydrological models. Water stored in aquifer is isolated from the total water storage measured by GRACE by removing the contributions of both the surface reservoir, derived from satellite imagery and radar altimetry, and the root zone reservoir simulated by hydrological models.

## 1 Introduction

All climate scenarios agree on the high sensitivity of the northern regions to global change, with a stronger warming at these latitudes than globally. Average annual arctic temperatures have increased at almost twice the global rate over recent decades and are predicted to increase by an additional 4–7 °C by the end of the century (Serreze et al., 2000). Continued warming of the Arctic regions will likely have profound consequences, many of which may have regional and global implications (Peterson et al., 2002). For instance, an increase in air temperature is expected to intensify the Arctic hydrological cycle (Stocker and Raible, 2005; Wu et al., 2005) and important modifications of the terrestrial water cycle have been already observed in northern latitudes. One observable change that attracted considerable scientific attention is the increase over the last century by 7% in the average annual discharge of fresh water from the six largest Eurasian rivers to the Arctic Ocean (Peterson et al., 2002; Yang et al., 2002).

### Terrestrial water storage variations in the Lower Ob'

F. Frappart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Other recent evidence of changes in the Arctic hydrology was revealed with the analysis of optical remote sensing images by Smith et al. (2005) which showed a reduction between 1975 and 1997 in the number of lakes in West Siberia that could be related to permafrost thawing. These modifications could lead to significant repercussions on both the biogeochemical and hydrological cycles of northern high latitude systems, with further influence on the global climate.

Among the various reservoirs in which fresh water on land is stored (e.g., ice caps, glaciers, snowpack, soil moisture and groundwater), surface waters (rivers, lakes, reservoirs, wetlands and inundated areas) play a crucial role in the global biogeochemical and the hydrological cycles (de Marsily et al., 2005). Although wetlands and floodplains cover about 6% of the Earth surface (OECD, 1996), they have a substantial impact on flood flow alteration, sediment stabilization, water quality, groundwater recharge and discharge (Maltby, 1991; Bullock and Acreman, 2003). Moreover, floodplain inundation is an important regulator of river hydrology owing to storage effects along channel reaches. Reliable and timely information about the extent, spatial distribution, and temporal variation of wetlands and floods as well as the amount of water stored is crucial to better understand their relationship with river discharges, and also their influence on regional hydrology and climate.

Remote sensing techniques have been very useful for studying cold region climate and hydrology (Massom, 1995; Smith, 1997; Seidel and Martinec, 2004) as they provide a unique mean to observe large regions and are the only alternative to in situ data in remote areas. Good capabilities for monitoring snow extent and depth using passive microwave measurements (Yang et al., 2002; Grippa et al., 2005) or multispectral images (Hall et al., 2002; Robinson and Frei, 2004), water levels and discharges using radar altimetry (Kouraev et al., 2005), and inundation extent combining multisatellite information (Papa et al., 2007; 2008a) have been demonstrated. The Gravity Recovery And Climate Experiment (GRACE) mission, launched in 2002, detects tiny changes in the Earth's gravity field which can be related to the spatio-temporal variations of the terrestrial water storage (TWS) at monthly or 10-day time-scales (Tapley et al., 2004).

---

**Terrestrial water storage variations in the Lower Ob'**F. Frappart et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Previous studies provide important information on changes in TWS and snow mass at high latitudes (Muskett and Romanovsky, 2010; Frappart et al., 2010). TWS, representing an integrated measurement of the water stored in the different hydrological reservoirs and consequently the sum of the surface water, water stored in the root zone, snowpack and groundwater, can be considered as a good indicator of the changes that occur in the hydrological conditions globally and at basin-scale. Nevertheless, TWS is difficult to measure due to the lack of a complete network of in situ observations of the terrestrial hydrological components. The hydrologic regime of rivers is strongly influenced by the presence of permafrost at high latitudes. The river basins with high permafrost coverage have lower subsurface storage capacity, which causes a lower winter baseflow and a higher summer peak flow, than basins with low coverage (Woo, 1986; Kane, 1997; Smith et al., 2007).

Variations in the groundwater storage can be extracted from the TWS measured by GRACE using external information on the other hydrological reservoirs such as in situ observations (Yeh et al., 2006), model outputs (Rodell et al., 2009), or both (Leblanc et al., 2009). No similar studies have been yet undertaken for large river basins characterized by extensive wetlands or floodplains in high latitude regions.

In this study, we demonstrate that remote sensing and hydrological modelling can be combined to estimate water storage changes in the Lower Ob' basin using remote sensing and hydrological modelling by combining high-resolution imagery-derived inundation extents, altimetry-derived water level measurements and gravimetry from space products, and outputs from hydrological models. We first present the analysis of twelve years of surface water storage variations. Then, a new method, based on the combination of multisatellite-derived hydrological products and outputs from global hydrology models, is proposed to derive the spatio-temporal variations of water volume anomalies in the aquifer and the permafrost of the Lower Ob' basin. Water storage anomalies in the different hydrological reservoirs are removed from the TWS measured by GRACE to isolate the sum of the groundwater and permafrost anomaly storage over 2003–2004.

**Terrestrial water storage variations in the Lower Ob'**

F. Frappart et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## 2 Study area and datasets

### 2.1 The Lower Ob' basin

The Ob' is the largest watershed of all Arctic rivers (2 975 106 km<sup>2</sup>) and is the third largest contributors of freshwater to the Arctic ocean after the Yenisey and Lena rivers (Shiklomanov et al., 2000). This watershed is mostly underlain by discontinuous permafrost with portions of continuous permafrost in its northern part and sporadic permafrost in its southern part (Brown et al., 1998; Zhang et al., 1999) and during the winter, a large part of the basin is covered by snow, which persists for six months or more. The latitudinal extent of this basin results in the gradual snowmelting from south to north during spring/summer seasons and leads to large flood events. During the spring/summer season, the Ob' River basin is frequently described as the world's biggest swamp. The object of our study is the lower Ob', the part of the Ob' River comprised from the Yrtish mouth to the Ob' bay (Fig. 1).

### 2.2 The multisatellite inundation dataset

The multisatellite inundation dataset quantifies at global scale the monthly distribution of surface water extent and its variations at ~25 km of resolution. The methodology which captures the extent (with an accuracy of ~10%) of episodic and seasonal inundations, wetlands, rivers, lakes, and irrigated agriculture over more than a decade, 1993–2004, is based on a clustering analysis of a suite of complementary satellites observations, including passive (SSM/I) and active (ERS) microwaves, visible and near-IR (AVHRR) observations (Papa et al., 2007, 2008 a, b, 2010; Prigent et al., 2007). As the microwave measurements are also sensitive to snow cover, snow masks are used to edit the results and avoid any confusion with snow-covered pixels. The weekly North Hemisphere and South Hemisphere snow mask from the National Snow and Ice Data Center (University of Colorado; NSIDC) is adopted and averaged on a monthly basis (Armstrong and Brodzik, 2005).

## Terrestrial water storage variations in the Lower Ob'

F. Frappart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



This dataset has been intensively used for climatological and hydrological analyses, such as the evaluation of the methane surface emission models (Ringeval et al., 2010) or the validation of the river flooding scheme in land surface models (Decharme et al., 2008). In high latitudes regions, Papa et al. (2007) evaluated the consistency of the spatial and temporal variations of this inundation dataset over the Ob' basin using in situ snow depth and river runoff and Papa et al. (2008) investigated the response of river discharge to seasonal flood extent change in the large Siberian watersheds.

### 2.3 T/P-derived water levels

First devoted to ocean studies, radar altimetry is commonly used to monitor water levels over lakes, rivers and floodplains (Birkett, 1998). In this study, we used 90 (old orbit) and 92 (new orbit) time series of water levels derived from T/P measurements over the Lower Ob' drainage basin (see Fig. 1 for their locations), following the methodology proposed by Frappart et al. (2006a), for the period 1993–2004. For each intersection between the river (or the floodplain) and the satellite ground track, we define a so-called “altimetry station”, represented by a rectangular window. Outliers are deleted using a  $3\text{-}\sigma$  criterion over the whole time span of analysis. For each 10 day cycle, the water level at a given altimetry station is obtained by computing the median of all the high-rate data (10 Hz) included in the rectangular window. This process, repeated for each cycle, allows the construction of a water level time-series at the altimetry station. The uncertainty associated with the water level height ranges between 5–25 cm for high water season to 15–50 cm during low water season. Note that in high latitudes regions, the presence of ice and snow perturbs the altimetric signal during a large portion of the year (Kouraev et al., 2004).

### 2.4 GRACE-derived land water and snow mass solutions

The Gravity Recovery And Climate Experiment (GRACE) mission, launched in March 2002, provides measurements of the spatio-temporal changes in Earth's gravity field.

## Terrestrial water storage variations in the Lower Ob'

F. Frappart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Several recent studies have shown that GRACE data over the continents can be used to derive the monthly changes of the total land water storage (Ramillien et al., 2005, 2008; Schmitt et al., 2008) with an accuracy of  $\sim 1.5$  cm of water thickness equivalent when averaged over surfaces of a few hundred square-kilometres. Here we used the three latest land water solutions (RL04) produced by GFZ, JPL (for these two first products, January 2003, June 2003 and January 2004 are missing), and CSR (June 2003 and January 2004 are missing) with a spatial resolution of  $\sim 400$  km and the derived snow solutions (Frappart et al., 2006b, 2010). The Level-2 raw data consist of monthly estimates of geo-potential coefficients adjusted for each 30-day period from raw along-track GRACE measurements by different research groups (i.e., CSR, GFZ and JPL). These coefficients are developed up to a degree 60 (or spatial resolution of 333 km) and corrected for oceanic and atmospheric effects (Bettadpur, 2007) to obtain residual global grids of ocean and land signals corrupted by a strong noise. These data are available at: <ftp://podaac.jpl.nasa.gov/grace/>.

## 2.5 Water in the root zone from hydrology models

### The LaD model

The LaD model provides monthly  $1^\circ \times 1^\circ$  gridded time series of surface parameters. For each cell of the model, the total water storage is composed of three stores: a snowpack store, a root-zone store and a groundwater store and the total energy storage is equal to the sum of latent heat of fusion of the snowpack and the glacier and sensible heat content (Milly and Schmakin, 2002).

### The WGHM model

The Water GAP Global Hydrology Model (WGHM) was specifically designed to estimate river discharge for water resources assessments. It computes  $0.5^\circ \times 0.5^\circ$  gridded time series of monthly runoff and river discharge and is tuned against time series of

## Terrestrial water storage variations in the Lower Ob'

F. Frappart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



annual rivers discharges measured at 724 globally distributed stations. The soil water balance takes into account the water content of the soil within the effective root zone, the effective precipitation as the sum of throughfall and snowmelt, the actual evapotranspiration and the surface runoff. Surface runoff is computed from the water balance equation that takes into account the water content within the effective root zone, the effective precipitation and the evapotranspiration (Döll et al., 2003).

We use the water content within the root zone from these two models to simulate the soil moisture reservoir. It is important to note that WGHM takes into account the occurrence of permafrost in a grid cell for the groundwater recharge but LaD not.

## 2.6 Precipitation estimates from GPCP

These data quantify the distribution of precipitation over the global land surface (Adler et al., 2003). We use the monthly Satellite-Gauge Combined Precipitation Data product Version 2 data, available from January 1997 to present with a spatial resolution of 1° of latitude and longitude. Over land surfaces, the uncertainty in the rate estimates from GPCP is generally lower than over the oceans due to the in situ gauge input (in addition to satellite) from the GPCC (Global Precipitation Climatology Center). Over land, validation experiments have been conducted in a variety of location worldwide and suggest that while there are known problems in regions of persistent convective precipitation, non precipitating cirrus or regions of complex terrain, the estimates uncertainties range between 10%–30% (Adler et al., 2003).

## 2.7 In situ hydrological data from ArcticRIMS

Daily water levels from the Salekhard station between February 2000 and December 2004, and monthly river discharges from Salekhard station between January 1993 and December 2004, and Belogorje station between January 1993 and October 1999 (but with missing data over long time periods for Belogorje station) were obtained from the ArcticRIMS web site (ArcticRIMS, 2003).

## Terrestrial water storage variations in the Lower Ob'

F. Frappart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





### 3 Methodology

#### 3.1 Monthly maps of water levels

Monthly maps of water levels over the floodplains of the Lower Ob' basin have been determined using combined observations from the multi-satellite inundation dataset, T/P derived water levels. For a given month during the flood season, water levels were linearly interpolated over the flooded zones of the Lower Ob' basin. A pixel of 25 km×25 km is considered inundated when its percentage of inundated area is greater than 0. Maps of interpolated surface water levels with 25 km resolution have been constructed for each month between January 1993 and December 2004. The water levels are expressed with respect to the GGM02C geoid model (Tapley et al., 2005). The elevation of each pixel of the water level maps is given with reference to its minimum computed over the 1993–2004 period. This minimum elevation represents either the bathymetry or very low water stage of the floodplain. More details about the methodology can be found in (Frappart et al., 2005, 2006c, 2008).

#### 3.2 Rescaling of the GRACE data

The GRACE solutions suffer from the presence of important high-frequency noise (i.e., north-south striping) caused by orbit resonance in spherical harmonics determination and aliasing of short-time phenomena which are geophysically unrealistic. An optimum filter method was determined by analyzing the correspondance of GRACE basin-average water storage to the ensemble mean of hydrological models (WGHM, LaD) and by analyzing the error budgets (satellite/leakage errors) and amplitude and phase biases for the different filter types. For the Lower Ob' basin, the simple Gaussian filter with 500 km radius was the best filter. Only a very small bias in the seasonal phase of storage changes resulted due to filtering for the selected optimum filter methods. The GRACE products have been rescaled with a factor of 0.98 to account for amplitude smoothing due to filtering determined from smoothed and unsmoothed basin-average model ensemble time series of water storage (Werth et al., 2009).

## Terrestrial water storage variations in the Lower Ob'

F. Frappart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 3.3 Groundwater storage estimates

The time variations in Terrestrial Water Storage (TWS) are the sum of the contributions of the different reservoirs present in a drainage basin:

$$\Delta TWS = \Delta SW + \Delta RZ + \Delta SN + \Delta GW + \Delta P \quad (1)$$

where SW represents the total surface water storage including lakes, reservoirs, in-channel and floodplains water, RZ is the water contained in the root zone of the soil (representing a depth of 1 or 2 m), SN is the snow storage, GW is the total groundwater storage in the aquifers and P the permafrost storage. These terms are generally expressed in volume (km<sup>3</sup>) or mm of equivalent-water height.

The sum of GW and P anomalies over 2003–2004 is obtained in Eq. (1) by calculating the difference between the TWS anomaly estimated by GRACE and the SW level anomaly maps previously derived from remote sensing and the RZ anomaly derived from hydrological models outputs. The TWS and RZ monthly anomalies are the average anomalies of respectively the Level-2 GRACE CSR, GFZ and JPL destriped and smoothed solutions at 300 km of averaging radius, and the outputs from LaD and WGHM.

### 3.4 Water volume variations

For a given month  $t$ , the regional water volume of TWS, SW, RZ or GW+P storage  $\delta V(t)$  in a basin with surface area  $S$ , is simply computed from the water heights  $\delta h_j$ , with  $j = 1, 2, \dots$  (expressed in mm of equivalent water height) inside  $S$ , and the elementary surface  $R_e^2 \sin\theta_j \delta\lambda \delta\theta$  (and the percentage of inundation  $P_j$  for SW):

$$\delta V(t) = R_e^2 \sum_{j \in S} P_j \delta h_j(\theta_j, \lambda_j, t) \sin\theta_j \delta\lambda \delta\theta \quad (2)$$

where  $\lambda_j$  and  $\theta_j$  are co-latitude and longitude respectively,  $\delta\lambda$  and  $\delta\theta$  are the grid steps in longitude and latitude (generally  $\delta\lambda = \delta\theta$ ), and  $R_e$  the mean radius of the Earth (6378 km). The surface and total water volume variations are expressed in km<sup>3</sup>.

## Terrestrial water storage variations in the Lower Ob'

F. Frappart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 3.5 Annual cycle of water volumes

From the series of TWS, SW, RZ and GW+P anomaly grids, the temporal trend, seasonal and semi-annual amplitude were simultaneously fitted by least-square adjustment at each grid point. We assumed that, at 1st order, the changes  $\delta q(t)$  at each grid point are the sum of a linear trend, an annual sinusoid (with frequency  $\omega_{\text{ann}} = \frac{2\pi}{T_{\text{ann}}}$ , with  $T_{\text{ann}} \sim 1$  year), a semi-annual sinusoid (with frequency  $\omega_{\text{semi-ann}} = \frac{2\pi}{T_{\text{semi-ann}}}$ , with  $T_{\text{semi-ann}} \sim 6$  months) and water mass residuals  $\delta q^{\text{RES}}(t)$ :

$$\delta q(t) = At + B + C \cos(\omega_{\text{ann}} t + \varphi_{\text{ann}}) + D \cos(\omega_{\text{semi-ann}} t + \varphi_{\text{semi-ann}}) + \delta q^{\text{RES}} \quad (3)$$

The parameters which we adjusted for each grid point ( $\theta$ ,  $\lambda$ ) are the linear trend (i.e. slope  $A$  and y-intercept  $B$ ), the annual cycle (i.e. amplitude  $C$  and phase  $\varphi_{\text{ann}}$ ) and the semi-annual cycle (i.e. amplitude  $D$  and phase  $\varphi_{\text{semi-ann}}$ ). For this purpose, we merely used a error-free least-square fitting to solve the system:

$$\delta Q = \Phi \cdot X \quad (4)$$

where the vector  $\delta Q$  is the list of the SWE values,  $\Phi$  and  $X$  are the configuration matrix and the parameter vector, respectively. The latter two terms are:

$$\{\Phi_j\} = [t_j \ 1 \ \cos(\omega_{\text{ann}} t_j) \ \sin(\omega_{\text{ann}} t_j) \ \cos(\omega_{\text{semi-ann}} t_j) \ \sin(\omega_{\text{semi-ann}} t_j)] \quad (5)$$

$$X = [\alpha \ \beta \ \xi_{\text{ann}} \cos \varphi_{\text{ann}} \ -\xi_{\text{ann}} \sin \varphi_{\text{ann}} \ \xi_{\text{semi-ann}} \cos \varphi_{\text{semi-ann}} \ -\xi_{\text{semi-ann}} \sin \varphi_{\text{semi-ann}}] \quad (6)$$

for adjusting the temporal trend and for fitting the annual and semi-annual amplitude and phase.

According to the least-squares criteria, the solution vector of the system is:

$$X^{\text{SOL}} = (\Phi^T \Phi)^{-1} \Phi^T \delta Q \quad (7)$$

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 4 Results and discussion

### 4.1 Monthly inundation extent

The inundated area fractions for the Lower Ob' watershed were extracted from the multisatellite inundation dataset. The temporal evolution of the total inundated extent (km<sup>2</sup>) over the lower Ob River Basin for the period 1993–2004 is presented in Fig. 2. The mean extent during the snow-free months is only of  $29 \times 10^3$  km<sup>2</sup>. The inundation extent always reaches its maximum in July, except for 1996, 1998, and 1999 where the maximum occurred in August. The inundation exhibits a strong seasonal cycle, with the floods generally starting in June and lasting until October. It also presents an important interannual variability. In 1993, the maximum inundated extent reached  $67 \times 10^3$  km<sup>2</sup> whereas in 1995, it only reached  $32 \times 10^3$  km<sup>2</sup>.

The spatial pattern of the inundation over the Lower Ob' Basin is shown in Fig. 3. Figure 3a displays maps of the annual maximum of fractional inundation extent (in %, 100% of inundations equals to an area of 773 km<sup>2</sup>) from satellite estimates for each year of the period 1993–2004. Figure 3b shows the corresponding month during which the maximum inundation occurs. The inundation extent is clearly delineated over the river channel and the associated floodplains. The pixels with a very high percentage of inundation (greater than 50%) are located along the river channel. A large inundation area is present downstream the junction between the Ob' and the Irtysh Rivers around 60° N and 65° E.

Figure 3b presents a map of the month during which occurred the maximum inundation extent. As seen in the results from Papa et al., 2008, this map clearly shows a propagation from south to north resulting from the latitudinal dependence of the snow melting. The maximum inundation extent occurs from May to June in the southern part of the basin and from July to August on the northern region. The propagation of the maximum inundation extent from south to north generally takes two to three months (this rough estimate is limited by the monthly temporal resolution of the dataset).

HESSD

7, 6647–6676, 2010

## Terrestrial water storage variations in the Lower Ob'

F. Frappart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 4.2 Altimetry-derived water levels time series

The T/P altimetry stations where water level time series can be constructed are unevenly distributed across the basin (Fig. 1). An important point to notice is that this dataset gives valuable information on water levels especially for unmonitored regions of the Lower Ob' watershed, in particular over the floodplains.

The water level time series derived from radar altimetry for the stations close to Salekhard are presented in Fig. 4 along with in situ river heights time series. Comparisons have been made between the in situ gauge station of Salekhard at the mouth of the Ob' River and the closest altimetry stations on the river channel for both the old and the new orbit of T/P (Table 1). For the old orbit, two altimetry stations can be defined at locations around 65 km at the South of Salekhard. As previously observed by (Kouraev et al., 2005), a good agreement between altimetry-derived and in situ water levels is found. For the new orbit, the closest altimetry station is located around 100 km from Salekhard. As a consequence, the derived water levels do not agree as well as the ones from the old orbit.

## 4.3 Monthly maps of water levels

The monthly flood maps indicate that the flood period in the Lower Ob' generally ranges from June to October, whereas low-water period ranges from December to April, during the winter with a maximum occurring generally in July due to the snowmelt. Figure 5 shows a series of maps of interpolated water levels over the floodplains at the maximum of inundation for the period 1993–2004. The maximum difference of water levels between high and low waters is lower than 6 m in the Lower Ob' basin. Maximum differences of water levels between high and low stages are observed along the river stream. Large variability both in spatial patterns and interannual variability are also revealed by the series of maps.

The surface water interannual variability was compared with river discharges and precipitation. The annual maximum surface water storage, i.e., the difference between

HESSD

7, 6647–6676, 2010

### Terrestrial water storage variations in the Lower Ob'

F. Frappart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the annual maximum and minimum surface water volumes, was confronted to the volume of water that has flown in the lower Ob' basin during the year, and to the annual amount of precipitation (expressed in terms of annual average) in the whole Ob' basin. The results are presented in Fig. 6.

5 The annual water volume flowed in the Lower Ob' watershed has been estimated by time-integrated the difference between the monthly river discharges in the outlet of the basin in Salekhard and the monthly river discharges corresponding to the contributions of the Upper and Middle Ob', and of the Irtysh River in Belogorje. We completed the time series of monthly river discharges in Belogorje multiplying the river discharges in  
10 Salekhard by 0.795 as observed by Yang et al. (2004) in its climatology of the Ob' River basin.

Similar time variations can be observed between the annual water volume flow in the Lower Ob', the maximum storage and the average precipitation.

#### 4.4 Water volume variations

15 Using equations from Sect. 3.2 to 3.5 , Fig. 7 presents the time series of volume of TWS, and of the water volume stored in each compartment of the land water cycle, i.e., the surface water, the snow, the water contained in the root zone, and the sum of groundwater and permafrost over 2003–2004, i.e., the overlapping period of the different datasets.

20 These TWS and snow times series exhibit a similar temporal evolution with maxima occurring during winter. The soil moisture only represents a small fraction of the TWS and peaks when the snow starts melting in April or May. The peak of surface water appears during the flood period, in spring/summer, with maximum in June. It is caused by both the local snowmelt and snowmelt from upstream locations (Papa et al., 2007),  
25 and large rainfall occurring between May and September. When the flood decreases, the recharge of the groundwater reservoir begins and lasts until mid-winter and the beginning of the snowmelt.

---

## Terrestrial water storage variations in the Lower Ob'

F. Frappart et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The spatial pattern of the TWS and of the different components of the land water cycle is presented in Fig. 8 in terms of amplitude of the annual cycle over 2003–2004. The annual amplitude of TWS presents very low variations in the Lower Ob' basin, from 50 in the South-East to 70 mm in the north North-West. The snow annual amplitude map presents a meridian gradient and largest amplitudes on the western part of the basin (up to 50 mm). Both spatial pattern and amplitudes are similar to those can be found in climatologies (see Frappart et al., 2010). RZ has a very low annual amplitude, except in the eastern part of the basin with value reaching 25 mm. The map of annual amplitude of the groundwater and permafrost components exhibits a low signal in the whole basin, except just below the junction between the Ob and the Itysh where it reaches 30 mm, where the permafrost is sporadic. The recharge of the aquifer of the lower Ob' basin seems to be mainly related to the percolation of surface water in the large floodplains of the lower Ob' basin rather than the melt of the snow present in the basin. The recharge process seems to be strongly related to the fluxes of water coming from the Middle Ob' and the Irtysh. Permafrost acts as a barrier to the groundwater recharge, increasing the surface runoff and diminishing the subsurface, low annual amplitudes of the annual cycle orz and GW+P are consistent with the presence of sporadic and discontinuous permafrost in the Lower Ob' basin.

## 5 Conclusions

In this study, we estimated surface water extent and storage variations in the Lower Ob' for the 1993–2004 period. The combined use of altimetric water level observations from T/P and inundation patterns derived from multisatellite information to determine water volume variations provides valuable information on the inundation dynamics of river floodplains in the Arctic drainage basins. These surface water volume variations exhibit similar interannual variations as water flowed in the lower Ob' basin and precipitation Seasonal and interannual variabilities are consistent with precipitation and rivers discharges. Monitoring the variability of both flood extent and storage is crucial

### Terrestrial water storage variations in the Lower Ob'

F. Frappart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to better the relationships between snow, flood and surface runoff. These results are also of great interest for the preparation of the future wide swath altimetry mission SWOT (Surface Water and Ocean Topography) that will measure 2D water levels over a 120 km swath.

5 The complementarity between remote sensing derived hydrological products (altimetry, imagery and gravimetry from space) and hydrological modelling was used to determine the time variations of the water stored in the aquifers of the Lower Ob' watershed over 2003–2004. These preliminary results have important implications for the better understanding of the pan-Arctic hydrology, and especially the possible role of the  
10 groundwater in the increase in river discharge to the Arctic Ocean (Smith et al., 2007; Ye et al., 2009). Further work has to be done to investigate the possible increase of subsurface storage in the watersheds covered with permafrost by extending the common availability of these datasets to a longer time-period.

15 *Acknowledgements.* This work was partly supported by the foundation STAE in the framework of the CYMENT project and by NASA's NEWS Grant NNDX7AO90E managed by Jared K. Entin.



The publication of this article is financed by CNRS-INSU.

---

**Terrestrial water storage variations in the Lower Ob'**

F. Frappart et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## References

- Adler, R. F., Huffman, G. J., Chang, A., Ferraro, R., Xie, P., Janowiak, J., Rudolf, B., Schneider, U., Curtis, S., Bolvin, D., Gruber, A., Susskind, J., and Arkin P.: The Version 2 Global Precipitation Climatology Project (GPCP) monthly precipitation analysis (1979-present), *J. Hydrometeorol.*, 4, 1147–1167, doi:10.1175/1525-7541(2003)004<1147:TVGPCP>2.0.CO;2, 2003.
- ArcticRIMS: A Regional, Integrated Hydrological Monitoring System for the Pan-Arctic Land Mass <http://www.watsys.sr.unh.edu/arctic/RIMS/>, last access: October 2009, 2003.
- Armstrong, R. L. and Brodzik, M. J.: Northern Hemisphere EASE-Grid weekly snow cover and sea ice extent version 3., Boulder, Colorado USA, National Snow and Ice Data Center, Digital media, (see <http://nsidc.org/data/nsidc-0046.html>), 2005.
- Bettadpur, S.: CSR level-2 processing standards document for level-2 product release 0004, GRACE 327–742, Rev. 3.1, 18 pp., 2007.
- Birkett, C. M.: Contribution of the TOPEX NASA radar altimeter to the global monitoring of large rivers and wetlands, *Water Resour. Res.*, 34(5), 1223–1239, doi:10.1029/98WR00124, 1998.
- Brown, J., Ferrians Jr., O. J., Heginbottom, J. A., and Melnikov, E. S.: Circum-Arctic map of the permafrost and ground ice conditions, National Snow and Ice Data Center/World Data Center for Glaciology, Digital Media, Boulder, CO (see <http://nsidc.org/data/ggd318.html>), revised February 2001.
- Bullock, A. and Acreman, M.: The role of wetlands in the hydrological cycle, *Hydrol. Earth Syst. Sci.*, 7, 358–389, doi:10.5194/hess-7-358-2003, 2003.
- Decharme, B., Douville, H., Prigent, C., Papa, F., and Aires, F.: A new river flooding scheme for global climate applications: Off-line validation over South America, *J. Geophys. Res.*, 113, D11110, doi:10.1029/2007JD009376, 2008.
- Döll, P., Kaspar, F., and Lehner, B.: A global hydrological model for deriving water availability indicators: Model tuning and validation, *J. Hydrol.*, 270, 105–134, 2003.
- Frappart, F., Martinez, J.-M., Seyler, F., León, J.G., and Cazenave, A.: Floodplain water storage in the Negro River basin estimated from microwave remote sensing of inundation area and water levels, *Remote Sens. Environ.*, 99, 387–399, doi:10.1016/j.rse.2005.08.016, 2005.
- Frappart, F., Calmant, S., Cauhopé, M., Seyler F., and Cazenave A.: Preliminary results of ENVISAT RA-2 derived water levels validation over the Amazon basin, *Remote Sens. Environ.*,

## Terrestrial water storage variations in the Lower Ob'

F. Frappart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Terrestrial water storage variations in the Lower Ob'

F. Frappart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



100, 252–264, doi:10.1016/j.rse.2005.10.027, 2006a.

Frappart, F., Ramillien, G., Biancamaria, S., Mognard, N. M., and Cazenave, A.: Evolution of high-latitude snow mass derived from the GRACE gravimetry mission (2002–2004), *Geophys. Res. Lett.*, 33, L02501, doi:10.1029/2005GL024778, 2006b.

5 Frappart, F., Do Minh, K., L'Hermitte, J., Cazenave, A., Ramillien, G., LeToan, T., and Mognard-Campbell, N.: Water volume change in the lower Mekong basin from satellite altimetry and imagery data, *Geophys. J. Int.*, 167(2), 570–584, doi:10.1111/j.1365-246X.2006.03184.x, 2006c.

10 Frappart, F., Papa, F., Famiglietti, J. S., Prigent, C., Rossow, W. B., and Seyler, F.: Interannual variations of river water storage from a multiple satellite approach: a case study for the Rio Negro River basin, *J. Geophys. Res.*, 113, D21104, doi:10.1029/2007JD009438, 2008.

Frappart, F., Ramillien, G., and Famiglietti, J. S.: Water balance of the Arctic drainage system using GRACE gravimetry products, *Int. J. Remote Sens.*, doi:10.1080/01431160903474954, in press, 2010.

15 Grippa, M., Mognard, N. M., and Le Toan, T.: Comparison between the interannual variability of snow parameters derived from SSM/I and the Ob river discharge, *Remote Sens. Environ.*, 98, 35–44, 2005.

Hall, D. K., Kelly, R. E. J., Riggs, G. A., Chang, A. T. C., and Foster, J. L.: Assessment of the relative accuracy of hemispheric-scale snow-cover maps, *Ann. Glaciol.*, 34(1), 24–30, 2002.

20 Kane, D. L.: The impact of Arctic hydrologic perturbations on Arctic ecosystems induced by climate change, in *Global Change and Arctic Terrestrial Ecosystems*, *Ecol. Stud.*, 124, 63–81, edited by: Oerchel, W. C., Callaghan, T., Elling, H., et al., 61–83, Springer, New York, 1997.

25 Kouraev, A. V., Zakharova, E. A., Samain, O., Mognard, N. M., and Cazenave, A.: Ob' River discharge from Topex-Poseidon satellite altimetry (1992–2002), *Remote Sens. Environ.*, 93, 238–245, 2005.

Leblanc, M. J., Tregoning, P., Ramillien, G., Tweed, S. O., and Fakes, A.: Basin-scale, integrated observations of the early 21st century multiyear drought in southeast Australia, *Water Resour. Res.*, 45, W04408, doi:10.1029/2008WR007333, 2009.

30 Maltby, E.: Wetland management goals: Wise use and conservation, *Landscape Urban Plann.*, 20, 9–18, 1991.

- de Marsily, G.: Eaux continentales, C. R. Geosci., 337, 1–2, 2005.
- Massom, R: Satellite remote sensing of polar snow and ice: present status and future directions, Polar Res., 31(1), 99–114, 1995.
- Milly, P. C. D. and Shmakin, A. B.: Global modeling of land water and energy balances: 1. The Land Dynamics (LaD) model, J. Hydrometeorol., 3, 283–299, 2002.
- Muskett, R. R. and Romanovsky, V. E.: Groundwater storage changes in arctic permafrost watersheds from GRACE and in situ measurements, Environ. Res. Lett., 4(4), 045009, doi:10.1088/1748-9326/4/4/045009, 2010.
- Organisation for Economic Cooperation and Development: Guidelines for Aid Agencies for Improved Conservation and Sustainable Use of Tropical and Sub-tropical Wetlands, Guidelines Aid Environ., 9, Paris, 69 pp., 1996.
- Papa, F., Prigent, C., and Rossow, W. B.: Ob' River flood inundations from satellite observations: a relationship with winter snow parameters and river runoff, J. Geophys. Res., 112, D18103, doi:10.1029/2007JD008451, 2007.
- Papa, F., Prigent, C., and Rossow, W. B.: Monitoring Flood and Discharge Variations in the Large Siberian Rivers From a Multi-Satellite Technique, Surv. Geophys., 29, 297–317, 2008a.
- Papa, F., Güntner, A., Frappart, F., Prigent, C., and Rossow, W. B.: Variations of surface water extent and water storage in large river basins: A comparison of different global data sources, Geophys. Res. Lett., 35, L11401, doi:10.1029/2008GL033857, 2008.
- Papa, F., Prigent, C., Aires, F., Jimenez, C., Rossow, W. B., and Matthews, E.: Interannual variability of surface water extent at global scale, J. Geophys. Res., 115, D12111, doi:10.1029/2009JD012674, 2010.
- Peterson, B. J., Holmes, R. M., McClelland, J. W., Vörösmarty, C. J., Lammers, R. B., and Shiklomanov, A. I.: Increasing river discharge to the Arctic Ocean, Science, 298, 2171–2173, 2002.
- Prigent, C., Papa, F., Aires, F., Rossow, W. B., and Matthews, E.: Global inundation dynamics inferred from multiple satellite observations, 1993–2000, J. Geophys. Res., 112, D12107, doi:10.1029/2006JD007847, 2007.
- Ramillien, G., Frappart, F., Cazenave, A., and Güntner, A.: Time variations of the land water storage from an inversion of 2 years of GRACE geoids, Earth Planet. Sc. Lett., 235, 283–301, doi:10.1016/j.epsl.2005.04.005, 2005.

---

**Terrestrial water storage variations in the Lower Ob'**F. Frappart et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Terrestrial water storage variations in the Lower Ob'

F. Frappart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Ramillien, G., Famiglietti, J. S., and Wahr, J.: Detection of continental hydrology and glaciology signals from GRACE: A review, *Surv. Geophys.*, 29(4–5), 361–374, doi:10.1007/s10712-008-9048-9, 2008.
- 5 Ringeval, B., de Noblet-Ducoudré, N., Ciais, P., Bousquet, P., Prigent, C., Papa, F., and Rossow, W. B.: An attempt to quantify the impact of changes in wetland extent on methane emissions on the seasonal and interannual time scales, *Global Biogeochem. Cy.*, 24, GB2003, doi:10.1029/2008GB003354, 2010.
- Robinson, D. A. and Frei, A.: Seasonal variability of northern hemisphere snow extent using visible data, *The Professional Geographer*, 52(2), 307–315, 2000.
- 10 Roddel, M., Velicogna, I., and Famiglietti, J. S.: Satellite-based estimates of groundwater depletion in India, *Nature*, 460, 999–1003, doi:10.1038/nature08238, 2009.
- Seidel, K. and Martinec, J.: Remote sensing in snow hydrology: runoff modelling, effect of climate change, Springer Praxis books – Geophys. Sci., 150 pp., 2004.
- Serreze, M. C., Walsh, J. E., Chapin III, F. S., Osterkamp, T., Dyurgerov, M., Romanovsky, V., 15 Oechel, W. C., Morison, J., Zhang, T., and Barry R. G.: Observational evidence of recent change in the northern high-latitude environment *Climatic Change*, 46, 159–207, 2000.
- Schmidt, R., Flechtner, F., Meyer, U., Neumayer, K.-H., Dahle, Ch., Koenig, R., and Kusche, J.: Hydrological Signals Observed by the GRACE Satellites, *Surv. Geophys.*, 29, 319–334, doi:10.1007/s10712-008-9033-3, 2008.
- 20 Shiklomanov, I. A., Lammers, R. B., Peterson, B. J., and Vörösmarty, C. J.: The dynamics of river water inflow to the Arctic Ocean. In the freshwater budget of the Arctic Ocean, proc. of the NATO advanced research workshop, 281–296, Kluwer Academic Publishers, Norwell, 2000.
- Smith, L. C.: Satellite remote sensing of river inundation area, stage and processes: a review, *Hydrol. Process.*, 11, 1427–1439, 1997.
- 25 Smith, L. C., Sheng, Y., MacDonald, G. M., and Hinzman, L. D.: Disappearing Arctic lakes, *Science*, 308, 1429, doi:10.1126/science.1108142, 2005.
- Smith, L. C., Pavelsky, T. M., MacDonald, G. M., Shiklomanov, A. I., and Lammers, R. B.: Rising minimum daily flows in northern Eurasian rivers: a growing influence of groundwater in the high-latitude hydrologic cycle, *J. Geophys. Res.*, 112, G04S47, doi:10.1029/2006JG000327, 30 2007.
- Stocker, T. F. and Raible, C. C.: Water cycle shifts gear, *Nature*, 434, 1560–1563, 2005.



# HESSD

7, 6647–6676, 2010

## Terrestrial water storage variations in the Lower Ob'

F. Frappart et al.

**Table 1.** Comparisons between the gauge records from Salekhard and the water levels derived from T/P measurements.

	RMS (m)	Correlation
In situ/TP 112 (old orbit)	0.89	0.89
In situ/TP 187 (old orbit)	1.32	0.84
In situ/TP 214 (new orbit)	1.34	0.69

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Location of T/P altimetry stations in the Lower Ob basin



**Fig. 1.** Map of Lower Ob' basin and locations of the Topex/Poseidon water level stations. The line with stars shows the delineation of the watershed.

**Terrestrial water storage variations in the Lower Ob'**

F. Frappart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**Terrestrial water storage variations in the Lower Ob' River Basin**


F. Frappart et al.



**Fig. 2.** Monthly time variations of the inundated areas ( $\text{km}^2$ ) in the Lower Ob' River Basin from 1993 to 2004.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)





**Fig. 3.** (a) the 1993–2004 mean fractional inundation at yearly maximum (%) (b) the average month of the year (1993–2004) when the fractional inundation is maximum (in month).

## Terrestrial water storage variations in the Lower Ob'

F. Frappart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 4.** Time series of water levels at Salekhard from in situ gauge (black), T/P altimetry stations.

## Terrestrial water storage variations in the Lower Ob'

F. Frappart et al.

<a href="#">Title Page</a>	
<a href="#">Abstract</a>	<a href="#">Introduction</a>
<a href="#">Conclusions</a>	<a href="#">References</a>
<a href="#">Tables</a>	<a href="#">Figures</a>
<a href="#">◀</a>	<a href="#">▶</a>
<a href="#">◀</a>	<a href="#">▶</a>
<a href="#">Back</a>	<a href="#">Close</a>
<a href="#">Full Screen / Esc</a>	
<a href="#">Printer-friendly Version</a>	
<a href="#">Interactive Discussion</a>	






Fig. 5. Water level maps during the peak flood (July) for 1993–2004.

Terrestrial water storage variations in the Lower Ob'

F. Frappart et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion




**Terrestrial water storage variations in the Lower Ob'**

F. Frappart et al.



**Fig. 6.** Time series of annual maximum of surface water storage volume (light green), annual water volume flow in the Lower Ob' basin (blue), and annual water volume of precipitation.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




**Fig. 7.** Monthly variations of TWS volume (blue), surface water volume from T/P radar altimetry and multisatellite inundation dataset (black), soil moisture from hydrological model outputs (green), snow derived from GRACE measurements (yellow), sum of groundwater and permafrost by difference between TWS and the other compartments (red). Units:  $\text{km}^3$ .

## Terrestrial water storage variations in the Lower Ob'

F. Frappart et al.

<a href="#">Title Page</a>	
<a href="#">Abstract</a>	<a href="#">Introduction</a>
<a href="#">Conclusions</a>	<a href="#">References</a>
<a href="#">Tables</a>	<a href="#">Figures</a>
<a href="#">◀</a>	<a href="#">▶</a>
<a href="#">◀</a>	<a href="#">▶</a>
<a href="#">Back</a>	<a href="#">Close</a>
<a href="#">Full Screen / Esc</a>	
<a href="#">Printer-friendly Version</a>	
<a href="#">Interactive Discussion</a>	





**Fig. 8.** Amplitude of the annual cycle over 2003–2004: **(a)** TWS from GRACE, **(b)** Snow Water Equivalent (SWE) from GRACE, **(c)** Water in the Root zone from hydrological models, **(d)** Groundwater and permafrost. Units: mm.

**Terrestrial water storage variations in the Lower Ob'**

F. Frappart et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

