

This discussion paper is/has been under review for the journal Hydrology and Earth System Sciences (HESS). Please refer to the corresponding final paper in HESS if available.

Mapping surface soil moisture over the Gourma mesoscale site (Mali) by using ENVISAT ASAR data

F. Baup¹, E. Mougin², P. de Rosnay³, P. Hiernaux², F. Frappart², P. L. Frison⁴, and M. Zribi¹

¹CESBIO (Centre d'Etudes Spatiales de la Biosphère, UMR 5126, UPS-CNRS-CNES-IRD), 18 avenue Edouard Belin 31401 Toulouse cedex 9, France

²LMTG (Laboratoire des Mécanismes et Transferts en Géologie, UMR 5563 CNRS-UPS-IRD) 14 avenue Edouard Belin 31400 Toulouse, France

³ECMWF (European Centre for Medium-Range Forecasts), Reading, Berkshire, UK

⁴Laboratoire des Géomatériaux, Université Marne La Vallée, Institut Francilien des Sciences Appliquées, 5 bd Descartes, 77454 Marne la Vallée Cedex 2, France

Received: 23 July 2010 – Accepted: 13 August 2010 – Published: 28 September 2010

Correspondence to: F. Baup (frederic.baup@cesbio.cnes.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

HESSD

7, 7417–7450, 2010

**Mapping soil
moisture in Sahel
using ASAR data**

F. Baup et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

The potentialities of ENVISAT ASAR (Advanced Synthetic Aperture Radar) multi-angle data for mapping surface soil moisture (SSM) in Sahelian rangelands are investigated at medium scale (30 000 km²). The Wide Swath data are selected to take advantage of their high temporal repetitivity (about 8 days at the considered scale) associated to a moderate spatial resolution (150 m). In the continuity of previous studies conducted at a local scale in the same region, SSM maps are here processed over the whole AMMA Gourma mesoscale site at 1 km resolution scale. Overall, the generated maps are found to be in good agreement with field data, EPSAT-SG (Estimation des Pluies par SATellite – Second Generation) rainfall estimates and ERS (European Remote Sensing) Wind Scatterometer (WSC) SSM products. The present study shows that the spatial pattern of SSM can be realistically estimated at a kilometric scale. The resulting SSM maps are expected to provide valuable information for initialisation of land surface models and the estimation of the spatial distribution of radiative fluxes. Particularly, SSM maps could help to desaggregate low-resolution products such as those derived from WSC data.

1 Introduction

Soil moisture monitoring over the Sahel is a major issue of the agenda of the AMMA (African Monsoon Multidisciplinary Analysis) project which aims at a better understanding of the West African Monsoon and its physical, chemical and biological environments (GEWEX-news, 2006). Monitoring the spatio-temporal variability of soil moisture is also needed to better understand soil–vegetation–atmosphere interactions and to address the role of soil moisture on West African monsoon dynamics (e.g. Monteny et al., 1997; Clark et al., 2004; Taylor and Ellis, 2006). Moreover, surface soil moisture (SSM) is a key parameter for surface processes which govern the incident solar energy partitioning between surface heat and water fluxes (Taylor et al., 2006).

HESSD

7, 7417–7450, 2010

Mapping soil moisture in Sahel using ASAR data

F. Baup et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Mapping soil moisture in Sahel using ASAR data

F. Baup et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

For over 20 years, microwave remote sensing has been demonstrating capabilities to estimate SSM under a large variety of topographic and vegetation cover conditions. Several large-scale field experiments that included aircraft microwave radiometric measures were conducted within the framework of HAPEX-Sahel, FIFE and Monsoon'90 (Schmugge et al., 1992). So far, passive and active sensors have been used to map SSM over semi arid areas by using C- and L-band frequencies (e.g. Chanzy et al., 1997; Frison et al., 1998; Wagner and Scipal, 2000; Jarlan et al., 2002; Njoku et al., 2003; Zribi et al., 2008). These independent multi-frequency and multi-resolution sensors offer complementary information due to their different interactions with the surface components (soil and vegetation). L-band data are less influenced by the vegetation contribution due to a higher wave penetration into the vegetation and thus are more correlated with SSM. Although C-band data are more influenced by the vegetation canopy, they are also strongly linked to SSM in arid regions where the vegetation cover is low (Ulaby et al., 1982; Wagner and Scipal, 2000; Zine et al., 2005; Zribi et al., 2008).

The last generation of spaceborne radars, such as the Advanced Synthetic Aperture Radar (ASAR) onboard ENVISAT or RADARSAT, provides images combining a high spatial resolution (≤ 150 m) and a large spatial cover (up to 500 km) at C-band frequency. Particularly, the ASAR Wide Swath mode provides images with a high temporal sampling (3–4 days at local scale and 8 days at meso-scale) associated to a moderate spatial resolution (150 m). Others ASAR modes such as Alternating polarisation or Image modes could also be used for improving SSM estimates. However, for the time being, their use is constrained by their low spatial coverage that is lower than 100 km (ENVISAT ASAR product Handbook, 2007).

SSM estimation can take advantage of the multi-angular capabilities of SAR antenna. At low incidence angles and for low vegetation density (Leaf Area Index –LAI– < 2), the interferences due to the green vegetation can be either neglected or simply taken into account (Tansey et al., 1999; Moran et al., 2000). At higher incidence angles, the vegetation effects on the radar signal can be corrected using the angular properties of the backscattering coefficient (Wagner et al., 2000; Baup et al., 2007a).

Finally, the determination of SSM was found to be possible from ASAR data, for values higher than 4% (m^3/m^3). Below this threshold, the radar signal instability (± 0.6 dB for ENVISAT/ASAR) combined with unknown surface temporal changes prevent from an accurate SSM estimation (Moran et al., 2000; Mattia et al., 2006; Baup et al., 2007a; Zribi et al., 2008).

In the framework of the AMMA project, meso-scale sites have been instrumented in Mali, Niger and Benin, providing ground data along the latitudinal gradient between Sahelian and Sudanian bioclimatic regions (Lebel et al., 2009; Cappelaere et al., 2009; Mougouin et al., 2009; de Rosnay et al., 2009). The objective of the present study is to up-scale a local approach for estimating SSM (Baup et al., 2007a) to the Malian meso-scale site in the Gourma region by taking into account soil type and vegetation cover. To this end, we use HH-polarized ENVISAT/ASAR Wide Swath data which combine high spatial cover, moderate spatial resolution and high temporal sampling. The paper is organised as follows: the data used (ground and satellite) and the overall methodology are presented in Sect. 2. Then, the performance of the algorithm is evaluated using EPSAT-SG (Estimation des Pluies par SATellite – Second Generation) rainfall products, European Remote Sensing (ERS) wind scatterometer SSM products and local raingauges data (Sect. 3). Conclusions on the relevance of ASAR SSM maps and perspectives are given in Sect. 4.

2 Data and methods

2.1 The study site

The Gourma mesoscale site (1°W – 2°W , 14.5°N – 17.5°N) is entirely located within the Sahel bioclimatic zone (Mougouin et al., 2009). This region, mainly pastoral, is bracketed by the 150 and 500 mm isohyets. The rain distribution is strictly mono-modal with rainfall starting in June and ending in September with a maximum in August. This short rainy season is followed by a long dry season during which there is no green

Mapping soil moisture in Sahel using ASAR data

F. Baup et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



vegetation except for rare perennial herbaceous and the foliage of some of scattered trees and shrubs (Hiernaux et al., 2009). Bare soils are thus very much exposed during most of the dry season with contrasted colors, textures and roughness (Fig. 1a–c). A high cover of perennial vegetation is observed over the Bamba site as illustrated in Fig. 1c.

During the wet season, rangeland vegetation is composed of a low herbaceous layer dominated by annuals and a sparse woody plant population (mean <5% cover). Annual herbs growth is governed by soil moisture regime itself driven by the pattern and magnitude of the rain events, and by run-off function of rain intensity, topography and soil texture (Casenave and Valentin, 1989). Annual herbs germinate with the first marked rains, in June or July, and unless the plants wilt before maturity owing to a lack of rainfall, the senescence coincides approximately with the end of the rainy season, several days after the drying up of soils.

2.2 Ground data

2.3 SSM measurements

The objectives of the soil moisture stations network are to collect information about the spatio-temporal variation of soil moisture over the mesoscale window at different depths (de Rosnay et al., 2009). From April to December 2005, soil moisture was automatically recorded at 15 or 30 min time step in three sandy local sites, namely Agoufou (15.3° N, 1.5° W), Bamba (17.1° N, 1.4° W), Ebang Mallam (15.4° N, 1.3° W) and one hard pan (rocky) site: Eguerit (15.5° N, 1.4° W) (Fig. 2). Only measurements recorded at 5 cm depth are considered in accordance with the microwave soil penetration depth (Ulaby et al., 1981).

In all sites, soil moisture is measured by reflectometers (Campbell Scientific CS616, Campbell Scientific, 2002). Sensors are calibrated by using in situ gravimetric measurements and estimation of soil bulk density (de Rosnay et al., 2009). In this study, SSM is expressed in volumetric water content (m^3/m^3) with an accuracy of 2.5%

Mapping soil moisture in Sahel using ASAR data

F. Baup et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(Campbell Scientific, 2002). Figure 3a and b illustrate the temporal variation of SSM for the Agoufou sandy- and Eguerit rocky-sites.

During the dry season, SSM is close to 0.05% and reaches values as high as 18% at Agoufou during the wet season. At the Eguerit site, SSM values range between 3.5% and 15%. Local SSM measured by automatic stations typically range between 0.05 and 24% over the whole mesoscale site (de Rosnay et al., 2009).

2.3.1 Rainfall measurements

Rainfalls are recorded by 8 automatic rain gauges distributed along the latitudinal climatic gradient (Fig. 2). Rainfall data are summed over different time step depending of the site (Table 1). In the following, only the Agoufou (Ag) and the Tin Tadeini (TT) station data are detailed but all the stations are nevertheless used in the analyses.

2.3.2 Soil map

Over the Gourma region, nine soil types are mapped by visual interpretation of a mosaic of Landsat TM images at a resolution of 30 m and systematic field observations (Mougin et al., 2009). Soil surface types are defined according to topography, terrain morphology, top soil texture and soil surface features. Soil types (initially divided in 9 classes) are grouped in 4 dominant classes (clay, mixed, sandy and rocky) and presented in Fig. 4 at mesoscale and supersite scales. For best clarity, only main classes are presented for the mesoscale site. Each class is represented by its cover percentage varying from 0 to 100%. The sandy and rocky soil classes contribute more than 90% of the study site.

Mapping soil moisture in Sahel using ASAR data

F. Baup et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.4 Satellite data

ENVISAT ASAR

The ENVISAT satellite was launched by ESA (European Space Agency) on 1 March 2002. The ASAR instrument is a multi-mode sensor which operates at C-band (5.3 GHz) at several polarizations (HH, VV, HV and VH), incidence angles, and spatial/radiometric resolutions depending on the functioning mode (Desnos et al., 1999). At the C-band frequency, atmospheric perturbations, excepted originating from strong precipitation systems, can be considered negligible (Ulaby et al., 1981). ASAR data are provided by the ScanSAR Wide Swath (WS) mode at HH polarization, at a spatial resolution of 150 m and incidence angles ranged between 16° and 43° (ENVISAT handbook, 2004). The confidence interval is estimated at ± 0.6 dB at 1σ (where σ denotes the standard deviation), over the 1×1 km² sites (Baup et al., 2007b).

From July to December 2005, 23 images were acquired at various incidence angles over the AMMA mesoscale site. Only 7 images were analyzed: 5 for the wet season and 2 for the dry season (Table 2). The 16 other images have been acquired during the dry season and characterized by very low and almost constant SSM. The 7 selected images are calibrated with the B.E.S.T. (Basic ENVISAT SAR Toolbox) software algorithm described in Laur et al. (1998). Then, their geolocation is improved by using ground control points taken from a Landsat TM image at a spatial resolution of 30 m. Mean offsets of 1 pixel (75 m) in latitude and 7 pixels (525 m) in longitude have been observed and corrected for.

ERS Scatterometer data

The ERS WSC was initially designed to measure wind speed and direction at the sea surface. The instrument consists of three antennas, which transmit radar beams that are pointed 45° forward, 45° sideways, and 45° backward with respect to the satellite nadir. The incidence angle θ varies over the instrument swath from 18° to 47° for

HESSD

7, 7417–7450, 2010

Mapping soil moisture in Sahel using ASAR data

F. Baup et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the midbeam antenna and from 25° to 59° for the fore-beam and aft-beam antennas (Frison and Mougin, 1996). The sensor operates at 5.3 GHz and VV polarization, like the ERS/SAR instrument. Its spatial resolution is around 50 km, and measurements are repeated every three to four days. The data are processed using a grid with elementary 0.25° square cells. The methodology presented in Zribi et al. (2008) enables SSM products to be obtained at regional scale over West Africa. Angular normalisation is applied for each cell and local empirical surface soil moisture inversion functions are applied at larger scale. Vegetation and soil roughness variations are interpreted as a “dry signal” as presented in Zribi et al. (2008). The observed variations of the backscattering signal above the dry signal are attributed to SSM changes.

EPSAT-SG data

EPSAT-SG (Estimation des Pluies par SATellite – Second Generation) precipitation products are generated at LMD (Laboratoire de Météorologie Dynamique) within the framework of AMMA. Satellite and raingauge data are used to produce high resolution rainfall estimations. The EPSAT-SG algorithm is composed of two parts: the computation of rainfall probability and the actual rainfall estimation (Bergès et al., 2005). Rainfall probabilities are computed using a feed forward neural network from MSG (Meteosat Second Generation), multi-spectral images and TRMM PR (Tropical Rainfall Measuring Mission Precipitation Radar) data. For every 15 min MSG image, a rainfall probability map is produced (Bergès and Chopin, 2004). Then, potential rainfall intensity maps are computed merging GPCP-1dd (Global Precipitation Climatology Project at one degree of space resolution) information with the rainfall probability map. The estimated rainfall intensity is the product of the rainfall probability and the potential intensity (Chopin et al., 2005).

The AMMA project provides rain rates within gridded surface over West Africa (from 25° W to 25° E and from 20° N to 5° S) estimated with the EPSAT-SG algorithm at a spatial resolution of 0.1° and a temporal resolution of 30 min from June to September (mid-October in 2005) over the 2004–2006 period (<http://bddamma.ipsl.polytechnique>).

Mapping soil moisture in Sahel using ASAR data

F. Baup et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



fr/). For the need of the study, the rain rates are summed either over 1 or over 5 days.

2.5 Methodology of surface soil moisture mapping

The SSM maps are produced from satellite and ground data in a several steps process (Fig. 5).

1. The proposed methodology consists of an angular-normalization of ASAR images depending on the acquisition angle, soil type and vegetation. The normalization method consists in characterizing the angular variation function of the backscattering coefficient for different soil types (Baup et al., 2007b). Over sandy soils, vegetation effects are taken into account by using an average angular-normalization function (Baup et al., 2007a) that is applied only during the wet season after the start of herbaceous vegetation growth detected here using Normalized Difference Vegetation Index (NDVI) images acquired by the Moderate Resolution Imaging Spectroradiometer (MODIS). Here, a NDVI threshold of 0.14 is retained. The more the vegetation, the more the backscattering signal increases at high incidence angle ($>30^\circ$). In accordance with previous studies (Baup et al., 2007a, b), the angular variation function is used to normalise the radar data at 23° of incidence angle. At lower incidence angles, no significant correction is applied. Over rocky erosion and hard pan surfaces, vegetation effects are neglected since the vegetation cover remain generally very low, even during the wet season (Hiernaux et al., 2009).
2. Normalized images are filtered to reduce speckle effects using the spatial GAMMA filter of the ENVI 4.2 software (ENVI, 1998). The image resolution is thus degraded from 150 m to 1 km as a result of the filter application.
3. A change detection method is applied to reduce roughness effects. It consists of processing the difference between one dry season image chosen as a reference and the wet season image of interest (IOI) (Wagner et al., 2000; Zribi et al., 2006).

Mapping soil moisture in Sahel using ASAR data

F. Baup et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Mapping soil moisture in Sahel using ASAR data

F. Baup et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The reference image is selected to be representative of the soil roughness when no vegetation is present (here the 20 December). In situ measurements show that surface soil roughness does not strongly vary between dry and wet seasons over sandy soils. The instrument used is a 2 m long needle like profilometer with a 1 cm horizontal sampling distance, allowing the estimation of standard deviation of roughness heights and the autocorrelation length. Measurements were made in August 2005 (wet season) and February 2006 (dry season) at the Agoufou site (Table 3) with a total of 60 and 16 profiles for the wet and dry seasons, respectively. Results show that the roughness parameters for sandy soils do not significantly vary between the two contrasted seasons. Since the surface of sandy soils are the most sensitive to meteorological conditions (wind, rain) and animal trampling, it is assumed constant roughness values over time for the other soil types as well.

4. A mask of the Niger River is built from the soil map and applied to the whole set of ASAR images. Masks are also applied over mixed surfaces for which the relation between SSM and σ^0 is unknown.
5. Finally, SSM maps are established at kilometric scale over the mesoscale window by applying empirical inversion functions estimated for sandy and rocky soils. Because of residual negative SSM values, a minimal threshold of 0.5% is imposed.

3 Surface Soil Moisture Maps

3.1 Relationships between SSM and normalized σ^0

Figure 6 presents the temporal variation of the normalized backscattering coefficient for the two main soil types at the sandy Agoufou (Fig. 6a) and hard pan (rocky) Eguerit (Fig. 6b) sites.

As already indicated in Baup et al. (2007a, b), the backscattering coefficient, σ^0 , over the Agoufou sandy site displays a strong seasonal dynamics (>5dB), due both to the growth of annual herbs and to high SSM values reached during the wet season. In contrast, the seasonal dynamics of σ^0 recorded on the Eguerit rocky site is much weaker (<2 dB) due to both a low vegetation cover (Hiernaux et al., 2009) and overall low SSM.

Correlation coefficients between σ^0 and SSM for the four considered local sites are presented in Table 4. The highest values are observed for the sandy Agoufou and Bamba sites ($r \sim 0.8$). For this latter, only the data acquired during the wet season are considered as the backscattered radar signal is dominated by the response of perennial vegetation during the dry season (see Fig. 1c). The third sandy site, Ebang Mallam, exhibits a lower correlation coefficient ($r = 0.48$) due to a very high spatial heterogeneity in terms of vegetation cover and soils.

For the rocky site, Eguerit, backscattering coefficients are weakly correlated with SSM ($r = 0.53$), even for descending passes which show the highest σ^0 dynamics. This could be explained by two main reasons. The first one is associated to the difficulties of measuring soil moisture in such a medium composed of gravels and rocks for which the used SSM probes are not specifically designed. The second difficulty originates from the high spatial heterogeneity of soil texture within the site due to the presence of sparsely distributed sandy-loamy sheets. This justifies the exclusion of mixed soil types in the SSM mapping process. Finally, the soil and land cover types excluded from the analysis represent 28% of the mesoscale site; this includes the Niger River and seasonally flooded acacia forests.

3.2 SSM maps

Following the methodology described in Sect. 2.4, SSM maps are generated at meso-scale. On the resulting images, black pixels result from either the applied mask or the lack of radar data (see for example Fig. 8). At such scale, the evaluation of SSM maps

Mapping soil moisture in Sahel using ASAR data

F. Baup et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



is not a straightforward exercise. Here, the ASAR SSM maps are interpreted with the help of EPSAT-SG rainfall estimations, WSC soil moisture products and raingauges data. Because they differ in spatial resolution (1 km, 10 km and 25 km, respectively), their spatial patterns are compared in a first analysis.

Figure 7 shows the generated SSM map corresponding to the beginning of the 2005 rainy season (16 July, DoY 197). Patches of high SSM to the North of Tin Tadeini can be related to the precipitations that fell on the same day (a total of 2 mm estimated from EPSAT-SG) and to rain events during the previous 4 days totalling 10 to 12 mm (Fig. 7). Raingauge records at Tin Tadeini (6 mm) confirm the occurrence of rainfall events. The large strip of high SSM to the South of Agoufou can be related to 25 mm cumulative estimated rainfalls (EPSAT-SG) over the 4 previous days confirmed by the raingauge records in Agoufou (17 mm). Overall, most of the SSM values are still below the noise threshold (a SSM of 4%). This can be attributed both to speckle effects which are not completely removed at 1 km scale, and to local higher SSM values associated to topographic effects (depressions, small dunes) and to different soil textures. No WSC soil moisture product is available at this date over the whole meso-scale site.

During the wet season (Figs. 8 and 9), soil moisture maps of sandy soils display different patterns. On 4 August (DoY 216), the SSM map shows an overall good agreement with the daily rainfall EPSAT-SG map, especially in the Northern part (Fig. 8). Particularly, the SSM maps point out the high spatial heterogeneity of SSM and thus of the rainfall events that is not revealed by EPSAT-SG data. Besides, spots of high SSM values located in the South also indicate residual moisture due to past rainfall events supported by the cumulative EPSAT-SG product (about 35 mm of rain cumulated over the previous 4 days). This weak temporal signature of SSM confirms the limit of 4 days to detect the occurrence of rainfall events over sandy soils (Baup et al., 2007b). A high infiltration rate on sandy soils associated to high evaporation rates explain the rapid decrease of SSM at the soil surface. The finer spatial resolution of ASAR data provide information about the geographic distribution of rain events estimated by EPSAT-SG. Local SSM station data are in accordance with satellite products in the Southern part

Mapping soil moisture in Sahel using ASAR data

F. Baup et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

of the image, and with the recorded cumulative rainfall of about 39 mm at Agoufou.

WSC products show a strong agreement with ASAR products but with much higher SSM values estimated over dry areas (between 14.5° N to 16.5° N).

On 20 August (DoY 232), the SSM map is highly contrasted with a large swath of saturated SSM values (20%) in the Southern Gourma that corresponds to the occurrence of a large convective system a few hours before the radar image acquisition (EPSAT-SG product indicates 8 mm and Agoufou local station recorded 21 mm). High and medium SSM values are more scattered in Northern Gourma in relation with rain events that occurred during the previous days (20 mm EPSAT-SG cumulated rains in the North, confirmed by 11 mm recorded at Tin Tadeini over 3 days). The WSC product exhibit high values in the South and in the North-West part of the mesoscale site (up to 20%) in agreement with EPSAT-SG cumulative products.

On 3 September (DoY 246), the SSM map displays local patches that can be related to local rainfalls observed on EPSAT-SG rainfall products during the previous days (about 20 mm along a Northeast to Southwest stripe) (Fig. 9). The Agoufou raingauge station also recorded a rainfall event but of only 2.5 mm during on DoY 244.

On 8 September (DoY 251), the accumulation of medium SSM values in the Southern part of the site can be related to the occurrence of a convective system during the 4 previous days. In accordance with EPSAT-SG maps, no rains are recorded at Tin tadeini nor at Agoufou during the day corresponding to the ASAR acquisition. Local patches of high SSM values can also be observed in Northern and middle of the map. They are attributed to residual moisture in depressions as no rainfall has been monitored.

In contrast, soil moisture patterns derived from WSC data are not in accordance with ASAR and EPSAT products.

Mapping soil moisture in Sahel using ASAR data

F. Baup et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4 Evaluation of SSM ASAR maps

At meso-scale, the distribution function of SSM values drastically changes between the dry and the wet seasons (Fig. 10). Mean (standard deviation) SSM values are about 1% (1.8) and 4% (5.0) for the dry and wet season, respectively. The 19 December (dry season) and the 20 August (wet season) maps exemplify it well (Figs. 7 and 8). During the wet season, higher values reach about 20% in accordance with local automatic measurements. The dry season distribution indicates that SSM values mainly range between 0 and 4% showing the “noise” effect during this period.

Among the different products used in this study, only the ERS WSC ones can be used for a quantitative evaluation of the ASAR SSM maps. Furthermore, both data sets are derived from radar sensors operating at the same frequency (5.3 GHz). To enable a direct comparison, 1 km resolution ASAR maps are aggregated to the WSC spatial resolution (25 km). Then, the comparison is made by considering the ASAR pixels corresponding to at least 95% of a WSC pixel. Figure 11a and b shows the results obtained from SSM maps generated for the driest and wettest dates in 2005 (8 and 20 September, respectively). For each selected pixel, its latitudinal location is also indicated.

For the dry season with no rainfall event, WSC-derived SSM surprisingly range between 3% and 10% (Fig. 11a). The highest SSM values are located in the Northern part of the site where the WSC signal is mainly affected by the Niger River. The 3%–10% SSM dynamic range is explained by the “noise” effect of the radar signal whereas the initial offset of 3% is explained by the inversion method used (Zribi et al., 2009). In contrast, ASAR-derived SSM show a higher stability with only 2% of dynamic range and an offset of about 1%.

Results obtained for the rainy season indicate a high correlation ($r^2=0.91$) between the two data sets despite the small number of samples ($n = 11$) considered which results from both the small number of available ASAR images and to the small size of the Gourma site when compared to the spatial resolution of WSC product (25 km).

HESSD

7, 7417–7450, 2010

Mapping soil moisture in Sahel using ASAR data

F. Baup et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 Concluding remarks

A local empirical methodology inversion of SSM from ENVISAT/ASAR data has been up-scaled at meso-scale for the AMMA Malian site. Resulting soil moisture maps take into account surface roughness, vegetation density and angular variation of σ^0 . They offer both a moderate spatial resolution (1 km) and a large spatial coverage (30 000 km²) contrary to the maps derived from the Alternate Polarization (AP) high resolution (Zribi et al., 2007) and Global Monitoring (GM) low resolution modes (Wagner et al., 2005) that were previously used for that purpose.

During the 2005 wet season, five dates are available characterized by a large soil moisture dynamics. On the whole, there is a good agreement and consistency between ASAR derived SSM maps and EPSAT-SG rainfall fields and local rainfall measurements. Finally, when aggregated to 25 km resolution, the ASAR-derived SSM maps are found to be in a very good agreement with ERS WSC-derived SSM maps generated during the wet season. Furthermore, in contrast to ERS WSC dry season maps which show unrealistic high SSM values, ASAR derived maps exhibit low and constant SSM values during the whole dry season. These results demonstrate the great potentialities of ASAR WS images for mapping and monitoring SSM at moderate resolution (1 km) in semi-arid regions. Such SSM products could be advantageously used to de-aggregate lower resolution images, such as those derived from AMSR, SMOS or ERS WSC data (Wagner et al., 2000; Berger et al., 2002; Njoku et al., 2003).

In the future, temporal SSM monitoring in the Sahel could be improved by combining ASAR Wide Swath data to other existing microwave sensors. Indeed, the ASAR WS temporal resolution of about 8 days is still insufficient to enable fast varying surface variables like soil moisture to be accurately monitored. For instance, the combination with ALOS/PALSAR (L-Band) data should improve the sampling of SSM temporal variations while keeping a good spatial resolution.

Acknowledgements. This work was performed within the framework of the AMMA project. Based on a French initiative, AMMA has been constructed by an international group and is

HESSD

7, 7417–7450, 2010

Mapping soil moisture in Sahel using ASAR data

F. Baup et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



currently funded by large number of agencies, especially from France, the UK, the US and Africa. It has been the beneficiary of a major financial contribution from the European Community's Sixth Framework Research Programme. Detailed information on the scientific coordination and funding is available on the AMMA international web site (<https://www.amma-eu.org/>).

5 The authors thank ESA for providing the ENVISAT data used in the present study (Project ID 443, E. Mougin). The authors are grateful for all the help they received during the field measurement campaigns, especially from their colleagues and collaborators from the national institute for agronomic research in Mali, the "Institut d'Economie Rurale".



10 **INSU**
Institut national des sciences de l'Univers

The publication of this article is financed by CNRS-INSU.

References

15 Baup, F., Mougin, E., de Rosnay, P., Timouk, F., Lavenu, F., and Chênerie, I.: Surface soil moisture estimation over the AMMA sahelian site in Mali using ENVISAT/ASAR data, *Remote Sens. Environ.*, 109, 473–481, 2007a.

Baup, F., Mougin, E., Hiernaux, P., Lopes, A., De Rosnay, P., and Chênerie, I.: Radar signatures of Sahelian surfaces in Mali using ENVISAT-ASAR data. *IEEE T. Geosci. Remote Sens.*, 45, 2354–2363, 2007b.

20 Berger, M., Camps, A., Font, J., Kerr, Y., Miller, J., Johannessen, J., Boutin, J., Drinkwater, M. R., Skou, N., Floury, N., Rast, M., Rebhan, H., and Attema, E.: Measuring Ocean Salinity with ESA's SMOS Mission, *ESA Bulletin* 111, 113f, 2002.

Campbell Scientific: CS616 Water Content Reflectometer, user guide, Issued 6.3.02, 33 pp., 2002.

Bergès, J. C. and Chopin, F.: Satellite rainfall estimation using a feed forward network: improve-

HESSD

7, 7417–7450, 2010

Mapping soil moisture in Sahel using ASAR data

F. Baup et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1998.

GEWEX-news: Global Energy and Water Cycle Experiment, special issue 'AMMA west African monsoon studies are addressing water cycle issues', 16(1), 20 pp., 2006.

5 Hiernaux, P., Cissé, M. I., Diarra, L., and de Leeuw, P. N.: Fluctuations saisonnières de la feuillaison des arbres et des buissons sahéliens. Conséquences pour la quantification des ressources fourragères. *La Revue d'Élevage et de Médecine Vétérinaire des Pays Tropicaux*, 47, 1, 117–125, 1994.

10 Hiernaux, P., Mougin, E., Diarra, L., Soumaguel, N., Lavenu, F., Tracol, Y., Diawara, M., and Jarlan, L.: Rangeland response to rainfall and grazing pressure over two decades: herbaceous growth pattern, production and species composition in the Gourma, Mali, *J. Hydrol.*, 375, 114–127, 2008.

Jarlan, L., Mougin, E., Frison, P. L., Mazzega, P., and Hiernaux, P.: Analysis of ERS wind scatterometer time series over Sahel (Mali), *Remote Sens. Environ.*, 81, 404–415, 2002.

15 Laur, H., Bally, P., Meadows, P., Sanchez, J., Schaettler, B., Lopinto, E., and Esteban, D.: Derivation of the backscattering coefficient σ_0 in ESA SAR products, ESA publication, Document No: ES-TN-RS-PM-HL09 17, 2, Rev. 5d., 53 pp., 1998.

Lebel, T., Cappelaere, B., Galle, S., Hanan, N., Kergoat, L., Levis, S., Vieux, B., Descroix, L., Gosset, M., Mougin, E., Peugeot, C., and Seguis, L.: AMMA-CATCH studies in the Sahelian region of West-Africa: an overview, *J. Hydrol.*, 375, 3–13, 2009.

20 Mattia, F., Satalino, G., Dente, L., and Pasquariello, G.: Using a priori information to improve Soil Moisture Retrieval from ENVISAT ASAR AP data in semiarid regions, *IEEE T. Geosci. Remote Sens.*, 44, 900–912, 2006.

Monteny, B. A., Lhomme, J. P., Chehbouni, A., Troufleau, D., Amadou, M., Sicot, M., Verhoef, A., Galle, S., Said, F., and Lloyd, C. R.: The role of the Sahelian biosphere on the water and the CO₂ cycle during the HAPEX-Sahel Experiment, *J. Hydrol.*, 188–189, 516–535, 1997.

25 Moran, M. S., Hymer, D. C., Qi, J., and Sano, E. E.: Soil moisture evaluation using-temporal synthetic aperture radar (SAR) in semiarid rangeland, *Agr. Forest Meteorol.* 105, 69–80, 2000.

30 Mougin, E., Hiernaux, P., Kergoat, L., Grippa, M., de Rosnay, P., Timouk, F., Le Dantec, V., Demarez, V., Lavenu, F., Arjounin, M., Lebel, T., Soumaguel, N., Ceschia, E., Mougénot, B., Baup, F., Frappart, F., Frison, P. L., Gardelle, J., Gruhier, C., Jarlan, L., Mangiarotti, S., Sanou, B., Tracol, Y., Guichard, F., Trichon, V., Diarra, L., Soumaré, A., Koité, M., Dembélé, F., Lloyd, C., Hanan, N. P., Damesin, C., Delon, C., Serça, D., Galy-Lacaux, C., Seghier, J.,

HESSD

7, 7417–7450, 2010

Mapping soil moisture in Sahel using ASAR data

F. Baup et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Mapping soil moisture in Sahel using ASAR data

F. Baup et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Becerra, S., Dia, H., Gangneron, F., and Mazzega, P.: The AMMA Gourma observatory site in Mali: Relating climatic variations to changes in vegetation, surface hydrology, fluxes and natural resources, *J. Hydrol.*, 375, 14–33, 2009.

Mattia, F., Satalino, G., Dente, L., and Pasquariello, G.: Using a priori information to improve soil moisture retrieval from ENVISAT ASAR AP data in semiarid regions, *IEEE T. Geosci. Remote Sens.*, 44, 900–912, 2006.

Njoku, E. G., Jackson, T. J., Lakshmi, V., Chan, T. K., and Nghiem, S. V.: Soil moisture retrieval from AMSR-E, *IEEE T. Geosci. Remote Sens.*, 41, 215–229, 2003.

Schmugge, T., Jackson, T., Kustas, T. J., and Wang, J. R.: Passive microwave remote sensing of soil moisture: results from HAPEX, FIFE and MONSOON'90, *ISPRS Journal of Photogrammetry and Remote Sensing*, 47, 127–143, 1992.

Tansey, K. J., Millington, A. C., Battikhi, A. M., and White, K. H.: Monitoring soil moisture dynamics using satellite imaging radar in northeastern Jordan, *Appl. Geogr.*, 19, 325–344, 1999.

Taylor, C. M. and Ellis, R. J.: Satellite detection of soil moisture impacts on convection at the mesoscale, *Geophys. Res. Lett.*, 33, 3, L03404, doi:10.1029/2005GL025252, 2006.

Ulaby, F. T., Fung, A. K., and Moore, R. K.: *Microwave and remote sensing active and passive*. Norwood, MA: Artech House, 456 pp., 1981.

Ulaby, F. T., Fung, A. K., and Moore, R. K.: *Microwave and remote sensing: active and passive: surface scattering and emission theory*, Norwood, MA, Artech House, 1064 pp., 1982.

Wagner, W. and Scipal, K.: Large-scale Soil moisture mapping in western Africa using the ERS Scatterometer, *IEEE T. Geosci. Remote Sens.*, 38, 1777–1782, 2000.

Wagner, W., Scipal, K., Bartsch, A., and Pathe, C.: ENVISAT's Capabilities for Global Monitoring of the Hydrosphere, *Geosci. Remote Sens. Symposium 2005 (IGARSS '05)*, IEEE International, 8, 5678–5680, 2005.

Zine, S., Jarlan, L., Frison, P. L., Hiernaux, P., and Rudant, J. P.: Land surface parameter monitoring with ERS scatterometer data over the Sahel: A comparison between agro-pastoral and pastoral areas, *Remote Sens. Environ.*, 96, 438–452, 2005.

Zribi, M., Andre, C., Saux-Picard, S., Descroix, L., Otlle, C., Kallel, A.: Soil moisture mapping based on ASAR/ENVISAT radar data over a sahelian region, *Int. J. Remote Sens.*, 29, 3547–3565, 2007.

Zribi, M., André, C., and Decharme, B.: A method for soil moisture estimation in Western Africa based on ERS Scatterometer, *IEEE T. Geosci. Remote Sens.*, 46, 2, 438–448, 2008.

HESSD

7, 7417–7450, 2010

Mapping soil moisture in Sahel using ASAR data

F. Baup et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Table 1. Raingauges acquisitions time step.

Raingauge location	Tin Tadeini	Ekia	Eguerit	Bangui Mallam	Agoufou	Hombori	Kyrnia	Kobou
Acquisition time step	real time	real time	15 min	Real time	15 min	Daily	real time	real time

Mapping soil moisture in Sahel using ASAR data

F. Baup et al.

Table 2. Dates, time and orbit passes (Descending (D) or Ascending (A)) of radar image used in this study.

Season	ASAR				Orbit	WSC	
	Month	Day	Time	Month		Day	
wet	07	16	10:03:41	D	07	16	
wet	08	04	10:06:28	D	08	04	
wet	08	20	10:03:36	D	08	20	
wet	09	3	22:06:13	A	09	3	
wet	09	8	10:06:25	D	09	8	
dry	12	19	10:00:29	D	–	–	
dry	12	20	22:11:58	A	–	–	

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





Fig. 1. Views of the Agoufou **(a)**, Eguerit **(b)** and Bamba **(c)** sites during the dry season, respectively in April (a and b) and October 2005 (c).

Mapping soil moisture in Sahel using ASAR data

F. Baup et al.




Fig. 2. The AMMA mesoscale site in Gourma (Mali) (1°W – $2^{\circ}\text{W}/14.5^{\circ}\text{N}$ – 17.5°N) showing the location of the soil moisture stations and automatic raingauges in 2005.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 3. Seasonal variation of surface soil moisture (SSM) at 5 cm depth at the Agoufou sandy **(a)** and Eguerit rocky **(b)** sites in 2005.




Fig. 4. Soil types map of the AMMA mesoscale site in Mali. The 9 soil types (stacked bars) are grouped into 4 dominant soil texture classes (polygons).

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

Mapping soil moisture in Sahel using ASAR data

F. Baup et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Mapping soil moisture in Sahel using ASAR data

F. Baup et al.




Fig. 5. Chart of the surface soil moisture (SSM) mapping algorithm.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Mapping soil moisture in Sahel using ASAR data

F. Baup et al.




Fig. 6. Temporal variation of the normalized (23°) backscattering coefficient in 2005 for the Agoufou sandy (**a**) and the Eguerit rocky (**b**) sites.

Mapping soil moisture in Sahel using ASAR data

F. Baup et al.




Fig. 7. Soil moisture (SSM) maps established during the 2005 dry (December, 19) and wet (July, 16) seasons compared to EPSAT-SG rainfalls estimates on the same day and over the four previous days and with rainfall recorded at Agoufou and Tin Tadeini during the five same days.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Mapping soil moisture in Sahel using ASAR data

F. Baup et al.




Fig. 8. Comparisons of August soil moisture (SSM) maps (4 and 20) derived from ENVISAT-ASAR and ERS WSC SSM products and EPSAT-SG rainfalls estimates on the same day and over the four previous days and with rainfall recorded at Agoufou and Tin Tadeini during the five same days.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 9. Comparisons of September soil moisture (SSM) maps (3 and 8) derived from ENVISAT-ASAR and ERS WSC soil moisture products and EPSAT-SG rainfalls estimates on the same day and over the four previous days, and with rainfall recorded at Agoufou and Tin Tadeini during the five same days.

Mapping soil moisture in Sahel using ASAR data

F. Baup et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Mapping soil moisture in Sahel using ASAR data

F. Baup et al.




Fig. 10. Distribution of soil moisture (SSM) values ($\% \text{ m}^3/\text{m}^3$) at meso-scale site during the dry and the wet 2005 seasons.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Mapping soil moisture in Sahel using ASAR data

F. Baup et al.




Fig. 11. Comparisons of SSM values estimated from ENVISAT-ASAR and ERS-WSC data for the 2005 driest (**a**) and wettest (**b**) dates (2005-09-08 and 2005-08-20, respectively). The location of SSM pixels versus the latitude is also indicated.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)