

This discussion paper is/has been under review for the journal Hydrology and Earth System Sciences (HESS). Please refer to the corresponding final paper in HESS if available.

**Simulation of the
snowmelt runoff
contributing area**

C. M. DeBeer and
J. W. Pomeroy

Simulation of the snowmelt runoff contributing area in a small alpine basin

C. M. DeBeer and J. W. Pomeroy

Centre for Hydrology, University of Saskatchewan, 117 Science Place, Saskatoon,
Saskatchewan, S7N 5C8 Canada

Received: 21 January 2010 – Accepted: 28 January 2010 – Published: 8 February 2010

Correspondence to: C. M. DeBeer (cmd225@mail.usask.ca)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

Simulation of areal snowmelt and snow-cover depletion over time can be carried out by applying point-scale melt rate computations to distributions of snow water equivalent (SWE). In alpine basins, this can be done by considering these processes separately on individual slope units. However, differences in melt timing and rates arise at smaller spatial scales due to the variability in SWE and snowpack cold content, which affects the timing of melt initiation, depletion of the snow-cover and spatial extent of the snowmelt runoff contributing area (SRCA). This study examined the effects of variability in SWE, internal energy and applied melt energy on melt rates and timing, and snow-cover depletion in a small cold regions alpine basin over various scales ranging from point to basin. Melt rate computations were performed using a physically based energy balance snowmelt routine (Snobal) in the Cold Regions Hydrological Model (CRHM) and compared with measurements at three meteorological stations over a ridge within the basin. At the point scale, a negative association between daily melt rates and SWE was observed in the early melt period, with deeper snow requiring greater energy inputs to initiate melt. SWE distributions over the basin (stratified by slope) were measured using snow surveys and repeat LiDAR depth estimates, and used together with computed melt rates to simulate the areal snow-cover depletion. Comparison with observations from georeferenced oblique photographs showed an improvement in simulated areal snow-cover depletion curves when accounting for the variability in melt rate with depth of SWE in the early melt period. Finally, the SRCA was characterized as the product of the snow-covered area and the fraction of the SWE distribution undergoing active melt on each slope unit. Results for each slope were then aggregated to give the basin scale SRCA. The SRCA is controlled by the variability of melt amongst slope units and over individual SWE distributions, the variability of SWE and the resulting snow-cover depletion patterns over the basin.

HESSD

7, 971–1003, 2010

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

1 Introduction

The end-of-winter snow-cover over open alpine terrain is characterized by considerable spatial variability in its snow water equivalent (SWE) depth, primarily as a result of snow redistribution by wind (Elder et al., 1991; Winstral et al., 2002). Topographic depressions, areas of exposed alpine shrubs and vegetation, and leeward slopes tend to accumulate snow drifts up to several metres or more in depth over the winter, whilst adjacent and more exposed terrain is scoured by wind and may accumulate little or no snow. During the spring snowmelt period, a complex patchwork of snow-covered and snow-free areas emerges as a result of this heterogeneity and the spatial variation in applied melt energetics over the landscape. Both sources of variation control the timing and rate of areal snow-cover depletion (SCD), which is important hydrologically since the fraction of snow-covered area defines the maximum extent of the snowmelt runoff contributing area (SRCA) within the basin. Thus, studies dealing with alpine runoff require sufficient understanding and representation of this variability and the SCD to realistically derive the snowmelt hydrograph.

Previous work has shown that snowmelt dynamics in complex terrain can be represented at intermediate spatial scales in models reasonably well by disaggregating the landscape into terrain units defined by features such as slope and aspect, and treating coupled mass and energy balances separately over these units (DeBeer and Pomeroy, 2009; Dornes et al., 2008a, b; Pomeroy et al., 2003). This avoids having to rely on fully distributed melt representations, whilst at the same time utilizing an objective means of landscape stratification rather than an arbitrary grid system that does not conform to the natural variability of the terrain and the processes involved. In this regard, the approach is useful for explicitly representing the variability in applied melt energetics between slopes as a result of differences in irradiation. Further, areal SCD can be represented within individual slope units by stochastically accounting for the pre-melt distribution of SWE and applying slope-based melt rate computations to the SWE distribution. DeBeer and Pomeroy (2009) demonstrated how this approach yields improved

HESSD

7, 971–1003, 2010

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

results in simulated SCD curves from the case of uniform applied melt energetics and a spatially lumped distribution of pre-melt SWE over the landscape.

DeBeer and Pomeroy (2009) noted, however, that problems with this approach may arise in situations where melt rates and timing are non-uniform within individual terrain units. For example, small scale statistical associations between SWE and melt rates may accelerate or prolong areal SCD. Faria et al. (2000) and Pomeroy et al. (2004) attributed small scale negative associations to the effects of vegetation. In areas with a cold, redistributed and highly non-uniform snow-cover, such associations would also result from differences in the internal energy content of the snow. This may be particularly problematic in the early melt period as shallow snow becomes isothermal and begins to melt while adjacent deeper snow is still warming to 0 °C (Gray and O'Neill, 1974; Male and Gray, 1975). Essery and Pomeroy (2004) suggested integrating over the joint SWE and melt rate frequency distributions in the case of inhomogeneous melt to derive SCD curves, but noted that these integrals will generally be intractable. This is because both the frequency distributions of these variables and the association between them are not constant over time, and thus an analytical solution is not generally possible.

Other than studies using a fully distributed approach, there has been no attention given to the effects of small scale spatial variation of initial SWE and internal snowpack energy storage on the computation of melt rates over a SWE distribution. Physically-based (i.e. energy/mass balance) point-scale melt rate computations applied to these distributions have typically been based on a single initial snowpack state (depth, density, internal energy, etc.) (Liston, 1999; Luce et al., 1999), and have tended to neglect internal energy changes. At the same time, the SRCA has generally been assumed to be equal to the snow-covered area for estimating meltwater volume (Martinec et al., 1998). This is an oversimplification of reality for cold, redistributed snow-covers since not all areas will generate and release meltwater in the early stages of melt (e.g., Gray and O'Neill, 1974; Marsh and Pomeroy, 1996). Often, melt is first produced from the shallow snow, which may completely disappear before the areas with a deeper snow-

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

pack begin to produce melt (Male and Gray, 1975).

The purpose of this study is therefore to investigate the problems described above more thoroughly and to suggest a potential approach to resolve them in modelling applications. We first examine, at the point scale, how variability in initial SWE depth affects the computation of snowmelt rates and timing on variously orientated slopes over an alpine ridge within the Canadian Rocky Mountains. The effects of point-scale melt rate variation due to differences in internal energetics over a non-uniform snow-cover are then examined in terms of the areal SCD and the temporal evolution of the SRCA for two different slopes in an adjacent cirque basin. We finally explore these features at the scale of the small headwater basin, which encompasses multiple slope units of different aspect, gradient, and elevation. This analysis provides an initial basis for predicting the magnitude and timing of the snowmelt hydrograph here and in similar environments.

2 Study area and field methods

This study was conducted at Fisera Ridge (~2315 m) and the adjacent sparsely vegetated cirque basin within the Marmot Creek Research Basin, Kananaskis, Alberta, Canada (Fig. 1). Marmot Creek is located within the Front Ranges of the Canadian Rocky Mountains, where climatic conditions are dominated by continental air masses. Winters are long and cold, with an average temperature of -15°C for the months of January through March for Fisera Ridge. The spring season is generally cool and wet, often producing late season snowfall events at high elevations. Average temperatures at Fisera Ridge for May and June (during the primary snowmelt period) are $+2$ and $+8^{\circ}\text{C}$, respectively. Historically, annual precipitation at Marmot Creek has been observed to average about 900 mm, increasing to over 1140 mm at treeline near the Mt. Allan cirque (Storr, 1967); recent observations corroborate these values. Roughly 60–75% of the annual precipitation falls as snow at Marmot Creek with the percentage increasing with elevation.

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Continuous meteorological and snowpack condition observations were made for three consecutive years (2007, 2008, 2009) at a permanent meteorological station located on Fisera Ridge, and at two outlier stations on north and south-east slopes, either side of the ridge, for 2008 and 2009 (Fig. 1). Net radiation was measured at the ridge station by a Kipp & Zonen CNR1 net radiometer; the incoming shortwave component was measured with a CM3 pyranometer (directional error Max. 25 W m^{-2} at 1000 W m^{-2}), and the incoming longwave component was measured with a CG3 pyrgeometer (expected accuracy of $\pm 10\%$ for daily sums). Air temperature and relative humidity were measured at each of the sites with a Vaisala HMP45C212 hygrometer housed inside a Gill radiation shield. Wind speed and direction were measured at the main station using an R.M. Young propeller anemometer (Model 05103-10), and wind speed at both of the outlier stations was measured with Met-One 014A 3-cup anemometers. Rainfall was measured at Fisera Ridge using a Campbell Scientific TB4-L tipping bucket rain gauge, and in August 2008 an Alter-shielded Geonor T-200B weighing precipitation gauge was installed in a sheltered site on the ridgetop near the main station to measure total precipitation. Measurements of total precipitation for the entire study period were also available from a similar Geonor gauge located at a mid-elevation (1843 m) forest clearing site within the Marmot Creek Basin roughly 2 km away. Snow depths were recorded using Campbell Scientific SR50 sonic ranging sensors at each of the stations (at the two outlier stations, the sensors were oriented perpendicular to the sloping ground surface). To monitor the internal snowpack temperature, a series of fine-wire thermocouples was installed vertically at the two outlier stations, beginning at two cm above the ground surface and continuing up at 20-cm intervals.

A digital single lens reflex camera (Pentax model K110D) mounted to the main station on Fisera Ridge was used to take several pictures daily of the cirque. Photos taken at 12:00 p.m. LT each day were selected for analysis of the snow-covered area over the cirque, except in situations when low cloud cover or snowfall obscured the terrain. The images were projected orthogonally onto a 1-m resolution digital elevation model

(DEM) of the cirque derived from airborne LiDAR (Light Detection And Ranging) using the *Georeferencing Terrestrial Photography* software tool (Corripio, 2004; see DeBeer and Pomeroy (2009) for a description of this procedure). From the corrected images, the snow-covered area fraction was determined for the cirque and for various slopes within it using a threshold procedure in ESRI® ArcMap™ 9.1 GIS. Snow-covered area was determined as the ratio of the number of “snow” pixels to the total number of pixels over the relevant area (excluding pixels containing “no-data” values which were hidden from view of the camera).

Snow surveys were repeated a number of times throughout the winter and spring along fixed transects over the ridge and within the cirque (Fig. 1). These surveys were done to characterize the spatial variability of SWE on different slopes over time. Snow depth was measured using a graduated aluminium rod every three metres along each transect, and density measurements were taken roughly every fifth depth measurement where possible using an ESC-30 snow tube and scale. Several snow pits were located at various positions across the ridge and the within the cirque to examine the vertical snowpack structure and measure the density of the individual snow strata.

In late March, 2008 a second LiDAR DEM was obtained over Marmot Creek for the purpose of examining spatial patterns of snow accumulation over the basin (Hopkinson et al., 2010). The DEM representing snow-free conditions was subtracted from the snow-covered DEM to generate a 1-m spatial resolution raster image of snow depth values. Comparison of point values of snow depth extracted from this raster image with values measured along the Fisera Ridge survey transect on 28 March 2009 showed good correspondence ($R^2=0.94$; $n=137$; RMS error=0.17 m) over the range of measured snow depths from 0 to 2.88 m, and the LiDAR depths reproduced the observed spatial pattern over the ridge very well. There were no apparent biases in error due to slope or aspect, but LiDAR-derived snow depth had greater error (up to ~0.6 m) in locations with exposed alpine shrubs. Thus, the LiDAR snow depth raster appears to provide a useful tool for examining the spatial pattern and distribution of snow depth over most of the sparsely vegetated and open alpine cirque.

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

3 Snowmelt modelling and validation

Simulations of snowmelt rate and timing were carried out using the Cold Regions Hydrological Model (CRHM) platform (Pomeroy et al., 2007). CRHM is a flexible, object-oriented modelling system that can be used to develop, support, and apply dynamic hydrological process algorithms. These algorithms are applied over hydrological response units (HRU; i.e. homogeneous landscape units characterized by their terrain and surface properties), within which conditions and processes are represented by single sets of parameters, state variables, and energy and mass fluxes. Various component modules representing basin characteristics, observations, and hydrological processes are combined within CRHM to form an operational model of the system that has a level of complexity specified by the needs of the user.

Internal snowpack temperature and melt rates were computed using *Snobal* (snowmelt energy balance model; Marks et al., 1998, 1999), which was incorporated as a module within CRHM. This module approximates the snowcover as being composed of two layers: a surface active layer of fixed maximum thickness, and a lower layer representing the remaining snowpack. The module solves for the temperature ($^{\circ}\text{C}$) and the specific mass (kg/m^2) or water equivalent depth per unit area (mm) of each layer for each timestep. The energy balance of the snowpack at a point is expressed as:

$$Q_m = Q^* + Q_H + Q_E + Q_G + Q_P - dU/dt, \quad (1)$$

where Q_m is the energy available for snowmelt, Q^* is the net radiation comprised of both short- and longwave components, Q_H , Q_E , and Q_G are the sensible, latent, and ground heat fluxes, respectively, Q_P is the energy added to the snowpack by precipitation, and U is the internal energy of the snowpack. The melt energy can be expressed as a depth of melt, m , by:

$$m = \frac{Q_m}{\rho h_f \beta}, \quad (2)$$

where ρ is the density of the snow, h_f is the latent heat of fusion (0.334 MJ kg^{-1}), and

HESSD

7, 971–1003, 2010

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



β is the fraction of ice in snow (taken as 0.97). Melt is computed in either layer when the accumulated energy exceeds that required to bring the snowpack to 0°C , at which point positive values of Q_m result in snowmelt. A comprehensive description of the Snobal model is given by Marks et al. (1998, 1999).

The model was run at the point scale at 15-min time intervals in CRHM using meteorological observations from the various stations as the external forcing variables. Observations of short and long-wave radiation components were only available from the main station, so these had to be corrected for slope, aspect, and sky exposure to be applied at the two outlier stations. For shortwave radiation this was done using the modules *Global* and *Slope_Qsi* within CRHM, which use expressions from Garnier and Ohmura (1970) and List (1968) to determine clear sky direct and diffuse radiation on slopes. The ratio of measured shortwave radiation to the calculated theoretical clear sky direct and diffuse radiation on a horizontal plane is used to adjust the calculated clear sky shortwave radiation value to the slope. Incoming longwave radiation was adjusted for differences in sky exposure between the main station and the outlier stations using a modified version of the parameterization given by Sicart et al. (2006). More detailed descriptions of these procedures and the formulae used are given by DeBeer and Pomeroy (2009).

The model was run for the late winter and spring periods of 2007, 08, and 09. The first two years of observations were used to calibrate the model and select the most suitable parameter values; observations from the final year were used to evaluate the model performance. Model evaluation criteria included internal snowpack temperature, SWE, and daily melt rates.

Parameter selection focused on three parameters related to snowpack properties, and four related to the decay function for snow albedo, α . The first three parameters are the roughness height of the snow surface, z_0 , the maximum active layer thickness of the snowpack, $\max_{z,s0}$, and the maximum liquid water holding capacity of the snowpack, $w_{c,max}$. Roughness height was set as $z_0=2.0\times 10^{-3}$ m, as measurements of z_0 over seasonal snow-covers generally indicate small values on the order of 10^{-4} to 10^{-3} m

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

(Harding, 1986). The maximum active layer thickness was set as 0.25 m. Maximum water holding capacity, which is defined as the ratio of the volume of liquid water to the volume of pore space within the snowpack, was set as 0.01 following Marks et al. (1998). The snow albedo was parameterized as an exponential decay during the melt period using the following expressions from Essery and Etchevers (2004). For each timestep with snowmelt, the albedo is updated according to:

$$\alpha \rightarrow (\alpha - \alpha_{\min}) \exp\left(\frac{-\Delta t}{\tau}\right) + \alpha_{\min}, \quad (3)$$

where Δt is the timestep length, τ is a time constant applied to melting snow, and α_{\min} is the minimum albedo that the decay declines asymptotically toward. For time steps with snowfall, the albedo is increased by:

$$\alpha \rightarrow \alpha + (\alpha_{\max} - \alpha) \frac{S_f \Delta t}{S_{\min, \alpha}}, \quad (4)$$

where S_f is the snowfall amount during the timestep, and $S_{\min, \alpha}$ is the minimum snowfall amount required to refresh the albedo to α_{\max} . Values of $\tau=10^6$ s, $\alpha_{\min}=0.3$, $S_{\min, \alpha}=10$ mm, and $\alpha_{\max}=0.85$ were used in Eqs. (3) and (4).

Figure 2 provides an example of the model calibration and performance at a point for the south-east facing station. The model results are in reasonable agreement in terms of the timing and rate of snowmelt at this site and the others. Although simulations of internal snowpack temperature were problematic at times prior to the onset of melt, the timing of the pack reaching an isothermal state of 0°C at each site corresponded very well with the observations. Table 1 provides the root mean square (RMS) errors between simulation results and measurements of several variables at each of the sites. The success of the model for predicting snowmelt timing and rates at Fisera Ridge indicates that it can be used towards examining potential effects of differences in initial SWE on these variables at other sites in the Mt Allan cirque.

4 Simulated melt rate – SWE associations

To examine the sensitivity of snowmelt timing and melt rates to initial SWE and the associated differences in internal energy content, the model was run at the point scale for various initial depths of SWE. These simulations were carried out using the parameter values listed in the previous section and the observed 15-min meteorological conditions at Fisera Ridge and the two outlier stations for 2008. Outputs of SWE from the model each day at midnight were differenced to derive the simulated daily melt rates, which were then compared with SWE values at the beginning of each day.

Figure 3 shows plots of melt rates vs. SWE for the south-east and north facing sites at different times during the melt period. It is clear from these plots that there are considerable differences in the simulated rates and timing of snowmelt between slopes of different orientation. For example, in the early melt period shallow snow (i.e. SWE < 200 mm) at the south-east facing site began melting nearly one week before snow at the north facing site. Once snowmelt began at the north facing site, the rate was roughly an order of magnitude less than that simulated at the south-east facing site. Similar variation in melt rates is evident at other times in the melt period as well. Differences in melt rates amongst individual slope units have been previously reported (DeBeer and Pomeroy, 2009; Dornes et al., 2008a; Pomeroy et al., 2003) and are explained primarily by differences in short-wave radiation receipt.

The results in Fig. 3 also show that differences in initial SWE have a significant effect on the rate and timing of simulated melt under the same applied melt energetics. The effect is especially important early in the melt period. The figure shows an early phase of the snowmelt period in 2008, which ceased with cooler weather and snowfall beginning on 5 May. At the south-east facing site during this time, simulated shallow snow began melting nearly a week prior to deeper snow (i.e. ~300–700 mm SWE), while negligible melt was predicted for the deepest snow. The effect was even more pronounced at the north facing site, with only the shallowest snow beginning to melt by the end of this early phase of the melt period. A strong negative association between

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

simulated melt rates and SWE is clear at both sites during this time.

At later times in the snowmelt period, Fig. 3 shows these effects to change. A more pronounced phase of the snowmelt period began in mid-May, and within a few days, all snow began actively melting and the negative melt rate–SWE association disappeared at both sites. This was due to the fact that all snow depths were either at, or near to isothermal conditions. A negative association became evident again for a brief time in the late melt period in June at both sites following a snowfall event. Again, the deeper snow would take greater energy inputs to bring it back to 0 °C following the short period of colder weather. Although our model assumed a uniform ground heat flux, more rapid melt of shallow snow at later times in the melt period following snowfall events may also result from greater energy supplied to the snow from the ground surface in areas which were previously snow-free and heated by the sun. Interestingly, at other times a slight positive association between melt rates and SWE was predicted by the model (e.g. 18 May, 22 June). This is explained by the refreezing of liquid water within shallow snow overnight, which requires greater energy inputs than deeper snow (due to the greater thermal mass) to satisfy the latent heat requirement and restore melting conditions.

The variation in melt rates indicated by the model due to variable SWE depth alone is significant. Briefly, at times such as the early snowmelt period or following cold weather and snowfall events, it is of equal magnitude as the variation between slopes resulting from differences in radiation receipt. The association between melt rates and SWE at these times tends to be non-linear and changes over time, making the effects of the association difficult to parameterize in terms of a constant linear covariance term (e.g., Pomeroy et al., 2004). This is an important result that has not been previously considered in most snowmelt modelling applications, but which has significant implications for simulating melt rates over a cold, heavily redistributed snow-cover. These implications are considered in the following section.

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

5 Areal SCD and snowmelt runoff contributing area

Simulated snowmelt rates were used to model the decline in snow-covered area and the temporal evolution of the snowmelt runoff contributing area (SRCA) over two opposing slopes within the adjacent cirque. Our approach is based on the theoretical lognormal frequency distribution, which can be expressed for SWE values in the following linear form:

$$\overline{SWE} = \overline{SWE}(1 + KCV), \quad (5)$$

where \overline{SWE} and CV are the mean and the coefficient of variation (standard deviation/mean), respectively, and K is the frequency factor for each particular SWE value calculated from the rank of observation (Chow, 1954). If the underlying distribution is lognormal, values of SWE plotted against K should approximate a straight line with a slope equal to the standard deviation of SWE and an intercept at $K=0$ equal to \overline{SWE} . The theoretical framework for determining K values from observed data and for deriving SCD curves is described by DeBeer and Pomeroy (2009), Faria et al. (2000), Pomeroy et al. (1998) and Shook (1995). Essentially, the value of K corresponding to $SWE=0$ (i.e. K_{\min}) provides an index of the fraction of snow-covered area over an individual terrain unit, and by applying slope-based computed melt rates to the distribution, K_{\min} values can be continuously recalculated to generate the SCD curve.

Areal SCD was simulated using this framework on both a north- and a south-facing slope within the cirque basin (following DeBeer and Pomeroy, 2009). Measurements from the snow surveys and the LiDAR snow depth raster were used to define the parameters of the SWE distributions on these slopes (Table 2). Snowmelt modelling was carried out as described previously in this paper, with the short-wave and long-wave radiation inputs adjusted for the gradient, aspect, and sky view of the two slope units. To investigate the effects of the variation in computed melt rates due to differences in initial SWE on simulated SCD curves, daily melt rates were applied uniformly from simulations based on \overline{SWE} for each measured distribution, and were also applied across

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

the distribution using melt rates computed for different initial values of SWE depth. In the latter case, SCD was simulated by discretizing the linear form of the SWE distribution (Eq. 5) on each slope into a number of individual segments, which were defined by the initial SWE values used in the melt computations. As each of these points melted

5 over time, the segments were readjusted between the points, and that representing the shallowest part of the distribution was used to determine K_{\min} , and thus, snow-covered area. The effects of snowfall events during the melt period, which refresh the snow-cover until the new snow has melted, were handled using a rescaled depletion curve following Moore et al. (1999).

10 The SRCA on each slope was characterized as the product of the snow-covered area and the fraction of the remaining SWE distribution producing an appreciable melt quantity (i.e. non-zero SWE with melt rates $> \sim 5$ mm/day). To determine this fraction the relative proportion of the distribution within each of the initial segments of the linear representation of the SWE distribution must be known. For the lognormal distribution,

15 this can be found knowing the initial $\overline{\text{SWE}}$ and CV as follows:

1. For each value of initial SWE used to define the limits of these segments, determine the corresponding K value from Eq. (5). The value of the frequency factor K provides an index of the exceedence probability of that particular value of SWE in the distribution.
- 20 2. To determine the exceedence probability, calculate the value of the transformed frequency factor K_y (i.e. Eq. 3 in DeBeer and Pomeroy, 2009) corresponding to each value of K . Theoretically, K_y is calculated as:

$$K_y = \frac{\ln(1 + KCV)}{s_y} + \frac{s_y}{2}, \quad (6)$$

where s_y is the standard deviation of the log-transformed data, which can be

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



estimated from (Chow, 1954):

$$s_y = \sqrt{\ln(CV^2 + 1)}. \quad (7)$$

The probability of K_y being exceeded, $P(K_y)$, is equivalent to the exceedence probability of SWE, and can be determined as:

$$P(K_y) = \frac{1}{\sqrt{2\pi}} \int_{K_y}^{\infty} \exp\left(-\frac{K_y^2}{2}\right) dK_y. \quad (8)$$

3. The fraction of the total initial distribution represented by each segment can then be determined as the difference in exceedence probability between the lower and upper limits of each class.

At any time, the fraction of the remaining distribution represented by the individual segments in the plot of K vs. SWE can be found by multiplying the fraction of the total initial distribution represented between K values by the reciprocal of the snow-covered area. By this approach, the range of SRCA is limited in value between zero (in the case of no significant melt computed for any SWE depth) and the fraction of snow-covered area (in the case of all SWE actively melting).

Figure 4 shows the simulated and observed SCD curves, as well as the evolution of the SRCA on both the north and south facing slopes in the cirque during the early melt period of 2008. The results show an improvement in the simulated SCD curves when accounting for the variability of melt rates over different initial SWE depths, particularly on the south facing slope. This initial phase of SCD on the south facing slope was due mainly to the melt and disappearance of large areas of shallow snow, while adjacent deeper snow was still warming to 0°C . Thus, the simulation using melt rates based on a single melt rate for average SWE poorly represented the decline of snow-covered area in this early snowmelt phase. On the north facing slope at this time, SCD was initially due to wind scouring of the snow-cover (which is not accounted for in this

approach). Melt of the shallow snow did not begin until 4 May, at which time the model based on SWE classes predicted a small decline in snow-covered area, corresponding to the observations. The simulation based on melt rates computed for mean SWE predicted no decline in snow-covered area at this time.

5 The temporal evolution of the SRCA differs considerably between the two slopes during this early melt period as a result of differences in the SWE distributions and SCD curves, as well as the melt rates computed over these distributions. Simulated snowmelt runoff generation began on 26 April over the south facing slope, but the areal extent of the SRCA was limited since not all of the SWE distribution had begun actively
10 melting. Over the next several days this extent grew larger as melt occurred over an increasing portion of the SWE distribution. However, at the same time, the depletion of snow-covered area partially counteracted the effect of melt over an increasing fraction of the remaining distribution. The maximum extent of the SRCA occurred on 30 April (0.60) after which it declined due to further areal SCD. On the north facing slope, simulated snowmelt runoff generation did not begin until 2 May and reached its maximum
15 extent on 4 May (0.73). In this case, the maximum areal extent of the SRCA was larger than that over the south facing slope because of the limited areal SCD occurring on this slope. Subsequent snowfall occurring on 5 and 6 May refreshed the snowcover and temporarily ended snowmelt runoff generation on both slopes.

20 Figure 5 shows the results of this modelling on the two slopes carried out over the entire spring melt period of 2008. Following the initial period of melt and SCD, there was a period of cooler weather with several large snowfall events, resulting in a refreshed snowcover with high albedo and no snowmelt runoff generation. Melting conditions resumed on 14 May (south facing slope) and 15 May (north facing slope), and the value
25 of the SRCA rapidly approached the snow-covered area on both slopes as significant melt rates were computed for all SWE depths. Over the course of the spring there were several more snowfall events, with similar effects on areal snowcover, melt rates, and extent of the SRCA. Both approaches (uniform applied melt and melt computed for different SWE depths) yielded very similar SCD curves that closely matched the

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

observations after the initial melt phase in late April/early May. This is because the effects of variation in SWE on computed melt rates were less pronounced later in the snowmelt period, and the subsequent periods when a negative association was observed were too brief to have a major effect on areal SCD. This partly explains why the simulated SCD curves of DeBeer and Pomeroy (2009) were found in good agreement with observations later in the melt period using uniform slope-based melt rates.

This modelling framework was extended to all slopes within the upper Middle Creek basin to simulate areal SCD and SRCA at the basin scale. The basin was segregated into four slope-based HRUs, which include the north and south facing slopes. In addition, a broad east facing slope on Mt. Allan in the upper part of the basin was defined, and the small cirque floor area was distinguished from the other slope units. The floor is characterized by relatively level terrain with several small knolls and depressions and incised by several gulleys. Table 2 gives the statistical parameters of the SWE distributions on each HRU used to approximate the lognormal distribution. The basin contains some forested areas in the lower region, where the tundra snow modelling framework in this study is not expected to apply. These areas were therefore excluded from the analysis. Further, there are a few local areas of steep cliffs in the upper part of the basin, which are permanently snow-free. To distinguish these areas and exclude them from the analysis, a terrain classification similar to that of Blöschl et al. (1991) was carried out to identify areas with a gradient steeper than 50° . This value of the critical slope gave results that corresponded well with permanent snow-free areas identified in the corrected oblique photographs.

The simulation results on each slope unit were weighted according to the area of the slope and combined to give the basin scale SCD curve and SRCA, which are shown in Fig. 6. As on the individual slopes, the SRCA is initially zero and approaches the value of the snow-covered area as melt begins to occur over an increasing portion of the distributions on these slopes. Periods of new snowfall refresh the snowcover and reduce the snowmelt runoff generation area to zero until melt resumes following these events. These results show that the basin scale SCD curve and SRCA are controlled

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

by the relative extent of the various slope units and the changing snow-covered area and snowmelt dynamics over them with time.

As an example of the spatial variability of the SCD and SRCA within the basin, Fig. 7 shows the extent of snow-covered area and melting snow over several dates in the early melt period of 2008. These patterns were mapped from the LiDAR-derived snow depth raster image by accounting for the melt and depletion of different SWE depths over time on each of the slopes. Initially, simulated runoff generation began over limited parts of the south facing slope and the cirque floor. These areas then expanded and subsequently became depleted of snow as more parts of the other slopes began to produce melt. By the time the SRCA had expanded to include most of the area on the north facing and upper east facing slopes, the snow-covered area on the other slopes had declined considerably, thus limiting the runoff generating source area there. Thus, the differences in melt rates and SWE distributions amongst slope units controls not only the overall timing of the development of the SRCA, but the location of contributing areas as well.

6 Discussion and conclusions

This study has shown that differences in the warming and melt rates over a cold and highly redistributed snowcover have an important effect on areal SCD as well as the timing and development of the SRCA. The effects are most pronounced early in the melt period when only the shallow snow is actively melting. This leads to an initial acceleration of the SCD from the case of uniform applied melt rates based on the mean SWE depth, due to earlier and faster melt of areas with a thin snowcover. As the melt period progresses and the effects of variable SWE and internal energy differences become reduced, melt rates over a given slope (and SWE distribution over the slope) become more uniform and consideration of melt differences amongst SWE classes becomes less important. However, it is noted that this variability can be important again at other times such as following spring snowfall events (which are common in this environ-

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ment and refresh the snow-cover) or periods of cold weather. Even night-time cooling and longwave radiation losses (particularly on clear nights) can have a considerable effect on internal snowpack energetics and become manifested as differences in melt rates amongst different SWE depth classes the following day. Male and Gray (1975) showed the importance of the internal energy term in applying the energy balance principle to shallow snowcovers. Shallow snow tends to undergo large diurnal variations in internal energy due to overnight cooling and refreezing, in contrast to deeper snow, which takes longer to warm in spring, but exhibits relatively damped diurnal variation of internal energy content due to its greater thermal mass. Thus, it is important to consider differences in melt rates due to variation in SWE throughout the entire melt period in such environments.

Realistic simulation of the SRCA depends on proper representation of the areal SCD, which limits the maximum contributing area, as well as the fraction of the SWE distribution that is actively melting. Consideration of differences in melt timing and rates over a non-isothermal SWE distribution is therefore important in this regard. As described previously, earlier and initially faster melt of the shallow snow depletes the snow-cover and exposes the ground more rapidly in these sites. Thus, the effect of the earlier melt of shallow areas of snow, which initially expands the SRCA, is somewhat counteracted by the fact that these areas become more rapidly snow-free and the runoff contributing area is then limited by the snow-covered area. At the basin scale, which generally encompasses multiple slope units of different aspect, gradient, and elevation, the pattern is even more complex. Snowmelt and areal SCD begin first over areas with shallow snow on south facing slopes or level lower elevation sites, and then proceed to deeper snow within these areas and shallow snow on other slopes. By the time that most of the snow-cover on the remaining slopes begins to melt, much of the snow-cover on the south facing slopes may have disappeared. Subsequent spring snowfalls add even more complexity to these patterns as the snow-cover is re-established over the landscape and the SRCA must again evolve.

The approach used here was to apply slope-based melt energetics to different SWE

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



classes from a spatial frequency distribution of SWE over the different slope units in the basin. This yielded reliable prediction of areal SCD that corresponded well with observations, and it is reasonable to infer that the simulated SRCA was in good agreement with the actual areal extent and location of the snow-covered area producing appreciable melt quantities, based on the success of the model at the point scale and the known spatial distributions of SWE. The approach required far less data input and parameter information than a fully distributed approach, and yet was capable of simulating the complex spatial evolution of the snow-covered area and SRCA during the spring. This is because the processes and SWE distributions were represented at the appropriate spatial scales, areas and locations. This approach could easily be applied in other similar alpine basins with meteorological forcing adjusted for the slope/aspect and elevation, and information on the basic parameters describing the initial SWE distributions on the slopes (i.e. \overline{SWE} and CV for the lognormal distribution).

The approach is limited and may not be suitable in some situations. For example, it is not expected to work where other factors have a significant influence on the association between melt rate and SWE, unless those factors can be accounted for by the model. In this study, we neglected densely vegetated parts of the basin where a more complex relation between melt rates and SWE may exist (Faria et al., 2000; Pomeroy et al., 2001). When applying this framework over mountainous terrain it is necessary to account for cliffs, which are subject to snow avalanches and do not retain snowcover (Blöschl et al., 1991). The lognormal distribution generally provides a useful approximation to the SWE distribution over the remainder of the slope. However, if the terrain is highly complex with extensive cliffs comprising much of the slope unit area, then this approximation may fail. Further measurements are needed to determine whether and how the presence of cliffs over larger parts of the landscape might affect SWE distributions.

The findings and developments in this work can be used towards an improved framework for simulating the spatial variability of meltwater generation and release from the snowpack in open alpine terrain. Realistic, physically-based representation of these

processes must account for the spatial variability in melt rates and SWE at multiple scales (i.e. amongst individual slopes and within them). This leads to a further issue that should then be considered – the retention of meltwater within the snowpack and the transmission of meltwater fluxes through the pack. The rate of propagation of meltwater fluxes through snow is related to the magnitude of the flux (Colbeck, 1972; Colbeck and Davidson, 1973), while the travel time through the snowpack is a function of this rate as well as the depth of snow. This travel time can become important for some of the deepest snow found in this study site. Thus, the framework described in this study for simulating the areal SCD and SRCA should eventually be coupled with a variable meltwater routing scheme through snow that accounts for these principles. This will allow for a more realistic simulation of the meltwater release over the landscape, which can then be used as the input to hydrological models for streamflow simulation.

Acknowledgements. Funding for this work was provided by the Canadian Foundation for Climate and Atmospheric Sciences through the IP3 Network, the Natural Sciences and Engineering Research Council of Canada (NSERC), the University of Calgary Biogeoscience Institute, and the Canada Research Chairs program. Nakiska Ski Resort provided logistical support. We thank Michael Solohub, Chad Ellis, Matt MacDonald, Jim MacDonald, Anne Sabourin, Tom Brown, and Kevin Shook for their help in various aspects of the field work and modelling.

References

- Blöschl, G., Kirnbauer, R., and Gutknecht, D.: A spatially distributed snowmelt model for application in alpine terrain, in: Snow, Hydrology and Forests in High Alpine Areas, Proceedings of the Vienna Symposium, Aug 1991, edited by: Bergmann, H., Lang, H., Frey, W., Issler, D., and Salm, B., IAHS publication No. 205, IAHS, Wallingford, 51–60, 1991.
- Chow, V. T.: The log-probability law and its engineering applications, *P. Am. Soc. Civil Eng.*, 80, 1–25, 1954.
- Colbeck, S. C.: A theory of water percolation in snow, *J. Glaciol.*, 11, 369–385, 1972.
- Colbeck, S. C. and Davidson, G: Water percolation through homogeneous snow, in: *The Role*

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- of Snow and Ice in Hydrology, Proceedings of the Banff Symposia, Sep 1972, Unesco-WMO-IAHS, Geneva-Budapest-Paris, 242–257, 1973.
- Corripio, J. G.: Snow surface albedo estimation using terrestrial photography, *Int. J. Remote Sens.*, 25, 5705–5729, 2004.
- 5 DeBeer, C. M. and Pomeroy, J. W.: Modelling snow melt and snowcover depletion in a small alpine cirque, *Canadian Rocky Mountains, Hydrol. Process.*, 23, 2584–2599, 2009.
- Dornes, P. F., Pomeroy, J. W., Pietroniro, A., Carey, S. K., and Quinton, W. L.: Influence of landscape aggregation in modelling snow-cover ablation and snowmelt runoff in a sub-arctic mountainous environment, *Hydrolog. Sci. J.*, 53, 725–740, 2008a.
- 10 Dornes, P. F., Pomeroy, J. W., Pietroniro, A., and Verseghy, D.: Effects of spatial aggregation of initial conditions and forcing data on modeling snowmelt using a land surface scheme, *J. Hydrometeorol.*, 9, 789–803, 2008b.
- Elder, K., Dozier, J., and Michaelsen, J.: Snow accumulation and distribution in an alpine watershed, *Water Resour. Res.*, 27, 1541–1552, 1991.
- 15 Essery, R. H. and Pomeroy, J. W.: Implications of spatial distributions of snow mass and melt rate for snow-cover depletion: theoretical considerations, *Ann. Glaciol.*, 38, 261–265, 2004.
- Essery, R. H. and Etchevers, P.: Parameter sensitivity in simulations of snowmelt, *J. Geophys. Res.*, 109, D20111, doi:10.1029/2004JD005036, 2004.
- Faria, D. A., Pomeroy, J. W., and Essery, R. H.: Effect of covariance between ablation and snow water equivalent on depletion of snow-covered area in a forest, *Hydrol. Process.*, 14, 2683–2695, 2000.
- 20 Garnier, B. J. and Ohmura, A.: The evaluation of surface variations in solar radiation income, *Sol. Energy*, 13, 21–34, 1970.
- Gray, D. M. and O’Neill, A. J.: Application of the energy budget for predicting snowmelt runoff, in: *Advanced Concepts and Techniques in the Study of Snow and Ice Resources*, National Academy of Sciences, Washington D. C., 108–118, 1974.
- 25 Harding, R. J.: Exchanges of energy and mass associated with a melting snow pack, in: *Modelling Snowmelt-Induced Processes*, edited by: Morris, E. M., IAHS Publication No. 155, IAHS Press, Wallingford, 3–15, 1986.
- 30 Hopkinson, C., Collins, T., Pomeroy, J., Anderson, A., and DeBeer, C.: Investigating terrain and canopy cover controls on snowpack depth in a mountainous environment using airborne lidar, *Hydrol. Earth Syst. Sci.*, submitted, 2010.
- List, R. J.: *Smithsonian Meteorological Tables*, 6th edn., The Smithsonian Institute, Washing-

ton, D. C., 527 pp., 1968.

- Liston, G. E.: Interrelations among snow distribution, snowmelt, and snow cover depletion: implications for atmospheric, hydrologic, and ecologic modeling, *J. Appl. Meteorol.*, 38, 1474–1487, 1999.
- 5 Luce, C. H., Tarboton, D. G., and Cooley, K. R.: Sub-grid parameterization of snow distribution for an energy and mass balance snow cover model, *Hydrol. Process.*, 13, 1921–1933, 1999.
- Male, D. H. and Gray, D. M.: Problems in developing a physically based snowmelt model, *Can. J. Civil Eng.*, 2, 474–488, 1975.
- Marks, D., Domingo, J., Susong, D., Link, T., and Garen, D.: A spatially distributed energy
10 balance snowmelt model for application in mountain basins, *Hydrol. Process.*, 13, 1935–1959, 1999.
- Marks, D., Kimball, J., Tingey, D., and Link, T.: The sensitivity of snowmelt processes to climate conditions and forest cover during rain-on-snow: a case study of the 1996 Pacific Northwest flood, *Hydrol. Process.*, 12, 1569–1587, 1998.
- 15 Marsh, P. and Pomeroy, J. W.: Meltwater fluxes at an arctic forest-tundra site, *Hydrol. Process.*, 10, 1383–1400, 1996.
- Martinec, J., Rango, A., and Roberts, R.: Snowmelt Runoff Model (SRM) user's manual, *Geographica Bernensia P35*, Department of Geography, University of Berne, 84 pp., 1998.
- Moore, R. J., Bell, V. A., Austin, R. M., and Harding, R. J.: Methods for snowmelt forecasting in
20 upland Britain, *Hydrol. Earth Syst. Sci.*, 3, 233–246, 1999,
<http://www.hydrol-earth-syst-sci.net/3/233/1999/>.
- Pomeroy, J. W., Gray, D. M., Shook, K. R., Toth, B., Essery, R. H., Pietroniro, A., and Hedstrom, N. R.: An evaluation of snow accumulation and ablation processes for land surface modelling, *Hydrol. Process.*, 12, 2339–2367, 1998.
- 25 Pomeroy, J. W., Hanson, S., and Faria, D. A.: Small-scale variation in snowmelt energy in a boreal forest: an additional factor controlling depletion of snow cover?, in: *Proceedings of the Eastern Snow Conference*, 58th, 14–17 May 2001, Ottawa, Ontario, 85–96, 2001.
- Pomeroy, J. W., Toth, B., Granger, R. J., Hedstrom, N. R., and Essery, R. H.: Variation in surface energetics during snowmelt in complex terrain, *J. Hydrometeorol.*, 4, 702–716, 2003.
- 30 Pomeroy, J. W., Essery, R. H., and Toth, B.: Implications of spatial distributions of snow mass and melt rate for snow-cover depletion: observations in a subarctic mountain catchment, *Ann. Glaciol.*, 38, 195–201, 2004.
- Pomeroy, J. W., Gray, D. M., Brown, T., Hedstrom, N. R., Quinton, W. L., Granger, R. J., and

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Carey, S. K.: The cold regions hydrological process representation and model: a platform for basing model structure on physical evidence, *Hydrol. Process.*, 21, 2650–2667, 2007.
- Shook, K. R.: Simulation of the ablation of prairie snowcovers, Unpublished Ph. D. thesis, University of Saskatchewan, Saskatoon, 189 pp., 1995.
- 5 Sicart, J. E., Pomeroy, J. W., Essery, R. H., and Bewley, D.: Incoming longwave radiation to melting snow: observations, sensitivity and estimation in northern environments, *Hydrol. Process.*, 20, 3697–3708, 2006.
- Storr, D.: Precipitation variations in a small forested watershed, in: Proceedings of the Annual Western Snow Conference, 35th, 18–20 April 1967, Boise, Idaho, 11–17, 1967.
- 10 Winstral, A., Elder, K., and Davis, R.: Spatial snow modeling of wind-redistributed snow using terrain-based parameters, *J. Hydrometeorol.*, 3, 524–538, 2002.

HESSD

7, 971–1003, 2010

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy

Table 1. RMS errors between simulated and measured snow variables at the three snowpack/meteorological stations on Fisera Ridge for the 2009 snowmelt period. No measurements of internal snowpack temperature were available at the main ridge-top station.

	S-E facing	N facing	ridge-top
SWE	35 mm	41 mm	45 mm
T_{layer1}	0.95 °C	1.21 °C	–
T_{layer2}	0.39 °C	0.73 °C	–
melt rate	6.5 mm/d	7.3 mm/d	8.4 mm/d

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy

Table 2. Statistical parameters of pre-melt SWE distributions over individual slope units in 2008 based on snow surveys and the LiDAR snow depth raster.

Slope unit	$\overline{\text{SWE}}$ (mm)	st. dev. (mm)	CV
S. facing	121	157	1.30
N. facing	158	263	1.67
E. facing	155	187	1.21
Cirque floor	243	347	1.43

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy




Fig. 1. Shaded topographic map of Fisera Ridge and the Upper Middle Creek Basin (hatched area) within the Marmot Creek Research Basin, Alberta. Snow survey transects in 2008 are represented by the dashed lines and location of the detailed Fisera Ridge inset map is indicated. Other insets include an aerial photograph of the basin and a map showing the location of Marmot Creek within Western Canada.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy




Fig. 2. Comparison of simulated and observed SWE and internal snowpack temperature at the south-east facing outlier site on Fisera Ridge during the melt period in 2008 (calibration) and 2009 (validation).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy




Fig. 3. Plots of SWE vs. simulated melt rates at different times in the melt period of 2008 on two opposite slopes over Fisera Ridge.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy




Fig. 4. Simulated vs. observed SCD curves and the SRCA during the early melt period within the Mt. Allan cirque in 2008: **(a)** North-facing slope unit; **(b)** South-facing slope unit. SCD curves were derived based on uniform slope-based melt rates and melt rates obtained from computations based on different initial SWE values (inhomogeneous melt). Observed SCD curves were derived from the orthogonally corrected photographs of the cirque.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy




Fig. 5. Simulated vs. observed SCD curves and the SRCA over the entire spring period within the Mt. Allan cirque in 2008: **(a)** North-facing slope unit; **(b)** South-facing slope unit.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy




Fig. 6. Simulated basin-scale SCD curve and SRCA over the Upper Middle Creek Basin for the spring period of 2008.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Simulation of the snowmelt runoff contributing area

C. M. DeBeer and
J. W. Pomeroy




Fig. 7. Spatial snow-cover and SRCA patterns over the Upper Middle Creek Basin (showing the four different slope-based HRUs) in the early snowmelt period of 2008. Patterns were mapped based on the spatial distribution of snow depth and SWE derived from the repeat LiDAR data. Snow melt patterns were not simulated in the densely forested or cliff areas.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion