

Appendix 2 - References

Duffner, A., Ingwersen, J., Sangchan, W., Hugenschmidt, C., Streck, T., (in preparation): Identifying pesticide transport pathways from a sloped litchi orchard to an adjacent stream based on hydrograph separation.

Hodnett, M.G., and Tomasella, J., (2002): Marked differences between van Genuchten soil water-retention parameters for temperate and tropical soils: A new water-retention pedo-transfer function developed for tropical soils. *Geoderma* 108: 155–180.

Irwin R.R., (1976): Replacing shifting agriculture through intensive settled agriculture, crop diversification and conservation farming. Working Paper 9, UNEP/FAO Mae Sa integrated watershed and forest land use project. Food and Agriculture Organization of the United Nations, Chiang Mai, Thailand.

Kinner, D. A. and R. F. Stallard (2004): Identifying Storm Flow Pathways in a Rainforest Catchment Using Hydrological and Geochemical Modelling. *Hydrological Processes* 18: 2851-2875.

Ng, H. Y. F. and S. B. Clegg (1997): Atrazine and metolachlor losses in runoff events from an agricultural watershed: the importance of runoff components. *The Science of the Total Environment* 193: 215-228.

Schreinemachers, P., Praneetvatakul, S., Sirijinda, A., Berger, T., (2008): Agricultural statistics of the Mae Sa watershed area, Thailand, 2006. Report of the SFB 564 - The Uplands Program, University of Hohenheim, Stuttgart.

Stewart, M. K., Mehlhorn, J., Elliot, S., (2007): Hydrometric and natural tracer (oxygen-18, silica, tritium and sulphur hexafluoride) evidence for a dominant groundwater contribution to Pukemanga Stream, New Zealand. *Hydrological Processes* 21: 3340-3356.

Zimmermann, B., Elsenbeer, H., De Moreas, J. M., (2006): The influence of land-use changes on soil hydraulic properties: Implications for runoff generation. *Forest Ecology and Management* 222: 29-38.