

# Evaluation of Global Continental Hydrology as Simulated by the Land-surface Processes and eXchanges Dynamic Global Vegetation Model

S.J. Murray<sup>1</sup>, P.N. Foster<sup>1</sup> and I.C. Prentice<sup>1,2,3</sup>

<sup>1</sup> *QUEST, Department of Earth Sciences, Wills Memorial Building, University of Bristol, Queens Road, Bristol, BS8 1RJ, United Kingdom*

<sup>2</sup> *Department of Biological Sciences, Macquarie University, North Ryde, NSW 2109, Australia*

<sup>3</sup> *Grantham Institute of Climate Change, and Division of Biology, Imperial College, Silwood Park, Ascot, SL5 7PY, United Kingdom*

*Correspondence to: S.J. Murray (steve.murray@bristol.ac.uk)*

Response to reviewers' comments:

We would like to thank Dieter Gerten and Reviewer #2 for their constructive comments, which have helped us to improve this manuscript. We have carried out a major revision of the text in light of these comments, which are addressed here in turn.

## Reviewer #2's Comments

Main concerns:

- 1) "In order for the model to be used as a hydrologic analysis tool, it needs calibration to estimate reasonable streamflow volume (and also streamflow timing to some extent) for at least few selected river basins."

We believe this comment is wide of the mark, because such calibration would invalidate the main purpose of using a DGVM! In contrast to the classical approach in hydrological modelling, we intentionally do *not* calibrate streamflow to fit observed streamflow for any specific basin. If we did so, we would forfeit our claim to have a generic model applicable to all regions and environments, and under changing environments. The classical approach in hydrological modelling can produce a much better fit to observations under current conditions, but does so by calibration of parameters that could, in reality, vary under changes in climate and other relevant environmental conditions, including atmospheric CO<sub>2</sub> concentration. Which approach should be adopted depends on the time scale of interest. We would argue that at least for the century time scale typically adopted for climate change impact analysis, there is a strong case for using a DGVM, without calibration—even if we recognize that there is a distinct need for improvement in the hydrological process representations in DGVMs, as their application to water resources is a relatively new field.

- 2) “It is recommended to illustrate the differences in model parameterization or set up that were different from the study by Gerten et al. (2004). Further, the authors indicate that few basins improved in streamflow estimation while few became worse. Then what is the additional contribution of this paper in improving streamflow simulations?”

The differences in process representation and model setup relative to LPJ are presented in sections 2.1.3 and 3.1.6. Although a new and more realistic process-oriented fire regime is implemented in the model (Prentice *et al.*, in revision), and fire effects on vegetation certainly would be expected to impact on the partitioning of precipitation between evapotranspiration and runoff, there is no certainty that the demonstrated improvements in fire modelling would lead to improvements in simulated hydrology. The results presented here show that the differences in process representations lead to improvements in hydrology for some catchments, but not so for others, thus highlighting complexities in the controls of runoff.

We do *not* claim, therefore, that LPX (as presently configured) represents a major step forward in terms of runoff simulation; although we do establish that its performance in this respect is at least as good as that of earlier models. The purpose of the MS is rather to *critically evaluate* the hydrological realism of a model that represents the state of the art in DGVM development.

- 3) “Include a brief description how snow accumulation and ablation are represented in the LPX model (e.g. see Fassnacht and Soulis, 2002, ATMOSPHERE-OCEAN, 40 (4), 389-403).”

A description has now been included in section 2.1.1.

- 4) Regarding Dai *et al.* (2009) global river discharge observations: “Where were the gaps in time-series of observations? How were the data filled? Trends are sensitive to the infilling method when gaps are large.”

Data infilling is described in section 2.3 (second-last paragraph) including reference to Dai *et al.* (2009), who provide full details of the procedure. Gaps in the time-series and the values obtained through infilling are shown in Figure 5 of Dai *et al.* (2009) for 24 large catchments.

Minor comments:

We have dealt with all minor comments by implementing suggested changes, or including clarifications or corrections.

Selected points of note:

- 3) “Was the trend statistically significant?”

An error has been corrected and a *t* test has been applied.

- 4) “The model tends to perform well in replicating trends and is often within, or close to the periphery of the runoff envelope shown by the composite and monthly summed converted river discharge data (Fig. 4). This is not apparent from Figure 4.”

This issue has now been clarified in section 3.1.5. It is emphasised that while the model generally captures the interannual variability of runoff, it is subject to bias in some catchments. The bias is quantified at the global scale in section 3.1.3.

- 6) “Why LPX has a propensity to late prediction of peak intra-annual flow when the flow is available at the basin outlet within a given day?”

Difference of 0.068 months is converted to ~2 days. Reasons for slightly late runoff are likely to vary among catchments. In northern catchments, the simplistic snowmelt routine is likely to be an important issue, whereas elsewhere, the lack of explicit flow routing may be the critical point.

- 10) “Axes are not clear in Figure 5. Why summed monthly streamflow are higher than monthly summed precipitation (e.g. Amazon, Congo, Ob, and Lena) in Figure 5?”

Figure 5 has been redrawn and now includes clearer axes. It should be clearer now, in the new version of this Figure, that runoff is *not* simulated to exceed precipitation in the basins mentioned. Simply for clarity of presentation, however, we have used different y-axis scales for precipitation and runoff.

Steve Murray  
Pru Foster  
Colin Prentice

4<sup>th</sup> October 2010