

This discussion paper is/has been under review for the journal Hydrology and Earth System Sciences (HESS). Please refer to the corresponding final paper in HESS if available.

Turbulent flux modelling with a simple 2-layer soil model and extrapolated surface temperature applied at Nam Co Lake basin on the Tibetan Plateau

T. Gerken^{1,2}, W. Babel², A. Hoffmann¹, T. Biermann², M. Herzog¹, A. D. Friend³, M. Li⁴, Y. Ma⁵, T. Foken², and H.-F. Graf¹

¹Centre for Atmospheric Science, Department of Geography, University of Cambridge, UK

²Department of Micrometeorology, University of Bayreuth, Germany

³Department of Geography, University of Cambridge, UK

⁴Cold and Arid Region Environmental and Engineering Research Institute, Chinese Academy of Sciences, Lanzhou, China

⁵Institute of Tibetan Plateau Research, Chinese Academy of Sciences, Beijing, China

Received: 19 October 2011 – Accepted: 31 October 2011 – Published: 21 November 2011

Correspondence to: T. Gerken (tobias.gerken@uni-bayreuth.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

This paper introduces a surface model with two soil-layers for use in a high-resolution circulation model that has been modified with an extrapolated surface temperature, to be used for the calculation of turbulent fluxes. A quadratic temperature profile based on the layer mean and base temperature is assumed in each layer and extended to the surface. The model is tested at two sites on the Tibetan Plateau near Nam Co Lake during four days during the 2009 Monsoon season. In comparison to a two-layer model without explicit surface temperature estimate, there is a greatly reduced delay in diurnal flux cycles and the modelled surface temperature is much closer to observations. Comparison with a SVAT model and eddy covariance measurements shows an overall reasonable model performance based on RMSD and cross correlation comparisons between the modified and original model. A potential limitation of the model is the need for careful initialisation of the initial soil temperature profile, that requires field measurements. We show that the modified model is capable of reproducing fluxes of similar magnitudes and dynamics as the more complex methods chosen as reference.

1 Introduction

Turbulent fluxes of momentum, latent and sensible heat are some of the most important interactions between land surface and atmosphere. These fluxes are responsible for the development or modification of mesoscale circulations and the generation of clouds that, as a consequence, feed back on surface fluxes through the modification of solar radiation. The effects of vegetation influencing boundary layer structure and moisture are widely acknowledged (i.e. Freedman et al., 2001; van Heerwaarden et al., 2009), while the feedback from short-lived clouds is less understood, but important. Shallow cumulus-surface interactions were shown in an LES study to impact surface temperature and fluxes on very short time scales (Lohou and Patton, 2011). For improved process understanding, it is necessary to use: (1) atmospheric models with sufficiently

A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



high resolution to resolve boundary layer processes as well as clouds and (2) surface models capable of reproducing the system's surface flux dynamics.

Our research focuses on surface-atmosphere interactions on the Tibetan Plateau (TP) in the Nam Co Lake region. With more than 4700 m a.s.l., a semi-arid climate and with a highly adapted *Kobresia pygmaea* alpine steppe (Miehe et al., 2011), the TP proves to be a difficult environment for surface models (Yang et al., 2003, 2009). Specific problems include large temporal and spatial variability in soil moisture (Su et al., 2011), large diurnal variations of surface temperature from surface freezing before sunrise to more than 30 °C at noon. Ma et al. (2009) give an overview about the TP surface-atmosphere processes. On the TP, the fraction of diffuse solar radiation is very small, making cloud feedbacks especially important for the surface-atmosphere system. The model studies with a regional model of Cui et al. (2007) imply that some of the precipitation events on the TP are predominantly local and therefore not captured by coarser resolution models.

In this paper we present results of a rather simple two-layer flux algorithm based on a two-layer soil model that is part of a vegetation dynamics and biosphere model (Friend, 2010). The original model produces a substantial delay in the diurnal turbulent flux cycle due to the low responsiveness of the upper model layer to changes in atmospheric forcing and fails to capture important dynamics. We therefore introduce an extrapolated surface temperature and show that this is capable of reproducing diurnal flux dynamics for two vegetation covered surfaces near Nam Co Lake. These sites are representative for the basin, but show very different dynamics. In our future studies, the same surface-model version will also be coupled to a spatially and temporally high resolution atmospheric model including radiation, cloud microphysics and active tracer transport. As simulations of the high-resolution model will be run for approximately 24 h we tested the surface model in column mode forced with standard atmospheric measurements for the same period of time with initialisation at 00:00 h Beijing Standard Time (BST). We acknowledge that this approach is different from most surface model studies that are run for longer periods, but it is necessary for the planned study

A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

of the coupled surface-atmosphere system. Such a surface flux algorithm is generally suitable for high-resolution atmospheric modelling studies of different ecosystems as it does not have built in assumptions about horizontal scales.

It is our objective to test the suitability of a simple two-layer soil model with an improved surface or “skin” temperature estimated from the mean temperature of the uppermost layer that shall subsequently be used for driving an atmospheric circulation model for the Nam Co region on the TP. Therefore, fluxes derived from the surface flux algorithm with and without a specific formulation for “skin” temperature are compared to measured fluxes by eddy covariance technique and to fluxes derived by a more complex Soil-Vegetation-Atmosphere Transport (SVAT) Model, with five soil layers.

2 Site description and experimental data

Nam Co Lake is located on the Tibetan Plateau at approximately 4730 m a.s.l., circa 150 km north of Lhasa. During the monsoon season of 2009 a joint measurement campaign was conducted from 27 June to 8 August by the University of Bayreuth (UBT) and the Institute of Tibetan Plateau Research – Chinese Academy of Sciences (ITP). Data from 2 locations are used. Site 1, both referred to and operated by UBT is an eddy covariance complex close to the shore of a small lake next to Nam Co lake. UBT has a fairly constant soil moisture below circa 60 cm depth due to the influence of a water table. Additionally, the atmospheric measurements are influenced by a land lake breeze. Site 2 (operated by and referred to as ITP) is at the Nam Co Station for Multisphere Observation and Research (Li et al., 2009; Cong et al., 2009), approximately 300 m distant from both UBT and the lake with a sandy soil and a very low field capacity (FC = 5 %) compared to overall pore volume (39 %). The vegetation at both sites is grassland (Metzger et al., 2006) with UBT having a small bare soil fraction (0.1) compared to 0.4 at ITP). Small FC and the generally low volumetric top soil water contents (θ_v) at ITP, lead to large sensible energy fluxes compared to latent heat fluxes ($Q_H \gg Q_E$). After rain events however, θ_v may exceed FC by a factor of up to 3 leading

A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to a similar flux regime at the two stations with $Q_E > Q_H$. Due to the generally drier conditions at ITP, surface temperature frequently drops below 0°C in the early morning hours. At UBT there were soil temperature sensors installed at 2.5, 5, 10, 20, 30 and 50 cm depths. At ITP no accurate soil temperatures were available at depths less than 20 cm. Comprehensive information of ITP and UBT surface and soil properties is found in Table 1.

2.1 Forcing data

The 24-h model runs are initialised with the soil temperature profile and soil moisture at 00:00 h BST ($\sim 22:00$ in local solar time). Four days were selected from the campaign representing different weather conditions. 10 July was a complex day with rain in the morning and sunshine in the afternoon. 27 July was a cloudy day without rain. 5 August was a radiation day after a period of rain leading to moist conditions and large Q_E at ITP. 6 August was similar to the previous day, but with some of the water drained from the soil at ITP and developing clouds in the afternoon. During 10 July and 5 August, the station close to the lake (UBT) came under the influence of a lake breeze during which the forcing data (except for radiation measurements) correspond rather to the nearby lake than the land surface. Due to the overcast sky on 27 July the lake breeze was severely weakened as described in Zhou et al. (2011), so that there was only limited influence of the lake surface onto the atmospheric measurements.

The model is forced with measured atmospheric data from two sites on the TP providing air temperature, water vapour mixing ratio, wind speed, air pressure, precipitation and downwelling long and shortwave radiation. These data are displayed in Fig. 1 for UBT and Fig. 2 for ITP. In general 30-min mean values were linearly interpolated to the surface model time step that was the same as a typical time step of an atmospheric model ($\Delta t = 2.5\text{ s}$). The only selected day with precipitation (rain) during day-time was 10 July 2009. However, there was also rain recorded at UBT from about 22:00 h BST on 6 August 2009. Half hourly precipitation was scaled down to the model time step assuming a constant precipitation rate per 30-min interval. There was little difference

A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



between the data measured at ITP and UBT, as expected due to the proximity of the sites. However there was an offset of approx. 5 hPa between the recorded pressures, that was not corrected for as this is likely within the uncertainty of the sensors and the model should not be too sensitive to such a pressure difference. Unlike UBT where rain 30-min precipitation was available, there were only daily sums recorded for ITP, which had to be downscaled to 30-min values by scaling them linearly with UBT observations.

3 Modelling approach

The surface model Hybrid (Friend et al., 1997; Friend and Kiang, 2005) is currently coupled to the high-resolution Active Tracer High-resolution Atmospheric Model (ATHAM) by Oberhuber et al. (1998) and Herzog et al. (1998) for the investigation of feed-backs between atmospheric processes and surface fluxes.

Our high-resolution modelling approach aims at a spatial and temporal resolution in the order of 500 m and 2.5 s, respectively. As our focus is on diurnal surface-atmosphere interactions, the surface model must capture the magnitude of the fluxes and must be able to react quickly to changes in atmospheric forcing. Therefore, a surface model that is capable of reproducing realistic turbulent energy and water vapour fluxes at a sufficiently high temporal resolution and at reasonable computational costs is needed. We decided against a model with more than two soil-layers due to higher computational cost and instead modified the original Hybrid model to meet these requirements.

3.1 The surface model

The modified version of Hybrid which is a process based terrestrial ecosystem and surface model, incorporates a simple two-layer representation of the soil and uses the turbulent transfer parameterisations taken from the GISS model II (Hansen et al., 1983). The transfer equations in Hybrid are described in Friend and Kiang (2005). Bare

A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



soil parameterisation follows the approach of SSiB (Xue et al., 1996) that is based on Camillo and Gurney (1986) and Sellers et al. (1986). Turbulent fluxes are calculated using a bulk approach for the sensible heat flux (Q_H):

$$Q_H = c_p \rho C_H u(z) (T_0 - T(z)) \quad (1)$$

5 with air specific heat capacity (c_p [$\text{Jkg}^{-1}\text{K}^{-1}$]), the Stanton number (C_H) which is calculated as a function of roughness length z_0 and Bulk Richardson Number, air density (ρ [kgm^{-3}]), measured wind speed ($u(z)$ [ms^{-1}]) and air temperature ($T(z)$) at measurement height z [m] and with T_0 being the surface temperature. All temperatures used are in K. The latent heat flux (Q_E) is derived in a more complex manner from bulk soil
10 evaporation and a canopy resistance approach estimating plant transpiration. Transfer coefficients are modified from Deardorff (1968). Plant physiology and stomatal conductance are included via generalised plant types (GPT). As an ecosystem model Hybrid is designed to work on hourly to climate scales (Friend, 2010) and should therefore be capable of reproducing diurnal flux cycles as well as ecosystem changes on climate
15 scales. It was originally developed as a biosphere-surface component for the GISS GCM. A “thin” upper layer with 10 cm thickness follows the daily cycle of surface temperatures, whereas a lower layer with 4 m thickness acts as the memory for the annual cycle. However, an upper layer of such thickness imposes a substantial dampening on the diurnal temperature cycle and will effectively act as a lowpass filter for events
20 of short durations such as cloud shading that, especially under the conditions found at the TP, has a substantial immediate impact on surface temperatures and on fluxes as well. This is also seen in Fig. 12 of Hansen et al. (1983), where a time delay of approximately 2 h is visible for surface temperature in the diurnal cycle. A similar behaviour of the original Hybrid is discussed in Sect. 5.4. Shortcomings with the representation of
25 diurnal cycles may also impact on longer term studies as the model drifts away from a realistic state. As we plan to apply the coupled model for high-resolution simulations with a time step in the order of seconds, we focus in this work on the accuracy of the diurnal flux cycles that can be achieved with such a model.

A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.2 The modified soil model in Hybrid

Two main changes that lead to a better representation of the actual soil state have been introduced to Hybrid. They address its main problems of a delay in diurnal flux evolution and its weak responsiveness to sudden short-term changes in atmospheric forcing.

3.2.1 Diagnostic surface temperature

An extrapolated surface temperature (T_0) is being introduced that is then subsequently used for the calculation of atmospheric stability through the Bulk Richardson number as well as for Q_H and Q_E . This approach is somewhat similar to the “force-restore method” (Blackadar, 1979) that also aims at providing a realistic surface temperature imitating the behaviour of real soils. However, while “force-restore” uses an oscillating heat source as forcing term and a heat-flux into the ground as restoring term (Yee, 1988), our method is not dependent on a periodic heating function and uses the concept of layer heat storage. T_0 is derived from a set of assumptions that were already included in Hybrid going back to Hansen et al. (1983). For both layers denoted with the subscripts 1 and 2 from the model top, we assume a quadratic temperature profile $T(z)$ (Fig. 3):

$$T_{1,2}(z_{\text{rel}}) = a_{1,2} (z_{\text{rel}} - d_{1,2})^2 + T_{\text{base}_{1,2}} \quad (2)$$

with a [Km^{-2}] being a constant, z_{rel} [m] the depth below the top of the layer, d [m] the layer thickness and T_{base} as temperature at the base of the respective layers. There is no transfer of heat through the lower boundary of the model so that T_{base_2} is constant and equal to annual mean temperature. The relationship between layer heat content E [J] and temperature profile is given by:

$$E = c_p \int_{z_L}^{z_U} T(z) dz \quad (3)$$

A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where z_L and z_U are the lower and upper boundaries of the layer and c_{p_s} [$\text{Jm}^{-3}\text{K}^{-1}$] is the total soil heat capacity. Hence, with a known heat content for each layer it is possible to solve for

$$a_2 = \frac{\frac{E_2}{c_{p,2}} - d_2 T_{\text{base}_2}}{\frac{d_2^3}{3}}, \quad (4)$$

5 by integrating Eq. (3) with Eq. (2) and solving for a_2 , thus deriving the base temperature of the first layer:

$$T_{\text{base}_1} = T_{\text{base}_2} + a_2 d_2^2. \quad (5)$$

In a similar fashion a_1 and T_0 can be approximated:

$$a_1 = \frac{\frac{E_1}{c_{p,1}} - d_1 T_{\text{base}_1}}{\frac{-z_1^3}{3}} \quad (6)$$

10 and

$$T_0 = T_{\text{base}_1} + a_1 d_1^2. \quad (7)$$

3.2.2 Heat diffusion estimation

In the original heat diffusion algorithm of Hybrid (Hansen et al., 1983), the heat flux from the first to the second soil layer $F(z)$ is dependent on the difference between mean surface layer temperatures (\bar{T}), the soil heat flux calculated as residual of turbulent and radiation fluxes ($F(0)$), layer thickness and thermal resistances r ,

$$F(z) = \frac{3\bar{T}_1 - 3\bar{T}_2 - 0.5F(0)r_1}{r_1 + r_2} \times \Delta t, \quad (8)$$

where Δt is model time-step. This leads to unrealistic modelled heat fluxes $F(z)$ as $F(z)$ is largely dominated by $F(0)$, which is positive during nighttime and negative during

A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



daytime, thus leading to a net transfer of heat from a cold to a warm layer. With the assumption of a subgrid temperature profile the heat flux between the two layers Eq. (8) was modified with a heat diffusion approach and integration of

$$\frac{\partial T}{\partial t} = D \frac{\partial^2 T}{\partial z^2} \quad (9)$$

with D being the diffusion constant for heat. We assume ∂z to be approximated by the diffusion length $L = 2\sqrt{\Delta t D} = \Delta z$. As the model is run with the short time-step of the atmospheric model, such a formulation becomes valid. A rough calculation for L with $D = 10^{-6} \text{ m}^2 \text{ s}^{-1}$, which is close to the determined value, and $\Delta t = 30 \text{ min}$ gives $L = 0.08 \text{ m}$, which is close to d_1 , posing an upper limit on Δt for this method.

3.3 Surface temperature profile initialisation

Due to the quadratic nature of the soil layer temperature profiles and their potential kink at the layer interface (see Fig. 3), the modified model depends on careful initialisation that full-fills two requirements: (1) a realistic estimate of surface temperature and (2) an appropriate estimate of ground heat storage (E) allowing the upper layer to react in a realistic way. In this study soil temperature measurements at several depths were used in order to accomplish both requirements. Surface temperature was estimated from upwelling longwave radiation according to the Stefan-Boltzmann law with a longwave emissivity of $\epsilon = 0.97$. The 10 cm temperature at UBT was used as T_{base_1} and the temperature curve for the upper soil layer was fitted to measurements by minimising the squared mean error. Due to the lacking 10 cm temperature at ITP, this temperature had to be estimated from the 20 cm measurement. It should be noted that the assumed quadratic temperature profile in the lower soil layer clearly underestimated the vertical temperature gradient in the soil as estimated UBT temperatures at 50 cm were always higher than measured temperatures. This difference is reduced from July to August as the summer warming reaches lower layers. This is a limitation due to fixed layer depths.

A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2 shows the initial temperatures for each day. From the span of layer temperatures \bar{T}_1 and \bar{T}_2 , the theoretical parameter space of T_0 for a constant T_{base_2} (Fig. 4) can be derived. A random combination of the initial temperatures given in Table 2 would yield T_0 in the range of -10 to 30°C . In contrast, the actual model layer temperatures, indicated by the crosses in Fig. 4c, occupy a much smaller area and are, with the exception of one day, clustered closely. This highlights the importance of a careful initialisation of the soil temperature profile requiring knowledge about subsurface temperatures that are difficult to estimate without field measurements.

4 Flux comparison

Surface fluxes derived with any method contain inaccuracies such as measurement errors or theoretical limitations. Therefore we are not comparing our modelling results to the absolute truth, but to two flux references.

4.1 EC and SEWAB reference fluxes

Fluxes estimated by both versions of Hybrid are compared with observed fluxes derived by eddy covariance (EC) method and fluxes modeled by the SVAT model SEWAB (Surface Energy and Water Balance model – Mengelkamp et al., 1999), which has been calibrated to the two sites for gap-filling and up-scaling of flux measurements. Both flux references yield fluxes averaged over 30-min intervals. Unlike many SVAT models that derive the soil heat flux from the flux residual, SEWAB is solving the surface energy balance equation ($Q_E + Q_H + Q_{\text{Rad}} + Q_{\text{Soil}} = 0$) iteratively for T_0 by Brent's method (Mengelkamp et al., 1999), hence closing the energy balance locally (Kracher et al., 2009). In contrast, the surface energy balance closure derived by EC is only in the order of 0.7 at Nam Co Lake (Zhou et al., 2011). This means that 30% of the net radiation is not captured by surface flux measurements. However, energy balance closure must not be used as a quality measure for flux measurements (Aubinet et al., 1999)

A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

as surface heterogeneity leads to organised low frequency structures and mesoscale circulations (Panin et al., 1998; Kanda et al., 2004) that are mainly responsible for the lack of closure (Foken, 2008). This is especially true for sea (lake) breeze systems, where a significant portion of the energy fluxes is transported horizontally (Kuwagata et al., 1994). Therefore, SEWAB (and Hybrid) fluxes are comparatively larger than measured ones. When the energy balance is closed artificially by redistributing the residual to fluxes according to the Bowen ratio (Twine et al., 2000), the resulting fluxes are in much better agreement with SEWAB (Babel et al., 2011). Therefore we use, when possible, energy balance corrected fluxes ($Q_{E_{EC,EBC}}$ and $Q_{H_{EC,EBC}}$). EC data were collected at 3 m height and calculated using the TK3 package (Mauder et al., 2008; Mauder and Foken, 2011). Quality checks were performed according to Foken et al. (2004). A detailed description of the instrumentation can be found in Biermann et al. (2009). The rain event of 10 July leads to the exclusion of fluxes due to quality concerns. Both Hybrid and SEWAB produce fluxes during rain, but their quality is unknown as they cannot be compared to measurements.

Measuring in close proximity to the lake also means that depending on wind direction the fluxes measured at UBT are originating from land, water or a mixture of both as the footprint of the EC system and the forcing data is located upwind of the site. This leads to problems in the energy balance closure and the integration of fluxes. The development of a lake breeze system at Nam Co means that during most days there are no flux measurements available from the late morning or early afternoon until the lake breeze ceases. The days of 27 July and 6 August are the only days during which the field campaign provides data that do not have a full lake breeze influence at UBT. Therefore it is beneficial to compare not only to measured fluxes, but also to SEWAB ($Q_{E_{SEWAB}}$ and $Q_{H_{SEWAB}}$).

For completeness, fluxes over the lake calculated by the TOGA-COARE algorithm (Fairall et al., 1996a,b) that is also part of the coupled surface-atmosphere model are given during lake breeze events.

4.2 Statistical evaluation measures

In order to assess the model quality we use Root Mean Square Deviation (RMSD) and Cross Correlation (Eq. 11):

$$\text{RMSD} = \sqrt{\frac{1}{N} \sum_{i=1}^N (P_p - P_r)_i^2} \quad (10)$$

$$R^2(j) = \left(\frac{\text{cov}(P_p(1+j:N), P_r(1:N-j))}{\sigma_{P_p(1+j:N)} \sigma_{P_r(1:N-j)}} \right)^2 \quad (11)$$

with $R^2(j)$ being the coefficients of determination for the predicted (P_p) and reference (P_r) flux time series shifted by j elements, the total number of elements in each time series (N) and σ as their respective standard deviations. Both SEWAB and EC measurements produce 30-min flux averages, whereas Hybrid was set to 10-min averaged fluxes. Therefore the reference fluxes were linearly interpolated to Hybrid's output times before statistical evaluation. Periods when no energy balance corrected EC measurements were available were excluded from the calculation of the statistical measures.

5 Results and discussion

The following section presents and discusses the improvements that are achieved for a simple two-layer model when the formulation for surface temperature is included.

The original two-layer model Hybrid fails to reproduce the diurnal dynamics observed at UBT due to the thermal inertia of the top-layer. The delayed response in surface temperature leads to a shift in the resulting turbulent surface fluxes. This causes an underestimation of Q_E and Q_H until $\sim 18:00$ h BST and later to an overestimation due to delayed surface cooling. The improvement of the modified Hybrid over the original formulation is discussed in more detail in Sects. 5.1 and 5.4.

A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5.1 Discussion of turbulent fluxes

The original two layer model reacts only slowly to the atmospheric forcing, delaying the fluxes' response to atmospheric forcing. Such a time lag leads to a shift in the diurnal cycle and is problematic for the coupling to atmospheric models since surface fluxes are one of the main drivers of regional and local circulation as well as cloud development. These will certainly be affected by erroneous surface flux dynamics. In our specific case, the dampening of the diurnal temperature cycle and the delay in surface fluxes may reduce the intensity of the land-lake breeze or may delay its development through a reduction of differential heating between land and lake surface. However, there is still a minor delay visible in the modified Hybrid as the surface temperature is purely diagnostic and dependent on \bar{T}_1 . This is discussed in more detail in Sect. 5.4.

Table 3 shows the results of the RMSD between the modelled results and the reference quantities. With the modified Hybrid model there is a 40–60 % improvement in the RMSDs compared to the original Hybrid, when both are compared against SEWAB. The only notable exception for this is 6 August at ITP, where a strong deviation of turbulent fluxes derived by SEWAB and measured fluxes was encountered. This is due to a problem with SEWAB for this day as it falls into a dry interval between rainy periods, where SEWAB underestimates the soil water content. The picture is more diverse for the comparison between the energy balance corrected EC fluxes and Hybrid. There is reduction in the error for all instances, except Q_H on 6 August at ITP, but the reductions cover a much larger range from less than 1 % to 80 %. Due to data quality concerns the number of comparable elements is much lower (N given in Table 3) and probably too small for meaningful statistics in case of UBT. This is especially true as the daytime lake breeze influence coincides with the times when one would expect the best quality of EC fluxes. As flux qualities are usually lower during conditions with limited vertical exchange (stable stratification), EC fluxes at ITP mainly reflect the daytime model performance whereas the comparison with SEWAB also takes into account the night-time,

A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



This may either be due to errors in soil moisture, surface emissivity (ϵ) or due to the surface temperature extrapolation function used in this work.

5.3 Soil moisture variation and evapotranspiration

After a 24 h run the moisture content of the first model layer (last two columns of Table 2) is smaller than measurements suggest. For the case UBT, measured soil moisture content does hardly vary on a day to day scale and is kept well above FC due to groundwater influence. This is not reflected by the model as it lacks the the capability to include groundwater tables. The true soil moisture at ITP has a much larger variation due to its low FC and comparatively large pore volume. During the dry day of 27 July the upper soil layer loses 1.5 mm of water whereas during the moist days of August there is a total loss from layer one of 6.7 and 5.3 mmd^{-1} , respectively. Comparing $\theta_{1,\text{end}}$ of 5 August with $\theta_{1,\text{obs}}$ of the next day shows that the model would perform considerably worse if it were not restarted every day. This is caused by a very limited soil hydrology included in Hybrid. Hu et al. (2008) have estimated the summer evapotranspiration on a central Tibetan grassland site to be in the order of 4–6 mmd^{-1} . An experiment conducted within the framework of TiP has estimated bare soil evaporation and evapotranspiration of a very dry soil at Kema in 2010 (~150 km northeast of Nam Co Lake) at 2 mmd^{-1} rising to at least 6 mmd^{-1} and possibly more for a vegetated *Kobresia* pasture during an irrigation experiment (H. Coners – University of Göttingen, personal communication, 27 June 2011). Even though the soils are not directly comparable this suggests similar dynamics in Q_E to the ITP site. One factor likely to play a role in the local water cycle that is not included is dew fall in the early morning hours. Direct absorption of atmospheric moisture on bare soil (Agam (Ninari) and Berliner, 2004) and dew fall are often considered a significant moisture input for semi-arid environments (Agam and Berliner, 2006). Heavy dewfall in the vicinity of Nam Co Lake is frequently observed, but has, at least to our knowledge, never been quantified. This additional source of water and the associated local recycling of water may account for a significant fraction of the missing water. In addition to this, the too simplistic representation

A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of soil hydrology is very likely responsible for the remaining water deficit in the upper layer of the soil model.

5.4 Cross correlation of turbulent fluxes

A different way of looking at the model performance is cross correlation of the modelled surface fluxes against EC measurements and SEWAB (Fig. 7). These measures give an insight into the reasons for the delayed response of the surface model and the amount of flux-variance explained, but does not yield information whether the model and the reference fluxes show a true one-to-one correlation. As with RMSD the quality of the analysis is limited by the number of data points that can be correlated, which is comparatively small for the energy balance corrected EC measurements at ITP and even smaller at UBT due to lake breeze influences (Fig. 7a–e). Hence, it is very difficult to interpret the cross correlations for EC. It is probably fair to say that there is a tendency for smaller time lags during the time series with higher number of elements, notably UBT 6 August and all days of ITP and that the total explained variances are at the same level of determination, when comparing the maximum $R^2(j)$. A notable exception is ITP 5 August.

For the comparison with SEWAB (Fig. 7f–h), it becomes notable that for many cases the maximum $R^2(j)$ of the modified Hybrid approach $R^2 \rightarrow 1$ and that their maxima are usually found at lags of 10–30 min ($j = 1 - 3$). Solar radiation rapidly modifies the skin temperature that is governing turbulent fluxes. As SEWAB has an instantaneous surface temperature solver for each model time step, one would expect a direct response of SEWAB to changes in solar radiation. This may even be faster than in reality, especially for Q_E flux that is not only dependent on the actual skin temperature, but also on the vegetation's response. Including negative values of j into Fig. 7 would show a gradual decrease of correlations with decreasing j , showing that the flux dynamics of Hybrid never precede EC measurements or SEWAB.

A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

5.5 Natural variability of fluxes

Atmospheric quantities and turbulent surface fluxes have a large natural variability that is difficult to measure or to model. The EC approach is dependent on averaging procedures and most standard measurements will yield mean values. In order to use high-frequency measurements for flux estimation, less common techniques such as conditional sampling or wavelet-spectra have to be used. Even if models are capable of reproducing variability on realistic scales it is difficult to supply forcing data with similar resolution. The forcing data used in this study, sampled and averaged 10 or 30 min means, are used for SEWAB. Running Hybrid at time steps comparable to a high-resolution mesoscale model requires interpolation of the forcing data and therefore potentially causes a smoothing of the model's response compared to the actual weather forcing as it would be provided by a coupled model. As surface models share a similar approach to the parameterisation of surface fluxes and close the surface energy balance locally, SEWAB and Hybrid fluxes are more similar to each other than they are to field measurements.

6 Conclusions

The accurate generation of surface fluxes is a necessary pre-requisite for studies of surface-atmosphere interactions and local to mesoscale circulations. In order to gain a better process understanding of the interaction between atmospheric circulation, clouds, radiation and surface fluxes, the generated diurnal flux cycles have to be of realistic magnitude and without temporal shift. The original two-layer surface model without a specific formulation for T_0 produced both a considerable time lag and failed to capture the full diurnal dynamics due to its unresponsiveness.

We have demonstrated that the introduction of an extrapolated surface temperature enables even a quite simplistic soil model to realistically estimate turbulent surface fluxes and skin temperatures. The delay of fluxes during the daily cycle was greatly

HESSD

8, 10275–10309, 2011

A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



reduced, making the model usable for diurnal process studies. The total magnitude of fluxes is also much improved, when few and computationally cheap additional physically based processes are introduced. Comparing SEWAB with Hybrid, the RMSD for both fluxes and surface temperature is decreased by generally 40–60%. The improvement in quality was somewhat more varied in comparison to EC measurements, as comparison of models and measurements is not straight forward. The improved $R^2(j)$ for smaller values of j shows that temporal shifts of the flux time series have been greatly reduced and the overall correlations are high. As with any natural system it is impossible to obtain complete data sets that capture the full amount of natural variability. However, the modified model has been tested for a larger spectrum of environmental conditions on the TP and produced reasonable results for both dry and moist conditions.

We have shown that a rather simple soil surface model can efficiently calculate turbulent fluxes at a high temporal resolution when driven by realistic atmospheric conditions. Nevertheless, it is quite clear that such an approach with extrapolated surface temperature needs careful model initialisation. The initial soil heat contents and therefore knowledge of soil temperature profiles is necessary. Due to the fact that the surface temperature in this study is a purely diagnostic quantity, there may still be some limitations such as a delayed or smoothed response to atmospheric forcing on very short timescales, such as the feedback between passing boundary-layer clouds and the surface fluxes. The influence of surface fluxes and their dynamics to regional circulation will be investigated in a future study.

Acknowledgements. This research was funded by the German Research Foundation (DFG) Priority Programme 1372 “Tibetan Plateau: Formation, Climate, Ecosystems” as part of the Atmosphere – Ecology – Glaciology – Cluster (TiP-AEG). ITP data was provided through CEOP-AEGIS, which is a EU-FP7 Collaborative Project/ Small or medium-scale focused research project – Specific International Co-operation Action coordinated by the University of Strasbourg, France under the call ENV.2007.4.1.4.2: “Improving observing systems for water resource management.” The authors would like to thank everyone who has contributed to the collection of field data in one of the world’s most remote regions. ADF acknowledges

A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



support from the European Community's Seventh Framework Programme (FP7/2007-2013) under grant agreement no. 238366. AH acknowledges funding by the Fonds National de la Recherche (FNR-Luxembourg), under the grant BFR07-089 and support by the Cambridge European Trust (CET-UK).

5 References

Agam (Ninari), N. and Berliner, P. R.: Diurnal Water Content Changes in the Bare Soil of a Coastal Desert, *J. Hydrometeorol.*, 5, 922–933, doi:10.1175/1525-7541(2004)005;0922:DWCCIT;2.0.CO;2, 2004. 10291

Agam, N. and Berliner, P. R.: Dew formation and water vapor adsorption in semi-arid environments – A review, *J. Arid Environ.*, 65, 572–590, doi:10.1016/j.jaridenv.2005.09.004, 2006. 10291

Aubinet, M., Grelle, A., Ibrom, A., Rannik, U., Moncrieff, J., Foken, T., Kowalski, A., Martin, P., Berbigier, P., Bernhofer, C., Clement, R., Elbers, J., Granier, A., Grünwald, T., Morgenstern, K., Pilegaard, K., Rebmann, C., Snijders, W., Valentini, R., Vesala, T., Fitter, A., and Raffaelli, D.: Estimates of the Annual Net Carbon and Water Exchange of Forests: The EUROFLUX Methodology, *Adv. Ecol. Res.*, 30, 113–175, doi:10.1016/S0065-2504(08)60018-5, 1999. 10285

Babel, W., Huneke, S., and Foken, T.: A framework to utilize turbulent flux measurements for mesoscale models and remote sensing applications, *Hydrol. Earth Syst. Sci. Discuss.*, 8, 5165–5225, doi:10.5194/hessd-8-5165-2011, 2011. 10286

Biermann, T., Babel, W., Olesch, J., and Foken, T.: Mesoscale Circulations and Energy and Gas Exchange over the Tibetan Plateau – Documentation of the Micrometeorological Experiment, Nam Tso, Tibet – 25 June–8 August 2009, *Arbeitsergebnisse* 41, University of Bayreuth, ISSN 1614-8616, Bayreuth, 2009. 10286

Blackadar, A.: High Resolution Models of the Planetary Boundary Layer, in: *Advances in Environmental Science and Engineering*, edited by: Pfafflin, J. and Ziegler, E., vol. 1, 50–85, Gordon and Breach, New York, 1979. 10282

Camillo, P. J. and Gurney, R. J.: A resistance parameter for bare-soil evaporation models, *Soil Sci.*, 141, 95–105, 1986. 10281

Cong, Z., Kang, S., Smirnov, A., and Holben, B.: Aerosol optical properties at

HESSD

8, 10275–10309, 2011

A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Nam Co, a remote site in central Tibetan Plateau, Atmos. Res., 92, 42–48, doi:10.1016/j.atmosres.2008.08.005, 2009. 10278

Cui, X., Langmann, B., and Graf, H.: Summer Monsoonal Rainfall Simulation on the Tibetan Plateau with a Regional Climate Model Using a One-way Double-nesting System, SOLA, 3, 49–52, 2007. 10277

Deardorff, J. W.: Dependence of Air-Sea Transfer Coefficients on Bulk Stability, J. Geoph. Res., 73, 2549–2557, 1968. 10281

Fairall, C. W., Bradley, E. F., Godfrey, J. S., Wick, G. A., Edson, J. B., and Young, G. S.: Cool-skin and warm-layer effects on sea surface temperature, J. Geoph. Res., 101, 1295–1308, 1996a. 10286

Fairall, C. W., Bradley, E. F., Rogers, D. P., Edson, J. B., and Young, G. S.: Bulk parameterization of air-sea fluxes for Tropical Ocean-Global Atmosphere Coupled-Ocean Atmosphere Response Experiment, J. Geophys. Res., 101, 3747–3764, 1996b. 10286

Foken, T.: The parameterisation of the energy exchange across the air-sea interface, Dynam. Atmos. Ocean, 8, 297–305, doi:10.1016/0377-0265(84)90014-9, 1984. 10288, 10307

Foken, T.: The Energy Balance Closure Problem: An Overview, Ecol. Appl., 18, 1351–1367, 2008. 10286

Foken, T., Göckede, M., Mauder, M., Mahrt, L., Amiro, B., and Munger, J.: Post field data quality control, in: Handbook of Micrometeorology: A guide for surface flux measurements and analysis, edited by: Lee, X., Massman, W., and Law, B., 181–208, Kluwer, Dordrecht, 2004. 10286, 10307

Freedman, J. M., Fitzjarrald, D. R., Moore, K. E., and Sakai, R. K.: Boundary Layer Clouds and Vegetation-Atmosphere Feedbacks, J. Clim., 14, 180–197, doi:10.1175/1520-0442(2001)013;0180:BLCAVA;2.0.CO;2, 2001. 10276

Friend, A. D.: Terrestrial plant production and climate change, J. Exp. Bot., 61, 1293–1309, doi:10.1093/jxb/erq019, 2010. 10277, 10281

Friend, A. D. and Kiang, N. Y.: Land Surface Model Development for the GISS GCM: Effects of Improved Canopy Physiology on Simulated Climate, J. Clim., 18, 2883–2902, 2005. 10280

Friend, A. D., Stevens, A. K., Knox, R. G., and Cannell, M. G. R.: A process-based, terrestrial biosphere model of ecosystem dynamics (Hybrid v3.0), Ecol. Model., 95, 249–287, 1997. 10280

Hansen, J., Russell, G., Rind, D., Stone, P., Lacis, A., Lebedeff, S., Ruedy, R., and Travis, L.: Efficient Three-Dimensional Global Models for Climate Studies: Models I and II, Mon.

- Weather Rev., 111, 609–662, 1983. 10280, 10281, 10282, 10283
- Herzog, M., Graf, H., Textor, C., and Oberhuber, J. M.: The effect of phase changes of water on the development of volcanic plumes, *J. Volcanol. Geoth. Res.*, 87, 55–74, doi:10.1016/S0377-0273(98)00100-0, 1998. 10280
- 5 Hu, Z., Yu, G., Fu, Y., Sun, X., Li, Y., Shi, P., Wang, Y., and Zheng, Z.: Effects of vegetation control on ecosystem water use efficiency within and among four grassland ecosystems in China, *Glob. Change Biol.*, 14, 1609–1619, doi:10.1111/j.1365-2486.2008.01582.x, 2008. 10291
- Kanda, M., Inagaki, A., Letzel, M. O., Raasch, S., and Watanabe, T.: LES Study of the Energy Imbalance Problem with Eddy Covariance Fluxes, *Bound.-Lay. Meteorol.*, 110, 381–404, doi:10.1023/B:BOUN.0000007225.45548.7a, 2004. 10286
- 10 Kracher, D., Mengelkamp, H., and Foken, T.: The residual of the energy balance closure and its influence on the results of three SVAT models, *Meteorolog. Z.*, 18, 647–661, doi:10.1127/0941-2948/2009/0412, 2009. 10285
- 15 Kuwagata, T., Kondo, J., and Sumioka, M.: Thermal effect of the sea breeze on the structure of the boundary layer and the heat budget over land, *Bound.-Lay. Meteorol.*, 67, 119–144, doi:10.1007/BF00705510, 1994. 10286
- Li, M., Ma, Y., Hu, Z., Ishikawa, H., and Oku, Y.: Snow distribution over the Namco lake area of the Tibetan Plateau, *Hydrol. Earth Syst. Sci.*, 13, 2023–2030, doi:10.5194/hess-13-2023-2009, 2009. 10278
- 20 Lohou, F. and Patton, E. G.: Land-surface response to shallow cumulus, EGU General Assembly, Vienna, Austria, 3–8 April 2011, EGU2011-10280-1, 2011. 10276
- Ma, Y., Wang, Y., Wu, R., Hu, Z., Yang, K., Li, M., Ma, W., Zhong, L., Sun, F., Chen, X., Zhu, Z., Wang, S., and Ishikawa, H.: Recent advances on the study of atmosphere-land interaction observations on the Tibetan Plateau, *Hydrol. Earth Syst. Sci.*, 13, 1103–1111, doi:10.5194/hess-13-1103-2009, 2009. 10277
- 25 Mauder, M. and Foken, T.: Documentation and instruction manual of the eddy covariance software package TK3, *Arbeitsergebnisse* 46, University of Bayreuth, ISSN 1614-8916, Bayreuth, 2011. 10286
- 30 Mauder, M., Oncley, S. P., Vogt, R., Weidinger, T., Ribeiro, L., Bernhofer, C., Foken, T., Kohsiek, W., Bruin, H. A. R., and Liu, H.: The energy balance experiment EBEX-2000. Part II: Intercomparison of eddy-covariance sensors and post-field data processing methods, *Bound.-Lay. Meteorol.*, 123, 29–54, doi:10.1007/s10546-006-9139-4, 2006. 10290

A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



- Mauder, M., Desjardins, R. L., and MacPherson, I.: Scale analysis of airborne flux measurements over heterogeneous terrain in a boreal ecosystem, *J. Geophys. Res.*, 112, D13112, doi:10.1029/2006JD008133, 2007. 10290
- Mauder, M., Foken, T., Clement, R., Elbers, J. A., Eugster, W., Grnwald, T., Heusinkveld, B., and Kolle, O.: Quality control of CarboEurope flux data - Part 2: Inter-comparison of eddy-covariance software, *Biogeosciences*, 5, 451–462, doi:10.5194/bg-5-451-2008, 2008. 10286
- Mengelkamp, H., Warrach, K., and Raschke, E.: SEWAB – a parameterization of the Surface Energy and Water Balance for atmospheric and hydrologic models, *Adv. Wat. Resour.*, 23, 165–175, doi:10.1016/S0309-1708(99)00020-2, 1999. 10285
- Metzger, S., Ma, Y., Markkanen, T., Göckede, M., Li, M., and Foken, T.: Quality assessment of Tibetan Plateau Eddy Covariance measurements utilizing Footprint modeling, *Adv. Earth Sci.*, 21, 1260–1267, X–XI, 2006. 10278
- Miehe, G., Miehe, S., Bach, K., Nölling, J., Hanspach, J., Reudenbach, C., Kaiser, K., Wesche, K., Mosbrugger, V., Yang, Y., and Ma, Y.: Plant communities of central Tibetan pastures in the Alpine Steppe/Kobresia pygmaea ecotone, *J. Arid Environ.*, 75, 711–723, doi:10.1016/j.jaridenv.2011.03.001, 2011. 10277
- Oberhuber, J. M., Herzog, M., Graf, H., and Schwanke, K.: Volcanic plume simulation on large scales, *J. Volcanol. Geoth. Res.*, 87, 29–53, doi:10.1016/S0377-0273(98)00099-7, 1998. 10280
- Panin, G. N., Tetzlaff, G., and Raabe, A.: Inhomogeneity of the Land Surface and Problems in the Parameterization of Surface Fluxes in Natural Conditions, *Theor. Appl. Climatol.*, 60, 163–178, doi:10.1007/s007040050041, 1998. 10286
- Panin, G. N., Nasonov, A. E., and Foken, T.: Evaporation and heat exchange of a body of water with the atmosphere in a shallow zone, *Izvestiya, Atmos. Ocean. Phys.*, 42, 337–352, doi:10.1134/S0001433806030078, 2006. 10288, 10307
- Ruppert, J., Thomas, C., and Foken, T.: Scalar Similarity for Relaxed Eddy Accumulation Methods, *Bound.-Lay. Meteorol.*, 120, 39–63, doi:10.1007/s10546-005-9043-3, 2006. 10290
- Sellers, P. J., Mintz, Y., Sud, Y. C., and Dalcher, A.: A Simple Biosphere Model (SIB) for Use within General Circulation Models, *J. Atmos. Sci.*, 43, 505–531, 1986. 10281
- Su, Z., Wen, J., Dente, L., van der Velde, R., Wang, L., Ma, Y., Yang, K., and Hu, Z.: The Tibetan Plateau observatory of plateau scale soil moisture and soil temperature (Tibet-Obs) for quantifying uncertainties in coarse resolution satellite and model products, *Hydrol. Earth Syst. Sci.*, 15, 2303–2316, doi:10.5194/hess-15-2303-2011, 2011. 10277

- Twine, T. E., Kustas, W. P., Norman, J. M., Cook, D. R., Houser, P. R., Meyers, T. P., Prueger, J. H., Starks, P. J., and Wesely, M. L.: Correcting eddy-covariance flux underestimates over a grassland, *Agric. For. Meteorol.*, 103, 279–300, doi:10.1016/S0168-1923(00)00123-4, 2000. 10286
- 5 van Heerwaarden, C. C., de Arellano, J. V., Moene, A. F., and Holtslag, A. A. M.: Interactions between dry-air entrainment, surface evaporation and convective boundary-layer development, *Q. J. Roy. Meteorol. Soc.*, 135, 1277–1291, doi:10.1002/qj.431, 2009. 10276
- Xue, Y., Zeng, F. J., and Schlosser, C. A.: SSiB and its sensitivity to soil properties – a case study using HAPEX-Mobilhy data, *Glob. Planet. Change*, 13, 183–194, doi:10.1016/0921-10281(95)00045-3, 1996. 10281
- 10 Yang, K., Koike, T., and Yang, D.: Surface Flux Parameterization in the Tibetan Plateau, *Bound.-Lay. Meteorol.*, 106, 245–262, doi:10.1023/A:1021152407334, 2003. 10277
- Yang, K., Chen, Y.-Y., and Qin, J.: Some practical notes on the land surface modeling in the Tibetan Plateau, *Hydrol. Earth Syst. Sci.*, 13, 687–701, doi:10.5194/hess-13-687-2009, 2009. 10277
- 15 Yee, S. Y. K.: The force-restore method revisited, *Bound.-Lay. Meteorol.*, 43, 85–90, doi:10.1007/BF00153970, 1988. 10282
- Zhou, D., Eigenmann, R., Babel, W., Foken, T., and Ma, Y.: The study of near-ground free convection conditions at Nam Co station on the Tibetan Plateau, *Theor. Appl. Climatol.*, 105, 217–228, doi:10.1007/s00704-010-0393-5, 2011. 10279, 10285
- 20

A simple two-layer soil model with extrapolated surface temperatureT. Gerken et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 1. Description of the two sites near Nam Co lake used in this work.

Site	UBT	ITP
Coordinates	30°46.50′ N 90°57.61′ E	30°46.44′ N 90°57.72′ E
Soil	sandy–loamy	sandy
Porosity	0.63	0.393
Field Capacity	0.184	0.05
Wilting Point	0.115	0.02
Heat capacity (c_p) [$\text{Jm}^{-3}\text{K}^{-1}$]	2.5×10^6	2.2×10^6
Thermal		
Conductivity [$\text{Wm}^{-1}\text{K}^{-1}$]	0.53	0.20
Surface Albedo (α)	0.2	0.2
Surface Emissivity (ϵ)	0.97	0.97
Vegetated Fraction	0.9	0.6
LAI [m^2m^{-2}]	0.9	0.6
Vegetation height [m]	0.07	0.15

A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.

Table 2. Initial soil temperatures used in study (\bar{T}_2 and \bar{T}_1 are estimated from the respective base and top temperatures of the layer according to a quadratic temperature profile), change of layer 1 mean temperature ($\Delta\bar{T}_1$) over the modified Hybrid run, soil moisture content of layer 1 at beginning of the modified model run ($\theta_{1,obs}$) and at the end of the simulation ($\theta_{1,end}$). The values in parenthesis are expressed as θ_1/FC [-].

Site	Date	\bar{T}_2 [°C]	$T_{1,base}$ [°C]	\bar{T}_1 [°C]	T_0 [°C]	$\Delta\bar{T}_1$ [°C]	$\theta_{1,obs}$ [%]	$\theta_{1,end}$ [%]
UBT	10 July	3.9	11.8	10.9	9.3	-1.6	26.9 (1.47)	41.1(2.24)
	27 July	4.5	13.4	12.5	10.6	-1.6	20.8 (1.14)	17.0 (0.92)
	05 August	4.8	14.4	13.4	11.2	-3.0	26.9 (1.47)	19.1 (1.04)
	06 August	4.75	14.3	12.8	9.8	-1.4	25.4 (1.39)	34.0 (1.85)
ITP	10 July	5.4	16.2	13.2	7.2	-1.2	6.0 (1.1)	25.1 (5.02)
	27 July	7.2	21.6	17.8	10.2	-1.7	3.0 (0.6)	1.6 (0.32)
	05 August	5.7	17.1	11.1	-0.8	0.2	11.0 (2.2)	4.3 (0.86)
	06 August	5.6	16.8	11.6	1.1	1.9	9.0 (1.8)	3.7 (0.73)

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 3. Root mean square deviation (RMSD) between the modelled quantities of the original and modified Hybrid and reference values. The reference quantities used are either measured by EC and corrected for energy balance closure (EC,EBC) or modelled with SEWAB for fluxes or taken from longwave outgoing radiation for T_0 . The values in parenthesis (N) correspond to the number of elements used for calculation of RMSD and R^2 ($l = 0$) in Fig. 7.

Site	Date	Run	RMSD				
			Q_E EC,EBC [Wm^{-2}]	Q_H [Wm^{-2}]	Q_E SEWAB [Wm^{-2}]	Q_H [Wm^{-2}]	T_0 [$^{\circ}C$]
UBT	10 July	orig	318	117 (8)	94	74 (94)	4.3 (139)
	27 July		97	58 (19)	60	59 (139)	4.5 (143)
	05 August		168	139 (11)	90	64 (110)	4.3 (143)
	06 August		159	84 (52)	87	71 (128)	3.7 (143)
ITP	10 July	orig	182	93 (25)	97	69 (143)	3.7 (143)
	27 July		43	64 (72)	58	75 (143)	3.8 (143)
	05 August		224	103 (64)	179	68 (143)	8.3 (143)
	06 August		118	80 (52)	130	119 (143)	5.1 (143)
UBT	10 July	mod	214	43 (8)	51	36 (94)	2.3 (139)
	27 July		79	44 (19)	32	28 (139)	2.9 (143)
	05 August		93	62 (11)	36	26 (110)	3.4 (143)
	06 August		78	57 (52)	39	32 (128)	3.2 (143)
ITP	10 July	mod	74	73 (25)	42	32 (143)	1.6 (143)
	27 July		42	58 (72)	55	36 (143)	2.6 (143)
	05 August		44	80 (64)	64	30 (143)	2.6 (143)
	06 August		68	82 (52)	113	77 (143)	3.5 (143)
UBT	all	orig	170	92 (90)	83	67 (471)	4.2 (568)
		mod	100	54 (90)	39	31 (471)	3.0 (568)
ITP	all	orig	152	84 (213)	125	86 (572)	5.6 (572)
		mod	54	73 (213)	74	48 (572)	2.7 (572)

A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.




Fig. 1. Forcing data measured at UBT used for model runs: **(a)** downward shortwave radiation (SW [Wm^{-2}]); **b)** downward longwave radiation (LW [Wm^{-2}]); **(c)** air temperature (T [$^{\circ}C$]); **(d)** water vapour mixing ratio(q [gkg^{-1}]); **(e)** wind speed (U [ms^{-1}]); **(f)** surface pressure (P [hPa]) and precipitation [$mm(0.5h)^{-1}$]) Height c–e is 3 m.

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶


◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.




Fig. 2. Same as Fig. 1, but with forcing data measured at ITP.

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 3. Conceptual drawing of the assumed quadratic subgrid soil temperature profile and the associated parameters. In order to derive \bar{T}_1 and \bar{T}_2 geometrically the areas A_1 and A_2 must be equal.

A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 4. Dependency of soil temperature parameters: **(a)** relationship between mean temperature of layer 2 (\bar{T}_2) and bottom temperature of layer 1 ($T_{base_1} - a_2$) is calculated according to Eq. (4); **(b)** surface temperature (T_0) contour plot as function of T_{base_1} and layer 1 mean temperature (\bar{T}_1) and **(c)** contours of (T_0) as function of \bar{T}_2 and \bar{T}_1 . The black rectangle at the intersection of the layer temperature ranges (yellow) indicates the theoretical parameter space given by the temperature values used in this study and the black crosses mark the actual configurations.

A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.




Fig. 5. Model results for the modified Hybrid at UBT for 10 July 2009 (a–b), 27 July 2009 (c–d), 5 August 2009 (e–f) and 6 August 2009 (g–h). Left column: latent heat flux (Q_E); right column: sensible heat flux (Q_H) and surface temperature T_s [$^{\circ}\text{C}$]. L and W refer to “land” and “water” as origin of the fluxes. *All* is the complete available time series. The subscript *Hybrid* refers to fluxes from Hybrid and *COARE* are fluxes from the lake derived by TOGA-COARE whereas *SEWAB* is a SVAT model and *HM* refers to a hydrodynamic multi-layer lake model after Foken (1984) and Panin et al. (2006). *EC* and *EC,EBC* refer to measurements by eddy covariance method where in the latter the energy balance has been closed by distributing the residual according to Bowen-ratio (this requires good data quality and fluxes and can only be done for fluxes that are attributed to land). The circles indicate poor data quality of the EC system according to Foken et al. (2004). Yellow shading indicates times where the flux footprint of UBT was over the lake.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.




Fig. 6. Same as Fig. 5, but for ITP. There are no contributions from the lake.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A simple two-layer soil model with extrapolated surface temperature

T. Gerken et al.




Fig. 7. Correlation of simulated fluxes against flux reference shifted by t_{lag} as multiples of 10 minutes (Cross Correlation – $R^2(t)$) for each of the four days simulated and for the original and modified Hybrid. The maximum number of elements used in the calculation of R^2 for each curve can be taken from Table 3.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)