

This discussion paper is/has been under review for the journal Hydrology and Earth System Sciences (HESS). Please refer to the corresponding final paper in HESS if available.

# Stream recession curves and storage variability in small watersheds

N. Y. Krakauer<sup>1</sup> and M. Temimi<sup>2</sup>

<sup>1</sup>Department of Civil Engineering, The City College of New York, USA

<sup>2</sup>NOAA-CREST, The City College of New York, USA

Received: 19 January 2011 – Accepted: 28 January 2011 – Published: 10 February 2011

Correspondence to: N. Y. Krakauer (nkrakauer@ccny.cuny.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

**HESSD**

8, 1827–1860, 2011

## Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

The pattern of streamflow recession after rain events offers clues about the relationship between watershed runoff (observable as river discharge) and water storage (not directly observable) and can help in water resource assessment and prediction. However, it has not been systematically analyzed across flow rates or related to independent assessments of terrestrial water storage. We characterized the streamflow recession pattern in 61 relatively undisturbed small watersheds (1–100 km<sup>2</sup>) across the coterminous United States with multiyear records of hourly streamflow from automated gauges. We used the North American Regional Reanalysis (NARR) to help identify periods where precipitation, snowmelt, and evaporation were small compared to streamflow. The order of magnitude of the recession timescale increases from 1 day at high flow rates (~1 mm/h) to 10 days at low flow rates (~0.01 mm/h), leveling off at low flow rates. There is significant variability in the recession timescale at a given flow rate between basins, correlated with climate and geomorphic variables such as the ratio of mean streamflow to precipitation and soil water infiltration capacity. Stepwise multiple regression was used to construct a six-variable predictive model that explained some 80% of the variance in recession timescale at high flow rates and 30–50% at low flow rates. Seasonal and interannual variability in storage shows similar time evolution to but is up to a factor of 10 smaller than regional-scale water storage variability estimated from GRACE satellite gravity data and from land surface modeling forced by observed meteorology. The discrepancy may point to a “disconnection” between the conceptual pool that supplies streamflow during dry periods and other dynamic pools such as soil moisture and deep groundwater.

## 1 Introduction

The observation that river flow gradually decreases after a rainstorm was modeled mathematically by the beginning of the 20th century with the concept of a “recession curve” that describes the characteristic decay of flow rate with time during rainless

# HESSD

8, 1827–1860, 2011

## Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



periods. Such recession curves have been used to forecast flows, estimate the probabilities of low flows, infer groundwater storage or aquifer characteristics, and detect change in watershed characteristics over time. Analytical expressions for the form of the recession curve could be derived for idealized basin shapes and subsurface flow properties, or curves could be fit empirically from streamflow measurements. Early applications were held back in part by the lack of a systematic procedure for determining an appropriate functional form and parameters for the recession curve shape from river discharge measurements (for reviews, see Hall, 1968; Tallaksen, 1995).

Brutsaert and Nieber (1977) developed a procedure for visualizing the recession curve for a given river that has been widely used and adapted. A family of functions describing river recessions is given by the power law

$$\dot{Q} = -aQ^b \quad (1)$$

where  $Q$  is river discharge,  $\dot{Q}$  is its rate of change, and  $a$  and  $b$  are parameters.  $b = 1$  corresponds to an exponential-decay recession curve. For a river recession that follows this pattern, a scatter log plot of  $\dot{Q}$  vs.  $Q$  should approximate a straight line, since from Eq. (1),

$$\log(-\dot{Q}) = \log(a) + b\log(Q), \quad (2)$$

when  $Q > 0$ . This is convenient because values for  $a$  and  $b$  can be fitted to recession data by linear regression. Further, the appropriateness of the assumed functional form Eq. (1) can be assessed by how well the points in the scatter plot are described by a straight line. In practice, Brutsaert and Nieber (1977) took time series of daily streamflow for several streams in New York state and plotted  $\dot{Q}$  as the difference in stream flow between consecutive days (that were at least 5 days after the most recent rain) against the corresponding  $Q$  (estimated as the average of the two days).  $a$  and  $b$  were then estimated by drawing lines to match the lower envelope of the cloud of  $(\log(Q), \log(-\dot{Q}))$  points. The slope of this lower envelope, corresponding to  $b$ , tended to be higher than 1 – around 1.5 at low flows, possibly increasing to about 3 at high

flows. Wittenberg (1999) fit a power law to data from German watersheds, also finding that  $b$  was around 1.5.

A number of studies have determined or assumed that, at least well after rain events,  $b = 1$  to acceptable accuracy, so that only one parameter,  $a$ , must be estimated from measured streamflow (e.g., Vogel and Kroll, 1992; Brandes et al., 2005; Eng and Milly, 2007; van Dijk, 2010). In this case the reciprocal of  $a$  is the recession timescale  $\tau$ , corresponding to the ratio  $-Q/\dot{Q}$ . In general, for other shapes of the recession curve (functional relationships between  $Q$  and  $\dot{Q}$ ), this recession timescale  $\tau$  would vary as a function of the flow rate  $Q$ . Brutsaert (2008) argued that the recession timescale is fairly constant not only for a given stream but also across streams, at least during summer low flows and for large basins, at  $45 \pm 15$  days.

Kirchner (2009) further refined the estimation of recession curves based on measured  $Q$  and  $\dot{Q}$ . In the approach of Kirchner (2009), the time series of hourly streamflow  $Q$  is sorted into bins and the average  $\dot{Q}$  is determined for each bin, using only time periods when both measured precipitation and estimated potential evaporation are small compared to streamflow, and parameters describing the relationship between  $Q$  and  $\dot{Q}$  are fit for the binned data. Excluding time periods when evaporation might be a substantial part of the water budget avoids possible bias in the recession curve due to evaporation, which would be expected to increase the streamflow recession rate  $-\dot{Q}$  at a given  $Q$  (Weisman, 1977; Wittenberg, 2003). It adds the complication of requiring hourly, rather than daily, streamflow data, since in most tropical and temperate watersheds, potential evaporation under rainless conditions is only likely to be much lower than streamflow at night; until recently, only daily streamflow data have generally been made available to the hydrological community. Kirchner (2009) showed that for two small watersheds in Wales, an empirically chosen quadratic functional form

$$\log(-\dot{Q}) = \log(a) + b\log(Q) + c\log^2(Q) \quad (3)$$

fits the binned hourly data well.  $b$  was found to be close to 2, and the departure from a log-linear power-law relationship (expressed by the quadratic term) was found to be small but significantly different from zero.

## Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A large number of studies have examined the variability in recession timescale across streams, most often in small regions (e.g., Bingham, 1986; Vogel and Kroll, 1992; Brutsaert and Lopez, 1998; Wittenberg, 1999; Zhang et al., 2009; Zhu et al., 2010), but also on continental and larger scales (van Dijk, 2010; Peña-Arancibia et al., 2010), and related inter-stream variability in the recession timescale to climate, topographic, or geologic factors. However, these studies have tended to concentrate on low-flow periods and fit a simple functional form of the recession curve, generally the power law (Eq. 1) (often with  $b = 1$  so that the recession timescale is taken to be constant), to derive the recession timescale, so it is not clear how the recession timescale and its spatial distribution varies across flow rates. Thus, one purpose of the current study is to examine the characteristics of recession curves derived with uniform procedures from hourly streamflow data across a range of climate and terrain.

We adopt the approach of Kirchner (2009) as a starting point because it has the advantage of making use of all hours for which streamflow data is available, excluding only those where other water fluxes such as precipitation and evaporation are likely to be significant. Other common selection criteria, such as fitting the lower envelope of  $\log(-\dot{Q})$  or excluding streamflow records from a certain number of days after rain events, involve arbitrary thresholds and make it difficult to estimate the error of the fitted recession timescale. Using hourly, as compared to daily, streamflow data enables the selection of low-evaporation periods and avoids bias in recession time estimates at higher flow rates when the recession timescale  $\tau = -Q/\dot{Q}$  is of order 1 day (Rupp and Selker, 2006). In this study, we consider only small watersheds ( $<100 \text{ km}^2$ ), so that the timescale for streamflow generated within the watershed to reach the gauge is not much more than an hour and the measured discharge gives a reasonable estimate of hourly runoff.

By making some assumptions, the recession curve, expressed as the function  $\tau(Q)$ , can relate the streamflow  $Q$  to the basin water store  $S$ . The rate of change of storage,  $\dot{S}$ , is the sum of the water fluxes in and out of the basin, namely streamflow,

## Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



precipitation  $P$ , and evaporation  $E$ :

$$\dot{S} = P - E - Q. \quad (4)$$

When the streamflow  $Q$  is the major flux of water in or out of the basin (precipitation and evaporation are slight), we can make the approximation

$$\dot{S} = -Q \quad (5)$$

and therefore

$$\frac{dQ}{dS} = \frac{\dot{Q}}{\dot{S}} = \frac{\dot{Q}}{-Q} = \tau^{-1}. \quad (6)$$

Having estimated  $\tau$  as a function of  $Q$  for periods with low  $P$  and  $E$  from recession curve analysis, one can therefore estimate the change in storage  $S - S_0$  corresponding to observed streamflows  $Q$ :

$$S(Q) - S_0 = \int_{Q_0}^Q dS = \int_{Q_0}^Q \tau(Q) dQ, \quad (7)$$

where the reference streamflow  $Q_0$  and reference storage level  $S_0 = S(Q_0)$  are arbitrary. Kirchner (2009) further observed that by invoking water balance, one can estimate not only the rate of change in storage but also the net flow into the watershed  $F = P - E$ :

$$F = Q + \dot{S} = Q + \tau \dot{Q}. \quad (8)$$

The fitted function  $\tau(Q)$  is derived from the recession curve over hours with negligible precipitation and evaporation; estimating  $S$  and  $F$  for other conditions from the above equations requires assuming that the deduced relationship between  $Q$  and  $S$  continues to hold. Kirchner (2009) argued that this is indeed the case for his two study watersheds, as evidenced by the ability of the basin storage-discharge relationship estimated using Eq. (6) to successfully infer rainfall using only streamflow observations, and to

## HESSD

8, 1827–1860, 2011

### Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



predict the evolution of streamflow using measured precipitation and estimated evaporation. If so, streamflow time series could be used to infer other water flows (precipitation, evaporation) and stores at the catchment scale, which Kirchner (2009) termed “doing hydrology backward” compared with the conventional hydrology approach of deriving streamflow from meteorological forcing and basin characteristics, which are often quite uncertain.

One recent application of “doing hydrology backward” is by Palmroth et al. (2010), who apply recession curve analysis to construct storage-discharge relationships that they used to estimate evapotranspiration over parts of North Carolina state, although correlation with independent (eddy covariance) measurements of evapotranspiration was found to be poor. Brutsaert (2010) quantified changes in summer terrestrial water storage across the central United States in recent decades based on changes in summer streamflows along with an assumed recession timescale  $\tau$  based on previous studies. The inferred changes in water storage were found to be consistent with groundwater observations in Illinois state (Brutsaert, 2008). To utilize this approach more widely, however, the validity of water storage changes inferred from recession-curve analysis warrants further testing and comparison with available watershed-scale hydrometeorological data.

Multi-year time series of streamflow measurements at high (sub-hourly) temporal resolution are now freely available for many streams. Here, we employ these data to construct recession curves across a range of topography, geology, and climate in order to answer the following questions:

1. What is the variability across streams of the recession timescale at different flow rates? How much of this variability is correlated with factors such as climate regime and topography?
2. How does the variability in basin water storage inferred from streamflow recession curve analysis compare to basin water storage variability inferred from other, independent methods?

## Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2 Methods

2.1 Streamflow data

The Hydroclimatic Data Network (HCDN) includes about 1500 stream gauge records from the United States Geological Survey (USGS) stream gauge network chosen to represent streams with long monitoring histories and whose flow has experienced minimal human disturbance (Slack and Landwehr, 1992). For this study, we chose HCDN gauges draining small watersheds (area under 100 km<sup>2</sup>) in the coterminous United States (USA) which had daily records over at least 2/3 of the period 1979–2008. High-resolution streamflow measurements for these streams were obtained from the USGS Instantaneous Data Archive (IDA, <http://ida.water.usgs.gov/ida/>; Showstack, 2007). The selection criteria yielded 75 streams, of which 61 had flow records at hourly or better resolution available through IDA. The median basin area for these 61 streams was 59 km<sup>2</sup> (range: 6.1–98 km<sup>2</sup>), and basin locations covered a wide range of climate as well as topography, although most were in the moister regions of the country, near the Atlantic and Pacific coasts (Fig. 1). Median streamflow per unit basin area was 428 mm/year (range: 51–1499 mm/year), compared to an average of about 160 mm/year for the coterminous USA over the same period (Krakauer and Fung, 2008).

The IDA streamflow records, typically at 15-min resolution, were averaged to generate hourly streamflow series. Streamflow values within one minute of the turn of the hour were assigned half-weight for estimating both hours' streamflow. Only hours with at least one streamflow measurement (or two measurements at their borders) were used in the analysis. The 61 streams in this study had an average of 148 thousand usable hours (equivalent to 17 years; range: 87–232 thousand). While these hours were not always consecutive (records frequently had gaps), this is not a problem for our analysis procedures.

Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## 2.2 Meteorology

Precipitation and evaporation records for the study watersheds were used to isolate hours with low/no precipitation or evaporation to use for estimating recession curves. Since sub-daily field measurements of precipitation and evaporation for each of the watersheds were not in general available, precipitation and evaporation were obtained from the National Centers for Environmental Prediction (NCEP) North American Regional Reanalysis (NARR) (Mesinger et al., 2006) for 1979–2008, which uses the Eta Model to simulate regional atmospheric circulation at relatively high horizontal resolution (32 km). For each watershed, meteorological fields were taken from the closest grid cell to the stream gauge – because the watersheds are all  $<100 \text{ km}^2$  while a NARR grid cell is  $\sim 10^3 \text{ km}^2$ , the entire watershed is likely to lie within a single grid cell.

Precipitation in NARR assimilates rain gauge and satellite observations and is therefore much more accurate than previous reanalyses. Evaporation in NARR is simulated by the Noah land surface model and is only indirectly tied to observations (Mesinger et al., 2006).

## 2.3 Binning streamflow recession data

Following Brutsaert and Nieber (1977), hourly  $-\dot{Q}$  was estimated as  $\frac{1}{\Delta t}(Q_h - Q_{h+1})$ , the difference between streamflow in adjacent hours, while the corresponding hourly  $Q$  was the average for those hours,  $Q = \frac{1}{2}(Q_h + Q_{h+1})$ . Pairs of hours were selected for which both precipitation plus snowmelt and evaporation were less than 10% of average streamflow (Fig. 2).

The NARR output fields include snowmelt, but the precipitation field is not divided into rain vs. snow. For determining rain-free hours, we required that the precipitation plus snowmelt be less than 0.1 of streamflow. For determining hours with low evaporation, we required that evaporation be less than 0.1 of streamflow.

NARR output fields are available at 3 h time resolution and were matched to the corresponding hourly streamflow data. To account for any delay between runoff generation and streamflow at the gauge location, we followed Kirchner (2009) in estimating this

## Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



lag for each basin from the position of the maximum lagged cross-correlation between precipitation and  $\dot{Q}$ . The streamflow time series were then shifted relative to the precipitation and evaporation time series by that amount before deciding what hours to exclude from the recession curve analysis as affected by precipitation or evaporation.

The small size of the basins we analyzed kept this lag small (0–2 h for 50/61 basins and 0–5 h for 59/61, ranging up to 11 h).

Because the NARR grid size is bigger than the areas of our watersheds, there remains the concern that localized bursts of precipitation are not reflected in the NARR precipitation record. To reduce the effect that such unrecorded precipitation might have on the estimated recession curve, we further excluded periods of two or more consecutive positive hourly  $\dot{Q}$  (rising streamflow), on the assumption that these correspond to precipitation or snowmelt events not necessarily captured in NARR. Altogether, exclusion based on NARR precipitation, snowmelt, and evaporation and on observed rising flow left 0.6–27% (median 7.3%, or 10 thousand hours) of the original number of hours for constructing the streamflow recession curve, with the lower percentages found in arid basins where streamflow was often very low and the higher percentages found in more humid basins.

The selected values of  $\dot{Q}$  (i.e. those when the lagged precipitation and snowmelt were small, and excluding periods of rising streamflow) were then averaged over ranges of  $Q$ . These ranges were selected as follows (cf. Kirchner, 2009): (1) begin with the top 1% of the logarithmic range in  $Q$ ; (2) compute the mean and standard error of  $\dot{Q}$  for all  $Q$  in that range; (3) if the number of values in the range is less than 9 or the mean  $\dot{Q}$  is nonnegative or the standard error in  $\dot{Q}$  is more than half its absolute mean value, expand the bin by another 1% of the logarithmic range; (4) otherwise, keep the mean  $Q$  and  $\dot{Q}$  of the bin and continue with the next 1% of the logarithmic range. This resulted in typically 20–70 bins (median 58 bins), each with a mean  $Q$ , mean  $\dot{Q}$ , and standard error of  $\dot{Q}$ . Regression on the binned values is basically equivalent to weighted regression on the original hourly data, where weighting is by the inverse variance of  $\dot{Q}$  over a small range in  $Q$ .

## Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Occasionally (1 of our 61 sites) this procedure did not converge (i.e. the scatter in  $\dot{Q}$  was large enough that the standard error did not drop sufficiently even when the bins were expanded). In such cases, the range in  $Q$  was divided into a number of bins with roughly equal numbers of elements (approximately the square root of the number of usable hours), and the mean and the standard error of  $\dot{Q}$  in each bin was calculated; bins with nonnegative  $\dot{Q}$  were simply discarded.

## 2.4 Fitting recession curves

We experimented with different functional forms for the recession curve, as fit to the binned  $\log(Q)$  and  $\log(-\dot{Q})$ , including the linear-in-logs (power law) relationship (Eq. 2) and the quadratic-in-logs relationship (Eq. 3). The goodness of fit of different functional forms was assessed by fitting the functional form to one half of each streamflow record and calculating the misfit between the fitted values and the binned values found for the other half of the same streamflow record. We found that for our sample of watersheds, a nonparametric functional form corresponding to locally-weighted least squares linear regression (similar to LOWESS, Cleveland, 1979) gave the best fit (Fig. 3a). Given that the slope of the relationship between  $\log(-\dot{Q})$  and  $\log(Q)$  reflects complex water flows in different parts of the watershed (Harman et al., 2009) and might be expected to vary, for example, in the transition from surface to subsurface runoff (Mizumura, 2005), it makes sense that a particular shape with few adjustable parameters would be less suitable for modeling the streamflow recession curve as compared to a flexible locally smooth function. At each bin's value of  $\log(Q)$  ( $\log(Q_{\text{bin}})$ ), a smoothed value of  $\log(-\dot{Q})$  is obtained by weighted linear regression with weights that favor adjacent bins, namely

$$w_{\text{bin},i} = \bar{w}_i \times \exp(-|\log(Q_i) - \log(Q_{\text{bin}})|/\alpha) \quad (9)$$

where  $\bar{w}_i$  is weighting based on the standard error of each bin (equal to the reciprocal of the square of  $E_i/\log(-\dot{Q}_i)$ ), where  $E_i$  is the standard error of the binned  $\dot{Q}_i$ ) and  $\alpha$  is a parameter that sets the size of the neighborhood that is considered in the locally

## Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



weighted linear regression. Between the points  $\log(Q_i)$ , the function was taken to be piecewise linear in  $\log(Q)$ .

We used generalized cross validation (Craven and Wahba, 1979; Krakauer et al., 2004) to estimate a suitable value for  $\alpha$  for each stream, which typically was around 0.3 log units. At the limit  $\alpha \rightarrow \infty$ , the result is the same as the linear relationship (Eq. 2), while as  $\alpha \rightarrow 0$  the fitted function becomes less smooth and approaches linear interpolation.

Given the above weights, uncertainties for the fitted function value were then calculated using standard linear regression methodology. If  $E_i/\log(-\dot{Q}_i)$  in fact reflect the error standard deviation of the binned values and if this error is normally distributed, then  $r$ , the weighted sum of squares of the residuals from the fitted function,

$$r = \sum_i \left( \frac{\log(-\dot{Q}_i) - \log(-\hat{\dot{Q}}_i)}{E_i/\log(-\dot{Q}_i)} \right)^2 \quad (10)$$

(where  $\log(-\hat{\dot{Q}}_i)$  is the value of the fitted function at  $\log(Q_i)$ ), should be close to the effective number of degrees of freedom  $n - m$ , where  $n$  is the number of bins and  $m$  the effective number of fitted parameters (which will increase as  $\alpha$  decreases). In practice, we found that the  $r$  was often somewhat larger than this, presumably reflecting non-lognormal errors in streamflow measurements that propagate to the calculated  $-\dot{Q}$ ; the median ratio  $r/(n - m)$  across our sample was 1.5. Therefore, calculated uncertainties were multiplied by  $\sqrt{\frac{r}{n - m}}$ . This adjusted uncertainty for the fitted recession curve was

found to be realistic: the difference between the binned  $\log(-\dot{Q})$  and  $\log(Q)$  derived from one half of a streamflow record and the fitted recession curve function derived from the other half was consistent with the adjusted uncertainty in the fit.

Given the fitted recession curve, which gives  $\log(-\dot{Q})$  as a piecewise linear function of  $\log(Q)$  with associated uncertainty, the recession time  $\tau(Q)$  was found as  $-Q/\dot{Q}$ , with fractional uncertainty equal to that in  $\dot{Q}$  (Fig. 3b). A lookup table was generated for the estimated storage  $S$  corresponding to various streamflow levels in the observed range

## Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



by numerically integrating  $\tau(Q)dQ$  (Eq. 7). Monthly mean storage was then computed as the mean of hourly  $S$  calculated from the observed  $Q$ , and the variability of this monthly watershed storage was compared with satellite and model estimates of water storage variability.

## 2.5 Explaining inter-stream variability in recession curves

For understanding the relationship between the recession curve and watershed hydrological processes, analysis of factors correlated with variability in the recession curve, as expressed by the function  $\tau(Q)$ , is helpful. For each stream, the function  $\tau(Q)$  was expressed as 31 values for  $\tau$  logarithmically spaced in  $Q$  for  $Q$  ranging between 0.0048 and 1.3 mm/h, corresponding to the median range of the streamflow bins used to fit the recession curves. Potential predictor variables used were mean streamflow (calculated from the streamflow time series); gauge longitude, latitude, and elevation, basin area, precipitation, climatological January minimum temperature, mean elevation, percent forest cover, percent lake cover, and soil water infiltration capacity, and stream length and slope (taken from the HCDN data files); and precipitation, snowfall, and evaporation (taken from NARR). The nonparametric (Spearman) correlation coefficient of each variable against  $\log(\tau)$  at each of the 31 flow rates was calculated, and the mean square correlation coefficient across the 31 flow rates was compared to that obtained from regressions with 1000 random permutations of the predictor values to assess whether this variable is a significantly correlated to  $\tau(Q)$  at the 0.05 level. Stepwise multiple linear regression against  $\log(\tau)$  (weighted by its site-specific estimated uncertainty) was also performed, where the predictor variable was added whose inclusion most increases the weighted mean  $R^2$  of the regression model. The procedure was terminated when the increase in  $R^2$  from one more variable being added to the regression model was not significant at the 0.05 level, as quantified by comparing to the increase in  $R^2$  when values for that variable were randomly permuted before being added to the regression. Any missing values for predictor variables in the HCDN

## Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



data file were filled in with the average value for that variable, to minimize bias in the estimated regression coefficients.

## 2.6 Water storage data

Monthly terrestrial water storage anomalies on a  $1^\circ \times 1^\circ$  grid for 2002–2008, estimated from gravitational anomalies reflected in the GRACE satellite positions, were obtained from the Jet Propulsion Laboratory (JPL). The water storage anomalies were derived from gravitational fields estimated from the satellite orbits by CSR (U. Texas/Center for Space Research), destriped to minimize processing artifacts, and rescaled based on a land surface model to minimize error in the storage anomaly magnitudes due to contributions from adjacent areas to the measured gravity anomaly (Swenson and Wahr, 2006; Swenson, 2011). Another set of estimates of monthly terrestrial water storage anomalies on a  $1^\circ \times 1^\circ$  grid for 2001–2008 from the Noah land surface model run using observed meteorological forcing as part of the Global Land Data Assimilation System (Ek et al., 2003; Rodell et al., 2004) was also obtained from JPL.

Measures of water storage variability calculated from the streamflow recession curves and for the corresponding grid cells in the GRACE and Noah datasets included the seasonal cycle amplitude (the standard deviation of the mean seasonal cycle) and interannual variability (the standard deviation of monthly storage anomalies once the mean seasonal cycle has been removed). Additionally, we calculated correlation coefficients to assess to what extent the spatial pattern of seasonal to interannual variability magnitudes was consistent between the streamflow, GRACE, and Noah storages, and to compare the temporal variation in water storage during the period of overlap of the different estimates.

### Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 3 Results

### 3.1 Recession curves

Recession time as a function of streamflow  $Q$  showed broadly similar patterns across the sample of watersheds, characteristically decreasing from  $\sim 10$  days at the lowest streamflow rates resolvable with our binning method ( $\sim 0.005$  mm/h) to  $\sim 1$  day at high streamflow rates seen soon after rain or snowmelt ( $\sim 1$  mm/h) (Fig. 4). These timescales are similar to the pattern seen in stream recession curves constructed in previous studies (Brutsaert and Nieber, 1977; Vogel and Kroll, 1992; Brandes et al., 2005; Kirchner, 2009; van Dijk, 2010; Peña-Arancibia et al., 2010), although notably smaller than the  $45 \pm 15$  day timescale for low-flow conditions seen in the studies cited by Brutsaert (2008), perhaps because of differences in the estimation procedure. Note that the function  $\tau(Q)$  is on average almost flat at low streamflows (corresponding to  $b \approx 1$  in the power-law relationship (Eq. 1)), while it decreases with increasing  $Q$  at higher streamflows (corresponding to  $b \approx 1.6$ ); thus, no single power-law relationship can represent accurately the average recession curve.

We see from Fig. 4a that there is across-stream variability of an order of magnitude in the recession time  $\tau$  for any given flow rate  $Q$ . Comparing with Fig. 3b suggests that this variability is larger than the uncertainty in the fitted recession time curve for any one stream. This is confirmed by comparing the weighted mean fit uncertainty (spread of lines in Fig. 4b) with the total standard deviation across streams (error bars in Fig. 4b), which is much greater across flow rates.

We performed two sensitivity analyses on the impact of the criteria for choosing suitable hours on the recession curve. In one analysis, we did not exclude hours with high evaporation (as most previous analyses did not). This typically resulted in little change in the recession curve at high flows (when evaporation was likely small compared to the flow), but lowered the recession timescale by up to 40% during low flow, qualitatively similar to the finding of Weisman (1977) that flow diminished faster during periods of high evaporation than during periods of low evaporation (Fig. 5). In a second

HESSD

8, 1827–1860, 2011

## Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





analysis, we did not exclude hours with rising streamflow (but little precipitation according to NARR). This resulted in a longer recession timescale at all flow rates, with the recession timescale almost doubling at low flow rates (Fig. 5).

Given the variability across streams in recession timescales, it is of interest to determine what basin-specific factors could influence the recession timescale. Longer recession timescales were significantly correlated (in decreasing order of significance) with higher ratio of streamflow to precipitation, higher channel slope, higher elevation, more forest cover, higher basin soil infiltration capacity, lower longitude (i.e. western as compared to eastern USA), lower temperature, higher latitude (i.e. northern as compared to southern USA), and higher cold-season precipitation fraction. Recession timescales were not significantly correlated with lake cover, channel length, watershed area, channel length, or mean precipitation, streamflow, or evaporation. Many of these predictor variables were highly correlated with each other (for example, slope and elevation, or latitude and temperature). Stepwise multiple regression analysis yielded a model with six predictor variables, in the order they were added the model: (1) longitude, (2) soil infiltration capacity, (3) latitude, (4) channel length, (5) forest cover, (6) HCDN precipitation. (Of these, (4) and (6) were not found to be significant predictors in the univariate analysis, cf. Fig. 6a.) The relationship of the predictor variables to the recession timescale varied across flow rates: for example, the soil infiltration capacity showed a significant positive association with recession timescale only at low and moderate flow rates, while the correlation of recession timescale with latitude is positive only at high flow rates (Fig. 6a). The multivariate model best predicted recession timescales at high flow rates, where  $R^2$  was around 0.8, while at low flow rates  $R^2$  was 0.3–0.5; the average  $R^2$  across the range of flow rates was 0.57 (Fig. 6b). Spatial semi-variograms of the recession timescale at particular flow rates showed no evidence of spatial clustering (not shown), beyond the continental-scale east–west and north–south trends captured by the linear relationship with longitude and latitude.

## Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





### 3.2 Basin storage

Seasonal variability in storage inferred from streamflow and the recession curve showed good coherence with the variability in terrestrial water storage inferred from GRACE, with a median (across sites) coefficient of determination  $R^2$  between the two of 0.69, compared with a median  $R^2$  of 0.41 between the modeled (Noah) seasonal cycle and GRACE over the same grid cells. This good performance of the recession curve technique for measuring storage is particularly impressive because of the scale mismatch between the watershed size and the GRACE data (tens of  $\text{km}^2$  versus  $\sim 10^4 \text{ km}^2$ ), as compared to the similarity in scale between GRACE and the Noah simulations. Interannual variability in storage (computed for each site over months when the two data sets overlapped) was less coherent between the streamflow and GRACE approaches, with a median  $R^2$  of 0.22, but this was still better than the performance of Noah compared to GRACE, where the median  $R^2$  for interannual variability was only 0.06.

Both the seasonal and interannual variability in storage as inferred from the recession curves were generally lower by around a factor of 10 than those derived from GRACE (Fig. 7). The median ratio between streamflow-inferred and GRACE standard deviation in storage was 0.081 for the annual cycle and 0.106 for interannual variability. Storage variability in the Noah model was lower than the results from GRACE, because Noah does not represent variability in groundwater and surface water (Syed et al., 2008), but still generally higher than the variability inferred from the recession curves. The median ratio between streamflow-inferred and Noah standard deviation in storage was 0.389 for the annual cycle and 0.227 for interannual variability. Also, there was little correlation across watersheds between storage amplitude inferred from streamflow and that inferred from GRACE (Fig. 7).

HESSD

8, 1827–1860, 2011

#### Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4 Discussion

4.1 What accounts for variability in recession timescales across streams?

In our sample, inter-stream variability in the recession curve was correlated with mea-  
sures of climate (ratio of streamflow to precipitation, forest cover, temperature) and  
geomorphology (elevation, channel slope, soil infiltration capacity). The geographic  
patterns observed (with higher recession timescales in the west and north) are proba-  
bly related to the continental gradient in climate and geomorphology. The influencing  
variables and the direction of correlations between them and streamflow are largely  
consistent with those found in previous studies.

In studies by van Dijk (2010) in Australia and Peña-Arancibia et al. (2010) across the  
tropics and subtropics, basin aridity was found to be a dominant control on recession  
time, with more arid areas having shorter recession timescales. In the current sample  
of temperate-zone watersheds, we found a similar pattern: a low ratio of streamflow to  
precipitation correlated with short recession timescale.

High forest cover, typically associated with moist conditions, was associated with  
longer recession time, as also found by Peña-Arancibia et al. (2010). Vegetation cover  
and density has a major impact on the spatial organization of soil moisture (Mohanty  
et al., 2000; Gómez-Plaza et al., 2000, 2001; Qiu et al., 2001; Cantón et al., 2004;  
Temimi et al., 2010). The presence of vegetation fosters the retention of water in the  
canopy, litter layer, and root zone, which leads to slower drainage and therefore longer  
recession timescale. Roering et al. (2010) found that trees modify the topography  
around them by promoting soil formation and porosity and reducing erosion, which  
would tend to enhance the slow percolation of precipitation. The reduction of surface  
evaporation by vegetation shading (Hébrard et al., 2006) would also tend to increase  
recession timescale, although this impact should be less pronounced in our analysis  
because we excluded periods with high evaporation when computing recession curves.

Stream recession  
and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



We found that channel slope and basin elevation were positively correlated with the recession timescale. Peña-Arancibia et al. (2010) also found a positive correlation between basin slope and recession timescale. This contradicts the theoretical expectation of a negative correlation: if the contributing aquifer has a larger slope, it would be expected to drain faster, all things being equal (Brutsaert and Nieber, 1977; Vogel and Kroll, 1992). A negative correlation is also seen in some observational studies covering smaller spatial scales (Zecharias and Brutsaert, 1988; Brandes et al., 2005). One possible explanation for this discrepancy is that local or watershed-level attributes like channel slope and elevation do not necessarily correspond to aquifer properties, which would depend more on regional topography and geology (Temimi et al., 2010). The positive correlation between soil infiltration capacity and recession timescale is more intuitive, although Peña-Arancibia et al. (2010) found no correlation between the recession timescale and mapped soil infiltrability and drainage indices.

An original contribution of this study is the attempt to quantify the factors controlling the recession timescale at different flow rates, rather than estimating a single recession timescale for each watershed. We found that some variables were significant predictors of the recession timescale only at high or low flow rates, showing the value of explicitly including flow rate in this sort of regression analysis. Particularly at low flow rates, there also appeared to be substantial inter-stream variability in the recession timescale not captured by the set of predictor variables we used. Studies of variability in recession timescales across smaller spatial scales of tens to hundreds of km point to important geological controls associated with indicators such as bedrock porosity, drainage density, and soil group (e.g., Bingham, 1986; Brandes et al., 2005) that were not available for our sample of watersheds. van Dijk (2010) found that, after controlling for aridity, variability in recession timescales in Australia was spatially correlated over distances of 100–150 km, presumably reflecting geologic or topographic controls on soil and bedrock properties that was not reflected in the set of predictor variables used. Intensive studies of flow pathways in research watersheds as well as studies of large samples of small gauged watersheds with watershed properties estimated from remote

## Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



sensing and other distributed data sets can help characterize the link between watershed geology and morphology, on the one hand, and stream hydrology as reflected in the recession timescale, on the other, on a regional to global scale.

4.2 Why are storage amplitudes inferred from the recession curve so small?

We found that recession timescales derived from the recession curve constructed for periods of low precipitation, evaporation, and snowmelt to estimate a watershed storage-discharge relationship can be used to estimate monthly storage fluctuations that are coherent with those inferred from GRACE – more so, in fact, than those estimated by a state-of-the-art land surface model forced by observed meteorology (Noah/GLDAS). Thus, recession curve analysis is promising for extending the GRACE record of terrestrial water storage variability for higher spatial and temporal resolution or over longer time periods. Additional study is required to extend discharge-based storage estimates to larger basins with concentration times of days or weeks, where the method used here to construct recession curves probably would not work, for direct comparisons with GRACE and with models over a regional spatial scale.

The annual cycle and the interannual variability estimated from streamflow recession curve analysis prove to be similar in time evolution to that measured by GRACE, but are typically smaller by a factor of 10. This suggests that quantitative estimates of basin storage based on streamflow fluctuations and recession analysis should be treated with caution. We see several possible reasons for the small dynamic storage found with the recession curve approach. No single reason appears sufficient to explain the full magnitude of the disparity with GRACE, but several of them taken together may do so.

- 1. Because of the scale mismatch between GRACE and our study watersheds, the storage variability being compared is over quite different spatial scales. It is possible that our sample of watersheds represents a subset of the coterminous USA with particularly low water storage capacity, perhaps because these

Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



## Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



are disproportionally mountainous watersheds with high hydraulic gradients and limited soil profile development. However, we see low storage amplitude, as compared to GRACE, even in watersheds with relatively little topographic relief. Further, we would expect that if all things are equal, storage amplitude measured over a small watershed would be larger than the regional mean sensed by GRACE, because storage variations in adjacent watersheds within a region partly cancel out in the regional mean to the extent that they are not completely in phase.

2. It is possible that our assumption is false that during periods with low precipitation, evaporation, and snowmelt (according to the reanalysis), streamflow is the dominant flux of water in or out of the watershed. If so, then the streamflow-storage relationship we construct would be biased. It is clear from the streamflow record that the reanalysis frequently misses periods of heavy snowmelt in high-mountain basins, partly because the large topographic relief in these basins (which impacts the periods of snow accumulation and melt) is not captured by the coarse ~32 km spatial scale of the reanalysis. The atmospheric model used in NARR tends to underestimate mountain snowfall, so that snow is added to the model land surface in the analysis steps to nudge NARR toward observed snow cover (Luo et al., 2007). However, this would not explain the low storage amplitudes compared to GRACE observed in basins where snow is not a major part of the water budget.
3. It is conjectured that groundwater movement out of headwater basins that is not reflected in streamflow could be an important term in basin-scale water balance over the USA (Schaller and Fan, 2009), which would also seriously bias the storage-discharge relationships we constructed. However, given that basins with net groundwater outflow must be largely balanced on the continental scale by those with net groundwater inflow, it is difficult to see how this effect could be large and consistent enough to account for the systematic underestimate of the storage amplitude seen in almost our entire sample.

4. We assumed that when flow drops to zero, basin storage remains constant at a minimum value extrapolated from the storage-discharge relationship for periods of positive flow. This unquestionably results in an underestimate of the storage variability, since evaporation will result in decreasing storage over periods of zero flow. However, this cannot explain the disparity seen, since only 15/61 of our sample basins have any recorded hours of zero flow.
5. Finally, the assumption that the storage-discharge relationship as determined for hours of low precipitation, evaporation, and snowmelt is valid for other periods, which is necessary for computing monthly-mean storage, may not hold. For example, it is likely that the pool of water contributing to evaporation (largely soil moisture) and the pool of water contributing to streamflow (largely groundwater) are partly decoupled, so that changes in basin storage due to evaporation would not be immediately reflected in streamflow. Similarly, when there is precipitation, streamflow generation is likely to be qualitatively different than during periods without precipitation, with overland flow and greater contributing area, resulting in a different storage-discharge relationship compared to that inferred here. This would mean that “doing hydrology backward” based on the storage-discharge relationship inferred from recession curve analysis using Eq. (8) does not work for the majority of small watersheds, although the concept may nevertheless have heuristic value if it is approximately valid, for example, over periods of light precipitation. Note, however, that assuming that a non-constant storage-discharge relationship explains the majority of the observed discrepancy would require this relationship to undergo very large fluctuations over period of high precipitation, evaporation, and/or snowmelt – enough to increase the mean recession timescale by a factor of 10; cf. Eq. (6), – compared to the relatively modest modifications seen when the subset of hours chosen for constructing the recession curve is modified (Fig. 5).

**Stream recession  
and basin storage**

N. Y. Krakauer and  
M. Temimi

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

# 5 Conclusions

We have outlined a systematic method for constructing recession curves for small watersheds based on high-frequency streamflow measurements combined with reanalysis meteorology. Recession curve analysis appears to provide valuable information on basin hydrological characteristics and should be studied further for its possible utility in forecasting floods and low flows. We found that for the selected continent-wide sample of small, undisturbed watersheds, recession curves, as constructed by a uniform method intended to minimize the impacts of precipitation, snowmelt, and evapotranspiration, had broad similarities, with recession timescales typically increasing by a factor of 10 going from high flows as seen immediately after storms to flows near the median level, and showing indications of leveling off at low flows. We were able to quantify the uncertainty in each recession curve, and linked variability in the recession timescale across watersheds to known climate and geomorphological factors, but with a component of small-scale variability (particularly at low flow rates) which needs to be investigated in larger samples or with more explanatory variables. Storage variations inferred from the recession curve agree in terms of timing, but not amplitude, with independent gravimetric estimates. Study of the discrepancy in the inferred storage amplitude may provide clues to the range of validity of the recession curve constructed according to the method used here.

*Acknowledgement.* GRACE land data were processed by Sean Swenson, supported by the NASA MEASURES Program, and are available at <http://grace.jpl.nasa.gov>. This study was supported by the National Oceanic and Atmospheric Administration (NOAA) under Grant Number NA06OAR4810162. The statements contained in this article are not the opinions of the funding agency or government, but reflect the views of the authors.

HESSD

8, 1827–1860, 2011

## Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## References

- Bingham, R. H.: Regionalization of Low-Flow Characteristics of Tennessee Streams, Water Resources Investigations Report 85-4191, USGS, 1986. 1831, 1845
- Brandes, D., Hoffmann, J. G., and Mangarillo, J. T.: Base flow recession rates, low flows, and hydrologic features of small watersheds in Pennsylvania, USA, *J. Am. Water Resour. As.*, 41, 1177–1186, doi:10.1111/j.1752-1688.2005.tb03792.x, 2005. 1830, 1841, 1845
- Brutsaert, W.: Long-term groundwater storage trends estimated from streamflow records: climatic perspective, *Water Resour. Res.*, 44, W02409, doi:10.1029/2007WR006518, 2008. 1830, 1833, 1841
- Brutsaert, W.: Annual drought flow and groundwater storage trends in the eastern half of the United States during the past two-third century, *Theor. Appl. Climatol.*, 100, 93–103, doi:10.1007/s00704-009-0180-3, 2010. 1833
- Brutsaert, W. and Lopez, J. P.: Basin-scale geohydrologic drought flow features of riparian aquifers in the Southern Great Plains, *Water Resour. Res.*, 34, 233–240, doi:10.1029/97WR03068, 1998. 1831
- Brutsaert, W. and Nieber, J. L.: Regionalized drought flow hydrographs from a mature glaciated plateau, *Water Resour. Res.*, 13, 637–643, 1977. 1829, 1835, 1841, 1845
- Cantón, Y., Solé-Benet, A., and Domingo, F.: Temporal and spatial patterns of soil moisture in semiarid badlands of SE Spain, *J. Hydrol.*, 285, 199–214, doi:10.1016/j.jhydrol.2003.08.018, 2004. 1844
- Cleveland, W. S.: Robust locally weighted regression and smoothing scatterplots, *J. Am. Stat. Assoc.*, 74, 829–836, 1979. 1837
- Craven, P. and Wahba, G.: Smoothing noisy data with spline functions: Estimating the correct degree of smoothing by the method of generalized cross-validation, *Numer. Math.*, 31, 377–403, doi:10.1007/BF01404567, 1979. 1838
- van Dijk, A. I. J. M.: Climate and terrain factors explaining streamflow response and recession in Australian catchments, *Hydrol. Earth Syst. Sci.*, 14, 159–169, doi:10.5194/hess-14-159-2010, 2010. 1830, 1831, 1841, 1844, 1845
- Ek, M. B., Mitchell, K. E., Lin, Y., Rogers, E., Grunmann, P., Koren, V., Gayno, G., and Tarp-ley, J. D.: Implementation of Noah land surface model advances in the National Centers for Environmental Prediction operational mesoscale Eta model, *J. Geophys. Res.*, 108, 8851, doi:10.1029/2002JD003296, 2003. 1840

# HESSD

8, 1827–1860, 2011

## Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





- Eng, K. and Milly, P. C. D.: Relating low-flow characteristics to the base flow recession time constant at partial record stream gauges, *Water Resour. Res.*, 43, W01201, doi:10.1029/2006WR005293, 2007. 1830
- Gómez-Plaza, A., Alvarez-Rogel, J., Albaladejo, J., and Castillo, V. M.: Spatial patterns and temporal stability of soil moisture across a range of scales in a semi-arid environment, *Hydrol. Process.*, 14, 1261–1277, doi:10.1002/(SICI)1099-1085(200005)14:7<1261::AID-HYP40>3.0.CO;2-D, 2000. 1844
- Gómez-Plaza, A., Martínez-Mena, M., Albaladejo, J., and Castillo, V. M.: Factors regulating spatial distribution of soil water content in small semiarid catchments, *J. Hydrol.*, 253, 211–226, doi:10.1016/S0022-1694(01)00483-8, 2001. 1844
- Hall, F. R.: Base-Flow Recessions – A Review, *Water Resour. Res.*, 4, 973–983, doi:10.1029/WR004i005p00973, 1968. 1829
- Harman, C. J., Sivapalan, M., and Kumar, P.: Power law catchment-scale recessions arising from heterogeneous linear small-scale dynamics, *Water Resour. Res.*, 45, W09404, doi:10.1029/2008WR007392, 2009. 1837
- Hébrard, O., Voltz, M., Andrieuxa, P., and Moussa, R.: Spatio-temporal distribution of soil surface moisture in a heterogeneously farmed Mediterranean catchment, *J. Hydrol.*, 329, 110–121, doi:10.1016/j.jhydrol.2006.02.012, 2006. 1844
- Kirchner, J. W.: Catchments as simple dynamical systems: Catchment characterization, rainfall-runoff modeling, and doing hydrology backward, *Water Resour. Res.*, 45, W02429, doi:10.1029/2008WR006912, 2009. 1830, 1831, 1832, 1833, 1835, 1836, 1841
- Krakauer, N. Y. and Fung, I.: Mapping and attribution of change in streamflow in the coterminous United States, *Hydrol. Earth Syst. Sci.*, 12, 1111–1120, doi:10.5194/hess-12-1111-2008, 2008. 1834
- Krakauer, N. Y., Schneider, T., Randerson, J. T., and Olsen, S. C.: Using generalized cross-validation to select parameters in inversions for regional carbon fluxes, *Geophys. Res. Lett.*, 31, L19108, doi:10.1029/2004GL020323, 2004. 1838
- Luo, Y., Berbery, E. H., Mitchell, K. E., and Betts, A. K.: Relationships between land surface and near-surface atmospheric variables in the NCEP North American Regional Reanalysis, *J. Hydrometeorol.*, 8, 1184–1203, doi:10.1175/2007JHM844.1, 2007. 1847
- Mesinger, F., DiMego, G., Kalnay, E., Mitchell, K., Shafran, P. C., Ebisuzaki, W., Jović, D., Woollen, J., Rogers, E., Berbery, E. H., Ek, M. B., Fan, Y., Grumbine, R., Higgins, W., Li, H., Lin, Y., Manikin, G., Parrish, D., and Shi, W.: North American Regional Reanalysis, *B. Am.*

## Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Meteorol. Soc., 87, 343–360, 2006. 1835
- Mizumura, K.: Analyses of flow mechanism based on master recession curves, J. Hydrol. Eng., 10, 468–476, doi:10.1061/(ASCE)1084-0699(2005)10:6(468), 2005. 1837
- Mohanty, B. P., Famiglietti, J. S., and Skaggs, T. H.: Evolution of soil moisture spatial structure in a mixed vegetation pixel during the Southern Great Plains 1997 (SGP97) Hydrology Experiment, Water Resour. Res., 36, 3675–3686, doi:10.1029/2000WR900258, 2000. 1844
- Palmroth, S., Katul, G. G., Hui, D., McCarthy, H. R., Jackson, R. B., and Oren, R.: Estimation of long-term basin scale evapotranspiration from streamflow time series, Water Resour. Res., 46, W10512, doi:10.1029/2009WR008838, 2010. 1833
- Peña-Arancibia, J. L., van Dijk, A. I. J. M., Mulligan, M., and Bruijnzeel, L. A.: The role of climatic and terrain attributes in estimating baseflow recession in tropical catchments, Hydrol. Earth Syst. Sci., 14, 2193–2205, doi:10.5194/hess-14-2193-2010, 2010. 1831, 1841, 1844, 1845
- Qiu, Y., Fu, B., Wang, J., and Chen, L.: Soil moisture variation in relation to topography and land use in a hillslope catchment of the Loess Plateau, China, J. Hydrol., 240, 243–263, doi:10.1016/S0022-1694(00)00362-0, 2001. 1844
- Rodell, M., Houser, P., Jambor, U., Gottschalck, J., Mitchell, K., Meng, C., Arsenault, K., Cosgrove, B., Radakovich, J., Bosilovich, M., Entin, J., Walker, J., Lohmann, D., and Toll, D.: The Global Land Data Assimilation System, B. Am. Meteorol. Soc., 85, 381–394, 2004. 1840
- Roering, J. J., Marshall, J., Booth, A. M., Mort, M., and Jin, Q.: Evidence for biotic controls on topography and soil production, Earth Planet. Sc. Lett., 298, 183–190, doi:10.1016/j.epsl.2010.07.040, 2010. 1844
- Rupp, D. E. and Selker, J. S.: Information, artifacts, and noise in  $dQ/dt - Q$  recession analysis, Adv. Water Res., 29, 154–160, doi:10.1016/j.advwatres.2005.03.019, 2006. 1831
- Schaller, M. F. and Fan, Y.: River basins as groundwater exporters and importers: Implications for water cycle and climate modeling, J. Geophys. Res., 114, D04103, doi:10.1029/2008JD010636, 2009. 1847
- Showstack, R.: Online database for instantaneous streamflow data, EOS, 88, 523, 2007. 1834
- Slack, J. R. and Landwehr, J. M.: Hydro-Climatic Data Network (HCDN): A U.S. Geological Survey Streamflow Data Set for the United States for the Study of Climate Variations, 1874–1988, Tech. rep., USGS Open-File Report 92-129, 1992. 1834
- Swenson, S.: Restoring Signal Loss in GRACE Terrestrial Water Storage Estimates, online at: [ftp://podaac.jpl.nasa.gov/tellus/grace\\_monthly/swenson\\_destripe/ss201008/doc/swenson.grace.scaling\\_description\\_doc\\_draft.pdf](ftp://podaac.jpl.nasa.gov/tellus/grace_monthly/swenson_destripe/ss201008/doc/swenson.grace.scaling_description_doc_draft.pdf), last access: 7 February, 2011. 1840

## Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Swenson, S. and Wahr, J.: Post-processing removal of correlated errors in GRACE data, *Geophys. Res. Lett.*, 33, L08402, doi:10.1029/2005GL025285, 2006. 1840
- Syed, T. H., Famiglietti, J. S., Rodell, M., Chen, J., and Wilson, C. R.: Analysis of terrestrial water storage changes from GRACE and GLDAS, *Water Resour. Res.*, 44, W02433, doi:10.1029/2006WR005779, 2008. 1843
- Tallaksen, L. M.: A review of baseflow recession analysis, *J. Hydrol.*, 165, 349–370, 1995. 1829
- Temimi, M., Leconte, R., Chaouch, N., Sukumal, P., Khanbilvardi, R., and Brissette, F.: A combination of remote sensing data and topographic attributes for the spatial and temporal monitoring of soil wetness, *J. Hydrol.*, 388, 28–40, doi:10.1016/j.jhydrol.2010.04.021, 2010. 1844, 1845
- Vogel, R. M. and Kroll, C. N.: Regional geohydrologic-geomorphic relationships for the estimation of low-flow statistics, *Water Resour. Res.*, 28, 2451–2458, doi:10.1029/92WR01007, 1992. 1830, 1831, 1841, 1845
- Weisman, R. N.: The effect of evapotranspiration on streamflow recession, *Hydrolog. Sci. J.*, 22, 371–377, doi:10.1080/02626667709491731, 1977. 1830, 1841
- Wittenberg, H.: Baseflow recession and recharge as nonlinear storage processes, *Hydrol. Process.*, 13, 715–726, doi:10.1002/(SICI)1099-1085(19990415)13:5<715::AID-HYP775>3.0.CO;2-N, 1999. 1830, 1831
- Wittenberg, H.: Effects of season and man-made changes on baseflow and flow recession: case studies, *Hydrol. Process.*, 17, 2113–2123, doi:10.1002/hyp.1324, 2003. 1830
- Zecharias, Y. B. and Brutsaert, W.: Recession characteristics of groundwater outflow and base flow from mountainous watersheds, *Water Resour. Res.*, 24, 1651–1658, doi:10.1029/WR024i010p01651, 1988. 1845
- Zhang, L., Chen, Y. D., Hickel, K., and Shao, Q.: Analysis of low-flow characteristics for catchments in Dongjiang Basin, China, *Hydrogeol. J.*, 17, 631–640, doi:10.1007/s10040-008-0386-y, 2009. 1831
- Zhu, R., Zheng, H., and Liu, C.: Estimation of groundwater residence time and recession rate in watershed of the Loess Plateau, *J. Geogr. Sci.*, 20, 273–282, doi:10.1007/s11442-010-0273-z, 2010. 1831

## Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Stream recession  
and basin storage**N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Fig. 1.** Locations of sample watersheds (markers,  $n = 61$ ) within the coterminous United States.



**Fig. 2.** Scatter of hourly  $\dot{Q}$  vs.  $Q$ , excluding periods with precipitation, high evaporation, or rising flow, at an example site (USGS gauge 01384500, Ringwood Creek, New Jersey; nonnegative  $\dot{Q}$  values not shown).

# HESSD

8, 1827–1860, 2011

## Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Stream recession  
and basin storageN. Y. Krakauer and  
M. Temimi

**Fig. 3.** (a) Binned mean hourly  $-dQ/dt$  vs.  $Q$  for sample site, together with the smoothed functional fit (middle line). Error bars are standard errors for each bin; upper and lower lines show uncertainty of fit ( $\pm 1$  standard error). (b) Binned mean recession time  $\tau$  vs.  $Q$  for sample site, together with the smoothed fit of the relationship (error bars are standard errors for each bin; upper and lower lines show fit uncertainty as  $\pm 1$  standard error).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 4. (a)** Fitted recession time  $\tau$  vs.  $Q$  across the 61 streams. **(b)** Inverse-variance weighted mean recession timescale  $\tau$  (curve, with weighted within-site uncertainty given by upper and lower envelope) along with variability across streams (weighted standard deviation given by error bars). Variability in recession curves across streams is consistently much greater than the uncertainty associated with recession-curve estimation for individual streams, meaning that most of the variability in recession times seen across streams is real.

## Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 5.** Inverse-variance weighted mean recession timescale  $\tau$  across sites (as in Fig. 4b) along with the mean recession timescale for cases where the hours considered (1) included periods of high evaporation, (2) included periods of rising flow.

# HESSD

8, 1827–1860, 2011

## Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion





Stream recession  
and basin storageN. Y. Krakauer and  
M. Temimi

**Fig. 6.** (a) Nonparametric (Spearman) correlation coefficient with the recession timescale at different streamflows of the six variables (1) longitude, (2) soil infiltration capacity, (3) latitude, (4) channel length, (5) forest cover, (6) precipitation;  $|r| \geq 0.3$  is significantly different from zero at the 0.05 level. (b) Coefficient of determination  $R^2$  for multivariate linear regression with these six predictor variables;  $R^2 \geq 0.3$  is significant at the 0.05 level.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



# Stream recession and basin storage

N. Y. Krakauer and  
M. Temimi



**Fig. 7.** Amplitude of water storage variability inferred from the recession curve vs. that inferred from GRACE gravimetry, showing the standard deviation of the **(a)** seasonal cycle and **(b)** interannual variability. The solid lines are 1–1 relationships. The amplitudes inferred from recession curve analysis are generally low compared to GRACE.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

