

This discussion paper is/has been under review for the journal Hydrology and Earth System Sciences (HESS). Please refer to the corresponding final paper in HESS if available.

# Hydrogeology and hydrogeochemistry of an alkaline volcanic area: the NE Mt. Meru slope (East African Rift – Northern Tanzania)

G. Ghiglieri<sup>1</sup>, D. Pittalis<sup>2</sup>, G. Cerri<sup>3</sup>, and G. Oggiano<sup>3</sup>

<sup>1</sup>Department of Earth Sciences, University of Cagliari, Italy, Desertification Research Group-NRD, University of Sassari, Italy

<sup>2</sup>Department of Territorial Engineering, Desertification Research Group-NRD, University of Sassari, Italy

<sup>3</sup>Department of Botanical, Ecological and Geological Sciences, Desertification Research Group-NRD, University of Sassari, Italy

Received: 22 July 2011 – Accepted: 12 August 2011 – Published: 1 September 2011

Correspondence to: G. Ghiglieri (ghiglieri@unica.it)

Published by Copernicus Publications on behalf of the European Geosciences Union.

**HESSD**

8, 8255–8289, 2011

**Hydrogeology and  
hydrogeochemistry  
of an alkaline  
volcanic**

G. Ghiglieri et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Abstract

The objective of this study is to analyze the geochemical conditions associated with the presence of fluoride ( $F^-$ ) in the groundwater of an area of Northern Tanzania. The studied aquifers are composed of volcanic rocks such as phonolitic and nephelinitic lavas, basalts, lahars of various ages and mantling ash. Sedimentary rocks consisting of fine-grained alluvial and lacustrine deposits occur as well. Samples collected from springs, borehole and surface water, during two monitoring surveys, were analyzed for the various physico-chemical and isotopic parameters. The geochemical composition of water is typically sodium bicarbonate. High values of  $F^-$  (up to  $68 \text{ mg l}^{-1}$ ) were recorded. The highest values of fluoride agreed with the highest values of pH, sodium and bicarbonate. Dissolution of major ions, exchange processes and precipitation of  $\text{Ca}^{2+}$  from super-saturated solutions joined with the local permeability and hydraulic gradients, control the fluoride mobilization and the contamination of the area.

## 1 Introduction

The pollution of groundwater has become a major environmental issue, particularly where groundwater represents the main source of drinking water. This situation is so common in many lesser developed countries that the security of drinking water supply has been chosen as one of the ten Millennium Development Goals. The natural contamination of groundwater resulting from its surrounding geological environment can be an important factor determining the quality of drinking water. One of the most severe such natural contaminants is fluoride, especially in the East African Rift Valley, where malformed bones and certain neurological ailments are known to be caused by fluorosis, and where the presence of fluoride has some influence on the ecology of the region. Sources of fluoride derives largely from rock minerals, air and seawater, but anthropogenic activity can also make a contribution (Fuge and Andrews, 1988). In Tanzania, for example, the level of fluoride in drinking water frequently exceeds the

**HESSD**

8, 8255–8289, 2011

## Hydrogeology and hydrogeochemistry of an alkaline volcanic

G. Ghiglieri et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

WHO guideline of  $1.5 \text{ mg l}^{-1}$  and, from time to time, even the  $4.0 \text{ mg l}^{-1}$  limit set by the Tanzanian Government. The alkaline volcanism in the Rift Valley has been implicated for the high fluoride concentration in the local groundwater (Clarke et al., 1990; Davies, 1996; Deocampo, 2003), but neither the behaviour of fluoride in relation to the rock composition of the aquifer, nor the residence time, nor the concentration of other ions have been widely researched. The present study sought to explore the chemistry of the groundwater in the Arusha region of Tanzania, using a combination of hydrochemical, minero-chemical and isotopic analyses, with a view to developing a set of criteria for identifying low fluoride carrying sources of groundwater (Ghiglieri et al., 2010).

## 2 Study area and methodology

### 2.1 Study area

The study area lies within the Meru District, which administratively comprises six divisions, 37 wards and 133 villages (Fig. 1).

It is bounded by Mt. Meru and the Arusha National Park, occupies about  $370 \text{ km}^2$  and includes nine villages within the Oldonyo Sambu and Ngarenanyuki wards. The area is part of the Maasai Steppe, which extends from Lake Turkana in Kenya to central Tanzania. The natural vegetation is typically savannah. The topography of the study region is dominated by the Mt. Meru volcanic cone of Pleistocene to recent origin. The local climate is temperate Afro-Alpine, with an annual precipitation of 450 mm (Hijmans et al., 2005) and mean daily temperature minima and maxima of, respectively,  $20.6^\circ\text{C}$  and  $28.5^\circ\text{C}$ . The rainfall is irregularly distributed between a main wet season from February to mid May (which contributes  $\sim 70\%$  of the annual precipitation), and a minor one from September to November which provides much of the remainder.

## Hydrogeology and hydrogeochemistry of an alkaline volcanic

G. Ghiglieri et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## 2.2 Geological and hydrogeological setting

The relationship between the local geology and hydrogeology has been described elsewhere (Ghiglieri et al., 2010). The volcano-sedimentary succession is Cenozoic, with some features dating back to the Miocene-Pliocene, and more recent ones to the Olocene. Volcanic rock dominates, with some recent alluvial deposits. There are no crystalline basement outcrops, rather this material lies at a shallow depth a few kilometers to the north of the study area. Major rift faults are present on the NW margin (Matuginigi and Matisiwi Escarpment). Linear features and benches are commonplace on the flanks of Mt. Meru, and it is highly probable that the early volcanic structure has been block-faulted. In the central area, the faults trend either N-S or NNE-SSW (Uwiro graben), while in the NW section, the trend is NW-SE (parasitic cone in Lassarkartarta). The two dominant hydrogeological features are the volcanic phonolitic and nephelinitic lavas (as well as basalts, lahars of various ages and mantling ash) and sedimentary material of fine-grained alluvial and lacustrine origin. Groundwater recharge, transmission and discharge are determined by a combination of geomorphology, geology and structural patterns (Ghiglieri et al., 2010), and both shallow and deep circulating groundwater can be distinguished. Shallow groundwater hosted in unconsolidated or semi-consolidated saturated sediments are referred to as local systems. Intermediate and deep groundwater circulation occurs where the permeability of the aquifer and a sufficient elevation difference between recharge and discharge area allow deep infiltration. Deep infiltration is also promoted where widespread fracturing and faulting affects the rock. Where these circumstances prevail, substantial precipitation levels can support productive wells and springs (e.g. the main cone group complex Nvm). The groundwater is characterized by a multi-directional flow, dominated by movement from the higher elevation southern part of the region towards the lower lying area in the north. Recharge is via both rainfall infiltration and lateral connections to other hydrogeologic units. The latter affects the Mkuru area in particular, where an aquifer lying in weathered and scoriaceous basalt at a depth of 40–60 m is fed by groundwater infiltrated from a high elevation area in the main cone group.

### Hydrogeology and hydrogeochemistry of an alkaline volcanic

G. Ghiglieri et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 2.3 Material and methods

### 2.3.1 Collection of field data and analytical procedures

Water was sampled at three different times, determined by a survey conducted in February 2007, with subsequent monitoring (Ghiglieri et al., 2010). A set of 58 sites was established, including 46 springs (30 in the Ngarenanyuki ward and 16 in Oldonyosambu) and six surface water sites (Fig. 1). Portable devices (Hanna Instrument models HI 98130 pH/EC/T and HI 93739 photometer), were employed to record pH, electrical conductivity, temperature and fluoride concentration. Two monitoring surveys were implemented, the first in March–April 2007, referred to as Masika, involved 34 samples (25 spring waters, nine surface waters), and the second in January 2008 (pre-Masika) 31 samples (25 spring waters, six surface waters). Two samples from each site were filtered (0.45  $\mu\text{m}$ ) into 1l polythene bottles thoroughly pre-washed with distilled water. The samples were transported in low temperature thermal bags and stored under refrigeration.

Standard methods (APHA, 1992) were applied to quantify the presence of major ions, and the analysis of nitrate, nitrite, ammonia and fluoride was performed at the AUWSA laboratory in Arusha. Other chemical determinations were undertaken at the University of Sassari Department of Territorial Engineering. Cation presence ( $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ) was determined by atomic absorption spectrometry (Perkin Elmer model AAnalyst 200), and that of anions ( $\text{F}^-$ ,  $\text{Cl}^-$ ,  $\text{SO}_4^{2-}$ ) by ion chromatography (anion column Alltech model allsep anion 7 $\mu\text{m}$ , 100 mm). Carbonate and bicarbonate contents were obtained by titration, and silicate by colorimetry. For the sample from spring 26 ENG, elemental analysis was performed by inductively coupled plasma mass spectrometry at the University of Barcelona. The ion balance errors for the analyses were generally within  $\pm 5\%$ . The saturation index (hereafter, SI) for fluorite, fluorapatite, calcite, villiaumite and the chemical facies were computed using PHREEQC v2.1 software (Parkhurst and Appelo, 1999) and AQUACHEM v3.1 (SigmaStat software). Isotopic analyses of  $^{18}\text{O}$ ,  $^2\text{H}$  and  $^3\text{H}$  were carried out at the CNR Laboratory (Pisa) for the set

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of Masika samples, along with a sample from Ichnusa Well1 collected in January 2008, applying analytical methods described, respectively, by Epstein and Mayeda (1953), Coleman et al. (1982) and the US Department of Energy (1997). Isotope content values ( $\delta^{18}\text{O}$ ,  $\delta\text{D}$ ) are expressed in ‰ relative to the Vienna Standard Mean Ocean Water (VSMOW) defined by Craig (1961), and  $^3\text{H}$  concentrations in terms of tritium units (TU). Mineralogical analyses were performed on a set of 12 samples at the Department of Botanical, Ecological and Geological Sciences, University of Sassari using a SIEMENS D5000 X-ray diffractometer (Bragg-Brentano geometry) equipped with a Cu tube and a graphite monochromator on the diffracted beam. The following operating conditions were applied: 40 kV, 30 mA,  $2\theta$  range 2–70°, step size 0.02°, 2 s step<sup>-1</sup>. Wet milling was avoided due to the possible presence of highly soluble phases; instead powders were prepared by hand in an agate mortar. Minerals were identified using Bruker EVA v14.2 software (Bruker AXS, 2008) and the PDF-2 database (ICDD, 2003). All statistical analyses were based on R v2.7.0 software (<http://www.r-project.org/>).

## 3 Results

### 3.1 Chemistry

The physico-chemical properties of the groundwater samples varied markedly (Table 1). The groundwater pH varies from 5.9 to 8.1. The Electrical conductivity (EC) varies from 190 to 5070  $\mu\text{S cm}^{-1}$ . Sodium and  $\text{HCO}_3^-$  are the dominant ions ranging, from 24.5 to 1100  $\text{mg l}^{-1}$  and from 89.5 to 2143  $\text{mg l}^{-1}$  respectively. The concentration of fluoride in the groundwater varies from 0.90 to 68.00  $\text{mg l}^{-1}$ . In the most of groundwater samples (84%),  $\text{F}^-$  exceeds the WHO limit (1.5  $\text{mg l}^{-1}$ ), whereas the 70% are above the Tanzanian limit (4  $\text{mg l}^{-1}$ ).

## Hydrogeology and hydrogeochemistry of an alkaline volcanic

G. Ghiglieri et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The ternary diagram analysis identified a unique water type, namely bicarbonate-alkaline-earth (Fig. 2).

Chloride concentrations were relatively low, as is the case for most natural water systems (Davies, 1996), while the concentration of the other major anions was rather variable. The positive correlation between the concentration of fluoride with those of bicarbonate and sodium (Figs. 3 and 4) suggested that in bicarbonate-alkaline-earth waters, the presence of bicarbonate and sodium ions may exert an influence over the fluoride concentration.

The river waters also shows a bicarbonate-alkaline hydrofacies, in particular at the three collection sites along the Ngarenanyuki river (river samples 1, 24 and 28), where the concentration of major ions and fluoride was higher than in the other river water samples (Fig. 5).

### 3.2 Mineralogy

The outcome of the X-ray analysis applied to the sample subset is given in Table 2. The more (ENG6; OLD11) or less (OLD9, ENG1; ENG31) pronounced Gaussian shape of the background in the  $\approx 15\text{--}40^\circ 2\theta$  region showed that, in addition to the crystalline phases, a glassy fraction was also present in all samples. The phonolite rock samples (ENG1 and ENG31) contain alkali feldspar and feldspathoids, along with riebeckite, augite, fluoroapatite; these phases also occur, with only minor exceptions, in all samples. Among the authigenic minerals, the zeolites phillipsite and chabazite (which are ubiquitous in lake deposits and frequently found in lahars) were formed because of the interaction between volcanic glass and alkaline solutions; cancrinite (limited to lahars and lake deposits) can be considered as a newly-formed phase, while analcime can be a magmatic as well as a secondary-formed phase. Nepheline and leucite are magmatic in this context, and represent detrital minerals in sedimentary lithotypes. Clay minerals, belonging to the illite/smectite group, were found in paleosoils interbedded within various lahar (ENG6, OLD9 and OLD11). Trona, natron, natrite, sylvite, apthitalite and calcite were present, though never all together. These phases can be directly

## Hydrogeology and hydrogeochemistry of an alkaline volcanic

G. Ghiglieri et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



precipitated from super-saturated solutions and frequently form, sometimes in conjunction with zeolites, crusts, hardpans and calcrete, as in the calcrete OLD9, in breccias (OLD3, 4 and 5) and in the crust of a lake deposit (2474 C). Trona is referred to locally as “scooped magadi” when it forms an efflorescent crust on the soil surface in association with mixtures of halite, quartz, villiaumite, kogarkoite and thermonatrite (Nielsen, 1999). This assemblage was not present in samples in which trona was associated with either natrite, sylvite, apthitalite, goethite, phillipsite, chabazite and analcime of probably secondary formation (ENG8b), or calcite, cancrinite, phillipsite, chabazite and secondary analcime (2474 C). The respective underlying samples (ENG8 and 2474 S, just below the crust) were richer in primary minerals and contained neither trona, calcite nor apthitalite. Only traces of natron and natrite were identified in ENG8.

### 3.3 Isotopic data

The majority of groundwater samples lay between GMWL and LMWL in the isotope diagram (Fig. 6), which implies relatively depleted levels of  $^{18}\text{O}$  and  $^2\text{H}$ , relatively unaffected by evaporation, and readily diluted by the infiltration of rainwater, as can be expected given the high hydraulic conductivity within the Mt. Meru recharge zone.

Nevertheless, some groundwater samples do show a higher  $^{18}\text{O}$  and  $^2\text{H}$  content (Table 3), and this is thought to reflect a stronger influence of evaporation, along with a slower rate of infiltration or an appreciable return flow of irrigation water (particularly in sediment and lahar formations), as well as some leaching from the rock into the groundwater. At higher altitudes, with only few exceptions, the water had a high  $^3\text{H}$  content and a moderately depleted stable isotope content (Fig. 7). Here, the groundwater was likely to have been largely recharged via the infiltration of rainwater through fractures, in line with hydrogeologic evidence (Ghiglieri et al., 2010). On the slope and in the lowland area, the waters were characterized by a rather low  $^3\text{H}$  content, suggesting a longer residence time. Most of these samples plot below the LMWL, indicating the importance of evaporation during recharge, and suggesting pounding effects in less permeable soil horizons. These same samples showed the greatest degree of  $^{18}\text{O}$

## Hydrogeology and hydrogeochemistry of an alkaline volcanic

G. Ghiglieri et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





depletion, and some originated from the Uwiro graben, the main tectonic depression in the area. In the high eastern part of the study area, the groundwater temperature was more elevated, and its stable isotope and  $^3\text{H}$  contents were both depleted. So here, recharge could have occurred either through rainwater carrying a depleted isotope content, and/or the current meteoric water has been circulating at a deep level, during which time its isotopic content has become altered by isotope exchange with volcanic  $\text{CO}_2$ .

### 3.4 Fluoride distribution

The distribution of fluoride was determined by a combination of altitude and aquifer lithology. Within phonolite and basalt aquifers, the fluoride level was lowest on the phonolite outcrops at higher altitudes ( $1.6 \text{ mg l}^{-1}$ ), whereas the concentration in water emerging from basalt at the base of the Mt. Meru cone reached  $7.2 \text{ mg l}^{-1}$ .

The highest contents were in the water emerging from lahar hosted aquifers in the flat land within the Uwiro graben. Figure 8 shows the flow route from the recharge zone of the NW side Mt. Meru (phonolite) to the Oldonyo Sambu valley (lahar), and documents the fluoride content of a number of local springs. The 26 ENG spring (along with 28 ENG and 29 ENG) were the exceptions to the trend of fluoride content decreasing with altitude. Its water had a constant temperature of  $22.4^\circ\text{C}$  and a depleted  $^3\text{H}$  content, which together suggested that the spring is of hydrothermal origin; this would account for its showing the highest fluoride content so far detected in the study area. 26 ENG feeds the Engarenanyuky river at a rate of  $6 \text{ l s}^{-1}$ , and thus represents a source of pollution for both the river and any aquifers hydraulically connected it, such as in the area lying on the Uwiro graben lowland.

## Hydrogeology and hydrogeochemistry of an alkaline volcanic

G. Ghiglieri et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 4 Insights gained from the physico-chemical analysis

### 4.1 The dissolution of major ions

Typically, silicate minerals and glass present in alkaline lava and ash are weathered by hydrolysis, producing  $\text{NaHCO}_3$  rich and  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  depleted groundwater (Jones et al., 1977). Here, this process affected the degradation of phonolite, tephrite-phonolite, Na-K-feldspar and Na-K-feldspatoid, which occurred in the lahar formations. Groundwaters, particularly those containing dissolved  $\text{CO}_2$ , react readily with alkaline silicate such as albite to release sodium and bicarbonate ions. This process appears to be commonplace in the NE-SW-trending fault system on the eastern flank of Mt. Meru, where the most suitable hydrothermal conditions occur. The alkalinity of groundwaters is strictly associated with the presence of alkali metals (Jalali, 2007), as demonstrated by the positive correlation noted between alkalinity (bicarbonate content) and the sum of alkali and earth alkaline metals (Fig. 9).

### 4.2 Exchange processes of the major ions

The exchange capacity process affects secondary mineral associations such as zeolite and other sodium-hydrate silicates characteristically formed in an high evaporation environment (Sebag et al., 2001). These minerals are able not only to concentrate cations, but also to release them in response to changes in water availability and temperature which occur as a result of normal seasonal cycling.

By influencing the concentration of cations in the water, the zeolites can indirectly play a role on fluorine distribution, permitting or preventing the precipitation of F-bearing phases, namely fluorides. As shown in Fig. 10, there is a marked negative correlation between  $\text{Na}^+ + \text{K}^+ - \text{Cl}^- - \text{F}^-$  and  $\text{Ca}^{2+} + \text{Mg}^{2+} - \text{HCO}_3^- - (\text{SO}_4)^{2-}$ . According to Su et al. (2007), samples lying furthest from the origin have experienced the greatest extent of ionic exchange.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion



### 4.3 Fluoride content

Since the presence/absence of certain minerals shows which minerals have been dissolved/ precipitated, saturation index calculations were derived for each of the ground-water samples. The fluoride content of the groundwaters was verified by the positive relationship existing between the saturation index (SI) and the ionic strength (*I*). When SI < 1, minerals tend to dissolve, while at SI > 1, they tend to be precipitated. The SI values for fluorite (−1.89 – +1.25 in the *Masika* samples, and 1.55–1.61 in the *pre-Masika* samples) and those for *I* (0.002–0.05) suggested that the phonolite and basalt groundwaters were sub-saturated with respect to fluorite, while those emerging from lahar and sedimentary rocks were at equilibrium between the mineral phase and the solution (Fig. 11). The data can also be used to infer the residence time of the groundwater, and these inferences largely supported the <sup>3</sup>H-derived conclusions, such that residence time of the groundwater in the phonolite formations was short, while that in the lahars was somewhat longer.

Although these data may be predictive with respect to fluorite super-saturation, this phase was not identified in the study area, presumably because of the scarcity of dissolved calcium as scavenged by CaCO<sub>3</sub> precipitation according to:



Calcrete and hardpan were noted at the base of these soil profiles, and travertine concretion on the porous lava flows. Calcium removal by the precipitation of calcite allows the fluoride concentration to increase (Kim and Young Jeong, 2005), as supported by the negative correlation obtaining between calcium and fluoride, particularly in the waters derived from phonolite aquifers (Fig. 12).

Ionic exchange can also enhance calcium capture and sodium release. The variation in the 26 ENG SI with respect to various minerals at various temperatures is shown in Fig. 13. Sub-saturation values for fluorite were not reached at temperatures >150 °C. At the temperature of the emerging spring water (22 °C), this phase is insoluble. Thus

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion



it may be that fluorapatite was the source of fluoride in the water.



As shown in the plot its behaviour is antithetic with respect to the  $\text{CaF}_2$ . At a temperature  $>36^\circ\text{C}$ , fluorapatite precipitates, so the expectation is that this mineral would remain sub-saturated in the groundwater. In supergene condition, a further source of fluoride can be provided by secondary mineral associations (Nielsen, 1999; Vuhahulaa et al., 2008), which are favoured by the high level of evaporation typical of this climatic region (Lahermo et al., 1991).

An increased  $\text{Na}^+/\text{Ca}^{2+}$  ratio could theoretically favour the fractionation of fluoride by villiaumite, a mineral phase which is extremely soluble (Gao et al., 2007). However, the groundwaters were uniformly strongly sub-saturated with respect to villiaumite (Fig. 14). Even though villiaumite may be present in evaporative lakes (Kilham and Hecky, 1973), it was not detected here either in saline crusts or in association with scooped magadi. A ready source of alkali and fluoride (up to 3700 ppm) is also presented by the scooped magadi and crusts of the sediment cover. Thus, the fluoride rich Ngarenanyuki river water, in combination with leachate from the scooped magadi, may well have contributed to the increased fluoride content in the lahar aquifer.

A further important source of fluoride is represented by the ubiquitous volcanic ash (not present on the higher slopes of Mt. Meru). A leaching test conducted at ambient temperature on a representative sample of slightly altered, yellowish, powder-like ash showed that a considerable amount of fluoride can be readily released (Fig. 15), explaining the high fluoride content of alkaline magmas. The melting of Na-phonolite shows a higher diffusivity of fluoride (Balcone-Boissard et al., 2009) with respect chloride, and as opposed to  $\text{H}_2\text{O}$  and S hardly degass from the melt so giving rise to fluoride-rich glassy ejecta (Signorelli et al., 1999).

**Hydrogeology and  
hydrogeochemistry  
of an alkaline  
volcanic**

G. Ghiglieri et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 5 Conclusions

The chemistry of the groundwaters showed that a variety of natural processes must have been responsible both for the diversity of the hydrochemical facies and for the presence of fluoride contamination. Chief among these processes are dissolution, exchange capacity and precipitation from super-saturated solutions. A fluoride contamination model demonstrated that an interaction between the groundwater and certain fluoride-rich minerals could explain the variation in fluoride concentration observed in the recharge and flow-through area of northern Mt. Meru, while in the discharge area, evaporation and exchange capacity made a greater contribution to the fluoride presence in the surface- and groundwater. The groundwater emerging from phonolite had a low content of both alkaline cations and fluoride, due to the relatively low reactivity of silicate. The phonolite hosted-aquifers are highly fractured, and are thus highly permeable (Ghiglieri et al., 2010). Water carried through a scoriaceous, highly permeable, basalt aquifer showed an even lower concentration of fluoride and alkaline cations. The elevated fluoride content of lahar-hosted groundwater, on the other hand, derived from both the high permeability of the rock and the contribution of zeolites which both contain a high level of exchangeable cations and have an ash matrix. Volcanic ash is a major source of fluoride release. The dissolution process (and particularly, the cyclic dissolution/precipitation of the fluoride-rich trona which occurs as a seasonal encrustation in low-lying river valleys and ponds at lake margins), on the other hand, is an important factor behind the high fluoride content of the saturated superficial sediments. Hydrothermal springs, such as those which feed into the Ngarenanyuki river, make a further contribution to the level of fluoride pollution in the groundwater.

The geo-lithological characteristics of an aquifer, the recharge altitude and the residence time of the groundwater together determine the extent of the water ionic charge. Three major types of aquifer can be distinguished. The first is the high altitude phonolite-hosted aquifer, in which high transmissivity, high elevation and low temperature (16°C) together serve to inhibit the rate of dissolution of both mineral and

**HESSD**

8, 8255–8289, 2011

### Hydrogeology and hydrogeochemistry of an alkaline volcanic

G. Ghiglieri et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

## Hydrogeology and hydrogeochemistry of an alkaline volcanic

---

G. Ghiglieri et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

glassy groundmass, giving rise to a low fluoride groundwater. The second is the sedimentary (lahar and alluvial/lacustrine) hosted aquifer, frequently found at the base of Mt. Meru. Their relatively low altitude means that high temperatures prevail near the surface, favouring the precipitation of  $\text{CaCO}_3$ , and thereby inhibiting the precipitation of  $\text{CaF}_2$  and producing a higher fluoride content. The occurrence of CEC-endowed neoformed phases, along with the magadi, enhances the seasonal cycle of fluoride entrapment/release. Finally, the buried basalt hosted type of aquifer, associated with an elevated recharge area, shows a high transmissivity (Ghiglieri et al., 2010) which is optimal for maintaining a low ionic content. Infiltration into these aquifers occurs under rather cool conditions, the water tends to have a low residence time and no feeding occurs from the flatter areas where calcrete formation can deplete the level of calcium and thereby raise the fluoride concentration in the water.

*Acknowledgements.* This research was done as part of a project funded by OIKOS Institute (Italy), Charity and Defence of Nature Fund (private foundation) and Sardinia local Government (Italy) (Regional Law 19/96: cooperation with developing countries). Thanks are due also to Fondazione Banco di Sardegna for the financial support to D. Pittalis. For the purpose of the research, thanks are due to OIKOS EAST-AFRICA for technical and logistic support in Tanzania.

## References

- APHA: Standard methods for examination of water and wastewater. America Public Health Association, Washington D.C.-APHA, 1992.
- Balcone-Boissard, H., Baker, D. R., Villemant, B., and Boudon, G.: *F* and *Cl* diffusion in phonolitic melts: Influence of the Na/K ratio, *Chemical Geology*, 263, 89–98, 2009.
- Bruker AXS: Diffrac<sup>plus</sup> Evaluation package EVA 14. Release 15-07-2008, Bruker AXS GmbH Karlsruhe, Germany, 2008.
- Clarke, M. C. G., Woodhall, D. G., Allen, D., and Darling, G.: Geological, volcanological and hydrological controls on the occurrence of geothermal activity in the area surrounding Lake Naivasha, Kenya, Ministry of Energy Report, Nairobi, Kenya, 138 pp., 1990.

## Hydrogeology and hydrogeochemistry of an alkaline volcanic

G. Ghiglieri et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Coleman, M. C., Shepherd, T. J., Durham, J. J., Rouse, J. D., and Moore, G. R.: Reduction of water with zinc for hydrogen isotope analysis, *Anal. Chem.*, 54, 993–995, 1982.
- Craig H.: Standards for reporting concentrations of deuterium and oxygen-18 in natural waters, *Science*, 133, 1833–1834, 1961.
- 5 Davies, T. C.: Chemistry and pollution of natural waters in western Kenya, *J. Afr. Earth Sci.*, 23(4), 547–563, 1996.
- Deocampo, D. M.: Hydrogeochemistry in the Ngorongoro Crater, Tanzania, and implications for land use in a World Heritage Site, *Appl. Geochem.*, 19, 755–767, 2003.
- Dettman, D. L., Palacios-Fest, M. R., Nkotagu, H., and Cohen A. S.: Paleolimnological investigations of anthropogenic environmental change in Lake Tanganyika: VII. Carbonate isotope geochemistry as a record of riverine runoff, *J. Paleolimnol.*, 34, 93–105, 2005.
- 10 Epstein, S. and Mayeda, T.: Variation of O<sup>18</sup> content of waters from natural sources, *Geochim. Cosmochim. Ac.*, 4, 213–224, 1953.
- Fuge, R. and Andrews, M. J.: Fluoride in the UK environment, *Environ. Geochem. Health*, 10, 96–104, 1988.
- 15 Gao, X., Wang, Y., Li, Y., and Guo, Q.: Enrichment of fluoride in groundwater under the impact of saline water intrusion at the salt lake area of Yuncheng basin, northern China, *Environ. Geol.*, 53(4), 795–803, doi:10.1007/s00254-007-0692-z, 2007.
- Ghiglieri, G., Balia, R., Oggiano, G., Ardaù, F., and Pittalis, D.: Hydrogeological and geophysical investigations for groundwater in the Arumeru District (Northern Tanzania). Presentation at the 84° National Meeting of the Italian Geologic Society. Sassari 15–17 September 2008 (Conference Proceedings, 431–433, Vol. 2), 2008.
- 20 Ghiglieri, G., Balia, R., Oggiano, G., and Pittalis, D.: Prospecting for safe (low fluoride) groundwater in the Eastern African Rift: the Arumeru District (Northern Tanzania), *Hydrol. Earth Syst. Sci.*, 14, 1081–1091, doi:10.5194/hess-14-1081-2010, 2010.
- 25 Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., and Jarvis, A.: Very high resolution interpolated climate surfaces for global land areas, *Int. J. Climatol.*, 25, 1965–1978, 2005.
- ICDD: International Centre for Diffraction Data. PDF-2. Release 2003. Newton Square, Pennsylvania, USA, 2003.
- 30 Jalali, M.: Hydrochemical identification of groundwater resources and their changes under the impacts of human activity in the Chah Basin in western Iran, *Environ. Monit. Assess.*, 130, 347–364, 2007.
- Jones, B. F., Eugster, H. P., and Rettig, S. L.: Hydrochemistry of the Lake Magadi basin, Kenya,

## Hydrogeology and hydrogeochemistry of an alkaline volcanic

G. Ghiglieri et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Geochim. Cosmochim. Ac., 41, 53–72, 1977.

Kilham, P. and Hecky, R. E.: Fluoride: Geochemical and Ecological significance in East African waters and sediments, *Limnol. Oceanogr.*, November, 18(6), 932–945, 1973.

Nanyaro, J. T., Aswathanarayana, U., Mungure, J. S., and Lahermo, P. W.: A geochemical model for the abnormal fluoride concentrations in waters in parts of northern Tanzania, *J. African Earth Sci.*, 2, 129–140, 1984.

Nielsen, J. M.: East African magadi (trona): fluoride concentration and mineralogical composition, *J. Afr. Earth Sci.*, 29, 423–428, 1999.

Parkhurst, D. L. and Appelo, C. A. J.: User's guide to PHREEQC (version 2) – a computer program for speciation, batch-reaction, one-dimensional transport, and inverse geochemical calculations. United States Geological Survey, Water Resources Investigations Report 99–4259, Washington, p. 326, 1999.

Sebag, D., Verrecchia, E. P., Leed, S. J., and Durand, A.: The natural hydrous sodium silicates from the northern bank of Lake Chad: occurrence, petrology and genesis, *Sediment. Geol.*, 139(1), 15–31, 2001.

Signorelli, S., Vaggelli, G., and Romano, C.: Pre-eruptive volatile (H<sub>2</sub>O, F, Cl and S) contents of phonolitic magmas feeding the 3550-year old Avellino eruption from Vesuvius, southern Italy, *J. Volcanol. Geoth. Res.*, v. 93, 237–256, 1999.

Su, Y. H., Feng, Q., Zhu, G. F., Si, J. H., and Zhang, Y. W.: Identification and Evolution of Groundwater Chemistry in the Ejin Sub-Basin of the Heihe River, Northwest China, *Pe-dosphere*, 17(3), 331–342, ISSN 1002-0160, doi:10.1016/S1002-0160(07)60040-X, June 2007.

US DEPARTMENT OF ENERGY, Environmental Measurements Laboratory: Tritium in water – acid electrolysis – 3H – 01 – RC, HASL – 300, 28th Edition, 1997.



**Table 1.** Major ion composition of groundwater.

ID sample	T		pH		Conductivity		TDS		Ca <sup>2+</sup>		Mg <sup>2+</sup>		Na <sup>+</sup>		K <sup>+</sup>		HCO <sub>3</sub> <sup>-</sup>		Cl <sup>-</sup>
	Apr 2007	Jan 2008	Apr 2007	Jan 2008	Apr 2007	Jan 2008	Jan 2008	Apr 2007	Jan 2008	Apr 2007	Jan 2008	Apr 2007	Jan 2008	Apr 2007	Jan 2008	Apr 2007	Jan 2008	Apr 2007	
1old	14.6	14.4	8.0	8.4	750	790	603.08	1.15	0.72	0.27	0.26	4.26	4.78	0.92	0.64	6.14	5.86	18.05	
2old	23.6	n.a.	6.4	n.a.	210	n.a.	n.a.	0.41	n.a.	0.13	n.a.	1.07	n.a.	0.47	n.a.	1.62	n.a.	0.50	
3old	14.4	14.2	6.2	6.5	480	470	355.13	0.35	0.07	0.02	0.02	3.09	3.22	0.71	0.52	3.24	2.75	0.14	
4old	14.2	15.9	6.4	7.2	540	540	410.72	0.59	0.52	0.28	0.26	3.00	3.09	0.85	0.71	3.89	3.24	0.20	
5old	17.7	17.2	7.4	7.4	610	640	514.34	0.27	0.14	0.04	0.04	4.35	4.61	0.87	0.74	4.39	4.24	0.21	
6old	13.1	14.7	6.0	6.0	690	660	587.75	0.36	0.24	0.20	0.19	5.22	5.22	0.50	0.42	6.29	5.82	0.25	
8old	10.7	12.3	6.7	7.0	190	190	185.83	0.07	0.04	0.01	0.01	1.35	1.39	0.24	0.14	1.46	1.26	0.13	
10old	14.5	14.5	7.2	7.1	480	470	370.05	0.19	0.10	0.03	0.03	3.18	3.26	0.54	0.47	3.16	2.79	0.06	
13old	14.1	15.7	7.3	7.1	580	540	442.59	0.39	0.30	0.17	0.13	3.52	3.22	1.21	0.64	4.98	4.00	0.15	
16old	11.9	16.0	7.2	7.0	200	170	155.82	0.24	0.14	0.08	0.04	1.26	0.67	0.47	0.35	1.79	1.22	0.15	
1eng	18.2	15.4	7.2	7.1	620	600	513.39	0.67	0.41	0.23	0.41	5.22	4.35	0.59	0.42	5.15	4.91	0.07	
2eng	15.2	n.a.	5.9	n.a.	490	n.a.	n.a.	0.79	n.a.	0.42	n.a.	2.52	n.a.	0.59	n.a.	3.72	n.a.	0.14	
2beng	n.a.	15.5	n.a.	7.1	n.a.	770	632.89	n.a.	1.27	n.a.	0.35	n.a.	4.35	n.a.	0.89	n.a.	5.99	n.a.	
3eng	23.6	24.0	7.4	7.3	720	670	577.40	0.49	0.39	0.14	0.12	4.70	5.22	0.77	0.63	6.22	5.72	0.17	
22.1	20.5	7.0	6.7	980	650	502.14	0.75	0.37	0.31	0.16	8.26	3.91	1.10	0.63	8.25	4.53	0.62		
8eng	22.7	21.5	7.0	7.0	1220	1340	945.64	0.95	0.74	0.40	0.31	5.92	9.35	1.25	0.95	7.50	8.63	0.40	
16eng	17.0	16.5	6.4	6.3	340	340	285.53	0.13	0.08	0.04	0.04	2.52	2.39	0.27	0.23	2.55	2.30	0.15	
18eng	18.5	18.1	6.5	6.4	470	450	384.42	0.24	0.20	0.12	0.10	3.18	3.22	0.36	0.36	3.53	3.04	0.20	
19eng	16.9	16.2	7.4	7.2	400	390	333.91	0.21	0.14	0.05	0.05	2.78	2.70	0.45	0.38	3.16	2.77	0.18	
22eng	12.7	14.0	5.9	6.9	250	230	213.87	0.55	0.31	0.28	0.16	1.30	1.17	0.43	0.25	2.06	1.55	0.12	
24eng	18.3	17.7	7.2	7.0	1070	1170	961.22	1.30	2.05	0.49	0.48	7.83	7.18	1.20	0.95	9.83	9.93	0.08	
26eng	22.4	22.3	7.7	7.6	5070	4730	3927.60	0.40	0.67	0.27	0.94	47.85	35.67	1.20	4.60	35.12	36.60	5.35	
27eng	13.9	12.2	6.8	6.9	330	320	289.63	0.03	0.03	0.00	0.01	2.04	2.57	0.20	0.14	2.08	2.39	0.06	
28eng	17.8	17.6	7.6	7.7	1500	1160	896.46	0.27	0.24	0.12	0.10	9.05	8.48	2.51	1.20	9.53	7.88	0.84	
29eng	17.3	17.5	7.8	7.7	1390	1870	1537.35	0.08	0.04	0.02	0.05	7.92	15.01	1.69	2.35	7.96	14.04	0.70	
30eng	24.8	24.7	8.1	7.9	3740	3500	2808.44	0.60	0.34	0.29	0.24	26.75	30.45	3.43	2.25	18.72	22.29	2.59	
lchnusa well 1	n.a.	21.5	n.a.	6.4	n.a.	620	526.46	n.a.	0.72	n.a.	0.23	n.a.	3.65	n.a.	0.57	n.a.	5.15	n.a.	

\* with 0.0 mg l<sup>-1</sup> are represented the values out of instrument sensitivity.

Chemical composition of groundwater in the study area in April (Apr) 2007 and January (Jan) 2008; all concentrations are expressed in meq l<sup>-1</sup> except pH, conductivity (μS cm<sup>-1</sup>), temperature (°C) and RESIDUE (mg l<sup>-1</sup>); n.a. = not available.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

⏴

⏵

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Hydrogeology and hydrogeochemistry of an alkaline volcanic

G. Ghiglieri et al.

**Table 1.** Continued.

ID sample	Cl <sup>-</sup>		SO <sub>4</sub> <sup>2-</sup>		NO <sub>3</sub> <sup>-</sup>		NO <sub>2</sub> <sup>-</sup>		NH <sub>4</sub> <sup>+</sup>		F <sup>-</sup>		SiO <sub>2</sub>	RESIDUE 110°C	RESIDUE 180°C	FACIES
	Jan 2008	Apr 2007	Jan 2008	Apr 2007	Jan 2008	Apr 2007	Jan 2008	Apr 2007	Jan 2008	Apr 2007	Jan 2008	Apr 2007	Jan 2008	Jan 2008	Jan 2008	
1old	0.27	0.28	0.31	0.05	0.23	0.004	0.003	0.001	0.001	0.23	0.25	0.803	460.00	420.00	Na-HCO <sub>3</sub>	
2old	n.a.	0.17	n.a.	0.07	n.a.	0.016	n.a.	0.002	n.a.	0.05	n.a.	n.a.	n.a.	n.a.	Na-HCO <sub>3</sub>	
3old	0.18	0.21	0.24	0.17	0.18	0.005	0.002	0.012	0.001	0.65	0.63	0.826	260.00	220.00	Na-HCO <sub>3</sub>	
4old	0.20	0.22	0.23	0.34	0.44	0.007	0.006	0.002	0.001	0.13	0.10	0.879	340.00	340.00	Na-HCO <sub>3</sub>	
5old	0.23	0.28	0.31	0.02	0.30	0.006	0.004	0.001	0.001	0.68	1.05	0.909	360.00	340.00	Na-HCO <sub>3</sub>	
6old	0.13	0.12	0.14	0.02	0.08	0.004	0.001	0.001	0.001	0.13	0.16	1.144	380.00	360.00	Na-HCO <sub>3</sub>	
8old	0.06	0.04	0.04	0.02	0.03	0.005	0.002	0.007	0.001	0.24	0.27	0.848	80.00	80.00	Na-HCO <sub>3</sub>	
10old	0.17	0.23	0.23	0.04	0.20	0.005	0.003	0.005	0.001	0.93	1.05	0.902	240.00	220.00	Na-HCO <sub>3</sub> -F	
13old	0.13	0.13	0.13	0.11	0.28	0.005	0.004	0.002	0.001	0.21	0.22	0.977	360.00	340.00	Na-HCO <sub>3</sub>	
16old	0.05	0.05	0.04	0.02	0.20	0.007	0.003	0.003	0.001	0.08	0.11	0.492	140.00	80.00	NaK-HCO <sub>3</sub> -NO <sub>3</sub>	
1eng	0.11	0.18	0.19	0.26	0.27	0.005	0.004	0.001	0.001	0.16	0.16	0.848	380.00	380.00	Na-HCO <sub>3</sub>	
2eng	n.a.	0.15	n.a.	0.05	n.a.	0.004	n.a.	0.001	n.a.	0.07	n.a.	n.a.	n.a.	n.a.	Na-HCO <sub>3</sub>	
2beng	0.33	n.a.	0.24	n.a.	0.24	n.a.	0.003	n.a.	0.001	n.a.	0.09	0.970	460.00	440.00	Na-HCO <sub>3</sub>	
3eng	0.18	0.19	0.14	0.02	0.10	0.003	0.002	0.003	0.001	0.26	0.28	0.818	440.00	420.00	Na-HCO <sub>3</sub>	
5eng	0.26	0.34	0.33	0.05	0.18	0.015	0.002	0.004	0.001	0.28	0.26	0.985	360.00	320.00	Na-HCO <sub>3</sub>	
8eng	0.57	1.35	1.08	0.15	0.16	0.005	0.002	0.001	0.001	0.53	0.53	0.924	740.00	720.00	Na-HCO <sub>3</sub>	
16eng	0.15	0.16	0.17	0.02	0.16	0.003	0.002	0.001	0.001	0.28	0.31	0.811	220.00	160.00	Na-HCO <sub>3</sub>	
18eng	0.20	0.33	0.33	0.04	0.16	0.004	0.002	0.001	0.001	0.27	0.30	1.098	300.00	280.00	Na-HCO <sub>3</sub>	
19eng	0.21	0.11	0.13	0.05	0.20	0.005	0.003	0.002	0.001	0.18	0.22	0.894	200.00	200.00	Na-HCO <sub>3</sub>	
22eng	0.10	0.05	0.03	0.02	0.23	0.004	0.003	0.006	0.001	0.07	0.10	0.902	180.00	180.00	Na-HCO <sub>3</sub> -NO <sub>3</sub>	
24eng	0.08	0.23	0.27	0.31	0.42	0.006	0.006	0.007	0.001	0.37	0.38	0.939	700.00	660.00	Na-HCO <sub>3</sub>	
26eng	5.17	6.92	7.63	0.00	0.19	0.003	0.003	0.000	0.001	3.11	3.58	0.652	3140.00	3120.00	Na-HCO <sub>3</sub>	
27eng	0.08	0.05	0.09	0.01	0.16	0.004	0.002	0.000	0.001	0.20	0.24	0.917	220.00	180.00	Na-HCO <sub>3</sub>	
28eng	0.78	1.57	1.38	0.03	0.18	0.003	0.002	0.000	0.001	1.48	1.05	0.697	720.00	660.00	Na-HCO <sub>3</sub>	
29eng	1.31	1.14	2.30	0.00	0.14	0.008	0.002	0.002	0.001	0.90	1.20	0.886	1200.00	1120.00	Na-HCO <sub>3</sub>	
30eng	2.83	10.67	9.90	0.18	0.15	0.005	0.002	0.001	0.001	1.63	1.57	0.583	2380.00	2320.00	Na-HCO <sub>3</sub> -SO <sub>4</sub>	
lchnusa well 1	0.18	n.a.	0.15	n.a.	0.21	n.a.	0.003	n.a.	0.002	n.a.	0.16	0.977	420.00	320.00	Na-HCO <sub>3</sub>	

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Table 2.** Mineral composition of sediments and rocks samples.

	Paleo-soil Lahar (Nzd2) (Nzd2)	Lahar (Nzd2)	Lake deposits (l)	Crusts of Lake deposits (l) (l)	Lake deposits (l)	Crusts of Lake deposits (l) (l)	Breccias on Mantling Ash (Nvf) (Nvf)	Lahar (Nzd1)	Phonolite (Nvm)	Calcrete on Mantling Ash (Nvf) (Nvf)	Silt Layer in soil	Phonolite (Nvm)
	ENG6	ENG7	ENG8	ENG8b	2474 S	2474 C	OLD 3,4,5	OLD 10	ENG1	OLD9	OLD11	ENG31
Phillipsite		x	x	x	x	x		x				
Chabazite		x	x	x	x	x		x				
Analcime		x	x	x	x	x			x			
Nepheline	x	x	x	x	x	x	x	x	x		x	x
Leucite	x	x	x	x	x	x			tr			tr
Anorthoclase	x	x	x	x	x	x	x	x	x	x	x	x
Sanidine	x	x	x		x		x	x	x	x	x	x
Albite		x	x								x	
Riebeckite		x	x			x		x			x	x
Augite	x	x	x	x	x	x	x	x	x		x	x
Biotite		x	x	x		x	x					
Illite/smectite	x									x	x	
Trona				x		x						
Natron			tr									
Natrite	x		tr	x					x			
Calcite						x	x			x		
Cancrinite		x	x			tr		x				
Sylvite			x	x	x							
Fluorapatite		x	x		x	x		x	x			x
Aphthalite				x								
Goethite				x								

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion

**Table 3.** Isotopic water analysis and composition.

ID sample	Date	$\delta^{18}\text{O}\text{‰}$		$\delta\text{D}$		$^3\text{H}$	
		V-SMOW	V-SMOW	(U.T.)	+/- (U.T.)	(U.T.)	+/- (U.T.)
1old	19/02/2007	-5.29	-27.6	1.2	0.6		
2old	21/02/2007	-2.79	-8.3	2.6	0.7		
3old	21/02/2007	-6.39	-40.1	1.6	0.6		
4old	22/02/2007	-4.76	-24.4	1.7	0.6		
5old	22/02/2007	-6.54	-39.07	0.9	0.6		
6old	17/04/2007	-6.25	-34.5	0.4	0.3		
8old	17/04/2007	-5.47	-28.5	1.1	0.4		
10old	17/04/2007	-6.70	-39.5	0.8	0.3		
13old	16/04/2007	-5.40	-28.8	1.1	0.4		
16old	16/04/2007	-5.19	-27.5	2.2	0.5		
1eng	18/02/2007	-4.68	-23.3	1.1	0.6		
2eng	18/02/2007	-5.14	16.50	n.a.			
3eng	21/02/2007	-5.78	-35.5	1.1	0.6		
5eng	22/04/2007	-5.34	-27.3	1.0	0.4		
8eng	17/02/2007	-5.55	-33.9	0.8	0.6		
16eng	22/04/2007	-5.88	-31.2	0.7	0.4		
18eng	18/04/2007	-5.75	-30.2	0.8	0.3		
19eng	18/04/2007	-5.47	-27.8	1.0	0.4		
22eng	20/02/2007	-5.03	-26.5	2.4	0.7		
26eng	05/04/2007	-6.62	-35.3	1.5	0.5		
27eng	05/04/2007	-4.62	-23.9	2.2	0.5		
28eng	05/04/2007	-5.15	-24.5	1.7	0.5		
29eng	05/04/2007	-4.66	-29.9	2.2	0.5		
Ich. well 1	18/01/2008	-5.64	-29	1.5	0.6		
24 river	23/04/2007	-5.24	-25.4	1.5	0.4		
Big M.L.	17/02/2007	3.88	19.3	1.8	0.7		
Small M.L.	17/02/2007	2.65	13.5	2.1	0.7		
Rain	22/04/2007	-0.73	5.8	2.8	0.9		

## Hydrogeology and hydrogeochemistry of an alkaline volcanic

G. Ghiglieri et al.



**Fig. 1.** Study area and related hydrogeological map indicating the location of water points.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Hydrogeology and hydrogeochemistry of an alkaline volcanic

G. Ghiglieri et al.



**Fig. 2.** Ternary plot showing the study area groundwater composition (= freshwater in the *Masika* period; = freshwater in the *pre-Masika* period; = hydrothermal water in *Masika* period; = hydrothermal water in the *pre-Masika* period). The Ca-Na-K ternary plot is shown on the *left*, and the  $\text{HCO}_3\text{-SO}_4\text{-Cl}$  one on the *right*.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Hydrogeology and hydrogeochemistry of an alkaline volcanic

G. Ghiglieri et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper




Fig. 3. Sodium vs. fluoride scatter diagram of all groundwater samples (symbols as in Fig. 3).



## Hydrogeology and hydrogeochemistry of an alkaline volcanic

G. Ghiglieri et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper



**Fig. 4.** Bicarbonate vs. fluoride scatter diagram of all groundwater samples (symbols as in Fig. 3).






Fig. 5. Distribution of the major ions in river water samples.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶


Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 6.** Plot of  $^2\text{H}$  vs.  $^{18}\text{O}$  content; between bracket  $^3\text{H}$  values. The Local Meteoric Water Line (LMWL) was defined following Dettman et al. (2005).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Hydrogeology and hydrogeochemistry of an alkaline volcanic

G. Ghiglieri et al.



**Fig. 7.** Distribution map of the study area, indicating the location of isotopic water points classified according to their  $^3\text{H}$  content.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶


◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Fig. 8.** Cross section of the NW side of Mt. Meru, showing the location of the springs and the fluoride content of the spring waters.

Hydrogeology and hydrogeochemistry of an alkaline volcanic

G. Ghiglieri et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper





Fig. 9. Plot of total cation content vs. alkalinity (symbols as in Fig. 3).



**Fig. 10.** A Ca-Mg-alkalinity-SO<sub>4</sub> vs. Na-K-Cl scatter diagram illustrates the occurrence of reverse ion exchange. The dotted line represents the 1:1 ratio (symbols as in Fig. 3).

## Hydrogeology and hydrogeochemistry of an alkaline volcanic

G. Ghiglieri et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper




Fig. 11. Plot of fluorite saturation index vs. ionic strength (symbols as in Fig. 3).

Hydrogeology and hydrogeochemistry of an alkaline volcanic

G. Ghiglieri et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper





Fig. 12. Plot of fluoride vs. calcium in phonolite groundwater samples (symbols as in Fig. 3).



**Hydrogeology and  
hydrogeochemistry  
of an alkaline  
volcanic**

G. Ghiglieri et al.



**Fig. 13.** Plot of the saturation indices of several mineral phases vs. temperature (°C) for the water of spring 26 ENG.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Hydrogeology and hydrogeochemistry of an alkaline volcanic

G. Ghiglieri et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





Fig. 14. Plot of villiaumite saturation index vs. ionic strength (symbols as in Fig. 3).

## Hydrogeology and hydrogeochemistry of an alkaline volcanic

G. Ghiglieri et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Fig. 15.** A leaching test of a mantling ash sample conducted at ambient temperature.