

This discussion paper is/has been under review for the journal Hydrology and Earth System Sciences (HESS). Please refer to the corresponding final paper in HESS if available.

Transient drawdown solution for a constant pumping test in finite two-zone confined aquifers

C.-T. Wang^{1,2}, H.-D. Yeh², and C.-S. Tsai²

¹Taiwan Branch, MWH Americas Inc., Taipei, Taiwan

²Institute of Environmental Engineering, National Chiao Tung University, Hsinchu, Taiwan

Received: 25 August 2011 – Accepted: 20 September 2011 – Published: 18 October 2011

Correspondence to: H.-D. Yeh (hdyeh@mail.nctu.edu.tw)

Published by Copernicus Publications on behalf of the European Geosciences Union.

HESSD

8, 9299–9321, 2011

Transient drawdown
solution for a
constant pumping
test

C.-T. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

The drawdown solution has been widely used to analyze pumping test data for the determination of aquifer parameters when coupled with an optimization scheme. The solution can also be used to predict the drawdown due to pumping and design the dewatering system. The drawdown solution for flow toward a finite-radius well with a skin zone in a confined aquifer of infinite extent in radial direction had been developed before. To our best knowledge, the drawdown solution in confined aquifers of finite extent so far has never before been presented in the groundwater literature. This article presents a mathematical model for describing the drawdown distribution due to a constant-flux pumping from a finite-radius well with a skin zone in confined aquifers of finite extent. The analytical solution of the model is developed by applying the methods of Laplace transforms and Bromwich contour integral. This solution can be used to investigate the effects of finite boundary and conductivity ratio on the drawdown distribution. In addition, the inverse relationship between Laplace- and time-domain variables is used to develop the large time solution which can reduce to the Thiem solution if there is no skin zone.

1 Introduction

The famous Theis solution (1935) was first introduced in the groundwater literature to describe the transient drawdown distribution induced by a constant pumping at a well of infinitesimal well radius in a homogeneous and isotropic confined aquifer of infinite extent. The radius of a well is in fact not zero in the real-world problems. It is well recognized that the solution developed based on the assumption of zero well radius can not give accurate drawdown predictions near the wellbore. Van Everdingen and Hurst (1949) developed the transient pressure solutions for the constant flow in finite and infinite confined reservoirs with considering the effect of well radius but neglecting the skin effect. Note that the term skin effect is used to reflect the increase or decrease of hydraulic conductivity caused by drilling practices in a region near the well.

Transient drawdown solution for a constant pumping test

C.-T. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



unconfined aquifer of infinite extent with a partially penetration well, finite-thickness skin. Yet, they adopted an approach such as a finite difference method to discretize the well screen for handling non-uniform wellbore flux problems.

The existing drawdown solutions for radial two-zone confined aquifers of infinite extent under constant-flux pumping were all developed in Laplace domain except the one given by Yeh et al. (2003) which was a time domain solution. Yeh et al.'s solution (2003) is in terms of an improper integral integrating from zero to infinity and its integrand comprises a singularity at the origin. In addition, the integrand is an oscillatory function with many product terms of the Bessel functions of the first and second kinds of zero and first orders. The numerical calculation of their solution is therefore time-consuming and very difficult to achieve accurate results. To avoid such numerical difficulties, we therefore propose an alternative analytical solution developed from a mathematical model similar to that of Yeh et al. (2003), except that the aquifer is of horizontally finite extent. The solution of the model is also developed by applying the methods of Laplace transforms and Bromwich contour integral. The integration of the contour integral in Yeh et al. (2003) results in a single branch point with no singularity at zero of the complex variable. Thus, a branch cut along the negative real axis of the contour should be chosen and thus a closed contour is produced. Such a procedure finally results in a complicated solution presented in Yeh et al. (2003). On the other hand, the integration of the contour integral arisen from the Laplace domain solution in our model has a simple pole at the origin and finite number of poles at other locations. The residue theorem is therefore adapted to obtain the time domain solutions for the skin zone or formation zone. These two solutions are in terms of a logarithmic function plus a summation term, rather than an integral, with the Bessel functions of the first and second kinds of orders zero and first. The solutions are much easier to calculate than those of Yeh et al. (2003) indeed involving a singularity in the integral. In addition, large-time solutions in simpler forms are also developed by employing the relationship of small Laplace-domain variable p versus large time-domain variable t , hereinafter referred to SPLT (Yeh and Wang, 2007), to the Laplace-domain solution.

Transient drawdown solution for a constant pumping test

C.-T. Wang et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
⏪	⏩
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



2 Mathematical model

2.1 Mathematical statement

The assumptions involved in the development of the mathematical model are: (1) the confined aquifer is homogeneous, isotropic, and of finite extent in radial direction; (2) the well fully penetrates the aquifer and has a finite well radius; (3) a skin zone is present around the pumping well as shown in Fig. 1; (4) the well discharge rate is maintained constant through out the entire pumping test.

The governing equations describing the drawdown distribution $s(r, t)$ in the skin zone and formation zone are, respectively,

$$\frac{\partial^2 s_1}{\partial r^2} + \frac{1}{r} \frac{\partial s_1}{\partial r} = \frac{S_1}{T_1} \frac{\partial s_1}{\partial t} \quad r_w \leq r < r_1 \quad (1)$$

$$\frac{\partial^2 s_2}{\partial r^2} + \frac{1}{r} \frac{\partial s_2}{\partial r} = \frac{S_2}{T_2} \frac{\partial s_2}{\partial t} \quad r_1 \leq r < R \quad (2)$$

where subscripts 1 and 2 express the skin zone and the formation zone, respectively, r is the radial distance from the central line of the pumping well, r_w is the well radius, r_1 is the outer radius of the skin zone, R is the distance from the central line of the well to the outer boundary, t is the pumping time, S is the storage coefficient, and T is the transmissivity.

Prior to pumping, there is no drawdown over the area that will be influenced by the test. Thus, the initial conditions for skin zone and formation can be written as

$$s_1(r, 0) = s_2(r, 0) = 0 \quad (3)$$

In addition, the drawdown at R is also considered to be zero. The flux along the well-bore is maintained at a constant rate Q . Thus, the outer and inner boundary conditions can be expressed, respectively, as

$$s_2(R, t) = 0 \quad (4)$$

Transient drawdown solution for a constant pumping test

C.-T. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$$\varphi = \frac{I_0(q_2 R) K_0(q_2 r) - I_0(q_2 r) K_0(q_2 R)}{I_0(q_2 R) K_0(q_2 r_1) - I_0(q_2 r_1) K_0(q_2 R)} \quad (13)$$

Note that p is the Laplace variable, $q_1 = \sqrt{pS_1/T_1}$, $q_2 = \sqrt{pS_2/T_2}$, I_0 and K_0 are the modified Bessel functions of the first and second kinds of order zero, respectively, and I_1 and K_1 are the modified Bessel functions of the first and second kinds of order first, respectively.

The values of γ and ω can reduce to $K_1(q_2 r_1)/K_0(q_2 r_1)$ and $K_0(q_2 r)/K_0(q_2 r_1)$, respectively, when R approaches infinity. Equations (8) and (9) are then equivalent to the solutions presented in Yeh et al. (2003, p. 750) as

$$\bar{s}_1 = \frac{Q}{4\pi T_2} \left[\frac{1}{p r_w T_1 q_1} \frac{\psi_2 K_0(q_1 r) + \psi_1 I_0(q_1 r)}{\psi_2 K_1(q_1 r_w) - \psi_1 I_1(q_1 r_w)} \right] \quad (14)$$

$$\bar{s}_2 = \frac{Q}{4\pi T_2} \left[\frac{1}{p r_w T_1 q_1} \frac{(\psi_2 K_0(q_1 r_1) + \psi_1 I_0(q_1 r_1)) K_0(q_2 r)}{(\psi_2 K_1(q_1 r_w) - \psi_1 I_1(q_1 r_w)) K_0(q_2 r_1)} \right] \quad (15)$$

with

$$\psi_1 = K_1(q_1 r_1) K_0(q_2 r_1) - \sqrt{\frac{S_2 T_2}{S_1 T_1}} K_0(q_1 r_1) K_1(q_2 r_1) \quad (16)$$

$$\psi_2 = I_1(q_1 r_1) K_0(q_2 r_1) + \sqrt{\frac{S_2 T_2}{S_1 T_1}} I_0(q_1 r_1) K_1(q_2 r_1) \quad (17)$$

2.3 Time-domain solution

The transient drawdown solution in time domain can be obtained by applying the Bromwich contour integral (Carslaw and Jaeger, 1959, p. 332) to the Laplace domain

Transient drawdown solution for a constant pumping test

C.-T. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

solution. Detailed development is shown in Appendix A and the result for the drawdown distribution in skin and formation zones is, respectively,

$$s_1 = \frac{Q}{2\pi T_1} \left\{ \ln \frac{r_1}{r} + \frac{T_1}{T_2} \ln \frac{R}{r_1} - \frac{\pi}{r_w} \sum_{n=1}^{\infty} \exp\left(-\frac{T_1}{S_1} \alpha_n^2 t\right) \cdot \frac{\alpha_n (J_1(\alpha_n r_w) Y_0(\alpha_n r) - Y_1(\alpha_n r_w) J_0(\alpha_n r))}{S_n^2 [B_n^2 + (\zeta (B_n C_n + A_n D_n) + \zeta A_n B_n / \alpha_n) / r_1 + \zeta^2 A_n^2] - \alpha_n^2} \right\} \quad (18)$$

5 and

$$s_2 = \frac{Q}{2\pi T_2} \left\{ \ln \frac{R}{r} - \frac{\pi}{r_w} \sum_{n=1}^{\infty} \exp\left(-\frac{T_1}{S_1} \alpha_n^2 t\right) \cdot \frac{\alpha_n (J_1(\alpha_n r_w) Y_0(\alpha_n r_1) - Y_1(\alpha_n r_1) J_0(\alpha_n r_w)) \cdot (Y_0(\xi \alpha_n R) J_0(\xi \alpha_n r) - Y_0(\xi \alpha_n r) J_0(\xi \alpha_n R))}{[S_n^2 B_n [B_n^2 + \zeta (B_n C_n + A_n D_n) / r_1 + \zeta A_n B_n / \alpha_n + \zeta^2 A_n^2] - \alpha_n^2]} \right\} \quad (19)$$

with

$$S_n = \frac{-\alpha_n J_1(\alpha_n r_w)}{-\zeta A_n J_0(\alpha_n r) - B_n J_1(\alpha_n r)} \quad (20)$$

$$A_n = [J_1(\xi \alpha_n r_1) Y_0(\xi \alpha_n R) - J_0(\xi \alpha_n R) Y_1(\xi \alpha_n r_1)] \quad (21)$$

$$10 B_n = J_0(\xi \alpha_n R) Y_0(\xi \alpha_n r_1) - J_0(\xi \alpha_n r_1) Y_0(\xi \alpha_n R) \quad (22)$$

$$C_n = -\xi R [J_1(\xi \alpha_n r_1) Y_1(\xi \alpha_n R) - J_1(\xi \alpha_n R) Y_1(\xi \alpha_n r_1)] - \xi r_1 B_n - \frac{A_n}{\alpha_n} \quad (23)$$

$$D_n = \xi R [J_1(\xi \alpha_n R) Y_0(\xi \alpha_n r_1) - J_0(\xi \alpha_n r_1) Y_1(\xi \alpha_n R)] - \xi r_1 A_n \quad (24)$$

Transient drawdown solution for a constant pumping test

C.-T. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

where J_0 and Y_0 are the Bessel functions of the first and second kinds of order zero, respectively, J_1 and Y_1 are the Bessel functions of the first and second kinds of order first, respectively, $\xi = \sqrt{T_1 S_2 / T_2 S_1}$, $\zeta = \sqrt{S_2 T_2 / S_1 T_1}$, and $\pm \alpha_n$ are the roots of

$$\left[\alpha J_1(\xi \alpha r_1) Y_0(\xi \alpha R) - \alpha J_0(\xi \alpha R) Y_1(\xi \alpha r_1) \right] \cdot \zeta \left[Y_1(\alpha r_w) J_0(\alpha r_1) - Y_0(\alpha r_1) J_1(\alpha r_w) \right] + \left[J_0(\xi \alpha r_1) Y_0(\xi \alpha R) - J_0(\xi \alpha R) Y_0(\xi \alpha r_1) \right] \cdot \left[\alpha Y_1(\alpha r_1) J_1(\alpha r_w) - \alpha J_1(\alpha r_1) Y_1(\alpha r_w) \right] = 0 \quad (25)$$

5 When ignoring the presence of skin zone, Eq. (19) can reduce to

$$s = \frac{Q}{2\pi T} \left\{ \ln \frac{R}{r} - \frac{\pi}{w} \sum_{n=1}^{\infty} \exp\left(-\frac{T}{S} \alpha_n^2 t\right) \cdot \frac{(J_1(\alpha_n r_w) Y_0(\alpha_n R) - Y_1(\alpha_n r_w) J_0(\alpha_n R))}{\alpha_n [J_1^2(\alpha_n r_w) - J_0^2(\alpha_n R)] / J_0^2(\alpha_n R)} \right\} \quad (26)$$

where α_n are the roots of $J_1(\alpha r_w) Y_0(\alpha R) - Y_1(\alpha r_w) J_0(\alpha R) = 0$. Note that Eq. (26) is exactly the same as the equation presented in Wang and Yeh (2008, Eq. 11). Additionally, the steady-state solution can be obtained from Eqs. (18) and (19) in the limit of time approaching infinity.

2.4 Large-time solution

The drawdown solution for two-zone confined aquifers of finite-extent at large times can be obtained by applying the SPLT technique and L'Hospital rule to Eqs. (8) and (9). Some limits of the Bessel functions with small arguments are given as $I_0(x) \sim 1$, $I_1(x) \sim x/2$, $K_0(x) \sim -\ln(x)$, and $K_1(x) \sim 1/x$ when x approaches zero (Abramowitz and Stegun, 1979, p. 375). The Laplace-domain drawdown distribution in skin zone and formation zone for small ρ can then be obtained, respectively, as

$$\overline{s_1}(r, \rho) = \frac{Q}{2\pi T_1 \rho} \left(\ln \frac{r_1}{r} + \frac{T_1}{T_2} \ln \frac{R}{r_1} \right) \quad (27)$$

$$\overline{s_2}(r, \rho) = \frac{Q}{2\pi T_2 \rho} \ln \frac{R}{r} \quad (28)$$

Transient drawdown solution for a constant pumping test

C.-T. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Accordingly, the large-time drawdown solutions can then be obtained by taking the inverse Laplace transform to Eqs. (27) and (28) as

$$s_1(r,t) = \frac{Q}{2\pi T_1} \left(\ln \frac{r_1}{r} + \frac{T_1}{T_2} \ln \frac{R}{r_1} \right) \quad (29)$$

$$s_2(r,t) = \frac{Q}{2\pi T_2} \ln \frac{R}{r} \quad (30)$$

- 5 The time invariant Eqs. (29) and (30) can also be obtained by applying the method of Tauberian theory (Sneddon, 1972) to Eqs. (8) and (9). This result indicates that the drawdown solution can reach its steady-state solution in confined aquifers of finite extent as declared in Wang and Yeh (2008). In addition, both Eqs. (29) and (30) can reduce to the Thiem solution if there is no skin zone, i.e. r_1 equals r_w and T_1 equals T_2 .

10 2.5 Dimensionless solution

Dimensionless variables are introduced as follows: $\kappa = T_2/T_1$, $\gamma = S_2/S_1$, $\tau = T_2 t / S_2 r_w^2$, $\rho = r/r_w$, $\rho_1 = r_1/r_w$, $\rho_R = R/r_w$, and $s_D = s(4\pi T_2)/Q$ where κ represents conductivity ratio, γ represents the ratio of storage coefficient, ρ represents dimensionless distance, ρ_1 represents dimensionless skin thickness, ρ_R represents dimensionless distance of the outer boundary, and s_D represents the transient distribution of dimensionless drawdown. The drawdown solutions in Eqs. (18) and (19) then becomes

$$s_1 = 2\kappa \left\{ \ln \frac{\rho_1}{\rho} + \frac{1}{\kappa} \ln \frac{\rho_R}{\rho_1} - \pi \sum_{n=1}^{\infty} \exp \left(-\frac{\gamma}{\kappa} \beta_n^2 \tau \right) \cdot \frac{\omega_n (J_1(\beta_n) Y_0(\beta_n \rho) - Y_1(\beta_n) J_0(\beta_n \rho))}{s_{Dn}^2 \left[b_n^2 + (\zeta (b_n c_n + a_n d_n) + \zeta a_n b_n / \beta_n) / \rho_1 + \zeta^2 a_n^2 \right] - \beta_n^2} \right\} \quad (31)$$

Transient drawdown solution for a constant pumping test

C.-T. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$$s_2 = 2K \left\{ \ln \frac{\rho_R}{\rho} - \pi \sum_{n=1}^{\infty} \exp \left(-\frac{\gamma}{\kappa} \beta_n^2 \tau \right) \right. \\ \left. \frac{\beta_n (J_1(\beta_n) Y_0(\beta_n \rho_1) - Y_1(\beta_n \rho_1) J_0(\beta_n)) \cdot (Y_0(\xi \beta_n \rho_R) J_0(\xi \beta_n \rho) - Y_0(\xi \beta_n \rho) J_0(\xi \beta_n \rho_R))}{[\zeta_{Dn}^2 [b_n^2 + (\zeta(b_n c_n + a_n d_n) + \zeta a_n b_n / \beta_n) / \rho_1 + \zeta^2 a_n^2] - \beta_n^2]} \right\} \quad (32)$$

where $\beta_n = r_w \alpha_n$ are the roots of

$$\left[\beta J_1(\xi \beta \rho_1) Y_0(\xi \beta \rho_R) - \beta J_0(\xi \beta \rho_R) Y_1(\xi \beta \rho_1) \right] \cdot \zeta \left[Y_1(\beta) J_0(\beta \rho_1) - Y_0(\beta \rho_1) J_1(\beta) \right] + \\ \left[J_0(\xi \beta \rho_1) Y_0(\xi \beta \rho_R) - J_0(\xi \beta \rho_R) Y_0(\xi \beta \rho_1) \right] \cdot \left[\beta Y_1(\beta \rho_1) J_1(\beta) - \beta J_1(\beta \rho_1) Y_1(\beta) \right] = 0 \quad (33)$$

and

$$s_{Dn} = \frac{-\beta_n J_1(\beta_n)}{-\zeta a_n J_0(\beta_n \rho_1) - b_n J_1(\beta_n \rho_1)} \quad (34)$$

$$a_n = J_1(\xi \beta_n \rho_1) Y_0(\xi \beta_n \rho_R) - J_0(\xi \beta_n \rho_R) Y_1(\xi \beta_n \rho_1) \quad (35)$$

$$b_n = J_0(\xi \beta_n \rho_R) Y_0(\xi \beta_n \rho_1) - J_0(\xi \beta_n \rho_1) Y_0(\xi \beta_n \rho_R) \quad (36)$$

$$c_n = -\xi \rho_R [J_1(\xi \beta_n \rho_1) Y_1(\xi \beta_n \rho_R) - J_1(\xi \beta_n \rho_R) Y_1(\xi \beta_n \rho_1)] - \xi \rho_1 b_n - \frac{a_n}{\beta_n} \quad (37)$$

$$d_n = \xi \rho_R [J_1(\xi \beta_n \rho_R) Y_0(\xi \beta_n \rho_1) - J_0(\xi \beta_n \rho_1) Y_1(\xi \beta_n \rho_R)] - \xi \rho_1 a_n \quad (38)$$

The numerical calculations for Eqs. (31) and (32) are achieved by finding the roots of Eq. (33) first using Newton's method and then adding the summation term for n up to 100. Generally, the results have accuracy to the fifth decimal place.

3 Advantages and applications of the present solution

3.1 Advantages over the existing solutions

The analytical solution developed herein has the following two advantages over Yeh et al.'s solution (2003). First, the present solutions can give the same predicted drawdowns as Yeh et al.' solution (2003) if the outer boundary distance in the present solution is very large. In other words, the solutions presented in Yeh et al. (2003) can be considered as a special case of the present solution. Second, the transient drawdown solution in Yeh et al. (2003) is given in terms of an improper integral integrating from zero to infinity and its integrand has a singularity at the origin. Due to the presence of singular point, the solution is very difficult to accurately calculate. In contrast, the present solution is composed of infinite series and can be easily calculated with accuracy to fifth decimal.

3.2 Potential applications

Due to the presence of skin zone, an aquifer system is characterized by five parameters, i.e. the outer radius of the skin zone and the transmissivity and storage coefficient for each of the skin and aquifer zones. Once the parameters are known, the presented solution can be used to predict the spatial or temporal drawdown distributions in both the skin and formation zones and explore the physical insight of the constant-flux test in two-zone aquifer systems. On the other hand, those five parameters can be determined via the data analyses if the parameter values are not available. The determination of unknown parameters is in fact a subject of inverse problems. Type-curve approach is common for the parameter estimation. However, it is almost impossible to develop type-curves for the parameter estimation because the unknown parameters of two-zone aquifers are too many. Alternative ways in determining those five unknown parameters are to use the presented solution in conjunction with the algorithm of extended Kalman filter (e.g. Leng and Yeh, 2003; Yeh and Huang, 2005) or a heuristic

Transient drawdown solution for a constant pumping test

C.-T. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the time becomes very large. On the other hand, the wellbore drawdown in infinite aquifers continuously increases with dimensionless time. Figure 2 also demonstrates the effect of skin property on the wellbore drawdown distribution. The aquifer with a positive skin has larger wellbore drawdowns than the one with a negative skin at the same pumping rate. Figure 3 presents the drawdown distributions at $\rho = 10$ for $\rho_1 = 3$ and $\kappa = 0.1, 1, \text{ and } 10$. It reveals that the drawdown in an aquifer with a negative skin is larger than that in the one with a positive skin. In other words, the effect of skin property on the drawdown distribution in the formation zone is opposed to that at the wellbore. The difference in drawdown distribution between aquifers with $\kappa = 1$ and 10 at $\rho = 1$ shown in Fig. 2 is significantly larger than those with $\kappa = 0.1$ and 1 at $\rho = 1$ (Fig. 2) and those with $\kappa = 1$ and 10 at $\rho = 10$ shown in Fig. 3, indicating that the two-zone aquifer system is rather sensitive to the pumping in positive skin cases.

5 Conclusions

A mathematical model has been developed to describe the drawdown distribution for a pumping test performed in a two-zone confined aquifer of finite extent. The Laplace-domain solution of the model for skin zone and formation zone is obtained by applying the method of Laplace transforms. The analytical solution (in time domain) is then developed by the Bromwich contour integral method. The drawdown distribution predicted from the analytical solution shows that the dimensionless drawdown distribution in a finite aquifer is significantly different from that in an infinite one at large pumping times. In other word, the drawdown solution of finite aquifer is applicable to predict that of an infinite aquifer only under the condition that the time is not large enough. In addition, the two-zone aquifer system is rather sensitive to the pumping in positive skin cases.

A large-time drawdown solution is also developed in this article based on the inverse relationship of Laplace- and time-domain variables. Surprisingly, this large-time solution is exactly the same as the steady-state solution, indicating that the solution of

Transient drawdown solution for a constant pumping test

C.-T. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



finite two-zone aquifers can reach its steady-state solution at large times. In addition, the large-time solution has been shown to reduce to the Thiem solution if neglecting the presence of skin zone.

Appendix A

5 Derivation of Eqs. (18) and (19)

The drawdown solution in time domain, denoted as $s(t)$ obtained by applying the Bromwich integral method (Carslaw and Jaeger, 1959) to the Laplace domain solution $\bar{s}(p)$ is expressed as

$$10 \quad s(t) = L^{-1} \{ \bar{s}(p) \} = \frac{1}{2\pi i} \int_{r-i\infty}^{r+i\infty} e^{pt} \bar{s}(p) dp \quad (A1)$$

where i is an imaginary unit and r_e is a real constant which is so large that all of the real parts of the poles are smaller than it. The graph of the Bromwich integral contains a close contour with a semicircle and a straight line parallel to the imaginary axis. According to Jordan's Lemma, the integration for the semicircle tends to be zero when
15 it radius approaches infinity. Based on the residue theorem, Eq. (A1) can be written as

$$s(t) = \sum_{n=1}^{\infty} \text{Res} \{ e^{pt} s(p); g_n \} \quad (A2)$$

where p_n are the poles in the complex plane. There are infinite singularities in $s(p)$ and obviously one pole at $p = 0$.

Introducing the following two variables

$$20 \quad \Delta = q_1 [\Phi_1 I_1(q_1 r_w) + \Phi_2 K_1(q_1 r_w)] \quad (A3)$$

$$\Psi = [\Phi_1 I_0(q_1 r) - \Phi_2 K_0(q_1 r)] \quad (A4)$$

Transient drawdown solution for a constant pumping test

C.-T. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Equation (8) can then be expressed as

$$\bar{s}_1 = \frac{-Q}{4\pi T_2} \frac{2T_2}{r_w T_1} \left[\frac{1}{\rho} \frac{\Psi(\rho)}{\Delta(\rho)} \right] \quad (\text{A5})$$

Let $\Delta = 0$, the roots α_n in $\rho = \rho_n = (-T_1 \alpha_n^2)/S_1$ can be then determined from Eq. (A3). Substituting $\rho = \rho_n = (-T_1 \alpha_n^2)/S_1$ into Eq. (A3) yields Eq. (25). From the following formula (Kreyszig, 1999), the residue of the pole at $\rho = 0$ is

$$\text{Res}\{e^{pt} s(\rho); 0\} = \lim_{\rho \rightarrow 0} s(\rho) e^{pt} (\rho - 0) \quad (\text{A6})$$

Substituting Eq. (A5) into Eq. (A6) and applying L'Hopital's rule results in

$$\text{Res}\{e^{pt} s(\rho); 0\} = \frac{Q}{2\pi T_1} \left[\ln \frac{r_1}{r} + \frac{T_1}{T_2} \ln \frac{R}{r_1} \right] \quad (\text{A7})$$

The other residues at the simple pole $\rho = \rho_n = -T_1 \alpha_n^2/S_1$ are expressed as

$$\text{Res}\{e^{pt} s(\rho); \rho_n\} = \lim_{\rho \rightarrow \rho_n} s(\rho) e^{pt} (\rho - \rho_n) \quad (\text{A8})$$

Applying L'Hopital's rule to Eq. (A8), the denominator term inside the brackets of Eq. (A5) becomes

$$\left[\rho \frac{d\Delta}{d\rho} \right]_{\rho = -T_1 \alpha_n^2/S_1} = \left[\frac{1}{2} q \frac{d\Delta}{dq} \right]_{q_1 = i\alpha_n, q_2 = i\kappa\alpha_n} = \frac{1}{2} q_1 \left\{ q_1 \left[\Phi_1' I_1(q_1 r_w) + \Phi_2' K_1(q_1 r_w) \right] + r_w q_1 \left[\Phi_1' I_0(q_1 r_w) - \Phi_2' K_0(q_1 r_w) \right] \right\} \quad (\text{A9})$$

where the variables Φ_1 and Φ_2 are defined in Eqs. (10) and (11), respectively, and Φ_1' and Φ_2' are the first differentiations of Φ_1 and Φ_2 , respectively.

To simplify Eq. (A8), a variable ζ_n is assumed based on Eq. (A3) and $\Delta = 0$:

$$\begin{aligned}\zeta_n &= \frac{q_1 I_1(q_1 r_w) K_0(\xi q_1 r_1)}{-\Phi_2 [(I_0(\xi q_1 R) K_0(\xi q_1 r_1) - I_0(\xi q_1 r_1) K_0(\xi q_1 R))]} \\ &= \frac{q_1 K_1(q_1 r_w) K_0(\xi q_1 r_1)}{\Phi_1 [(I_0(\xi q_1 R) K_0(\xi q_1 r_1) - I_0(\xi q_1 r_1) K_0(\xi q_1 R))]} \quad (A10)\end{aligned}$$

Two recurrence formulas (Carslaw and Jaeger, 1959, p. 490) are adopted to eliminate the imaginary unit in Eq. (8) as follows:

$$K_\nu(z e^{\pm \frac{1}{2} \pi i}) = \pm \frac{1}{2} \pi i e^{\mp \frac{1}{2} \nu \pi i} [-J_\nu(z) \pm i Y_\nu(z)] \quad (A11)$$

and

$$I_\nu(z e^{\pm \frac{1}{2} \pi i}) = e^{\pm \frac{1}{2} \nu \pi i} J_\nu(z) \quad (A12)$$

Substituting Eqs. (A11) and (A12) into Eq. (A10) yields

$$\frac{-\alpha_n J_1(\alpha_n r_w)}{-\zeta A_n J_0(\alpha_n r_1) - B_n J_1(\alpha_n r_1)} = \frac{\alpha_n Y_1(\alpha_n r_w)}{\zeta A_n Y_0(\alpha_n r_1) + B_n Y_1(\alpha_n r_1)} = \zeta_n \quad (A13)$$

The result of substituting Eq. (A13) into Eq. (A9) is

$$\begin{aligned}\left[p \frac{d\Delta}{dp} \right]_{p=-T_1 \alpha_n^2 / S_1} &= \left[\frac{1}{2} q \frac{d\Delta}{dq} \right]_{q_1 = i \alpha_n, q_2 = i \kappa \alpha_n} = \\ &= \frac{1}{2 \zeta_n} \left\{ \zeta_n^2 \left[B_n^2 + (\zeta (B_n C_n + A_n D_n) + \zeta A_n B_n / \alpha_n) / r_1 + \zeta^2 A_n^2 \right] - \alpha_n^2 \right\}\end{aligned} \quad (A14)$$

where the constants appeared on the right-hand side of Eq. (A14) are defined in Eqs. (20)–(24). Similarly, the numerator of Eq. (A5) can also be obtained as

$$\Psi = \frac{\pi \alpha_n}{2 \zeta_n} [J_1(\alpha_n r_w) Y_0(\alpha_n r) - Y_1(\alpha_n r_w) J_0(\alpha_n r)] \quad (A15)$$

The residues at the poles $p = p_n = -T_1 \alpha_n^2 / S_1$ are

$$\text{Res}\{e^{pt} s(p); p_n\} = \frac{-Q}{2\pi r_w T_1} \sum_{n=1}^{\infty} \exp\left(\frac{-\alpha_n^2 t T_1}{S_1}\right) \left\{ \frac{\alpha_n (J_1(\alpha_n r_w) Y_0(\alpha_n r) - Y_1(\alpha_n r_w) J_0(\alpha_n r))}{\xi_n^2 \left[(\xi A_n)^2 + B_n^2 + \xi ((B_n C_n + A_n D_n) + A_n B_n / \alpha_n) / r_1 \right] - \alpha_n} \right\} \quad (\text{A16})$$

Therefore, Eq. (A2) can be expressed as

$$h(t) = (\text{Res}\{e^{pt} s(p); 0\} + \text{Res}\{e^{pt} s(p); p_n\}) \quad (\text{A17})$$

Finally, the solution for the drawdown distribution in the skin zone can then be obtained as Eq. (18). The solution for the drawdown distribution in the formation zone can also be obtained in a similar way as Eq. (19).

Acknowledgements. This research is partially supported by the Taiwan National Science Council under the contract numbers NSC 99-NU-E-009-001, NSC 99-2221-E-009-062-MY3, and NSC 100-2221-E-009-106.

References

- Abramowitz, M. and Stegun, I. A.: Handbook of Mathematical Functions, Dover publications, New York, 1970.
- Baker, J. A. and Herbert, R.: Pumping test in patchy aquifer, Ground Water, 20, 150–155, 1988.
- Barry D. A., Parlange, J. Y., and Li, L.: Approximation for the exponential integral (Theis well function), J. Hydrol., 227, 287–291, 2000.
- Batu, V.: Aquifer Hydraulics, John Wiley & Sons, New York, 1998.
- Butler Jr., J. J.: Pumping tests in nonuniform aquifers-The radially symmetric case, J. Hydrol., 101, 15–30, 1988.

Transient drawdown solution for a constant pumping test

C.-T. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Transient drawdown solution for a constant pumping test

C.-T. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Butler Jr., J. J. and Liu, W. Z.: Pumping tests in nonuniform aquifers: The radially asymmetric case, *Water Resour. Res.*, 29, 259–269, 1993.
- Carlsaw, H. S. and Jaeger, J. C.: *Conduction of Heat in Solids*, 2nd edn., Oxford University Press, London, 1959.
- 5 Chen, C. S.: A reinvestigation of the analytical solution for drawdown distributions in a finite confined aquifer, *Water Resour. Res.*, 20, 1466–1468, doi:10.1029/WR020i010p01466, 1984.
- Hantush, M. S.: *Hydraulics of wells*, in: *Advances in Hydroscience*, Academic Press, New York, 1964.
- Huang, Y. C. and Yeh, H. D.: The use of sensitivity analysis in on-line aquifer parameter estimation, *J. Hydrol.*, 335, 406–418, 2007.
- 10 Ingersoll, L. R., Adler, F. T. W., Plass, H. J., and Ingersoll, A. G.: *Theory of earth heat exchangers for the heat pump*, HPAC Engineering, 113–122, 1950.
- Ingersoll, L. R., Zobel, O. J., and Ingersoll, A. C.: *Heat conduction with engineering, geological, and other applications*, 2nd edn., Madison, University Wisconsin Press, 1954.
- 15 Kreyszig, E.: *Advanced engineering mathematics*, John Wiley & Sons Inc., New York, 1999.
- Leng, C. H. and Yeh, H. D.: Aquifer parameter identification using the extended Kalman filter, *Water Resour. Res.*, 39, 1062, doi:10.1029/2001WR000840, 2003.
- Lin, Y. C. and Yeh, H. D.: Trihalomethane species forecast using optimization method: genetic algorithm and simulated annealing, *J. Comput. Civil Eng.*, 19, 248–257, 2005.
- 20 Liou, T. S. and Yeh, H. D.: Conditional expectation for evaluation of risk groundwater flow and solute transport: one-dimensional analysis, *J. Hydrol.*, 199, 378–402, 1997.
- Novakowski, K. S.: A Composite Analytical Model for Analysis of Pumping Tests Affected by Well Bore Storage and Finite Thickness Skin, *Water Resour. Res.*, 25, 1937–1946, 1989.
- Perina, T. and Lee, T. C.: General well function for pumping from a confined, leaky, or unconfined aquifer, *J. Hydrol.*, 317, 239–260, 2006.
- 25 Prodanoff, J. H. A., Mansur, W. J., and Mascarenhas, F. C. B.: Numerical evaluation of Theis and Hantush-Jacob well functions, *J. Hydrol.*, 318, 173–183, 2006.
- Sneddon, I. N.: *The Use of Integral Transforms*, New York, McGraw-Hill, 1972.
- Streltsova, T. D.: *Well Testing in Heterogeneous Formations*, John Wiley and Sons, Inc., New York, 45–49, 1988.
- 30 Theis, C. V.: The relation between the lowering of the piezometric surface and the rate and duration of discharge of a well using ground-water storage: *Trans American Geophysical Union*, 16, 519–524, 1935.

Transient drawdown solution for a constant pumping test

C.-T. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Tseng, P. H. and Lee, T. C.: Numerical evaluation of exponential integral: Theis well function approximation, *J. Hydrol.*, 281, 129–146, 1998.
- Van Everdingen, A. F. and Hurst, W.: The application of the Laplace transformation to flow problems in reservoirs, *Pet. Trans. AIME*, 186, 305–324, 1949.
- 5 Wang, C. T. and Yeh, H. D.: Obtaining the steady-state drawdown solutions of constant-head and constant-flux tests, *Hydrol. Process.*, 22, 3456–3461, 2008.
- Yeh, H. D. and Huang, Y. C.: Parameter estimation for leaky aquifers using the extended Kalman filter and considering model and data measurement uncertainties, *J. Hydrol.*, 302, 28–45, 2005.
- 10 Yeh, H. D. and Wang, C. T.: Large-time solutions for groundwater flow problems using the relationship of small p versus large t , *Water Resour. Res.*, 43, W06502, doi:10.1029/2006WR005472, 2007.
- Yeh, H. D., Chang, T. H., and Lin, Y. C.: Groundwater contaminant source identification by a hybrid heuristic approach, *Water Resour. Res.*, 43, W09420, doi:10.1029/2005WR004731, 2007.
- 15 Yeh, H. D., Yang, S. Y., and Peng, H. Y.: A new closed-form solution for a radial two-layer drawdown equation for groundwater under constant-flux pumping in a finite-radius well, *Adv. Water Resour.*, 26, 747–757, 2003.




Fig. 1. Schematic diagram of the pumping test in a finite-extent confined aquifer.

Transient drawdown solution for a constant pumping test

C.-T. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 2. The drawdown curves predicted at $\rho = 1$ (wellbore) for infinite aquifers denoted by the solid line and for finite aquifers having the outer boundary distances of 20, 30, and 50 represented by the dashed lines with the symbols of circle, diamond, and square, respectively, for $\rho_1 = 3$ and $\kappa = 0.1, 1$ and 10 .

Transient drawdown solution for a constant pumping test

C.-T. Wang et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	






Fig. 3. The drawdown curves predicted at $\rho = 10$ (in formation zone) for infinite aquifers denoted by the solid line and for finite aquifers having the outer boundary distances of 20, 30, and 50 represented by the dashed lines with the symbols of circle, diamond, and square, respectively, for $\rho_1 = 3$ and $\kappa = 0.1, 1$ and 10 .

Transient drawdown solution for a constant pumping test

C.-T. Wang et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

