

Interactive comment on “Land classification based on hydrological landscape units” by S. Gharari et al.

S. Gharari et al.

shervangharari@yahoo.com

Received and published: 10 August 2011

Reply to Comment by A. D. Nobre

Reply to brief history on development of HAND and A new landscape classification: the HAND Model

We appreciate the information provided by the reviewer on the background of the development of HAND. Although several scientists implicitly have been using this concept for quite a long time (e.g., Uhlenbrook et al., 2002; McGuire et al., 2005), they did not formalize it, i.e. they did not develop a unique algorithm for calculating

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Height Above Nearest Drainage (HAND). The reviewer claimed to have developed HAND, but he did not mention how we could understand this from the published literature, including Rennó et al. (2008). Although the author mentioned that Rennó et al. (2008) aimed at the remote sensing community and Nobre et al. (2011) refers to the hydrological aspects, HAND was clearly introduced the first time by Rennó et al. (2008). If HAND was already introduced in any previous work, we would kindly ask the reviewer to inform us so that we can appropriately refer to it.

We also note that a published paper is not a patent, and everybody can make use of concepts and ideas present in any published work. Our intention has not been to claim ourselves as owners of the concept. We just picked it up to investigate hydrological implications. The way to prevent these things from happening is simply not to publish.

References

McGuire, K. J., McDonnell, J. J., Weiler, M., Kendall, C., McGlynn, B. L., Welker, J. M., and Seibert, J.: The role of topography on catchment-scale water residence time, *Water Resources Research*, 41, W05 002, 10.1029/2004WR003657, 2005.

Nobre, A. D., Cuartas, L. A., Hodnett, M., Rennó, C. D., Rodrigues, G., Silveira, A., Waterloo, M., and Saleska, S.: Height above the Nearest Drainage, a hydrologically relevant new terrain model, *Journal of Hydrology*, 404(1-2), 13–29, 2011.

Rennó, C. D., Nobre, A. D., Cuartas, L. A., Soares, J. V., Hodnett, M. G., Tomasella, J., and Waterloo, M. J.: HAND, a new terrain descriptor using SRTM-DEM: Mapping terra-firme rainforest environments in Amazonia, *Remote Sensing of Environment*, 112(9), 3469–3481, 10.1016/j.rse.2008.03.018, 2008.

Uhlenbrook, S., Didszun, J., Tilch, N., McDonnell, J., and McGuire, K.: Breaking up is always difficult – Landscape discretization as a process-transfer approach for prediction in ungauged basins, *PUB-meeting-Brasilia*, 2002.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

Interactive comment on Hydrol. Earth Syst. Sci. Discuss., 8, 4381, 2011.

HESD

8, C3380–C3382, 2011

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

C3382

