

## ***Interactive comment on “Soil weathering rates in 21 catchments of the Canadian Shield” by D. Houle et al.***

**Anonymous Referee #1**

Received and published: 10 August 2011

The release of base cations (BC) from soil weathering to the soil solution and surface waters is an important issue considering acidification of soils and surface waters. The weathering related BC release maintains the natural buffering capacity of terrestrial and adjacent aquatic ecosystems up to a certain degree.

The here evaluated model PROFILE has been widely used to evaluate soil weathering related BC release and thus the natural buffering capacity mainly in boreal landscapes (in North America and Scandinavia) with rather poor soils (mainly shallow podsols) and softwater lakes.

The study by Houle et al. (the MS commented here) applies the PROFILE model to a rather large set of soil profiles (21 forested lake catchments with 3 profiles each) lo-

C3386

cated on the Canadian Shield within a 150 km wide strip parallel to the St. Lawrence River. For the parameterization of this model they apply several other models: the Uppsala model estimating mineralogical composition of a given soil horizon based on the bulk chemistry assessed by lab methods; the BioSIM model to generate monthly precipitation and temperature for each catchment from gauging data; FORHYM to simulate soil moisture, and FORSTEM to simulate soil temperature based on the BioSIM output.

Assuming that lake chemistry is determined by weathering within the soil profiles and the three soil profiles analyzed per lake catchment are representative for the whole catchment, Houle et al. evaluate the PROFILE-output for each base cation and the sum of BC by comparison with measured lake water concentrations. Further, they used the lake water concentration and FORHYM model output to estimate the stream export of BC from the catchments, assuming BC concentrations at the stream outlets equals that of the lakes.

They conclude that the PROFILE model output represents Ca stream exports quite well, while that of Mg is on average overestimated by about 50 %, and that of K (factor 6.9) and Na (factor 2.2) are even more overestimated. Further, they conclude that PROFILE is strong enough to reproduce geographical gradients in the weathering rates.

The presented research issue fits into the scope of HESS. However, before the manuscript can be considered for publications, I would like to suggest some major revisions to be done.

*\*General comments\**

First of all, I suggest to present the importance of that particular study in more detail:

-Why is this study important?

-What can the reader learn from this study? As this is mainly a technical paper: How

C3387

does this study help to improve existing methods for application in research and/or environmental survey programs?

-Why was this methodology chosen?

-What is the advantage of the here applied method compared to other approaches? I suggest to write a specific subsection on this issue in the result & discussion part.

-At the end of the conclusion the authors should present an outlook. What are important research questions left open?

As the PROFILE model has already been evaluated in previous studies (e.g. Kolka et al., 1996; Hodson et al., 1997; Ouimet and Duchesne, 2005; Whitfield et al., 2006) it might be more appropriate to evaluate the complete methodology presented in the MS, i.e. including the parameterization of PROFILE with the outputs of the UPPSALA, FORHYM, and FORSTEM models.

\*Comments on method section\*

The information on the study sites given in the MS is rather short. I suggest giving detailed information on each lake catchment as electronic supplement. This should include a more detailed description of the lithology, of the glacial till as well as of the bedrock. How thick is the till layer? Further, a description of the hydrologic setting would be of interest for the reader. Does every lake have a stream outflow? Are streams the major drainage ways from the lake catchments, or are groundwater exports from the catchments of higher importance? What is the mean groundwater table at the sites where the soil samples have been taken?

Further, the average hydrochemical properties for each lake should be listed in a table within the supplement. In the electronic supplement, a description of each soil profile should be given as well.

One of the parameters required by the PROFILE model is the DOC concentration within each soil horizon. Unfortunately, it is unclear how this information was derived for the

C3388

here presented study. Are the applied DOC values based on own measurements or based on literature values assumed to be representative?

In the method section it is said that for the clay fraction of the analyzed soil samples, mineralogy is assessed by X-ray diffraction. It is said that the results were used to validate the estimation of the mineralogy using the UPPSALA model. Unfortunately, the respective results are not given in the MS. I suggest to give these results as supplement information. The reader could get an idea how well the UPPSALA model really predicts the mineralogy, at least for the clay fraction.

\*Comments on results & discussion\*

In the MS, bulk chemistry and estimated mineralogical compositions are just given for the B-horizons of the soil profiles. It is argued that most of the BC release happens there, but it is not shown with data. For this, I suggest giving this information for the A, E, and C horizons as well, maybe in the electronic supplement.

The authors describe a spatial gradient and state that their estimation of BC release reproduces these gradients quite well. I suggest producing two small maps showing this spatial gradient: one map giving the spatial patterns of lake chemistry (BC-concentrations, observed data) and another map giving the spatial patterns of estimated BC release from soil weathering.

In the MS, studies comparing the PROFILE model with other approaches to assess BC release from catchments by weathering have been cited (Kolka et al., 1996; Ouimet and Duchesne, 2005; Whitfield et al., 2006), but yet not sufficiently discussed. Whitfield et al. (2006) generally compared approaches focusing on weathering within individual soil profiles, incl. the PROFILE model, and approaches addressing weathering within whole catchments. They pointed out two main problems related to the assessment of catchment weathering by approaches focusing on weathering rates within given soil profiles, like the PROFILE model does: 1) The chosen soil profiles might not be representative for the whole catchment, 2) Substantial weathering related BC release might

C3389

take place within the substrate below the developed soil horizons, which approaches like PROFILE model are neglecting. I strongly suggest to take these two points into account when discussing the results of this studies and associated uncertainties.

By assuming BC release is just happening within the soil profile it is implicated that weathering in the till/saprolite below the B-horizon would be negligible. If the authors want to stick to this assumption they should justify it.

When discussing other approaches, I suggest including the WITCH model (Probst et al., 2002; Godderis et al., 2006).

The authors stated that the ecosystems on the study sites are obviously not in a steady state. They used this as a probable explanation for the overestimation of K-exports. Is there a probable reason for this, e.g. a specific land use history. Are the forests growing (increasing biomass)? Is there any clue on this besides the K-flux overestimation?

There is still the problem of the high overestimation of Na fluxes. Is this related to wrong predictions by the UPPSALA model? I suggest stating some hypothesis why Na-fluxes are overestimated? Might this be due to wrongly estimated mineralogical composition by UPPSALA model or by wrongly estimated weathering rates of certain minerals by PROFILE model? Which minerals are of interest in this respect? I suggest to discuss such possibilities for K as well.

Spodic horizons of podsoles (B-horizons) are characterized by enrichment in amorphous organic substances and sesquioxides, forming coatings on the mineral grains. It is of interest in how far these coatings might influence the weathering rates of the minerals in these horizons. I suggest discussing this issue as well. Is this a source of uncertainty in the PROFILE model?

\*Other comments\*

p. 5745, line 10-13: cite Garrels and Mackenzie (1971), or earlier works of these authors

C3390

p. 5752, line 11-13: Especially as the authors refer to the B-Horizon of a podsol, I suggest to write that this mineralogical composition is typical for soils developed in such lithologies, rather than “are typical for Precambrian Shield geologies”.

Tables 1 & 2: In the captions, please indicate that it is weight-% to prevent any confusion.

Table 2: It is “Feldspar”, not “Feldspath”

Table 2: Authors distinguish albite from plagioclase. However, albite is a plagioclase. Please, correct this error.

In the MS ,base cations (BC) is used for the sum of Ca, Mg, K while Na is excluded. If the authors want to stick to this definition, I suggest giving an explanation why BC does not incorporate Na in this study.

Table 3: Here, BC is used as the sum of Ca, Mg, K, Na. This is in contrast to the rest of the MS, for which BC was defined as Ca+Mg+K, excluding Na.

p. 5751, line 26: “conservative” instead of “conservator”

\*References\*

Garrels, R.M. and Mackenzie, F.T., 1971. Evolution of Sedimentary Rocks. W.W. Norton, New York.

Godderis, Y., Francois, L.M., Probst, A., Schott, J., Moncoulon, D., Labat, D. and Viville, D., 2006. Modelling weathering processes at the catchment scale: The WITCH numerical model. *Geochimica Et Cosmochimica Acta*, 70(5): 1128-1147.

Hodson, M.E., Langan, S.J. and Wilson, M.J., 1997. A critical evaluation of the use of the PROFILE model in calculating mineral weathering rates. *Water Air and Soil Pollution*, 98(1-2): 79-104.

Kolka, R.K., Grigal, D.F. and Nater, E.A., 1996. Forest soil mineral weathering rates:

C3391

Use of multiple approaches. *Geoderma*, 73(1-2): 1-21.

Ouimet, R. and Duchesne, L., 2005. Base cation mineral weathering and total release rates from soils in three calibrated forest watersheds on the Canadian Boreal Shield. *Canadian Journal of Soil Science*, 85(2): 245-260.

Probst, A., Godderis, Y., Francois, L.M., Labat, D., Schott, J. and Viville, D., 2002. Modelling chemical weathering at river catchment scale: design and calibration of the WiTCh model. *Geochimica Et Cosmochimica Acta*, 66(15A): A615-A615.

Whitfield, C.J., Watmough, S.A., Aherne, J. and Dillon, P.J., 2006. A comparison of weathering rates for acid-sensitive catchments in Nova Scotia, Canada and their impact on critical load calculations. *Geoderma*, 136(3-4): 899-911.

---

Interactive comment on Hydrol. Earth Syst. Sci. Discuss., 8, 5743, 2011.