



## Abstract

Global land surface models (LSMs) such as the Joint UK Land Environment Simulator (JULES) are originally developed to provide surface boundary conditions for climate models. They are increasingly used for hydrological simulation, for instance to simulate the impacts of land-use changes and other perturbations on the water cycle. This study investigates how well such models represent the major hydrological fluxes at the relevant spatial and temporal scales – an important question for reliable model applications in poorly understood, data-scarce environments. The JULES-LSM is implemented in a 360 000 km<sup>2</sup> humid tropical mountain basin of the Peruvian Andes–Amazon at 12 km grid resolution, forced with daily satellite and climate reanalysis data. The simulations are evaluated using conventional discharge-based evaluation methods, and by further comparing the magnitude and internal variability of the basin surface fluxes such as evapotranspiration, throughfall, and surface and subsurface runoff, of the model with those observed in similar environments elsewhere. We find reasonably positive model efficiencies and high correlations between the simulated and observed streamflows, but high root-mean-square errors affecting the performance in smaller, upper sub-basins. We attribute this to errors in the water balance and JULES-LSM's inability to model baseflow. We also found a tendency to underrepresent the high evapotranspiration rates of the region. We conclude that strategies to improve the representation of tropical systems to be (1) addressing errors in the forcing (2) incorporating local wetland and regional floodplain in the subsurface representation.

## 1 Introduction

The humid tropics hosts extremely biodiverse ecosystems, which are subject to changing climate and land use patterns and potentially changing water cycles. With an area of approximately 6 million km<sup>2</sup> (Latrubesse et al., 2005), Amazonia hosts a significant part of the world's remaining rainforest and is an important supplier of atmospheric

**HESSD**

9, 12523–12561, 2012

### JULES-LSM hydrology for humid tropical environments

Z. Zulkafli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



moisture (Salati and Vose, 1984). Many hydrological studies have continued to focus on the continental and lower Amazon (e.g. Vörösmarty et al., 1989; D'Almeida et al., 2006; Paiva et al., 2011; Guimberteau et al., 2012), while the upper Andes–Amazon system receives far less attention despite being subject to increasing human impacts such as deforestation, oil exploitation, mining, and hydropower production. The potential impact of a changing climate and land use on the hydrological regime of the Amazon headwaters is a serious concern, not only because of its influence on the downstream basin, but also because of its link to local ecosystem services.

Hydrological models are a common approach to understanding and predicting the impact of change, but modellers of tropical environments face issues with process understanding and data availability for parameterization and validation of models (Giertz et al., 2006). Models for tropical basins in the literature tend to be conceptual models requiring few inputs (e.g. Darko, 2002; Campling et al., 2002; Bormann and Diekkrüger, 2004), but depending strongly on local calibration. This can be complicated by the low availability and high degree of uncertainty introduced by the streamflow data. This is especially true for the Amazon where the river morphology continuously change due to active erosion in the uplands (see Aalto et al., 2006). Another disadvantage of conceptual models is the unidentifiability of their parameters (Ebel and Loague, 2006), which complicates scenario analysis.

Distributed physics-based modelling has been attempted in smaller catchments with some success (Vertessy and Elsenbeer, 1999; Legesse et al., 2003; Bekoe, 2005). More recently, land surface models (LSMs) have been used for physics-based hydrological modelling at global scale (e.g. Arora and Boer, 2003; Alkama et al., 2011) and the continental scale Amazon (Decharme and Douville, 2006; Guimberteau et al., 2012). LSMs, also referred to in the literature as land surface schemes (LSS) and land surface parameterizations (LSP), were originally developed by the climate modelling community to provide the land-atmospheric boundary condition in operational weather forecasting and global climate simulations. An LSM operates in continuous time and fully-distributed mode, as it simulates the exchanges of energy, water and

## JULES-LSM hydrology for humid tropical environments

Z. Zulkafli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



carbon between the land surface and the atmosphere by accounting for processes in the ground and vegetation canopy. A particular strength of LSMs is the biophysical consistency between various modelled processes such as photosynthesis, carbon and nutrient cycles, irrigation, and crop growth. This provides an opportunity for hydrologists to study the impact of change on hydrology in interaction with the other land surface processes.

However, despite their sophistication, LSMs are newcomers to hydrological modelling and while there have been point-scale validation exercises in flux tower sites (Baldocchi et al., 2001) representing various environments (e.g. Blyth et al., 2011), none has thoroughly assessed their performance in tropical upland basins. First, there is the issue with model structure that stems from universalization of locally-observed processes. Such a simplifying assumption may not hold in topographically complex environments where there may be multiple interacting factors controlling the hydrological response. Additionally, there are established weaknesses in the LSMs even for temperate climates that may be more problematic over the humid tropics. For instance, LSMs are often criticized for an absence of a groundwater model and therefore unsuitability for basins with a shallow water table as it assumes free gravity drainage from the soil column (Yeh and Eltahir, 2005). Moreover, lateral runoff processes reported in the humid tropical literature (Dunne, 1978; Campling et al., 2002; Giertz et al., 2006; Chapell, 2010) such as saturation excess surface runoff, interflows in the organic layer, and natural pipes are seldom modelled explicitly due to the LSM vertical structure.

Secondly, parameter uncertainty is certainly an issue. In comparison with the temperate regions, the tropics has lagged in soil characterization with respect to runoff producing mechanisms (Giertz and Diekkrüger, 2003). In this context, an LSM offers an advantage in that its worldwide application means global datasets of soil and vegetation data are available. However, while these exist, most are not field based. For example, the soil parameters are derived from physical soil maps using pedotransfer functions (PTF), which are a product of regression analyses of soil water retention data of temperate soils (e.g. Cosby et al., 1984). Although the PTF for tropical soils

## JULES-LSM hydrology for humid tropical environments

Z. Zulkafli et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

(Tomasella and Hodnett, 1998; Tomasella et al., 2000; Hodnett and Tomasella, 2002) are better alternatives, they still lack representativeness for the upper Amazon Basins' younger volcanic soils.

The number of parameters required per LSM pixel certainly introduces issues of data collection and scaling. Consequently, the majority are effective parameters that are not well constrained and applied homogeneously in space. This may not be a serious issue in land-surface modelling, which is mainly concerned with representing fluxes at large spatial scales. However, it becomes problematic for hydrological modelling, where local heterogeneity and non-linear responses to perturbations need to be simulated, and especially in mountainous environment where a high degree of heterogeneity may be expected over a small scale.

In spite of the limitations, LSMs are increasingly used to simulate hydrological fluxes such as discharge at basin and global scales. This paper studies the implications for modelling humid tropical upland basins. The case study is the upper Amazon River (Marañón River) Basin in Peru which hosts half of the Pacaya Samiria National Reserve, the largest floodable forest reserve in the Peruvian Amazonia, where the hydrological system provides important ecosystem services for unique species of fish and freshwater turtles that are vulnerable to extinction, as well as approximately 10 000 km<sup>2</sup> aguaje (*Mauritia flexuosa*) palm forests of high economic importance (Kahn, 1988).

Our approach moves beyond the traditional model evaluation using strictly observational data to decide whether a model is acceptable. Instead, our aim is to evaluate a “fit for purpose” simulation system that “shadows” the natural system as closely as possible (Beven et al., 2012). This does not necessarily mean that internally all fluxes are correct at the pixel scale and at each time step, which is infeasible given the lack of observations. Rather, we intend to present a simulation that most closely resembles the reality in terms of the main statistical properties (especially the mean and variation of the basin's hydrological fluxes), under the assumption that such a system will be the most robust in representing the impact of perturbations at a basin level.

## JULES-LSM hydrology for humid tropical environments

Z. Zulkafli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 2 Materials and methods

### 2.1 The JULES land surface model

A full description of the model JULES can be found in Best et al. (2011), while a brief description is provided for completeness. Land surfaces are modelled as tiles consisting of vegetated and non-vegetated surfaces with distinct parameters for radiation balance, resistance to heat and momentum transfer, canopy interception, plant photosynthesis, respiration and growth, etc. Land-atmospheric heat and moisture exchanges for each grid are calculated by area-weighted averaging of the tile fluxes, and these are exchanged with a shared soil column.

The LSM requires time series of meteorological data i.e. incoming short wave and long wave radiation, temperature, specific humidity, wind speed, and surface pressure. These are used in a full energy balance equation that includes components of radiation, sensible heat, latent heat, canopy heat, and ground surface heat. The potential evaporation estimation is based on the Penman–Monteith (Penman, 1948) approach. Canopy evaporation is assumed to occur at the potential rate, while plant transpiration and bare soil evaporation are restricted by canopy resistance and the soil moisture state respectively.

In JULES, the local throughfall rate is proportional to the local precipitation rate by the fraction of occupied canopy storage,  $C/C_{\max}$ , where  $C_{\max}$  is a vegetation parameter and is a linear function of the leaf-area-index (LAI). On the ground surface, throughfall is partitioned into surface runoff and infiltration into the soil moisture pool based on the Hortonian infiltration excess mechanism, enhanced by a vegetation-specific factor to account for macroporosity in the soil.

In the subsurface, an instantaneous redistribution of moisture is assumed and water is exchanged between the soil layers using a finite difference approximation to the Darcy–Richards diffusion equation, with infiltration and gravity drainage as the upper and lower boundaries respectively, and root uptake as a sink. The soil water retention characteristics follow the model of Brooks and Corey (1964) or the alternative van

**HESSD**

9, 12523–12561, 2012

### JULES-LSM hydrology for humid tropical environments

Z. Zulkafli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Genuchten (1980) formulation. In the alternate soil hydrology model of JULES (Clark and Gedney, 2008), a grid-based implementation of TOPMODEL calculates the local saturation excess runoff based on a time-moving surface partial contributing area. In this configuration, the model applies an exponential decay to the soil hydraulic conductivity with depth, assumes a null flux lower boundary, and applies an anisotropic factor to generate lateral flows for the subsurface.

For the study basin, we evaluated both the basic JULES (JULES-BASE) and the JULES-TOPMODEL parameterisations in distributed mode. The study basin is divided into pixels of 0.125° latitude-longitude (~ 14 km) resolution. Each pixel is assigned a set of soil parameters, the distribution of the land cover types, and time-series of meteorological variables from global datasets. The model goes through a warming up period to initialise the internal states.

## 2.2 Runoff routing

The runoff generated by JULES consists of local surface and subsurface runoff that need to be routed for a meaningful assessment against streamflow measurement data. Our model is simple delay function, with the delay for each pixel being the distance between the pixel and the outlet divided by the flood wave velocity ( $C$ ). The flood wave velocity for the surface and subsurface runoff are the two parameters of the model that are optimized through a Monte-Carlo simulation.

The lag time  $t$  to the outlet from pixel  $i$  will vary by its distance  $d$  to the outlet along the stream network:

$$t_{i_1} = \frac{d_i}{C_1} \quad (1)$$

$$t_{i_2} = \frac{d_i}{C_2} \quad (2)$$

### JULES-LSM hydrology for humid tropical environments

Z. Zulkafli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where the subscripts 1 and 2 represent surface and subsurface components respectively. Finally, the simulated flow at the outlet ( $Q_{sim}$ ) is the sum of all contributing local hydrographs in the basin, lagged in time.

$$Q_{sim,t} = \sum_{i=1}^n (Q_{i_1(t-t_{i_1})} + Q_{i_2(t-t_{i_2})}) \quad (3)$$

## 2.3 Data

### 2.3.1 Study area

The study area is the Marañón (Peruvian Amazon) River Basin upstream of the confluence with the Ucayali River that forms the proper Amazon River (Fig. 1). On average, the yearly discharge in the Marañón River is  $14\,900\text{ m}^3\text{ s}^{-1}$ , as measured at San Regis station ( $74^\circ\text{ W}$ ,  $4.4^\circ\text{ S}$ ) (Espinoza-Villar et al., 2009a). The climate of the region has been discussed extensively by various works, see for example Kvist and Nebel (2001), Garreaud et al. (2009) and Espinoza-Villar et al. (2009b). The average yearly temperature in the Marañón River Basin is about  $23\text{--}27^\circ\text{ C}$ , and the austral winter (JJF) is warmer than austral summer (DJF), during which cloud formation and rainfall effectively reduces the temperature (Kvist and Nebel, 2001; Garreaud et al., 2009). Precipitation characteristics vary across latitudes as influenced by synoptic meteorological phenomena operating at varying time scales (see Table 1). Precipitation is also orographically controlled, with the wettest band found at an altitude of 1.3 ma.s.l. on the eastern face of the Andes (Bookhagen and Strecker, 2008). The Andean ranges, with elevations exceeding 4000 m are effective hydrometeorological controls separating their west and east and connecting regions at lower and higher latitudes (Garreaud et al., 2009).

The basin is underlain by silty Gleysols in the river floodplains, and young unstructured Cambisols in the Andean foothills and Leptosols and Regosols further upland (FAO, 2009). The tropical wet climate supports expansive lowland, montane, and floodable forests as well as wet grassland (locally known as páramos) on the Andes. In

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the southwest, shielding by the Eastern Andean range generates a drier climate and grassland above the tree line.

### 2.3.2 Land surface data

The watershed boundary for the study basin is delineated based on HydroSHEDS 90 m resolution hydrographic dataset (Lehner et al., 2008). A gridded map of mean and standard deviation of the topographic indices for TOPMODEL was also generated from this dataset.

A new set of land surface parameters governing the soil hydraulics were created using the pedotransfer functions developed by Tomasella and Hodnett (1998). These functions were derived using a database from the Brazilian Amazon and are considered the best available pedotransfer function for tropical soil. The functions take values of silt and clay fraction in the soil, which are available through the Harmonized World Soil Database (FAO, 2009) at 1 km resolution (FAO, 2009). The HWSD sources the ISRIC-World Soil Information Soil and Terrain (SOTER), which in South America is considered to have highest reliability (FAO, 2009). The global soil textural map provides information for the topsoil (top 30 cm) and subsoil that was assigned to the top 2 (total depth of 35 cm) and bottom 2 (total depth of 265 cm) layers in JULES.

The land cover was parameterized based on multiple land cover maps. The primary source is the Digital Ecological Systems Map of the Amazon Basin of Peru and Bolivia (Josse et al., 2007b). This is a 90 m resolution field-verified mapping based on satellite imagery. In areas outside of this map's coverage, the 1 km IGBP-DIS Land Cover Classification Map of Loveland et al. (2000) was used.

The model default vegetation parameters (see Tables 5 and 6 in Best et al., 2011) are applied uniformly in space with the exception of canopy height and LAI, whose spatial distribution were obtained from the UK Met Office version 7.7 Central Ancillary Program and are based on satellite derived normalized difference vegetation indices

## JULES-LSM hydrology for humid tropical environments

Z. Zulkafli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(NDVI). Additionally, canopy heights of broadtree leaves were mapped from the local digital ecological systems database (Josse et al., 2007a).

### 2.3.3 Meteorological data

The global land surface model driving data developed by Sheffield et al. (2006) were used and included meteorological variables such as long and short-wave radiation, temperature, pressure, specific humidity, wind, and precipitation. The dataset is the first generation NCEP (US National Center of Environmental Predictions Kalnay et al., 1996) climate reanalysis product merged with ground data (the data will be henceforth referred to as NCEP). The dataset at 1.0° latitude-longitude (111 km) grids were further disaggregated to 0.125° latitude-longitude (14 km) grids using the lapse rate interpolation method described in the same paper. The nearest neighbour interpolation method was used to disaggregate precipitation. An alternate precipitation dataset from the TRMM 3B42 (version 6, 0.25° resolution, Huffman et al., 2007) remote sensing product was bias-corrected with TRMM 2A25 climatology (0.1° resolution) of Nesbitt and Anders (2009) (the data will be henceforth referred to as TRMM). The simulation of JULES was performed over the entire basin for a period of 11 yr between 1998 and 2008 to coincidence with the periods of available data from NCEP and the TRMM precipitation.

### 2.3.4 Streamflow data

Daily streamflow data for four hydrological stations at San Regis, Borja, Santiago, and Chazuta, were obtained through HYBAM (geodynamical, HYdrological and Biogeochemical control of erosion alteration and material transport in the AMazon Basin) from the Servicio Nacional de Meteorología e Hidrología, Peru (SENAMHI) and the Nacional de Meteorología e Hidrología, Ecuador (INAMHI) monitoring networks (for

## JULES-LSM hydrology for humid tropical environments

Z. Zulkafli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



station locations and synthesis, refer to Fig. 1 and Table 2). Missing days from the time series were excluded from the analysis, and the 95th percentile lower and upper uncertainty bounds (*L.U.B.* and *U.U.B.*) were calculated adapting the method in Daren-Harmel and Smith (2007).

$$5 \quad L.U.B._t = Q_{\text{obs},t}(1 - \text{PER}/100) \quad (4)$$

$$U.U.B._t = Q_{\text{obs},t}(1 + \text{PER}/100) \quad (5)$$

For this study, the probable error ranges (PER) were estimated using the standard deviation of the errors between the gauged and field-measured discharges during the calibration campaigns. The water balance closure is assessed and 4 additional stations, i.e. Paute, Nueva Loja, Nuevo Rocafuerte, and San Sebastian from INAMHI are included for comparative analysis. The last three stations are located in the Napo River Basin in Ecuador, which is tributary to the Peruvian Amazon further downstream of San Regis.

## 15 2.4 Model evaluation

Streamflow simulations were assessed with the Nash Sutcliffe model efficiency (NSE), the root mean square error (RMSE), the relative bias and the Pearson correlation. The calculated deviations were modified using approach 1 described by (Daren-Harmel and Smith, 2007) to account for streamflow data uncertainty. The entire time series available from the modelling period of 1998–2008 were used.

The internal model fluxes (i.e. evapotranspiration, canopy throughfall, surface runoff, and subsurface runoff) are evaluated by calculating the statistics of the spatial variability over each major biomes – lowland forest and flood forest (below 1200 m a.s.l.), montane forest (between 1200 and 3500 m a.s.l.), and upland (above 3500 m a.s.l.) systems as shown in Fig. 2 – within the entire basin. In the absence of a dense network of local observations, the distribution of observations from the literature (Table 4) are taken as substitute for observations from a “real” system. The assumption is that the best

**JULES-LSM  
hydrology for humid  
tropical environments**

Z. Zulkafli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



simulation for the basin will produce similar natural variability, assessed in terms of the mean and spread of the distributions.

### 3 Results and discussion

#### 3.1 Uncertainties in the observations

Potential errors in the observations of precipitation and streamflow were assessed by analysing the runoff ratio, i.e. the total observed streamflow to the total precipitation input (Table 3). Ratios typical for humid tropical environments are in the range of 0.6–0.7 (Campling et al., 2002; Buytaert et al., 2006b; Rollenbeck and Anhuf, 2007), while the values found in the study basins exceed 0.80, with values up to 1.76. This suggests severe errors in either the streamflow measurements or precipitation products, or unaccounted sources of water. For the latter, cloud water input may be responsible, as this can be significant in montane forests can be significant, ranging between 22–1990 mm $\text{yr}^{-1}$  (see a comprehensive study by Bruijnzeel et al., 2011). However, a seasonality analysis of the water balance (results not shown) highlights that the largest overestimations of the runoff ratio occur during the austral winter. This is incompatible with cloud water input, which would occur during periods of persistent cloud cover (Zadroga, 1981) which in the case of the Mara $\tilde{\text{n}}$  Basin is the austral summer.

Previous studies that highlighted difficulties with water balance closure in similar magnitudes have attributed these to errors in precipitation because of the scarcity of gauge data for the upper basin (i.e. upstream of Brazil) (Guimberteau et al., 2012; Coe, 2002). Indeed, large errors have been reported in the literature with the TRMM and NCEP precipitation over the Amazon Basin. Ward et al. (2011) observed in Paute Basin both datasets underestimating precipitation during the dry season by 50 mm $\text{month}^{-1}$  on average. In the Brazilian Amazonia, Clarke et al. (2010) observed a negative bias in the maximum annual daily rainfall of up to 80 mm $\text{day}^{-1}$  in the NCEP reanalysis data,

## JULES-LSM hydrology for humid tropical environments

Z. Zulkafli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and an extension of the dry season when compared to gauged data. Both studies also found poor correlation to the gauged time series.

The most unrealistic runoff ratios were found at the Northern Andean Basins (Paute, Santiago, Nueva Loja, and San Sebastian, Table 3). Large, lowland basins such as Borja, Chazuta, San Regis, and Nuevo Rocafuerte have lower and more reasonable runoff ratios. This may highlight that the underestimation of precipitation by global precipitation products is most problematic at small scale and over mountainous regions. This is compatible with Ward et al. (2011), who found comparable, unrealistically high runoff ratio values over the Paute Basin in Ecuador and the Baker Basin in Chilean/Argentinean Andes, even with interpolated rain gauge data. The high data uncertainty will limit our ability to identify and reproduce the hydrological responses of these upper basins.

Several noteworthy trends can also be observed in the interannual variation in the runoff ratio (3). The ratios calculated with the TRMM product are higher and increasing between 1998 and 2004 but show a sharp decreasing trend after 2004. In contrast, the runoff ratios with NCEP precipitation are generally lower prior to 2004 but deteriorate after 2004, yielding values above 1. The significance of the year 2004 as a turning point for both datasets is not clear – in the case of TRMM data, it may be possible that this is linked to changes in the estimation algorithm for one of the contributing satellites in mid-2003 (Huffman et al., 2007), but in the case of NCEP, the reanalysis model has been held static throughout the time series. Nevertheless, there is the general tendency of a drier climate during the wet season (based on the trend of maximum annual flows at Borja, not shown) and the fact that calibration campaigns for the streamflow stations started in 2003–2004. Therefore, despite the strong case for precipitation uncertainty, the possibility of a high streamflow data uncertainty cannot be discounted.

### 3.2 Simulation of streamflow

Figure 4 illustrates several spatial and temporal trends in the model performance indicators that are in line with the observed trends in the water balance in Sect. 3.1.

## JULES-LSM hydrology for humid tropical environments

Z. Zulkafli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





Figure 6 is a comparison between the modelled and observed flow duration curves, which provide a better insight into the model performance over the entire flow regime. The slopes of the curves are reasonably well simulated, particularly with the TRMM precipitation as the driving variable. This is likely due to better estimates of precipitation intensities from the observation dataset, which also comes at a finer spatial resolution than the NCEP and therefore may be better at capturing local convective rainfall. However, there is an overall underestimation of the discharge in the low to mid-flow region and overestimation of peak flows, further confirming a missing flow attenuation component in the model.

For similar reasons, JULES-TOPMODEL is underperforming when compared to JULES-BASE. The flashiness of the response simulated by JULES-TOPMODEL is exaggerated and this is suspected to be due to errors in the parameter scale. Indeed, the mean topographic indices calculated over the coarse model grids were negatively skewed, and the time-series average of the grid saturated fractions (Fig. 7) clearly shows that the partial areas contributing to saturation excess runoff were unreasonably overpredicted. In the lower basin, 50 % gridbox is saturated on average, which is high even for the flood forests that goes through seasonal flooding.

### 3.3 Simulation of internal variability of surface hydrological fluxes

We further compared JULES' simulated internal variability of surface hydrological fluxes to the values published in the literature (Fig. 8) to assess its robustness in representing the overall hydrological balance of a humid tropical basin. We identified a general negative bias in the simulation of evapotranspiration (ET). The simulated mean for flood forest ET is even lower than that for the lowland forest, which is counter-intuitive given the higher water availability in floodplains. One possibility for this low bias is an underestimation of canopy interception, but the evidence for this is weak. Despite positive skews in the distribution, the simulated canopy throughfall (TF) largely resides within the literature ranges.

## JULES-LSM hydrology for humid tropical environments

Z. Zulkafli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

**JULES-LSM**  
**hydrology for humid**  
**tropical environments**Z. Zulkafli et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

A similar bias was observed by Blyth et al. (2011) (with JULES) and Guimberteau et al. (2012) (with ORCHIDEE-LSM) in Amazonia. Blyth et al. (2011) also underestimated primary production, which may be correlated as the canopy conductance is a function of photosynthesis rate within the model (Cox et al., 1999). Both these processes are dependent on the soil moisture, as is also the bare surface evaporation, and it is likely that the low bias with ET is due to the model overestimating soil water stress. Studies focusing on simulations of the soil moisture state have indeed shown negative biases particularly in the lowest layer in the soil profile, suggesting a weakness of the free gravity drainage assumption (Bakopoulou et al., 2012; Finch and Haria, 2006). This assumption prevents drawing up of soil moisture during dry periods when the water table dips below the maximum soil depth. The JULES-TOPMODEL implementation is an attempt to address this limitation with an underlying unconfined aquifer; however, the model is persistently saturated and loses the effective rainfall to the routing (as evident in the larger contribution to surface runoff with JULES-TOPMODEL), resulting in no observed improvement to the ET.

This reconfirms the need to simulate better the movement of lateral fluxes within the basin. Dadson et al. (2010) implemented a 2-D routing scheme based on the kinematic wave assumption that continuously estimates flood extents based on the simulated water level of each grid cell. The land cover tiles in JULES are updated in the subsequent timestep by converting the flooded fraction into an open water surface. The subsurface and consequently plant roots, however, do not gain access to this available moisture, as open water tiles in JULES do not infiltrate. Therefore, the improvement to ET estimates is solely due to the increase in the open water ET, which may not be sufficient for the flood forest system of the Marañón River Basin. A more optimal model may be the floodplain implementation with the ORCHIDEE land surface model by d'Orgeval et al. (2008), who model the surface area and volume of swamps and floodplain in order to calculate the water retention time, allowing reinfiltration into the subsurface during this period. However, their model splits each coarse LSM grid into smaller subbasins, likely requiring an instantaneous redistribution of soil moisture over the entire grid in

the subsequent timestep. This assumption may undermine the soil moisture accounting in flooded versus dry sections of the grid, although it may be less problematic over finer scale grids.

A final nonetheless important observation is that in the upland biome, ET is better simulated and because of errors in the water balance, this corresponds to poor runoff generation. This is a clear contrast to the simulations in the lowland forests, where the poor estimation of ET instead makes up for the water balance errors, resulting in well estimated runoff and consequently streamflow. In the latter case, data errors compensated for model and parameter errors and created a false impression of good modelling.

## 4 Conclusions

Our study has shown that hydrological predictions with the JULES-LSM's can be unreliable due to a large uncertainty in the driving data and the poor simulation of the baseflow component in the upper Andean Basins. In the peneplain, the model is unable to reproduce the well-regulated regime as it neglected the hydrological functions of the flood forest. Nevertheless, for a global model that is not purpose-built for hydrological modelling, JULES is capable of producing reasonable simulations of the flow regime at fine temporal scale.

In constructing a robust model for impact analysis of a resilient system such as the Amazon, it is important to represent the hydrological system holistically in terms of the internal states and fluxes, perhaps more than it is to score a near-perfect Nash Sutcliffe efficiency. We further assessed whether the model is capable of behaving as a mirror image of real systems elsewhere in terms of the basin's internal hydrological fluxes. We have identified the model weakness in the estimation of ET and suspect this to be due to errors in predicting soil water availability due to misrepresentation of the inundated areas of the Andean wetlands and in the Amazon floodplain. Future research will

## JULES-LSM hydrology for humid tropical environments

Z. Zulkafli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



explore adaptation of the model structure to better represent the hydrological functions of these natural features.

*Acknowledgements.* The study is funded by the Ministry of Higher Education, Malaysia, and grant NE/I004017/1 of the UK Natural Environment Research Council (NERC). We thank Richard Ellis and Douglas Clark (the Center of Hydrology) for assistance with JULES implementation, Grenville Lister (University of Reading) and Keir Bovis (UK Meteorological Office) for access to the Unified Model ancillary data, and Toby Marthews (the University of Oxford) for advice on pedotransfer functions.

## References

- 10 Aalto, R., Dunne, T., and Guyot, J. L.: Geomorphic controls on Andean denudation rates, *J. Geol.*, 114, 85–99, doi:10.1086/498101, 2006. 12525
- Alkama, R., Decharme, B., Douville, H., and Ribes, A.: Trends in global and basin-scale runoff over the late twentieth century: methodological issues and sources of uncertainty, *J. Climate*, 24, 3000–3014, doi:10.1175/2010JCLI3921.1, 2011. 12525
- 15 Arora, V. K. and Boer, G. J.: An analysis of simulated runoff and surface moisture fluxes in the CCCma coupled atmosphere land surface hydrological model, *Proc. of International Congress on Modelling and Simulation, MODSIM 2003*, vol. 1, Townsville, Queensland, Australia, 166–171, 2003. 12525
- Asdak, C., Jarvis, P., van Gardingen, P., and Fraser, A.: Rainfall interception loss in unlogged and logged forest areas of Central Kalimantan, Indonesia, *J. Hydrol.*, 206, 237–244, doi:10.1016/S0022-1694(98)00108-5, 1998. 12551
- 20 Bakopoulou, C., Bulygina, N., Butler, A., and McIntyre, N.: Sensitivity analysis and parameter identifiability of the land surface model JULES at the point scale in permeable catchments, *Proc. of British Hydrological Society National Symposium, Dundee, June 2012*. 12538
- 25 Baldocchi, D. D., Falge, E., Gu, L., Olson, R., Hollinger, D., Running, S., Anthoni, P., Bernhofer, Ch., Davis, K., Fuentes, J., Goldstein, A., Katul, G., Law, B.E., Lee, X., Mahli, Y., Meyers, T., Munger, W., Oechel, W., Paw U, K. T., Pilegaard, K., Schmid, H. P., Valentini, R., Verma, S., Vesala, T., Wilson, K., and Wofsy, S. W.: FLUXNET: a new tool to study the temporal and spatial variability of ecosystem-scale carbon dioxide, water

## JULES-LSM hydrology for humid tropical environments

Z. Zulkafli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



vapor, and energy flux densities, *B. Am. Meteorol. Soc.*, 82, 2415–2434, doi:10.1175/1520-0477(2001)082<2415:FANTTS>2.3.CO;2, 2001. 12526

Bekoe, E.: Application of a hydrological model in a data-poor tropical West African catchment: a case study of the Densu Basin of Ghana, PhD. thesis, Cranfield University, available at: <http://dspace.lib.cranfield.ac.uk/handle/1826/1102> (last access: 23 November 2011), 2005. 12525

Best, M. J., Pryor, M., Clark, D. B., Rooney, G. G., Essery, R. L. H., Ménard, C. B., Edwards, J. M., Hendry, M. A., Porson, A., Gedney, N., Mercado, L. M., Sitch, S., Blyth, E., Boucher, O., Cox, P. M., Grimmond, C. S. B., and Harding, R. J.: The Joint UK Land Environment Simulator (JULES), model description – Part 1: Energy and water fluxes, *Geosci. Model Dev.*, 4, 677–699, doi:10.5194/gmd-4-677-2011, 2011. 12528, 12531

Beven, K., Buytaert, W., and Smith, L. A.: On virtual observatories and modelled realities (or why discharge must be treated as a virtual variable), *Hydrol. Process.*, 26, 1905–1908, doi:10.1002/hyp.9261, 2012. 12527

Blyth, E., Clark, D. B., Ellis, R., Huntingford, C., Los, S., Pryor, M., Best, M., and Sitch, S.: A comprehensive set of benchmark tests for a land surface model of simultaneous fluxes of water and carbon at both the global and seasonal scale, *Geosci. Model Dev.*, 4, 255–269, doi:10.5194/gmd-4-255-2011, 2011. 12526, 12538

Bookhagen, B. and Strecker, M.: Orographic barriers, high-resolution TRMM rainfall, and relief variations along the Eastern Andes, *Geophys. Res. Lett.*, 35, L06403, doi:10.1029/2007GL032011, 2008. 12530

Borma, L. S., da Rocha, H. R., Cabral, O. M., von Randow, C., Collicchio, E., Kurzatkowski, D., Brugger, P. J., Freitas, H. Tannus, R., Oliveira, L., Rennó, C. D., and Artaxo, P.: Atmosphere an hyrological controls for the evapotranspiration over a floodplain forest in the Bananal Island region, Amazonia, *J. Geophys. Res.*, 114, G01003, doi:10.1029/2007JG000641, 2009. 12551

Bormann, H. and Diekkrüger, B.: A conceptual, regional hydrological model for Benin (West Africa): validation, uncertainty assessment and assessment of applicability for environmental change analyses, *Phys. Chem. Earth Pt. A/B/C*, 29, 759–768, doi:10.1016/j.pce.2004.05.003, 2004. 12525

Braud, I., Vich, A., Zuluaga, J., Fornero, L., and Pedrani, A.: Vegetation influence on runoff and sediment yield in the Andes region: observation and modelling, *J. Hydrol.*, 254, 124–144, doi:10.1016/S0022-1694(01)00500-5, 2001.

**JULES-LSM  
hydrology for humid  
tropical environments**

Z. Zulkafli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Brooks, R. and Corey, A.: Hydraulic properties of porous media, in: Hydrology Papers No. 3, Colorado State Univ., Fort Collins, 27 pp., 1964. 12528
- Bruijnzeel, L. A., Mulligan, M., and Scatena, F. N.: Hydrometeorology of tropical montane cloud forests: emerging patterns, *Hydrol. Process.*, 25, 465–498, doi:10.1002/hyp.7974, 2011. 12534, 12551
- 5 Buytaert, W. and Beven, K.: Models as multiple working hypotheses: hydrological simulation of tropical alpine wetlands, *Hydrol. Process.*, 25, 1784–1799, doi:10.1002/hyp.7936, 2011. 12536
- Buytaert, W., Iñiguez, V., Celleri, R., De Bièvre, B., Cisneros, F., Wyseure, G., Deckers, J., and Hofstede, R.: Human impact on the hydrology of the Andean paramos, *Earth-Sci. Rev.*, 79, 53–72, doi:10.1016/j.earscirev.2006.06.002 2006a. 12553
- 10 Buytaert, W., Iniguez, V., Celleri, R., De Bievre, B., Wyseure, G., and Deckers, J.: Analysis of the water balance of small paramo catchments in South Ecuador, in: *Environmental Role of Wetlands in Headwaters*, edited by: Krecek, J. and Haigh, M., Springer, 271–281, 2006b. 12534, 12553
- 15 Campling, P., Gobin, A., Beven, K., and Feyen, J.: Rainfall-runoff modelling of a humid tropical catchment: the TOPMODEL approach, *Hydrol. Process.*, 16, 231–253, doi:10.1002/hyp.341, 2002. 12525, 12526, 12534, 12551
- Carvalho, L. M. V., Jones, C., and Liebmann, B.: The South Atlantic convergence zone: intensity, form, persistence, and relationships with intraseasonal to interannual activity and extreme rainfall, *J. Climate*, 17, 88–108, doi:10.1175/1520-0442(2004)017<0088:TSACZI>2.0.CO;2 2004. 12548
- 20 Celleri, R.: Rainfall variability and rainfall runoff dynamics in the Paute River Basin – Southern Ecuadorian Andes, Ph. D. thesis, Katholieke Universiteit Leuven, Leuven, Belgium, 2007. 12553
- 25 Chappell, N.: Soil pipe distribution and hydrological functioning within the humid tropics: a synthesis, *Hydrol. Process.*, 24, 1567–1581, doi:10.1002/hyp.7579, 2010. 12526
- Clark, D. B. and Gedney, N.: Representing the effects of subgrid variability of soil moisture on runoff generation in a land surface model, *J. Geophys. Res.-Atmos.*, 113, D10111, doi:10.1029/2007JD008940, 2008. 12529
- 30 Clarke, R. T., Mendes, C. A. B., and Buarque, D. C.: A comparison of extreme rainfall characteristics in the Brazilian Amazon derived from two gridded data sets and a national rain gauge network, *J. Geophys. Res.-Atmos.*, 115, D13104, doi:10.1029/2009JD013217, 2010. 12534

---

**JULES-LSM**  
**hydrology for humid**  
**tropical environments**Z. Zulkafli et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**JULES-LSM  
hydrology for humid  
tropical environments**

Z. Zulkafli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Coe, M.: Long-term simulations of discharge and floods in the Amazon Basin, *J. Geophys. Res.*, 107, 1–17, doi:10.1029/2001JD000740, 2002. 12534
- Cosby, B. J., Hornberger, G. M., Clapp, R. B., and Ginn, T. R.: A statistical exploration of the relationships of soil moisture characteristics to the physical properties of soils, *Water Resour. Res.*, 20, 682–690, doi:10.1029/WR020i006p00682, 1984. 12526
- Cox, P. M., Betts, R. A., Bunton, C. B., Essery, R. L. H., Rowntree, P. R., and Smith, J.: The impact of new land surface physics on the GCM simulation of climate and climate sensitivity, *Clim. Dynam.*, 15, 183–203, doi:10.1007/s003820050276, 1999. 12538
- D’Almeida, C., Vörösmarty, C. J., Marengo, J. A., Hurr, G. C., Dingman, S. L., and Keim, B. D.: A water balance model to study the hydrological response to different scenarios of deforestation in Amazonia, *J. Hydrol.*, 331, 125–136, doi:10.1016/j.jhydrol.2006.05.027 2006. 12525
- d’Orgeval, T., Polcher, J., and de Rosnay, P.: Sensitivity of the West African hydrological cycle in ORCHIDEE to infiltration processes, *Hydrol. Earth Syst. Sci.*, 12, 1387–1401, doi:10.5194/hess-12-1387-2008, 2008. 12538
- Dadson, S. J., Ashpole, I., Harris, P., Davies, H. N., Clark, D. B., Blyth, E., and Taylor, C. M.: Wetland inundation dynamics in a model of land surface climate: evaluation in the Niger inland delta region, *J. Geophys. Res.*, 115, D23114, doi:10.1029/2010JD014474, 2010. 12538
- Daren-Harmel, R. and Smith, P. K.: Consideration of measurement uncertainty in the evaluation of goodness-of-fit in hydrologic and water quality modeling, *J. Hydrol.*, 337, 326–336, doi:10.1016/j.jhydrol.2007.01.043, 2007. 12533
- Darko, P. K.: Estimation of natural direct groundwater recharge in Southwest Ghana using water balance simulations, available at: <http://dlib.lib.cas.cz/6136/>, last access: 23 November 2011, *J. Hydrol. Hydromech.*, 50, 198–212, 2002. 12525
- Decharme, B. and Douville, H.: Introduction of a sub-grid hydrology in the ISBA land surface model, *Clim. Dynam.*, 26, 65–78, doi:10.1007/s00382-005-0059-7, 2006. 12525
- Dunne, T.: Field studies of hillslope processes, in: *Hillslope Hydrology*, edited by: Kirkby, M., John Wiley, 227–294, 1978. 12526
- Ebel, B. A. and Loague, K.: Physics-based hydrologic-response simulation: seeing through the fog of equifinality, *Hydrol. Process.*, 20, 2887–2900, doi:10.1002/hyp.6388, 2006. 12525
- Espinoza-Villar, J. C. E., Guyot, J. L., Ronchail, J., Cochonneau, G., Filizola, N., Fraizy, P., Labat, D., de Oliveira, E., Ordonez, J. J., and Vauchel, P.: Contrasting regional discharge evolutions in the Amazon Basin (1974–2004), *J. Hydrol.*, 375, 297–311, doi:10.1016/j.jhydrol.2009.03.004, 2009a. 12530

## JULES-LSM hydrology for humid tropical environments

Z. Zulkafli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Espinoza-Villar, J., Ronchail, J., Guyot, J. L., Cochonneau, G., Naziano, F., Lavado, W., De Oliveira, E., Pombosa, R., and Vauchel, P.: Spatio-temporal rainfall variability in the Amazon Basin countries (Brazil, Peru, Bolivia, Colombia, and Ecuador), *Int. J. Climatol.*, 29, 1574–1594, doi:10.1002/joc.1791, 2009b. 12530
- 5 FAO/IIASA/ISRIC/ISS-CAS/JRC: Harmonized World Soil Database, <http://hydrosheds.cr.usgs.gov/>, last access: 21 January 2011, FAO, Rome, Italy and IIASA, Laxenburg, Austria, 2009. 12530, 12531
- Finch, J. and Haria, A.: The representation of chalk soils in the JULES/MOSES soil hydrology model, 2006. 12538
- 10 Fleischbein, K., Wilcke, W., Goller, R., Boy, J., Valarezo, C., Zech, W., and Knoblich, K.: Rainfall interception in a lower montane forest in Ecuador: effects of canopy properties, *Hydrol. Process.*, 19, 1355–1371, doi:10.1002/hyp.5562, 2005. 12552
- Garreaud, R. D., Vuille, M., Compagnucci, R., and Marengo, J.: Present-day South American climate, *Palaeogeogr. Palaeoclimatol.*, 281, 180–195, doi:10.1016/j.palaeo.2007.10.032, 2009. 12530
- 15 Giertz, S. and Diekkrüger, B.: Analysis of the hydrological processes in a small headwater catchment in Benin (West Africa), *Phys. Chem. Earth Pt. A/B/C*, 28, 1333–1341, doi:10.1016/j.pce.2003.09.009, 2003. 12526
- 20 Giertz, S., Diekkrüger, B., and Steup, G.: Physically-based modelling of hydrological processes in a tropical headwater catchment (West Africa): process representation and multi-criteria validation, *Hydrol. Earth Syst. Sci.*, 10, 829–847, doi:10.5194/hess-10-829-2006, 2006. 12525, 12526
- Goller, R., Wilcke, W., Leng, M., Tobschall, H., Wagner, K., Valarezo, C., and Zech, W.: Tracing water paths through small catchments under a tropical montane rain forest in South Ecuador by an oxygen isotope approach, *J. Hydrol.*, 308, 67–80, doi:10.1016/j.jhydrol.2004.10.022, 2005. 12553
- 25 Guimberteau, M., Drapeau, G., Ronchail, J., Sultan, B., Polcher, J., Martinez, J.-M., Prigent, C., Guyot, J.-L., Cochonneau, G., Espinoza, J. C., Filizola, N., Fraizy, P., Lavado, W., De Oliveira, E., Pombosa, R., Noriega, L., and Vauchel, P.: Discharge simulation in the sub-basins of the Amazon using ORCHIDEE forced by new datasets, *Hydrol. Earth Syst. Sci.*, 16, 911–935, doi:10.5194/hess-16-911-2012, 2012. 12525, 12534, 12538
- 30 Harden, C. and Scruggs, P.: Infiltration on mountain slopes: a comparison of three environments, *Geomorphology*, 55, 5–24, doi:10.1016/S0169-555X(03)00129-6, 2003.

Hodnett, M. and Tomasella, J.: Marked differences between van Genuchten soil water-retention parameters for temperate and tropical soils: a new water-retention pedo-transfer functions developed for tropical soils, *Geoderma*, 108, 155–180, doi:10.1016/S0016-7061(01)00101-X, 2002. 12527

5 Huffman, G. J., Bolvin, D. T., Nelkin, E. J., Wolff, D. B., Adler, R. F., Gu, G., Hong, Y., Bowman, K. P., and Stocker, E. F.: The TRMM Multisatellite Precipitation Analysis (TMPA): quasi-global, multiyear, combined-sensor precipitation estimates at fine scales, *J. Hydrometeorol.*, 8, 38–55, doi:10.1175/JHM560.1, 2007. 12532, 12535

10 Josse, C., Navarro, G., Encarnación, F., Tovar, A., Comer, P., Ferreira, W., Rodríguez, F., Saito, J., Sanjurjo, J., Dyson, J., Rubin de Celis, E., Zárate, R., Chang, J., Ahuite, M., Vargas, C., Paredes, F., Castro, W., Maco, J., and Arreátegui, F.: Ecological Systems of the Amazon Basin of Peru and Bolivia, Classification and Mapping, Tech. rep., [http://www.natureserve.org/lacSite/sobreNosotros/documents/Sistemas\\_low\\_ENG.pdf](http://www.natureserve.org/lacSite/sobreNosotros/documents/Sistemas_low_ENG.pdf), last access: 3 February 2011, NatureServe, Arlington, Virginia, 2007a. 12532

15 Josse, C., Navarro, G., Encarnación, F., Tovar, A., Comer, P., Ferreira, W., Rodríguez, F., Saito, J., Sanjurjo, J., Dyson, J., Rubin de Celis, E., Zárate, R., Chang, J., Ahuite, M., Vargas, C., Paredes, F., Castro, W., Maco, J., and Arreátegui, F.: Digital Ecological Systems Map of the Amazon Basin of Peru and Bolivia, [http://www.natureserve.org/aboutUs/latinamerica/gis\\_data\\_downloads.jsp](http://www.natureserve.org/aboutUs/latinamerica/gis_data_downloads.jsp), last access: 25 January 2011, NatureServe, Arlington, Virginia, USA, 2007b. 12531

20 Kahn, F.: Ecology of economically important palms in Peruvian Amazonia, *Adv. Econ. Bot.*, 6, 42–49, 1988. 12527

Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Leetmaa, A., Reynolds, B., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K., Ropelewski, C., Wang, J., Roy, J., and Joseph, D.: The NCEP/NCAR 40-Year reanalysis project, *B. Am. Meteorol. Soc.*, 77, 437–472, doi:10.1175/1520-0477(1996)077<0437:TNYRP>2.0.CO;2, 1996. 12532

25 Kvist, L. and Nebel, G.: A review of Peruvian flood plain forests: ecosystems, inhabitants and resource use, *Forest Ecol. Management*, 150, 3–26, doi:10.1016/S0378-1127(00)00679-4, 2001. 12530

Latrubesse, E. M., Stevaux, J. C., and Sinha, R.: Tropical rivers, *Geomorphology*, 70, 187–206, doi:10.1016/j.geomorph.2005.02.005, 2005. 12524

## JULES-LSM hydrology for humid tropical environments

Z. Zulkafli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

**JULES-LSM  
hydrology for humid  
tropical environments**Z. Zulkafli et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Legesse, D., Vallet-Coulomb, C., and Gasse, F.: Hydrological response of a catchment to climate and land use changes in tropical Africa: case study South Central Ethiopia, *J. Hydrol.*, 275, 67–85, doi:10.1016/S0022-1694(03)00019-2, 2003. 12525

Lehner, B., Verdin, K., and Jarvis, A.: New global hydrography derived from spaceborne elevation data, available at: <http://hydrosheds.cr.usgs.gov>, last access: 3 May 2006, *EOS Trans. AGU*, 89, 93–94, 2008. 12531

Loveland, T. R., Reed, B. C., Brown, J. F., Ohlen, D. O., Zhu, Z., Yang, L., and Merchant, J. W.: Development of a global land cover characteristics database and IGBP DISCover from 1 km AVHRR data, *Int. J. Remote Sens.*, 21, 1303–1330, , 2000. 12531

Negrón-Juárez, R., Hodnett, M., Fu, R., Goulden, M., and von Randow, C.: Control of dry season evapotranspiration over the amazonian forest as inferred from observations at a southern amazon forest site., *J. Climate*, 20, 2827–2839, doi:10.1175/JCLI4184.1, 2007. 12551

Nesbitt, S. and Anders, A.: Very high resolution precipitation climatologies from the Tropical Rainfall Measuring Mission precipitation radar, *Geophys. Res. Lett.*, 36, L15815, doi:10.1029/2009GL038026, 2009. 12532

Paiva, R. C., Collischonn, W., and Tucci, C. E.: Large scale hydrologic and hydrodynamic modeling using limited data and a GIS based approach, *J. Hydrol.*, 406, 170–181, doi:10.1016/j.jhydrol.2011.06.007, 2011. 12525

Penman, H. L.: Natural evaporation from open water, bare soil and grass, *P. Roy. Soc. Lond. A*, 193, 120–145, 1948. 12528

Rasanen, M., Neller, R., Salo, J., and Jungner, H.: Recent and ancient fluvial deposition systems in the Amazonian foreland basin, *Peru Geol. Mag.*, 129, 293–306, 1992. 12536

Rollenbeck, R. and Anhof, D.: Characteristics of the water and energy balance in an Amazonian lowland rainforest in Venezuela and the impact of the ENSO-cycle, *J. Hydrol.*, 337, 377–390, doi:10.1016/j.jhydrol.2007.02.004, 2007. 12534, 12551

Salati, E. and Vose, P.: Amazon Basin: a system in equilibrium, *Science*, 225, 129–138, doi:10.1126/science.225.4658.129, 1984. 12525

Schmid, S., Burkard, R., Frumau, K. F. A., Tobón, C., Buijnzeel, L. A., Siegwolf, R., and Eugster, W.: Using eddy covariance and stable isotope mass balance techniques to estimate fog water contributions to a Costa Rican cloud forest during the dry season, *Hydrol. Process.*, 25, 429–437, doi:10.1002/hyp.7739, 2011.

## JULES-LSM hydrology for humid tropical environments

Z. Zulkafli et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)




[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


- Sheffield, J., Goteti, G., and Wood, E. F.: Development of a 50-year high-resolution global dataset of meteorological forcings for land surface modeling, *J. Climate*, 19, 3088–3111, doi:10.1175/JCLI3790.1 2006. 12532
- 5 Tomasella, J. and Hodnett, M. G.: Estimating soil water retention characteristics from limited data in Brazilian Amazonia, *Soil Sci.*, 163, 190–202, 1998. 12527, 12531
- Tomasella, J., Hodnett, M. G., and Rossato, L.: Pedotransfer Functions for the Estimation of Soil Water Retention in Brazilian Soils, *Soil Sci. Soc. Am. J.*, 64, 327–338, doi:10.2136/sssaj2000.641327x, 2000. 12527
- 10 van Genuchten, M. T.: A closed-form equation for predicting the hydraulic conductivity of unsaturated soils, *Soil Sci. Soc. Am. J.*, 44, 892–898, doi:10.2136/sssaj1980.03615995004400050002x, 1980. 12528
- Vertessy, R. A. and Elsenbeer, H.: Distributed modeling of storm flow generation in an Amazonian rain forest catchment: effects of model parameterization, *Water Resour. Res.*, 35, 2173–2187, doi:10.1029/1999WR900051, 1999. 12525
- 15 Vörösmarty, C. J., Moore, B. I., Grace, A. L., Gildea, M. P., Melillo, J. M., Peterson, B. J., Rastetter, E., and Steudler, P.: Continental scale models of water balance and fluvial transport: an application to South America, *Global Biogeochem. Cy.*, 3, 241–265, doi:10.1029/GB003i003p00241, 1989. 12525
- Ward, E., Buytaert, W., Peaver, L., and Wheatler, H.: Evaluation of precipitation products over complex mountainous terrain: a water resources perspective, *Adv. Water Resour.*, 34, 1222–1231, doi:10.1016/j.advwatres.2011.05.007, 2011. 12534, 12535
- 20 Yeh, P. J. F. and Eltahir, E. A. B.: Representation of water table dynamics in a land surface scheme, Part I: model development, *J. Climate*, 18, 1861–1880, 2005. 12526
- Zadroga, F.: The hydrological importance of a montane cloud forest area of Costa Rica, in: *Tropical Agricultural Hydrology*, edited by: Lal, R. and Russell, E. W., John Wiley and Sons Ltd., New York, 59–73, 1981. 12534
- 25



## JULES-LSM hydrology for humid tropical environments

Z. Zulkafli et al.

**Table 2.** Hydrological stations description

Station	River basin	Coordinates	Elevation (m)	Drainage Area (km <sup>2</sup> )	Period of available data	Mean $Q_{obs}$ (all available data)	Normalized mean $Q_{obs}$ (mm km <sup>-2</sup> )
San Regis	Marañón	73.9° W, 4.5° S	93	363 848	1986–2011	16 601	45.6
Borja	Marañón, upstream	77.5° W, 4.5° S	200	114 991	1986–2011	4539	39.5
Santiago	Santiago – northern tributary to Marañón upstream of Borja	78.0° W, 3.1° S	290	23 806	2001–2011	1585	66.6
Paute	Paute – tributary to Santiago	78.6° W, 2.6° S	1840	4917	1999–2004	109	22.2
Chazuta	Huallaga – southern tributary to Marañón downstream of Borja	76.1° W, 6.6° S	180	69 175	1998–2009	3042	44.0
Nuevo Rocafuerte	Napo – northern tributary to Marañón downstream of San Regis	75.4° W, 0.9° S	189	27 534	2001–2011	2176	79.0
San Sebastian	Coca – tributary to Napo	77.0° W, 0.3° S	290	5329	2000–2011	459	86.1
Nueva Loja	Aguarico – tributary to Napo	76.8° W, 0.0° N	299	4640	2001–2011	593	127.8

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## JULES-LSM hydrology for humid tropical environments

Z. Zulkafli et al.

**Table 3.** Data uncertainty and water balance closure calculated at the hydrological stations. PER = probable error range;  $Q_{\text{obs}}$  = observed streamflow;  $P$  = precipitation.

Station	PER (%)	$Q_{\text{obs}}/P$ -TRMM (1998–2008)	$Q_{\text{obs}}/P$ -NCEP (1998–2008)
San Regis	6.28	0.82	0.71
Borja	13	0.94	1.00
Santiago	7.05	1.36	1.14
Paute	–	1.10	0.39
Chazuta	1.31	0.81	1.11
Nuevo Rocafuerte	–	1.20	0.80
San Sebastian	–	1.47	1.09
Nueva Loja	–	1.76	1.88

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


**Table 4.** Summary of fluxes from humid tropical hydrology literature

Reference	Location	Precipitation (mm)	Evapo-transpiration (mm)	Throughfall (mm)	Surface Runoff (mm)	Subsurface Runoff (mm)
<b>Lowland forests</b>						
References in Bruijnzeel et al. (2011)	Central Amazonia, Ducke	2500	1310			
	Pasoh, Malaysia	1800	1448			
References in Rollenbeck and Anhof (2007)	Lambir Hills, Malaysia	2740	1303			
	Estado Amazonas, Venezuela	3244	1492	2595	195	1848
	French Guayana	3200	1346			
	Belem, Para, Brasilien	1819	1905			
	S Carlos De Rio Negro, Venezuela	3664	2065	3188		1759
	S Carlos De Rio Negro, Venezuela	3500	1502	2450		
	Reserva Ducke, Brazil	2209	1119	1966	67	640
	Lake Calado, Brazil	2870			115	1679
	Reserva Ducke, Brazil	2391				
	Reserva Ducke, Brazil	2391				
	Manaus, Brazil	3000				780
	Reserva Ducke, Brazil	2636	1318	2320		
Reserva Jaru, Brazil	3563					
Negrón-Juárez et al. (2007)		2291	1026			
Asdak et al. (1998)	Kalimantan, Indonesia	2199		1918		
		3563		3334		
Campling et al. (2002)	West Benin	1157	867			
<b>Flood forests</b>						
Borma et al. (2009)	Bananal Island, Brazil	1692	1332			
<b>Montane forests</b>						
References in Bruijnzeel et al. (2011)	Bolivia, Yungas	2310		1825		
	Columbia, Cordillera Central	3150		2394		
	Ecuador	2320		1554		
	Idem	2080		1477		
	Costa Rica, Monteverde	2500		1775		
	Costa Rica, Talamanca	2810		1967		
	Guatemala	2500		1625		
	Indonesia, Sulawesi	2900		2030		
	Peninsular Malaysia	2300		1426		
	Panama	3680		2318		
	Papua New Guinea	3800		2546		
	Peru, Central Cordillera	2220		1554		
	Tanzania, Usambara Mts	1230		972		
	Australia, Se Queensland	1350		1215		
	Australia, N Queensland, Ub Site	2985		2358		
	Idem, Mt Lewis	3315		2553		
	Idem, Mt Lewis	2610		2010		
	Bolivia, Yungas	3970		2938		
	China, Unnan, Ailao Mnt	1930		1679		
	Idem, Xishuangbanna	1485		1158		
	Colombia, Central Cordillera	2115		1851		
	Costa Rica Monteverde Windward Forest	6390		4473		
	Leeward Forest	2520		1638		
	Costa Rica, Monte De Los Olivos	3300				
	Windward Forest					
	Idem	3300		3498		
	Ecuador	2140		1819		
	Idem	2500		2275		
	Guatemala, Windward	2600		2106		
	Hawai, Maui, Leeward	1010		889		
	Honduras	3500		3325		
	Indonesia, West Java	3300		2640		
Mexico, Veracruz	3000		2475			
Tanzania, Mt Kilimanjaro	2480		2034			

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 4. Continued.

Reference	Location	Precipitation (mm)	Evapo- transpiration (mm)	Throughfall (mm)	Surface Runoff (mm)	Subsurface Runoff (mm)
	Idem, Dry Year	1960		1372		
	Thailand, Kog-Ma	2085		1856		
	Venezuela, San Eusebio	1575		1252		
	Venezuela, La Mucuy	3125		1688		
	Australia, Queensland, Bl	5300		6201		
	Colombia, Central Cordillera	1700		1513		
	Idem, Central Cordillera	1455		1273		
	Idem, Zipaquira	1615		1437		
	Costa Rica, Monteverde	4310		5086		
	Idem	6000		6480		
	Guatemala, Windward	2500		2825		
	Guatemala, Windward	2600		2288		
	Hawaii, Windward	2700		3537		
	Honduras Concave Slope	1500		1410		
	Honduras Convex Slope			1935		
	Honduras Ridge Top			2685		
	Jamaica, Pmull Forest	2850		2081		
	Malaysia, Peninsular	2115		1354		
	Peru, Central Cordillera	2750		2544		
	Philippines, Luzon	3910		3363		
	Puerto Rico, Palm Forest	4500		3105		
	Bolivia, Yungas	5150		3863		
	Hawaii	500		375		
	Jamaica, Mmor Forest	2855		1713		
	Puerto Rico, East Peak Elfin, Windward	5400		6210		
	Idem Reidge	4800		6000		
	Idem Leeward	6000		5760		
	Elfin, Windward	5400		8910		
	Idem, Sheltered	4500		4860		
	Reunion France	3000		3720		
	Spain, La Gomera	660		832		
	Madagascar, Perinet	2080	1290			
	Colombia, Sierra Nevada	1985	1270			
	Ecuador	2050	1271			
	Kenya, Kericho	2130	1342			
	Idem	2015	1249			
	Kenya, Kimakia	2305	1153			
	Tanzania, Mbeya	1925	1386			
	Queensland, Gambubal	1350	1256			
	Idem, Upper Barron	2985	1433			
	Idem, Mt St Lewis	3040	1459			
	Hawaii, Volcano Np	2500	1225			
	Indonesia, Jawa	3300	1155			
	Ecuador	2140	920			
	Mexico, Veracruz	3100	1426			
	Thailand, Kog-Ma	1768	813			
	Venezuela, San Eusebio	1465	982			
	Venezuela, La Mucuy	3125				
	Costa Rica, Monteverde	1450	783			
	Queensland, Bellenden Ker	1560	905			
	Puerto Rico, Luquillo Mountain	900	675			
	Puerto Rico, Luquillo Mountain	1010	566			
	La Gomera, Canary Islands	1270	533			
Fleischbein et al. (2005)	Loja, Ecuador, Site 1				1083	
	Loja, Ecuador, Site 2				1046	
	Loja, Ecuador, Site 3				1039	

**JULES-LSM**  
hydrology for humid  
tropical environments

Z. Zulkafli et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## JULES-LSM hydrology for humid tropical environments

Z. Zulkafli et al.

[Title Page](#)

[Abstract](#)   [Introduction](#)

[Conclusions](#)   [References](#)

[Tables](#)   [Figures](#)

[⏪](#)   [⏩](#)

[◀](#)   [▶](#)

[Back](#)   [Close](#)

[Full Screen / Esc](#)


[Printer-friendly Version](#)

[Interactive Discussion](#)

**Table 4.** Continued.

Reference	Location	Precipitation (mm)	Evapo- transpiration (mm)	Throughfall (mm)	Surface Runoff (mm)	Subsurface Runoff (mm)
Goller et al. (2005)	Loja, Ecuador, Site 1				1154	
	Loja, Ecuador, Site 2				680	
	Loja, Ecuador, Site 3				1191	
<b>Upper Andes</b>						
References in Buytaert et al. (2006a)	Machangara, Ecuador	1100				
	Ningar, Ecuador	950				
	Jima, Cuenca, Ecuador	1000				
	Chimborazo, Ecuador - Humid	900				
	- Dry	600				
	- Pantanal	900				
	Azoguez, Ecuador	1860				
	Cuenca, Ecuador	1270				
	Pichincha, Ecuador	1500				
		2000				
	El Angel, Ecuador	1150				
	Cotacachi, Ecuador	1500				
	Piedras Blancas, Venezuela	800				
	Andean paramos, from	700				
	to	3000				
	Columbian paramos	6000				
Buytaert et al. (2006b)	Huagrauma, Machanagra, Ecuador	1200	600		850	
	Soroche, Machanagra, Ecuador	800	450		500	
Celleri (2007)	Burgay	820				
	Duda	1120				
	Jadan	750				
	Matadero	1230				
	Mazar	1160				
	Yanuncay	1100				
	Tomebamba	980			200	600
	Paute	1030				





**Fig. 1.** The Peruvian Amazon Basin. The Marañón River merges with Ucayali River from the south and Napo River from the north beyond the downstream limits of the basin to form the Amazon proper river. The basin relief (in red) shows the location of the Andean ranges where the climate and physiography are highly variable

**JULES-LSM  
hydrology for humid  
tropical environments**

Z. Zulkafli et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶


Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 2.** The major biomes in the Marañón River Basin. The classification was performed using an ecosystem map and altitude and the evaluation of JULES was performed separately for each of the natural biomes.

## JULES-LSM hydrology for humid tropical environments

Z. Zulkafli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## JULES-LSM hydrology for humid tropical environments

Z. Zulkafli et al.




**Fig. 3.** Temporal and spatial trends in the runoff ratios calculated using observed streamflows and TRMM and NCEP precipitation. The expected value for humid tropical regions are between 0.6 and 0.7.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)



## JULES-LSM hydrology for humid tropical environments

Z. Zulkafli et al.



**Fig. 4.** Temporal and spatial trends in the hydrological performance indices. Blue: JULES-BASE and TRMM, magenta: JULES-TOPMODEL and TRMM, red: JULES-BASE and NCEP, orange: JULES-TOPMODEL and NCEP.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

**Fig. 5.** Modelled versus measured streamflows at all four gauging points for the modelling period 2006–2007. The overlying barplots are the basin-average precipitation time series. Black: observed flow, gray: 95 % confidence interval, blue: JULES-BASE and TRMM, magenta: JULES-TOPMODEL and TRMM, red: JULES-BASE and NCEP, orange: JULES-TOPMODEL and NCEP.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## JULES-LSM hydrology for humid tropical environments

Z. Zulkafli et al.



**Fig. 6.** Comparison of flow duration curves from observed and simulated streamflows.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## JULES-LSM hydrology for humid tropical environments

Z. Zulkafli et al.



**Fig. 7.** Mean area-saturated fraction of the gridbox from the JULES-TOPMODEL simulations.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## JULES-LSM hydrology for humid tropical environments

Z. Zulkafli et al.



**Fig. 8.** Internal basin's surface fluxes simulated by JULES compared with the variability found in the literature. Each subfigure is a kernel density plot of the surface hydrological fluxes in each biome, juxtaposed with a box plot of the distribution of published values from different world locations (Table 4). Blue: JULES-BASE and TRMM, magenta: JULES-TOPMODEL and TRMM, red: JULES-BASE and NCEP, orange: JULES-TOPMODEL and NCEP.