

This discussion paper is/has been under review for the journal Hydrology and Earth System Sciences (HESS). Please refer to the corresponding final paper in HESS if available.

Seasonal stratification and property distributions in a tropical estuary (Cochin estuary, west coast, India)

A. Shivaprasad, J. Vinita, C. Revichandran, P. D. Reny, M. P. Deepak, K. R. Muraleedharan, and K. R. Naveen Kumar

National Institute of Oceanography, Regional Centre, Dr. Salim Ali road, Kochi-18, Kerala, India

Received: 4 July 2012 – Accepted: 10 July 2012 – Published: 27 July 2012

Correspondence to: A. Shivaprasad (shivaprasadnio@gmail.com)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

The intratidal, spring-neap and seasonal variations in stratification were examined in Cochin estuary. The observations established a strong connection with the distribution of chemical and biological properties. The influence of tides and river discharge forcing in water column stability was quantified using potential energy anomaly (PEA) and stratification parameter. Partially mixed (neap) and well-mixed (spring) conditions during low river discharge (dry) period were altered in monsoon by the salt wedge intrusions. The ecological impact of salt wedge propagation on high tides bringing upwelled water to the system was evident from the bottom hypoxic, high chlorophyll *a* and nutrient-rich conditions. Phosphate and nitrite concentrations were higher at the bottom saline conditions but silicate and nitrate were clearly supplied by river water. However, during ebb tide this front was driven out of the estuary. The periodic advance and retreat of the salt wedge was inevitable in making the system immune from extended hypoxia/anoxia and maintaining the health of Cochin estuary. For the seasonally varying river flow in the estuary, salt intrusion receded with increasing river flow in monsoon and rebounded with decreasing river flow in dry season. During monsoon, the intense flushing and reduction in salinity field expansion seemed to be responsible for the limited chlorophyll *a* levels along the surface of Cochin estuary.

1 Introduction

The key of understanding stratification and de-stratification processes in different time scales (intra-tidal, spring-neap, seasonal) has gained the most attention during the last decades due to its tremendous relevance to the estuarine ecosystem. Estuaries can be classified into three types based on their longitudinal salinity distribution: partially mixed, vertically homogeneous or well-mixed and highly stratified or salt wedge type (Dyer, 1973). The type of the estuary essentially depends on river discharge and tidal regime of the estuary which have pronounced effect on the distributions of several

HESSD

9, 8979–9010, 2012

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

physical, chemical, and biotic processes within the estuary. The differential advection of salinity creates stratification which inhibits vertical mixing of momentum. With the increase in turbulent energy, stratification is reduced by mixing directly and indirectly by reduced shear (Nepf and Geyer, 1996). Stratification diminishes vertical fluxes of ecologically important variables (Uncles, 1990) like heat, salt, oxygen and nutrients. Physical dynamics play a critical role in estuarine biological production, material transport and water quality (Kasai et al., 2010).

The Cochin estuary is the largest among many extensive estuarine systems along the southwest coast of India. It has been identified as one of the most productive estuarine systems along the west coast of India by Menon et al. (2000). However due to anthropogenic interventions, the Cochin estuary has been reported to be on the brink of an ecological disaster by Dinesh Kumar et al. (1994). Further, the sediment heavy metal contamination of this estuary has placed the region among the impacted estuaries in the world (Balachandran et al., 2005). The run-off components like municipal waste discharge from the surrounding city, and riverine water carrying industrial and agricultural wastes are responsible for nutrient enrichment influencing the estuarine water quality (Joseph and Ouseph, 2010).

Numerous studies reveal the dynamics of the energetic environment of Cochin estuary. Tides in Cochin estuary are of mixed semidiurnal type with an average tidal range of 1 m (Qasim and Gopinathan, 1969). According to Srinivas et al. (2003b) the relative importance of the semi-diurnal and the diurnal components keeps changing throughout the month. Spring phase is dominated by semi-diurnal tides and neap phase by diurnal tides. There is a rapid decay in the amplitudes of the principal tidal constituents as they propagate upstream. However, tidal amplification is observed in the south estuary during pre-monsoon season (March) which is possibly caused by the closure of the hydraulic barrier at Thanneermukam (upstream) (Balachandran et al., 2008). In Cochin estuary, currents are dominated by tidal signals; semi-diurnal tidal regimes experience swift tidal currents than diurnal tidal regimes (Srinivas et al., 2003a). Balachandran et al. (2008) with the help of a model showed that strong currents prevail at the central

estuary (from Cochin inlet to 22 km south) whereas weak and slow currents are found in the north and south estuary. The strong ebb velocity of about 130 cm s^{-1} at Cochin inlet (Udaya Varma et al., 1981) which is comprised in the central estuary, maintains an effective flushing through the channel in this region (Balachandran et al., 2008).

For an ecological study, an interdisciplinary approach linking the physical phenomena with chemical and biological properties is essential and in Cochin estuary, this study is first of its kind. The physical processes (stratification, horizontal and vertical advection, flushing, etc.) that govern the ecological parameters have not been rigorously investigated in the region to date. For the efficient implementation of estuarine management plans, an imperative study of the impact of stratified systems on water quality and ecosystem ecology is essential. The objective of the present study is to assess (1) the intratidal and spring-neap variations in stratification of water column in dry and wet seasons (2) the horizontal extent of salt intrusion and the relation between salinity and property distributions.

1.1 Geomorphology

The Cochin estuary is the largest among many extensive estuarine systems along the southwest coast of India. It extends from Munambam ($10^{\circ} 10' \text{ N}$, $76^{\circ} 15' \text{ E}$) in the north, to Thannermukkam ($9^{\circ} 30' \text{ N}$, $76^{\circ} 25' \text{ E}$) in the south over a length of $\sim 80 \text{ km}$ (Fig. 1). The system is characterized by its long axis lying parallel to the coastline, with several small islands and interconnected waterways, and it covers an area of about 300 km^2 . The width of this estuarine system varies from 450 m to 4 km, and the depths range from 15 m at Cochin inlet to 3 m near the head with an average depth of 2.5 m (depths are reduced to chart datum). The estuary is separated from the sea by barrier spits interrupted by tidal inlets at Munambam, and Cochin. The openings at Cochin, known as the Cochin inlet, with a width of 450 m, and another at Munambam, provide perennial connections to the Arabian Sea. The Cochin Port, situated on the Willington Island, is near the Cochin inlet, which provides the main entrance channel to this harbour. The rivers that discharge freshwater into this estuarine system are Periyar in the north;

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Pampa, Achankovil, Manimala, and Meenachil in the south; and Muvattupuzha, midway between the two (Srinivas et al., 2003a). Munambam inlet is located further north, through which, 70 % of the Periyar River discharges into the Arabian Sea and the rest through the Cochin inlet (unpublished). Thannerermukkam barrier in the south was made functional in 1976 to prevent salt-water incursion and to promote cultivation in the low-lying fields. It remains closed from January to May every year.

1.2 Climatological setting

The domain experiences humid equatorial tropic climate with the mean daily temperatures ranging from 19.8 °C to 36.7 °C. Mean annual temperature ranges from 25.0–27.5 °C in the coastal lowlands (Government of Kerala, General features, 2005b). The average monthly rainfall for the year 1978 to 2002 at the different physiographic zones of the river basin is as shown in Fig. 2a. In most years, pre-monsoon (March–May) experiences the lowest recorded rainfall with a combined average of only 386 mm month⁻¹, thus defining the peak of the “dry” season. In contrast, southwest monsoon (June–September) receives the most rainfall with an average and maximum of 1400 mm month⁻¹ and 1891 mm month⁻¹, respectively; thus defining the peak of the “wet” season (Krishnakumar et al., 2009). The fresh water runoff data (Fig. 2b) for the year 2008 to 2009 was obtained from Central Water Commission. About 60 % to 70 % of the total discharge occurred during June–September and the least (6.82 %) occurred during December–February.

2 Materials and methods of observation

Intensive series of hydrographic surveys comprising three different time scales of observations were carried out in the Cochin estuary: (a) time-series at a single station positioned close to the main inlet of the system (temporal study of water column stability),

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

(b) synoptic (salt intrusion measurement) and (c) monthly observations (correlation analysis of environmental parameters) at ten stations.

2.1 Time series observations

Time series data were collected from a station, 5 km upstream of Cochin inlet during four surveys in 2007 (Fig. 1). Since the environmental behaviour and dynamics of the Cochin estuarine system is highly influenced by monsoonal rainfall and the associated runoff, we had ideally chosen both extreme conditions of seasons for data collection; pre-monsoon (dry) and monsoon (wet) seasons. Significantly, the observational coverage included two spring (2–3 May, 16–17 July 2007) and two neap (25–26 April, 24–25 July 2007) tidal phases. The measurements were done for dry (April, May) and wet (July) seasons for 30 h and 27 h respectively covering two consecutive semidiurnal cycles. A SBE Seabird 19 plus CTD was used for recording temperature (accuracy ± 0.001 C) and salinity (conductivity ± 0.001 S m^{-1}) profiles with a bin size 0.2 m for every 30 min interval. The water level measurements throughout the observations were collected using tide pole and it was crosschecked with the tide table (Published by Survey of India) and the tide predicted using TASK-2000 software (POL/PSMSL Tidal Analysis Software Kit 2000, Proudman Oceanographic Laboratory, UK). Water samples collected from surface, mid-depth (~ 4 m) and close to bottom at 3 h intervals were utilised to determine the nutrient and chlorophyll *a* concentrations.

As a convenient measure of water column stability, potential energy anomaly (PEA, ϕ) was calculated for the entire water column for each CTD profile using the equation:

$$\phi = \frac{1}{h} \int_{-h}^0 (\bar{\rho} - \rho) g z dz \quad \text{where } \bar{\rho} = \frac{1}{h} \int_{-h}^0 \rho z dz \quad (1)$$

Here g is gravitational acceleration ($m s^{-2}$), ρ is water density ($kg m^{-3}$), h is water depth (m) and z is depth interval (m).

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Simpson (1981) defined PEA as shown in equation above as the amount of mechanical energy (per m^3) required to instantaneously homogenize the water column completely. Thereafter, this method has proved crucial for quantifying the mixing efficiency in numerous stratification studies in coastal seas and estuaries (Nunes Vaz et al., 1989; Rippeth and Simpson, 1996; Lund-Hansen et al., 1996; Ranasinghe and Pattiaratchi, 1999). Although the meteorological phenomena like wind are possible sources of mixing energy, they are overshadowed by the constancy of tidal action (Blanton, 1969) and are therefore not treated in this study.

Water column stratification for each profile in time series observation was assessed using the stratification parameter, n_s defined as:

$$n_s = \frac{\delta S}{S'_m} \quad (2)$$

where $\delta S = S_{\text{bott}} - S_{\text{surf}}$, $S'_m = 1/2 (S_{\text{bott}} + S_{\text{surf}})$, with S_{surf} and S_{bott} the salinity at the surface and bottom of the water column, respectively. In case $n_s < 0.1$, then the water column is well mixed, when $0.1 < n_s < 1.0$ then partial mixing occurs, while if $n_s > 1.0$ stratification with the presence of salt-wedge is evident (Haralambidou et al., 2010).

2.2 Synoptic observations

The results of the time series measurements conducted in the most dynamical zone of Cochin estuary (Balachandran et al., 2008) inspired us to proceed our observations further along estuary to explore the longitudinal salinity dynamics. Hence, salinity intrusion surveys were undertaken from June 2008 to May 2009 during spring and neap tidal phases of each month by casting CTD-SBE 9¹¹ plus every 8 km intervals from northern arm to southern arm using a speed boat (40 km h^{-1}) covering ten stations along the estuary (Fig. 1). Morphology, tides and discharge were the deciding factors for the selection of sampling stations. Extensive data were gathered from stations 1 and 5, 6, 7 located at the proximity of northern and Cochin inlets, respectively, stations

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2, 3, 4 at the middle of northern arm and 8, 9, 10 comprising of southern arm. Occasionally, we had to face technical problems such that the measurements obtained with CTD did not reach bottom due to strong water currents in January and also a missing data of station 10 in June measurements.

5 The flushing time defined as the time taken to replace the existing freshwater in the estuary, at a rate equal to river discharge (Dyer, 1997), was calculated by Fresh water fraction method (Ketchum, 1983) i.e., $T = F/Q$ where F is the total volume of the fresh water in Cochin estuary and Q is the river discharge from the year 2008 to 2009.

2.3 Monthly observations

10 In order to relate the physical forcing (tide and river discharge) to chemical and biological property distributions, additional surveys were conducted at the mid of every month from June 2008 to May 2009 (similar to the period of synoptic survey). CTD-SBE 9¹¹ plus was used to measure temperature and salinity. Nutrients, chlorophyll *a* and dissolved oxygen (DO) concentrations were determined from the bottle samples collected
15 from the surface.

2.4 Chemical and biological parameters measurements

For the analysis of chlorophyll *a*, one litre of water sample from each depth was filtered through Whatmann GF/F filter and measured according to Strickland and Parsons (1972) using flourometer (Turner designs Instruments, Trilogy, USA). Pheophytin
20 (acidified 0.1 N HCL) concentrations were determined and deducted. Dissolved oxygen was analyzed by the Winkler's titrimetric method. Dissolved inorganic nutrients such as nitrite (NO₂⁻), nitrate (NO₃⁻), phosphate (PO₄³⁻) and silicate (SiO₄⁴⁻) were estimated following standard colorimetric techniques (Grasshoff et al., 1983).

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2.5 Statistical data analysis

The Pearson correlation coefficients were calculated using SPSS 17 statistical software to find out the linear relationship of salinity with all chemical and biological parameters. Two-way ANOVA was carried out on the monthly surface discrete samples to examine the difference in water quality variables among the along-estuary sampling stations (spatial) or through the different sampling months (temporal) with factors of interest being season, discharge and tidal activity.

3 Results

3.1 Temporal variations at single station

3.1.1 Tidal characteristics

Figure 3a–d gives the water level and predicted tide of spring and neap phases during the observation period. The maximum range of the spring tide was 1 m while the neap tides did not exceed 0.59 m. The diurnal tidal inequality was quite evident in all observations; it was prominent in dry neap with 0.20 m (Fig. 3a). Unlike all other tidal observations which were semi-diurnal in nature, the neap tide of wet season was diurnal (Fig. 3d).

3.1.2 Temperature

The vertical distribution of temperature is as shown in Figs. 4–7. A pronounced surface to bottom difference in temperature was absent during both spring and neap tidal phases in dry season. Even though, during the period when solar forcing during late afternoon was at its peak created a vertical difference of $\sim 1.5^\circ\text{C}$ (Figs. 4b, 5b). However, the high tides of both the tidal phases of monsoon season experienced typically

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

substantially lower temperature ($\sim 24.5^\circ\text{C}$) (Figs. 6b, 7b) at the bottom layers when compared with surface temperature ($\sim 27.5^\circ\text{C}$).

3.1.3 Salinity

The seasonally varying river flow and the tidal rhythm affected the vertical distribution of salinity as shown in Figs. 4–7. During the period April–May, low fresh water discharge (Figs. 4a, 5a) caused weaker salinity stratification (difference between the surface and near bottom salinities) relative to summer monsoon. In dry neap (April), the stratification was observed during flood with maximum of 16.35. With the arrival of ebb phase of the first tidal cycle, stratification declined to a remarkably low value of 0.19. Again, it increased with the upcoming flood tide. During spring phase of dry season (May), stratification of the entire water column over the tidal cycles was less and the depth averaged salinity ranged from 31.64 to 32.09. The stratification was higher in flood (12.5) than in ebb (0.15) either due to differential advection of salinity or vertical mixing. On the other hand, during both tidal phases of monsoon (Figs. 6a, 7a) large quantities of fresh water entered the estuary resulting in very low saline water at the surface and denser water at the bottom. Evolution of salinity stratification in water column (spring 33.24; neap 33.03) is associated with advection of salt wedge. The isopycnals were evenly spaced and flattened during highest high water (HHW). Notably the entire station was flushed with fresh water of salinity ~ 0.05 during low tides and lowest high water (LHW). The pycnocline became unstable during ebb and the saline wedge disappeared.

3.1.4 Stability factor and stratification parameter

The pattern of distribution of observed water column stability (Figs. 3e–h) corresponded well with the salinity distribution notwithstanding seasons and tidal cycles. The spring-neap cycles of stabilization and destabilization of water column were prominent which is having relative importance as far as tidal estuaries are concerned. During dry season, the average PEA value computed from all density profiles was 52.02 J m^{-3} in

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



neap which almost reduced to 25.9 Jm^{-3} during spring phase. This implies that the average energy required to mix the water column was about two-fold higher in neap than spring phase. It should also be noted that during dry neap the minimum PEA value of 5.0 Jm^{-3} was obtained during the ebb period. However, during dry spring the PEA values neared to zero for many of the density profiles over the tidal cycle. During wet the highest values of PEA ($> 200 \text{ Jm}^{-3}$) in spring and neap phases indicated stratification evolved due to the advancement of the salt wedge. However, for spring, on the arrival of ebb phase, PEA values gradually declined corresponding to the retreat of salt wedge and went as low as $< 1 \text{ Jm}^{-3}$, this low value sustained until the next tidal cycle and again the stratification began to develop at HHW. For wet neap, the water level was diurnal in nature. The observations of neap began with low PEA values ($\sim 10 \text{ Jm}^{-3}$) and the advancement of salt wedge occurred at the flood phase of the tidal cycle. Again, it began to retreat during ebb.

During dry season, stratification parameter fluctuated from 0.5 to 0.8 (partially mixed) in neap phase whereas for spring phase, most of the observations over the tidal cycle showed stratification number varying from 0 to 0.1 (well-mixed). During wet season, stratification parameter calculated from profiles during high tides showed values ranging from 1.3 to 1.9. This was due to the temporary stratification developed due to salt wedge intrusion. Once fresh water conditions prevailed and as the saline layer was pushed out of the estuary, stratification number became almost zero. The stratification parameter varies depending on the phase of tide and freshwater discharge indicating that Cochin estuary experiences a transition from partially mixed (neap) or well mixed (spring) in dry season to periodically stratified state during wet season.

3.1.5 Dissolved oxygen and nutrients

The distribution of chemical properties is shown in Figs. 4–7. During dry neap, the average surface DO value ($7.74 \pm 1.76 \text{ ml l}^{-1}$) was higher than bottom ($4.37 \pm 1.66 \text{ ml l}^{-1}$). In contrast, the average nutrient concentration (except for PO_4^{3-}) was higher at the bottom

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

than at the surface for neap phase during dry season; NO₂ (surface 0.55±0.41 μM; bottom 1.55 ± 0.21 μM), NO₃ (surface 1.79 ± 1.10 μM; bottom 4.10 ± 0.50 μM), PO₄³⁻ (surface 0.88 ± 0.30 μM; bottom 0.60 ± 0.13 μM) and SiO₄ (surface 12.02 ± 2.69 μM; bottom 15.73 ± 3.87 μM). Unlike neap, the water column property distributions were homogeneous over much of the tidal cycle in spring phase (Fig. 5c, e–h). During wet season, the incursion of hypoxic water (< 1.06 ml l⁻¹) through bottom layers (Figs. 6c, 7c) on flood tides was discernible during both spring and neap tidal phases. The near-bottom intruding water was characterised by high NO₂ (spring, 1.02 μM; neap, 1.84 μM), NO₃ (spring, 12.29 μM; neap, 17.5 μM), PO₄³⁻ (spring, 3.68 μM; neap, 3.24 μM) and low SiO₄ (spring, 39.65 μM; neap 38.03 μM) values. However, elevated levels of silicate (120.66 μM) which is a land derived nutrient were found at the surface. Silicate showed a negative relationship with salinity ($r^2 = 0.48$, $n = 31$) during wet season indicating that fresh water run-off is the principal source of silicate inputs. This is substantiated by the higher silicate concentrations in wet than during the low-run-off surveys.

3.1.6 Chlorophyll *a*

The distribution of chlorophyll *a* pigments which is a reliable measure of phytoplankton biomass is shown in Figs. 4–7. Seasonal variations in chlorophyll *a* distribution were observed with relatively higher values in dry seasons (on an average 12 mg m⁻³). The chlorophyll *a* concentration was higher at the surface with its peak (Figs. 4d, 5d) during 12:00 to 15:00 which may be attributed to the vertical migration of light-sensitive phytoplankton species to the surface. During dry season, the maximum surface chlorophyll *a* attained for neap and spring was 20.1 mg m⁻³ and 26.7 mg m⁻³, respectively. During salt wedge formation in both tidal phases of wet seasons (Figs. 6d, 7d), chlorophyll *a* maxima layer was formed at the bottom ocean water (salinity 33.29; chl *a* 11 mg m⁻³ (spring), salinity 33.04; chl *a* 6.8 mg m⁻³ (neap)) and minimum layer at the surface fresh water (salinity 0.05; chl *a* 1.1 mg m⁻³ (spring), salinity 0.01; chl *a* 0.9 mg m⁻³ (neap)).

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

3.2 Longitudinal salinity distributions (synoptic survey)

The results of the salinity intrusion survey for the months July 2008, August 2008, November 2008 and March 2009 are shown in Fig. 8. These months are characterised by south-west monsoon, north-east monsoon and dry period, respectively.

5 In the year 2008, with the onset of monsoon on 31 May, the river discharge peaked with 8.7 % of the total occurring in the month of June and flushing time calculated was 5.8 days. The depth averaged salinity of the southern arm stations (8, 9, 10) and of the stations (2, 3, 4) at the middle of northern arm presented low values averaging to 1.76-spring; 0.17-neap and 8.22-spring; 5.19-neap, respectively. Stations near Cochin
10 inlet (5, 6, 7) exhibited high salinity unlike the station 1 near northern inlet with tidal amplitudes having some influence. Stations 1, 2 are prone to fresh water influence due to Periyar river adjoining northern inlet which is the major river contributing to the total discharge of the estuary. Overall, the freshening of the estuary was initiated by the monsoon.

15 By the end of June, the monsoon peaked and July experienced the maximum fresh-water discharge amounting to 22.04 % of the total and flushing time decreased to 2.8 days. The salt wedge became prominent in July transforming the status of the estuary to that of a salt wedge type (Fig. 8a, b). A weaker salt wedge front originating from northern inlet, which was visible only in spring phase of July, persisted in all other monsoon months. Hence, higher salinities (18–34) were restricted to the bottom waters of the stations adjacent to the inlets (5, 6, 7 – Cochin inlet; 1, 2 – northern inlet). All the
20 other stations remained well mixed with depth averaged salinity reaching salinity values as low as 0.05.

25 The salt intrusion length during both spring and neap tides of the synoptic survey has been defined as the length from the river mouth (Cochin inlet) along the river channel to the point where the bottom salinity was 2 PSU. On an average, the intrusion length from the Cochin inlet to south averaged to 15 km and 11 km during spring and neap phases respectively of monsoon months. Distinctly, the results in August differed with

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

the intrusion greater in neap (15 km) than spring (13 km) since the neap observations were carried out during high tides whereas for spring it was during low tides (Fig. 8c, d). This entails the significance of the observation time with respect to tidal cycles denoting the relevance of time series measurements in the system. The depth of the interface (defined as the height of 2 PSU isohaline from the bottom) decreased gradually towards the outer end of the estuary. This depth which is also the thickness of the bottom saline water layer was greater in spring compared to neap phase at the mouth for the time series measurements (Figs. 6a, 7a) implying that saline intrusion was more during spring phase.

As the river discharge decreased from July to September (Fig. 2b), coincidentally the salt wedge advected further upstream suggesting an inverse relationship of salt intrusion with river flow. The withdrawal of monsoon concurred with the reduction in fresh water discharge. This resulted in the longitudinal dispersion of the salinity field. From 11.81 % in October, the river discharge was reduced to 3.34 % in December. Consequently, flushing time also increased from 3.5 days in October to 8.4 days in December. During the period, though the salinity at stations near Cochin inlet (5, 6, 7) were invariably high over the vertical, there was a consistent increase in the salinity at all other stations. However, the upper most station (10) remained far away from the influence of marine water irrespective of tidal phases. Later when the dry period (from January to April average discharge is only 1.4 % of total) commenced, the tidal actions dominated in the system. The flushing time was 9.3 days in January which increased to 13 days in April. The salinity field extended up to station 10 with maximum depth averaged salinity (15.12) attained in spring phase of March. In May, there was a slight increase in discharge to 2.5 % of the yearly discharge and the flushing time was 14.7 days. The aftermath of an anomalous rainfall in the catchment of Periyar during our spring observation caused station 1 at the northern inlet to be fresh water dominated.

3.3 Statistical analysis of monthly observation

The Pearson correlations between the surface values of ecologically important variables are shown in Table 1. Salinity showed significant positive correlation with P^H ($R = -0.524$, $P < 0.01$). Relevantly, it was negatively correlated with nitrate ($R = -0.308$, $P < 0.01$), DO ($R = -0.370$, $P < 0.01$) and silicate ($R = -0.650$, $P < 0.01$) indicating freshwater as their source of inputs. Chlorophyll *a* was highly correlated with phosphate ($R = 0.455$, $P < 0.01$), with no significant correlation with salinity. Also, there is no distinct correlation between salinity and phosphate. Although DO was inversely correlated with chlorophyll *a*, it was not significant indicating that biological processes are not the only factor influencing DO in the estuary. Negative correlations between salinity and DO were highly significant ($R = -0.370$, $P < 0.001$).

The two-way analysis of variance carried out on monthly samples is shown in Table 2. Salinity variations were found to be significant spatially ($F = 7.88$, $P < 0.001$) and temporarily ($F = 6.37$, $P < 0.001$). However, temperature varied more temporarily ($F = 2.29$, $P < 0.01$) than spatially ($F = 0.88$, $P < 0.01$) owing to the seasonal changes in the domain. Among the different nutrient species, only phosphate manifested fluctuations spatially ($F = 9.87$, $P < 0.001$) and temporarily ($F = 2.66$, $P < 0.005$). Chlorophyll *a* marked remarkably significant variations spatially ($F = 4.35$, $P < 0.001$) than temporarily ($F = 1.23$, $P < 0.5$).

4 Discussions and summary

Temperature has lesser influence in density stratification than salinity because the salinity range is larger than temperature in estuaries (Dyer, 1973). Thus, an attempt is made to analyze the variations in stratification considering salinity as the major determining factor. The data presented from this survey suggested that salinity fluctuates at different timescales, including intratidal, fortnightly of spring and neap tidal cycle, and seasonal wet and dry periods. It was evident from our study that Cochin estuary

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

experiences a transition from partially mixed (neap) or well mixed (spring) in dry season to periodically stratified state during monsoon.

The Arabian Sea, one of the major upwelling zones of the world, experiences upwelling from June to October (Banse, 1968; Naqvi et al., 2000). Srinivas et al. (2006) claims that the increased intensity of upwelling processes in off Cochin in July is instrumental in generating drop in sea level and surface temperature. Whilst this process aids in quicker flushing of the estuarine water through the bar mouth (Udayavarma, 1981), low tidal amplitudes and increasing number of oscillations in the south west coast may lead to small inter-tidal expanses, which reduce flushing (Qasim, 2003). Coastal upwelling in the Arabian Sea induces anomalies in the distribution of physiochemical properties characterized by a rapid decrease in temperature and DO concomitant with an increase in salinity and surface nutrients, particularly nitrates (Rotchord, 1975) which could foster enhanced primary productivity of the area (Mann and Lazier, 1996). The upwelled water area is situated just off the shelf break (Shetye et al., 1990), where the predominant tidal circulation is responsible in advecting these waters in to the estuaries. Significantly, the intruding water mass at the bottom layers during monsoon season was identified as upwelled water from the adjacent shelf with its peculiar characteristics of low temperature, high salinity, severely oxygen depleted, nutrient rich and high chlorophyll *a*.

The salt wedge plays an important role in the distribution of chemical and biological variables that have profound impacts on water quality (Haralambiduo et al., 2010). Coastal hypoxia (defined here as $< 1.42 \text{ ml l}^{-1}$) develops seasonally in many estuaries, fjords, and along open coasts as a result of natural upwelling (Levin et al., 2009). The upwelled water incursion and the stratification induced by fresh water discharge of monsoon season fueled the oxygen depletion at the bottom layer. Under persistent stratification, the probability of the shift of hypoxic to anoxic condition in the intruding frontal system cannot be ruled out. Our detailed hydrological analysis suggested that the dynamic and energetic environment of Cochin estuary hindered the sustainability of this front at the bottom. The onset of strong ebb phase and high flushing of monsoon

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



flushed the estuary from top to bottom with well oxygenated waters by pushing the saline wedge seaward. The duration of the existence of this front in our system was greatly determined by the tidal range and duration of flood-ebb cycles. The dependence of the salt wedge advancement on tidal range was clearly identifiable in spring phase as the tidal amplitude of 26.05 cm was not sufficient to force the transient stratified flow. The diurnal inequality of the neap phase contributed well to the flushing period (12 h) as when compared with the semi-diurnal spring phase (9 h 30 min). The time dependent nature of salt wedge has also been justified in Fraser estuary (Geyer et al., 1989), where during ebb the saline structure was eroded due to local enhancement of shear instability. Therefore, the periodic advance and retreat of the salinity wedge front is inevitable in preserving the ecosystem functioning and maintaining the health of the estuary.

Cochin estuary is reported to be a eutrophic estuary (Qasim et al., 2003). During high run-off surveys, an increase in the overall nutrient concentrations was noticed when compared to low run-off surveys. However, the salt wedge intrusion affected its distributions in the water column. Higher concentrations of phosphate and nitrite were restricted to the near bottom saline intrusions, but extreme levels of silicate and nitrate were observed in low salinities. Although the near-bottom seawater intruded with high nutrients, fresh water run-off was the principal source of silicate and nitrate supply. This is further substantiated by the statistical results which revealed that silicate and nitrate were highly negatively correlated with surface salinity whereas phosphate and nitrite which did not manifest a significant relationship. As the river flow weakened after monsoon, the flushing of the estuary diminished and the nitrite, nitrate, ammonia, phosphate and silicate loadings through anthropogenic activities (industries) and sediment re-suspension alter the nutrient stoichiometry substantially (Martin et al., 2008). Phosphorous showed both seasonal and spatial variability. The surface phosphate concentrations were moderately high in the stations with high salinity whereas concentration decreased in low saline regions. The previous work on the fractionation of phosphorous in Cochin estuary (Balchand et al., 1994) also concluded that enhanced amounts

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

of exchangeable P appeared in high saline waters, signifying the presence of biologically available nutrient phosphorus.

In Cochin estuary nutrients are not a limiting factor for the optimum phytoplankton growth at any time of the year (Balachandran et al., 2005) and also transient variations in the water quality play a significant role on phytoplankton behaviour (Madhu et al., 2009). Prior experimental studies had found that generally Cochin estuary exhibits high chlorophyll a when intermediate salinity (10–25) conditions prevail (Qasim and Sankaranarayanan, 1972). Our study substantiated the above statements such that increased chlorophyll a distribution was observed during dry season than wet season. The increased flushing during monsoon resulted in low chlorophyll concentration in surface layers where salinity was low. The effect of river discharge and surface salinity on the surface chlorophyll a distribution is clearly depicted in Fig. 2b such that the maximum chlorophyll concentrations were observed during lean-river flow months when the surface salinity was high. High riverine discharge leads to reduced residence time, leading to increased flushing of phytoplankton biomass out of the estuary (Lane et al., 2007). Further, the overall surface distribution of chlorophyll a during monsoon was affected when the increased river discharge restricted the salinity dispersion and hence the saline waters with biologically available phosphorous was limited in some areas along the estuary. Phosphorous is being attributed as a limiting nutrient in fresh water-dominated systems (Neil, 2005). The river pulses impede the salinity field expansion affecting the nutrient distributions. However, salinity could not be considered as the only determining factor for triggering phytoplankton growth. Unravelling these evidences is henceforth a major task to be accomplished through sustained research. Although the linking between temporal stratification of the water column with chlorophyll a distributions is a question of some subtlety, this study reveals the persuasive evidence of its critical importance in Cochin estuary.

The advancement in our understanding of time-dependent nature of water column stabilization and destabilization mechanisms can be achieved through continuous

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

monitoring of multiple stations along with supporting flow field to define various processes in spatial and time scales.

Acknowledgement. This study forms a part of project “Ecosystem modeling of Cochin estuary” funded by Integrated Coastal and Marine Area Management (ICMAM). We are indebted to the project staff in the small dinghies, who had overcome the adverse weather conditions for sample collection. We thank the Director of NIO-Goa, Scientist In-charge, NIO-Regional centre, Cochin for the support and encouragement provided towards the completion of this work. Shivaprasad Amaravayal is also grateful to CSIR, New Delhi, for financial support in the form of Senior Research fellowship. This work forms part of project “Ecosystem modeling of Cochin estuary” funded by ICMAM. This is NIO contribution no. XXXX.

References

- Balachandran, K. K., Laluraj, C. M., Nair M., Joseph, T., Sheeba, P., and Venugopal, P.: Heavy metal deposition in a flow-restricted, tropical estuary, *Estuar Coast Shelf S.*, 65, 361–370, 2005.
- Balachandran, K. K., Reddy, G. S., Revichandran, C., Srinivas, K., Vijayan, P. R., and Thottam, T. J.: Modelling of tidal hydrodynamics for a tropical ecosystem with implications for pollutant dispersion (Cochin estuary, Southwest India), *Ocean Dynam.*, 58, 259–273, 2008.
- Balchand, A. N. and Nair, S. M.: Fractionation of phosphorous in the sediments of a tropical estuary, *Environ Geol.*, 23, 284–294, 1994.
- Banse, K.: Hydrography of the Arabian Sea shelf of India and Pakistan and effects on demersal fishes, *Deep Sea Res.*, 15, 45–79, 1968.
- Blanton, J.: Energy dissipation in a tidal estuary, *J. Geophys. Res.*, 74, 5460–5466, 1969.
- Dinesh Kumar, P. K., Sankaranarayanan, V. N., and Saraladevi, K.: An introduction to the system, prior studies, historical trends and future implications, *Indian J. Environ. Protect.*, 14, 98–102, 1994.
- Dyer, K. R.: *Estuaries: a Physical Introduction*, 1st Edn., Wiley-Interscience, London, 1973.
- Dyer, K. R.: *Estuaries: a Physical Introduction*, 2nd Edn., John Wiley and Sons, Sydney, 1997.
- Geyer, W. R. and David, M. F.: Tide-induced variation of the dynamics of a salt wedge estuary, *J. Phys. Oceanogr.*, 19, 1060–1072, 1989.

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Government of Kerala General Features: available at: <http://en.wikipedia.org/wiki/Kerala#Climate> (last access: 23 July 2012), 2005.

Grasshoff, K., Ehrhardt, M., and Kremling, K.: *Methods of Seawater Analysis*, Verlag Chemie, Weinheim, p. 419, 1983.

5 Haralambidou, K., Sylaios, G., and Tsihrintzis, V. A.: Salt-wedge propagation in Mediterranean micro-tidal river mouth, *Estuar. Coast Shelf S.*, 90, 174–184, 2010.

Joseph, S. and Ouseph, P. P.: Assessment of nutrients using multivariate statistical techniques in estuarine systems and its management implications: a case study from Cochin estuary, India, *Water Environ. J.*, 24, 126–132, 2010.

10 Kasai, A., Kurikawa, Y., Ueno, M., Rober, D., and Yamashita, Y.: Salt-wedge intrusion of seawater and its implication for phytoplankton dynamics in the Yura Estuary, Japan, *Estuar. Coast Shelf S.*, 86, 408–414, 2010.

Ketchum, B. H.: *Estuaries and enclosed seas*, Elsevier Science Publication Comp, Amsterdam, p. 15, 1983.

15 Krishnakumar, K. N., Prasada Rao, G. S. L. H. V., and Gopakumar, C. S.: Rainfall trends in twentieth century over Kerala, India, *Atmos Environ.*, 43, 1940–1944, 2009.

Lane, R., Day, J., Marx, B. D., Reyes, E., Hyfield, E., and Day, J. N.: The effects of riverine discharge on temperature, salinity, suspended sediment and chlorophyll *a* in a Mississippi delta estuary measured using a flow-through system, *Estuar. Coast Shelf S.*, 74, 145–154, 2007.

20 Levin, L. A., Ekau, W., Gooday, A. J., Jorissen, F., Middelburg, J. J., Naqvi, S. W. A., Neira, C., Rabalais, N. N., and Zhang, J.: Effects of natural and human-induced hypoxia on coastal benthos, *Biogeosciences*, 6, 2063–2098, doi:10.5194/bg-6-2063-2009, 2009.

Lund-Hansen, L. C., Skyum, P., and Christiansen, C.: Modes of stratification in a semi-enclosed bay at the North Sea-Baltic Sea transition, *Estuar. Coast Shelf S.*, 42, 45–54, 1996.

25 Madhu, N. V., Balachandran, K. K., Martin, G. D., Jyothibabu, R., Thottathil, S. D., Nair, M., Joseph, T., and Kusum, K. K.: Short term variability of water quality and its implications on phytoplankton production in a tropical estuary (Cochin estuary – India), *Environ. Monit. Asses.*, 170, 287–300, doi:10.1007/s10661-009-1232-y, 2010.

30 Mann, K. H. and Lazier, J. R. N.: *Vertical Structure of the Open Ocean: Biology of the Mixed Layer. Dynamics of Marine Ecosystems-Biological-Physical Interactions in the Oceans*, 2nd Edn., Blackwell Science, Cambridge, MA, 56–97, 1996.

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Menon, N. N., Balchand, A. N., and Menon, N. R.: Hydrobiology of the Cochin estuary system a review, *Hydrobiologia*, 43, 149–183, 2000.
- Martin, G. D., Vijay, V. J., Laluraj, C. M., Madhu, N. V., Joseph, T., Nair, M., Gupta, G. V. M., and Balachandran, K. K.: Fresh water influence on nutrient stoichiometry in a tropical estuary, south west coast of India, *Appl. Ecol. Environ. Res.*, 6, 57–64, 2008.
- Naqvi, S. W. A., Jayakumar, D. A., Narvekar, P. V., Naik, H., Sarma, V. S., D'Souza, W., Joseph, T., and George, M. D.: Increased marine production of N₂O due to intensifying anoxia on the Indian continental shelf, *Nature*, 408, 346–349, 2000.
- Neil, M.: A method to determine which nutrient is limiting for plant growth in estuarine waters at any salinity, *Marine Pollut. Bull.*, 50, 945–955, 2005.
- Nepf, H. M. and Geyer, W. R.: Intratidal variations in stratification and mixing in the Hudson estuary, *J. Geophys. Res.*, 101, 12079–12086, 1996.
- Nunes Vaz, R. A., Lennon G. W., and De Silva Samarasinghe, J. R.: The negative role of turbulence in estuarine mass transport, *Estuar. Coast Shelf S.*, 28, 361–377, 1989.
- Qasim, S. Z.: *Indian Estuaries*, Allied Publication Pvt. Ltd., Heredia Marg, Ballard Estate, Mumbai, p. 259, 2003.
- Qasim, S. Z. and Gopinathan, C. K.: Tidal cycles and the environmental features of Cochin backwater (A tropical estuary), *Proceedings of the Indian Academy of Sciences*, Bangalore, 1969.
- Qasim, S. Z. and Sankaranarayanan, V. N.: Organic detritus of a tropical estuary, *Mar. Biol.*, 15, 193–199, 1972.
- Ranasinghe, R. and Pattiaratchi, C.: Circulation and mixing characteristics of a seasonally open tidal inlet: a field study, *Aust. J. Mar. Freshwater Res.*, 50, 281–290, 1999.
- Rippeth, T. P. and Simpson, J. H.: The frequency and duration of episodes of complete vertical mixing in the Clyde Sea, *Cont. Shelf Res.*, 16, 933–947, 1996.
- Rotchord, D. J.: Nutrient environment of Australian coastal waters, *Aust. J. Mar. Freshwater Res.*, 26, 223–243, 1975.
- Shetye, S. R., Gouveia, A., Shenoi, S. S. C., Sundar, D., Michael, G. S., Almeida, A. M., and Shatanam K.: Hydrography and circulation off the west coast of India during the southwest monsoon, *J. Marine Res.*, 48, 359–378, 1990.
- Simpson, J. H.: The shelf-sea fronts: implications of their existence and behaviour, *Philos. T. Roy. Soc. Lond.*, 302, 531–546, 1981.

- Srinivas, K. and Dinesh Kumar, P. K.: Atmospheric forcing on the seasonal variability of sea level at Cochin, southwest coast of India, *Cont. Shelf Res.*, 26, 1113–1133, 2006.
- Srinivas, K., Revichandran, C., Thottam, T. J., Maheswaran, P. A., Asharaf, M., and Murukesh, N.: Currents in the Cochin estuarine system during March 2000 (Southwest coast of India), *Indian J. Mar. Sci.*, 32, 123–132, 2003a.
- 5 Srinivas, K., Revichandran, C., Maheswaran, P. A., Asharaf, M., and Murukesh, N.: Propagation of tides in the Cochin estuarine system, southwest coast of India, *Indian J. Mar. Sci.* 32, 14–24, 2003b.
- 10 Strickland, J. D. H. and Parsons, T. R.: A Practical Handbook of Seawater Analysis, *Bull. Fish. Res. Board Canada*, 2, 167–310, 1972.
- Udaya Varma, P., Pylee, A., and Rama Raju, V. S.: Tidal influence on the seasonal variation in current and salinity around Willington Island, *Mahasagar*, 4, 225–237, 1981.
- Uncles, R. J., Ong, J. E., and Gong, W. K.: Observations and analysis of a stratification-destratification event in a tropical estuary, *Estuar. Coast Shelf S.*, 31, 651–665, 1990.

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.

Table 1. Results of Pearson correlation analysis (R) for the investigated physical and chemical variables during monthly observation.

Parameters	Salinity	Temp	DO	pH	Nitrate	Nitrite	Ammonia	Silicate	Phosphate	Chl <i>a</i>
Salinity	1									
Temp	0.028	1								
DO	-0.370**	0.375**	1							
pH	0.524**	0.690**	0.203*	1						
Nitrate	-0.308**	-0.027	0.173	-0.122	1					
Nitrite	0.106	0.059	-0.209*	-0.017	0.139	1				
Ammonia	-0.027	0.054	-0.092	0.064	-0.068	-0.204*	1			
Silicate	-0.650**	0.168	0.290**	-0.231*	0.251**	-0.186*	0.247**	1		
Phosphate	0.117	0.117	-0.127	0.168	0.125	-0.027	0.055	0.032	1	
Chl <i>a</i>	0.086	0.158	-0.038	0.16	0.014	-0.038	-0.048	-0.029	0.455**	1

* $P < 0.05$, ** $P < 0.01$, bold values indicate negative correlation.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.

Table 2. Two-way analysis of variance for differences of ecologically important parameters on sampling stations and sampling months for the survey (June 2008 to May 2009). *F* indicates the likelihood ratio; *P* indicates the probability. *** indicates statistically significant differences > 0.0001, ** indicates < 0.001 and * indicates < 0.01.

Variables	Station		Year	
	<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>
Salinity	7.883679	0.000065***	6.370352	0.00047**
Temperature	0.884074	0.541915	2.291627	0.014651
DO	3.411882	0.000984**	2.104926	0.025726
pH	3.836676	0.000301**	3.082916	0.001237*
Nitrate	0.419885	0.922091	12.94469	0.00002***
Nitrite	0.971728	0.4674	9.793611	0.00042**
Ammonia	1.006034	0.439704	8.134114	0.00032**
Silicate	1.843739	0.068204	7.809754	0.00008***
Phosphate	9.872694	0.0005**	2.662658	0.004654*
Chl <i>a</i>	4.357485	0.00007***	1.233476	0.273961

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 1. The Cochin estuary (west coast, India), showing stations and extent of backwaters, having two inlets to Arabian Sea Munambam (north) and Cochin inlet (middle of the extent of back waters). The time series station which is located 5 km way from Cochin inlet. Synoptic and monthly stations are discerningly marked in the backwaters.

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.




Fig. 2. (a) Climatology of rainfall (1978–2002) in the catchment of Cochin back waters. (b) Monthly river discharge, surface salinity, surface chlorophyll *a* starting from June 2008 and finishing May 2009.

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.




Fig. 3. (a–d) Semi-Diurnal variations of observed water level and predicted tide at the location. **(e–f)** Time series of potential energy anomaly (ϕ) (J m^{-3}) computed from the 30 min density profiles **(a, e)** 25 April 09:00 a.m. to 26 April 03:00 p.m. (neap-dry); **(b, f)** 2 May 09:00 a.m. to 3 May 03:00 p.m. (spring-dry); **(c, g)** 16 July 12:00 a.m. to 17 July 03:00 p.m. (spring-wet); **(d, h)** 24 July 12:00 a.m. to 25 July 03:00 p.m. (neap-wet).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.




Fig. 4. Depth-time contours of salinity, temperature DO, chlorophyll *a*, NO₂⁻, NO₃⁻, PO₄³⁻, SiO₄⁴⁻ for every three hours from surface, mid-depth and bottom during neap tides of dry season.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.




Fig. 5. Depth-time contours of salinity, temperature, DO, chlorophyll *a*, NO_2^- , NO_3^- , PO_4^{3-} , SiO_4^{4-} for every three hours from surface, mid-depth and bottom during spring tide of dry season.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.




Fig. 6. Depth-time contours of salinity, temperature DO, chlorophyll *a*, NO₂⁻, NO₃⁻, PO₄³⁻, SiO₄⁴⁻ for every three hours from surface, mid-depth and bottom during spring tide of wet season.

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.




Fig. 7. Depth-time contours of salinity, temperature DO, chlorophyll *a*, NO₂⁻, NO₃⁻, PO₄³⁻, SiO₄⁴⁻ for every three hours from surface, mid-depth and bottom during neap tide of wet season.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Seasonal stratification and property distributions in a tropical estuary

A. Shivaprasad et al.




Fig. 8. Longitudinal synoptic distribution of salinity measured twice monthly (during spring and neap tides) During July 2008, August 2008, November 2008 and March 2009. The Cochin inlet is at the coordinate origin. The northern/southern arm stations are at positive/negative distances, respectively. The insets show the tidal amplitude and the times (as X's) when each survey began and ended. Times of each station appear along the lower x-axis.