

Tree ring-based reconstruction of October to November runoffs in the Jiaolai River since 1826

L. Ma^{1,*}, T.X.Liu¹, Q.Wu²,L.H.Ji¹, R.Z.Gao¹

1. Water Conservancy and Civil Engineering College of Inner Mongolia Agricultural University, Hohhot, Inner Mongolia, 010018, China;

2. Land Surveying and Planning Institute of Inner Mongolia, Hohhot, Inner Mongolia, 010010, China;

*Corresponding author: malong4444333@163.com

Abstract

The Horqin Sandy Land is a typical desertification region in China hounded by ecological and environmental problems; ~~that which continue to~~ affect economic and social development. ~~Hence,~~ hydrological climate changes in this region need to be investigated. The current study reconstructed the runoff sequences ~~from in~~ the southwest edge of the LiaoHe River ~~into~~ the XiaWa station of the JiaoLai River during the months of October to November from 1826 to 2005. A comprehensive timeline for the regional tree wheel width of the Horqin Sandy Land was employed.

The timeline has been ~~in-used~~ for 183 years. For the past 180 years, the runoff has experienced six and four consecutive Feng and dry sections, respectively. From 1982 to 2005, the runoff reached the longest section of a continuous low-flow runoff, with ~~a~~ ~~the~~ mean average runoff amounting to only 63.58% of ~~that for~~ the entire period. ~~The~~ ~~R~~unoff has 3-, 11-, 15-, 24-, and 30-year quasi-periodic variations, consistent with changes in similar areas worldwide. ~~The change was gentler from~~ ~~The period of~~ 1826 to 1917 ~~presents a more gentle change~~. In 1956, the runoff increased; and then

significantly decreased for nearly 50 years. The drop rate is 1.7766 million m³/10 years, which shows a consistent downward trend with the precipitation (14.74 mm/10 years). The overall reduction in precipitation accounts for 29.86% of the initial value, which is far less than 75.58% ~~for of~~ the runoff. If the runoff and precipitation drops continue, more extensive and longer ecological and environmental problems are foreseen to occur.

Keywords: Northeast China; Horqin Sandy Land; Jiaolai River; Runoff; Response analysis; Sequence reconstruction

1. Introduction

China is one of several countries experiencing severe desertification. The rapid progression of dry land degradation has become an important ecological and socio-economic problem (Wang and Zhu, et al., 2002; Wang and Zhu, 2001). The Horqin Sandy Land is located between east longitude 117° 45' to 124° 06' and north latitude 42° 36' to 45° 20' (Figure 1). It belongs to the alluvial plain of the Liaohe River. In ancient times, the Horqin Sandy Land was a prairieland with lush vegetation ~~and beautiful sceneries~~. Now, it has become a typical Chinese desertification area, one of ~~the China's~~ four sandy lands ~~of China~~. Related research on the environmental changes in this area ~~has begun~~ in the 1950s, producing some substantive results (Wang and Zhu, 2001; Dong and Jin, et al., 1998; Wang and Wu, et al., 2004; Wang and Chen, et al., 2006; Li and Han, et al., 2007; Zhao and Zhou, et al., 2008) on the desertification process, evolution, structure, and drive mechanism. The results suggest that climate fluctuations could directly affect to some extent the

process of desertification via ~~through the~~ different periods of precipitation and temperature combinations (Wang and Wu, et al., 2004; Zhao and Zhou, et al., 2008).

The desertification is mainly subjected to millennial- and centennial-scale climate fluctuations (Dong and Jin, et al., 1998). Hence, a long-time-scale research on hydrological climate changes is particularly important.

Runoff and precipitation are closely related. The amount of runoff not only directly affects river ecology, but also has a profound impact on changes in river environments.

Consequently, the future trends ~~of~~ the ecological environment and past runoff variations need to be explored.

A number of studies have been conducted on runoffs in the Liaohe River. This river is located in the Horqin Sandy Land (Zhang and Wang, et al., 2007; Hao and Wang, et al., 2008; Yang and Liu, et al., 2009; Fang and Ren, et al., 2009; Jiang and Ren, et al., 2009; Wang and Jin, et al., 2011; Zhang and He, 2011; Gu and Wang, et al., 2011),

and the study periods ~~s~~ covers the years 1950s to 2008. ~~These works have research~~ involved the Liaohe River, ~~the~~ Liaohe tributaries, and other tributaries of the Laoha River.

Zhang et al. (2007) have shown that the measured runoff in the Liaohe Laoha River has significantly decreased since 1950. Fang et al. (2009) have revealed that

from the 1960s to the early 21st century, the variations in annual runoff and ~~the~~ annual runoff coefficient of the Laoha River ~~are were~~ more-less-less-more-less. The

decadal variation is much greater than the annual rainfall, and Wang et al. (2011) have ~~indicated pointed out~~ that the measured runoff in Liaohe Tieling show~~ed~~ a significant

reduction trend since the mid-1960s.

However, these studies are mainly based on ~~visual observational~~ data from hydrological observation stations, and the data life is generally 50 years. These data are clearly insufficient for studying longer-time-scale changes in runoff. ~~The~~ ~~h~~Hydro-climatic information revealed by tree-ring records is an effective means to address this limitation. Tree ring-based reconstructions present information that feature accurate positioning, good continuity, high resolution, and precise correlation with hydrological climate change (Fritts, 1976; Shao, 1997; Nicoletta, 2004). Such reconstructions are applied in research on hydro-climatic events abroad, such as runoff, drought, flood, rainfall, temperature, glaciation, and volcanic activity (Stockton and Meko, 1975; Hughes and Leggett, et al., 1978; Cook and Jacoby, 1983; Briffa and Jones, et al., 1995; Van and Stahle, et al., 1998; Clevel, 2000; Magda and Zelenova, et al., 2001; Meko and Therrell, et al., 2001; Woodhouse, 2001; Esper and Cook, et al., 2003; Law and Turner, et al., 2006; Michael and Loso, et al., 2006; Joseph and Mc, et al., 2006; Sophie and Heike, et al., 2007; Giovanna and David, et al., 2010; Samuli and Heikki, et al., 2010).

In China, ~~to study the chronology established, the tree ring and hydrological climate relationship, as well as the reconstruction,~~ the study objects of Sabina, spruce, pine, white stick, larch, and elm in cold and dry regions are examined to study the established chronology, relationship of the tree ring with the hydrological climate, and reconstruction~~were selected~~. Such regions included the Qinghai-Tibet Plateau, Qaidam Basin, Xinjiang, Qilian Mountain, Changbai Mountain, Qinling, and Inner Mongolia. The research has made great progress (Wu and Sun, et al., 1989; Shao and

Huang, et al., 2004; Liu and Qin, et al., 2004; Gou and Yang, et al., 2006; Ma and Liu, 2007; Shao and Wang, et al., 2007; Zhu and Zheng, et al., 2008; Eryuan and Shao, et al., 2008; Liu and Tian, et al., 2009; Li and Feng, et al., 2010; Liu and Sun, et al., 2010; Ma and Liu, et al., 2011). Part of the runoff is reconstructed in the Urumqi mountain basin, Black River, Tongtianhe, Huangshui, and Yellow River (Li and Yuan, et al., 1997; Kang and Chen, et al., 2002; Qin and Jin, et al., 2004; Wang and Chen, et al., 2004; Liu and Yang, et al., 2006; Kang and Wang, et al., 2008; Liu and Chen, et al., 2007; Sun and Liu, et al., 2011). The long sequence of changes in runoff characteristics ~~isare~~ analyzed.

Regardless of short time changes in runoff characteristics and the reconstruction of a long sequence, studies on the JiaoLai River are relatively limited. In the current paper, the 183-year Horqin sandy area elm tree ring width chronologies established by Ma et al. (2011) in a station in the JiaoLai River tributary during the months of October to November from 1826 to 2005 ~~were~~was used to study the runoff characteristics. The results provide some basic information on long-term changes in the Liaohe source runoff, ecological and environmental protection, as well as catchment economy progress.

Figure 1

2. Data used

2.1 Chronological data

The ring width chronology and reconstruction of elm in the Sandy Land were used (Ma and Liu, 2011). The ring width chronology ~~from for~~ 1826 to 2008 (183 years) is shown in Figure 2 (Ma and Liu, 2011).

Figure 2

2.2 Hydrological and meteorological data

The precipitation data selected were in accordance with the chronology of the establishment of meteorological stations. The 1951 to 2010 monthly (yearly) precipitation data were obtained from the eight meteorological stations closest to the sites. These stations were the Horqin Left Back Banner, TongLiao, Kailu, Horqin Left Middle Banner, Jarud Qi, Horqin Right Middle Banner, and Horqin Left Right Banner. Data from meteorological administration-integrated materials were also obtained. The uniformity of every meteorological element indicated that the precipitation data from the sites did not exhibit significant deviations as well as random changes, and that the data changes were relatively homogenous and consistent. Hence, the data were considered as reliable representations of the climate conditions in the region. Data from adjacent sites were used to interpolate measurements because some of the stations had incomplete information. The average data of the eight stations were used for area face value.

According to the same basin and proximity principle, the monthly (yearly) runoff data from 1957 to 2010 of the Xiawa station in the JiaoLai River were collected. This station is located at the edge of the Horqin Sandy Land and the import point to the

批注 [Ed1114 1]: Clarify this.

sand of the JiaoLai River. There is ~~low~~less human activity upstream, so the measured runoff data ~~were~~was closer to the natural values.

The ground precipitation in the Horqin Sandy Land is basically the same with ~~that~~the precipitation in the upstream watershed area. By combining all factors, the selected precipitation sites and hydrological stations were used in the present study.

3. Related analysis and explanation

3.1 Correlation of chronology and runoff

Tree ring climatology considers that rings formed in the year t are influenced not only by the climate at that time, but also by the climate of the previous ~~one~~ to ~~two~~ years (Fritts, 1976). Therefore, the standard chronological sequence and standard chronological $t + 1$, $t + 2$, and $t + 3$ sequences ~~in~~relat~~ed~~ion to the runoff in low-lying stations, as well as the correlation coefficients are shown in Table 1. The standard chronology had a significant relationship with the runoff in March, April, August, September, October, and November, as well as the annual runoff. The correlation in October and November was the highest.

Table 1

Figure 3

Table 2

The results of the correlation analysis between the annual and monthly runoffs show~~ed~~ that the runoff in July accounted for 30.19% of the annual runoff, and had the maximum correlation coefficient of 0.814. Although the runoff~~s~~ in Oct~~ober~~ and

November- were responsible for only 4.82% and 3.63% of the annual runoff, the annual runoff coefficient ranked only second to July; the coefficients were 0.755 and 0.699, respectively, ranking second and third. Based on the contrast curve among the annual runoff, runoff in October-, runoff in November-, and runoff in October- to November- (Figure 3), the four curves were concluded to have basically the same trend. Therefore, using a rebuilding chronology for the October to ~~November~~- runoff to analyze wet and dry season changes, cycle changes, which can represent the changes in annual runoff to some degree, were constructed. The reconstruction can also be carried out in August-, considering the better correlation coefficients in October to ~~November~~-, which has a more significant meaning in reconstruction.

3.2 Physiological mechanism explanation on the relationship between trees and runoff

The main supply source of runoff formation is precipitation. In different seasons ~~and~~ at different times, ~~the~~ groundwater comes from direct precipitation or snowmelt, and then supplies the runoff. According to the correlation between the precipitation sequence in the Horqin Sand Land and the runoff series in the XiaWa station (Table 2), the precipitation in July, August-, and September-, as well as the annual precipitation significantly correlated with the March to April, July to November-, and annual runoffs. Among them, the precipitation in July had a significant correlation with the March to April, July to November-, and annual runoffs, respectively. The runoff in July accounted for 30.19% of the annual runoff, whereas ~~the~~ precipitation in July was 32.18% of the total annual precipitation, which was the largest contribution

to the runoff. ~~The R~~rainfall in ~~August-~~ was significantly related to the ~~August-~~ to ~~October-~~ runoff, and the ~~August-~~ runoff accounted for 24% of the annual runoff. On the other hand, the precipitation in ~~August-~~ accounted for 21.46% of the annual precipitation, and ranked ~~eds~~ second in terms of runoff contribution. The precipitation in ~~September-~~ had a significant correlation with the ~~September-~~ to ~~November-~~ runoff, and the runoff in ~~September-~~ was 7.51% of the annual runoff. ~~ByIn~~ contrast, precipitation in ~~September-~~ accounted for 8.84% ~~ofin~~ the total annual precipitation. ~~The A~~annual precipitation was significantly related to the March to April, July to ~~November-~~, and annual runoffs. When precipitation transforms, some of it directly forms surface runoff; and some permeates the ground to supply surface runoff in the underground runoff form. Based on a combination of the above relationship, the formation of precipitation runoff can be concluded to have some lag. Therefore, the precipitation in July, ~~August-~~, ~~andas well as~~ ~~September-~~, was greater, and accounted for a significant proportion ~~ofin~~ the annual precipitation. The precipitation not only plays an important role in the formation of monthly surface runoff, but also affects the runoff information in ~~October-~~ to ~~NovNovember-~~ to some degree.

Precipitation also plays an important role in the growth of trees, especially the precipitation in July, ~~Aug-August~~, and ~~September~~, which were significantly correlated with the chronology (Ma and Liu, et al., 2011). Summer was the peak of elm growth, ~~and- Aat theseat months-time~~, elm growth mostly depended ~~most~~ on water. Good water conditions during this period effectively and rapidly promoted elm growth. This phenomenon was the same in ~~Sep-September~~

Considering the comprehensive situations in Tables 1 and 2, Figure 4 shows that precipitation played a major role in runoff formation and tree growth. In July, ~~Aug-August~~, and ~~Sep-September~~, the precipitation ~~not only~~ played an important role ~~not only~~ on tree growth, but also on runoff formation. Changes in the runoff also represented changes in the precipitation. Considering the lag effect, the precipitation in July, ~~Aug-August~~, and ~~Sep-September~~ had a significant impact on the runoff in ~~Oct-October~~ to ~~Nov-November~~. This phenomenon explained the good relationship between the chronology and the ~~Aug-August~~ to ~~Nov-November~~ runoff. The good relationship among the chronology sequences in years $t + 1$, $t + 2$, and $t + 3$ with the runoff also ~~completely fully~~ illustrated the delayed runoff response. ←

Figure 4

4. Reconstruction runoff

From the above response relationship, the standard sequence and standard chronology in years $t + 1$, $t + 2$, and $t + 3$ were used to reconstruct runoff in the XiaWa station of the JiaoLai River from ~~Oct-October~~ to ~~Nov-November~~ ~~from since~~ 1826 to 2005. The following equation was adopted:

$$R_t = -800.79 + 438.43 \times I_t + 98.01 \times I_{t+1} + 231.54 \times I_{t+2} + 818.15 \times I_{t+3}$$

$$-(N = 49, r = 0.74, R^2 = 0.54, R_{adj}^2 = 0.51, F(4,44) = 13.05, P < 0.001)$$

—where R_t is the runoff from ~~Oct-October~~ to ~~Nov-November~~ in year t , I_t denotes the index of standard tree ring chronology in year t (dimensionless), I_{t+1} is the index of standard tree ring chronology in year $t + 1$ (dimensionless), I_{t+2} is the index of

standard tree ring chronology in year $t + 2$ (dimensionless), and I_{t+3} is the index of standard tree ring chronology in year $t + 3$ (dimensionless).

Compared with the reconstructed sequence exhibiting the same trend as the corresponding sequence of the measured value (~~shown in~~ Figure 5), some extreme values cannot be completely consistent with the reconstructed values. In other words, part of the tree ring reconstruction results underestimates extreme hydrologically and climate events (Fritts, 1976). However, the reconstruction equation yielded good results for the reconstruction of the average minimum winter temperature. The reconstructed and measured values demonstrated good synchronization.

Figure 5

Additional tests on the stability and reliability of the reconstruction equation ~~are~~ necessary to ensure the credibility of the reconstruction value when a value generated beyond the calibration period ~~was~~ used. According to common international practice, the reduction error RE , S_1 (sign test), S_2 (first difference symbols for the test), and t (average test value for the product) were calculated. An RE of 0.54, ~~which was obviously greater than 0.3~~, was obtained. When ~~the~~ RE is higher than 0.3, the reconstruction value is considered credible (Li and Yuan et al., 2000). The values of the sign test results and first-difference symbols for the test results were 33/49 and 31/48, respectively. These ~~values~~ were significant at the 0.05 level, indicating that the reconstructed and measured value sequences were in good agreement with the changes in high-to-low frequencies. The average test value for the product was 2.91, which was significant at the 0.01 level, indicating a significant difference between the identical

and opposing serial number sequences. All parameters indicated that the reconstruction equation was stable and reliable. The reconstruction results for the runoff derived from this equation were also credible.

Consequently, the runoff series for the XiaWa Station of the JiaoLai River in ~~Oct.~~October to ~~Nov.~~November from 1826 to 2006 was rebuilt according to the reconstruction equation. The reconstruction sequence is shown in Figure 5.

5. Characteristics of runoff

5.1 Changes in wet and dry seasons

According to the reconstructed runoff series, the average runoff in the XiaWa station of the JiaoLai River in ~~Oct.~~October to ~~Nov.~~November from 1826 to 2006 was 9.7238 million m^3 . ~~with~~ ~~the~~ ~~maximum~~ ~~value~~ ~~of~~ ~~being~~ 18.5171 million m^3 ~~was~~ ~~found~~ ~~(occurred~~ ~~in~~ ~~the~~ ~~year~~ ~~1829)~~. ~~and~~ ~~the~~ ~~minimum~~ ~~value~~ ~~of~~ ~~was~~ 2.8576 million m^3 ~~was~~ ~~found~~ ~~(occurred~~ ~~in~~ ~~the~~ ~~year~~ ~~2005)~~. The extreme value was 6.48, which illustrated that the runoff in the JiaoLai River had greater amplitudes between years.

Reconstructing the sequence in mean runoff changes can reflect wet and dry season changes between years. The mean changes are shown in Table 3. These data showed the wet periods were more obvious in the 1820s, 1850s, 1900s, and 1950s during the era of scale changes, whereas the dry periods were remarkable in the 1880s, 1910s, and 1970s ~~to~~ ~~2000s~~.

Table 3

For further quantitative analysis of the reconstructed sequences in wet and dry season

changes, the coefficient ratio κ_p , which is the ratio of the runoff to the mean runoff for many years, was calculated. The following definitions were made for different κ_p values: more than 1.16, special wet years; between 1.06 and 1.16, partial wet years; between 0.95 and 1.06, flat water years; between 0.84 and 0.95, partial dry years; and less than 0.84, special dry years. The reconstruction sequence is shown in Figure 6.

This sequence indicated that the years of reconstructed sequences in the special wet, partial wet, flat water, partial dry, and special dry years were 40, 23, 40, 30, and 47, accounting for 22.22%, 12.78%, 22.22%, 16.67%, and 26.11% of the total number of years, respectively. The proportions of the special wet, flat water, and special dry years were large, with the ~~largest being that of~~ special dry years being the largest. This result revealed that the runoff in the JiaoLai River had a negative contribution to the reconstruction period.

According to the principle of at least ~~five~~ continuous years, the special and partial wet years were combined with the wet years, whereas the special and partial dry years were combined with the dry years. The results are shown in Figure 6. The consecutive wet years in the reconstruction sequences were 1826 to 1834, 1856 to 1861, 1872 to 1877, 1897 to 1905, 1938 to 1943, and 1946 to 1960. The durations ranged from 6 years to 15 years, and the total was 51 years. A total of 24 out of ~~Of~~ these 51 years, ~~24~~ occurred in the 19th century, and the remaining 27 occurred in the 20th century (mainly in the 1960s). The longest wet period occurred from 1946 to 1960, ~~a total of 15 years~~, with the mean annual runoff at 12.9962 million m^3 , which was 1.33 times the average runoff for the entire reconstruction period. There were four periods of continuous dry

years, namely, 1835 to 1840, 1880 to 1885, 1912 to 1922, and 1982 to 2005. The durations ranged from 6 years to 24 years, and the total was 47 years. Only 12 out of these 47 years, only 12 occurred in the 19th century, and the remaining 30 occurred in the 20th century (mainly from the 1980s to the late 20th century). The five years in the early 21st century were continuous dry years (from 1982 to 2005). The continued dry time reached 24 years, and the mean annual runoff was 6.1823 million m³, which was only 63.58% of the average runoff for the entire reconstruction period. The continuous dry periods occurred not only in the 1980s, but also from the early 1960s to 2005. The overall runoff trend was decreasing. From During the 1950s to the 2000s, the runoff decreased from 14.1031 million m³ to 3.4435 million m³, with a drop rate of 1.7766 million m³/10 years. The runoff significantly declined, which proved fully illustrated that the runoff in the JiaoLai River washad-been gradually decreasing for nearly 50 years, thereby forming the longest continuous dry season during the reconstruction period.

Figure 6

Figure 7

5.2 Periodic changes

An analysis of the runoff reconstruction with the power spectrum revealed that the runoff reconstruction sequence had 3-, 11-, 15-, 24-, and 30-year quasi-periodic variations under the 0.05 significance level. This result almost had basically the same cycle as the precipitation changes in the Horqin Sandy Land. This similarity signified that precipitation played an important role in the runoff formation.

5.3 Variation trends

The runoff sequence was reconstructed for a 10-year moving average, as shown in Figure 7. As can be seen from the trend line, the runoff change was gentler from 1826 to 1917 (92 years) in the JiaoLai River ~~for~~during the entire 180 years. Subsequently, the runoff increased in 1956, and then ~~for nearly 50 years,~~ significantly decreased for nearly 50 years. This finding was consistent with the aforementioned wet and dry seasonal changes.

6. Discussion and Conclusion

The present study used a comprehensive timeline for the regional tree wheel width of the Horqin Sandy Land to reconstruct the runoff sequences in the XiaWa station of the JiaoLai River during the months of ~~Oct.~~October to ~~Nov.~~November from 1826 to 2005. The reconstructed sequence exhibited the same trend as the corresponding sequence of the measured value. The calculated results of various tests showed that the reconstruction result was stable and reliable.

There were greater amplitudes in runoffs in the JiaoLai River between years. The reconstruction sequence had six continuous wet period years, or a total of 51 years, including 27 years that occurred in the early 20th century to the early 1960s. The multi-year average runoff in the wet period from 1946 to 1960 was 1.33 times the entire reconstruction period mean annual runoff. The runoff also went through four continuous dry periods or, a total of 47 years, ~~including which had~~ 30 years that occurred eding in the 1980s to the late 20th century. The multi-year average runoff from

1982 to 2005 was only 63.58% of the entire reconstruction period. Overall, the runoff gradually decreased for nearly 50 years; (~~from the early 1960s to the present~~), and the drop rate in runoff was 1.7766 million years $m^3/10$ years. This result indicated that the runoff significantly decreased; and the total reduction was 75.58% of the initial value.

Runoff in the JiaoLai River had 3-, 11-, 15-, 24-, and 30-year quasi-periodic variations. Among them, the 11-year variation had the same cycle with solar activity. The 24- and 30-year cycles were also consistent with the precipitation cycle. This result ~~fully demonstrated~~ explained that the runoff change in the JiaoLai River followed certain rules, with varying degrees of change similar ~~to~~with other areas around the globe.

Overall, ~~for~~during the ~~entire~~early 180 years in the JiaoLai River, the change in runoff was gentler from 1917 to 1826. Then, ~~in 1956~~, the runoff gradually decreased in 1956. ~~For~~In nearly 50 years, the runoff had a significantly decreasing trend. A comparison with the ten year sliding curve of the precipitation in the Horqin Sandy measured in July to ~~Nov~~November (Figure 7) revealed that the precipitation decreased for nearly 50 years, i.e., from 296.10 mm to 207.67 mm in the 1950s to the 2000s, with the decreased rate of 14.74mm/10 years. The overall decrease accounted for 29.86% of the initial value, which is a significant amount. ~~By~~In contrast, the proportional decrease in precipitation accounted for the initial value being much smaller than the proportion of runoff reduction, accounting for about half. In other words, a slight decrease in precipitation may lead to a significant reduction in runoff. Zhang et al. (2007) have drawn a relatively similar conclusion. They propose that in the past 50

years, the relationship between runoff and precipitation has ~~se~~ an overall downward trend. The results of the present study provide further ~~information~~~~referenees~~ on runoff changes in the Liaohe River, as well as possible strategies for ecological environmental protection and catchment economy progress.

Acknowledgments

This research was supported by ~~the~~ National ~~n~~Natural ~~s~~Science ~~f~~Funds ~~p~~Projects (Grant No: 50869005)-, Inner Mongolia ~~n~~Natural ~~s~~Science ~~f~~Fund ~~p~~Projects (Grant No: 2010BS0608), and ~~Inner Mongolia Agricultural University of the~~ "cold-dry water resources utilization" innovation team ~~for~~~~to~~ ~~planning~~ funding ~~of the Inner Mongolia Agricultural University~~.

References

- Briffa, K.R., Jones, P.D., and Schweinguber, F.H.: Unusual twentieth-century summer warmth in a 1000-year temperature record from Siberia, *Nature*, 376, 156-159, 1995.
- Cleveland, M.K.: A 963-year reconstruction of summer (JJA) streamflow in the White River, Arkansas, USA, from tree-rings, *The Holocene*, 10 (1), 33-41, 2000.
- Cook, E.R., and Jacoby, G.C.: Potomac River stream flow since 1730 as reconstruction tree rings, *Journal of Climate and Applied Meteorology*, 22, 1659-1672, 1983.
- Dong, G.R., Jin, H.L., and Chen, H.Z.: Genesis of desertification in semiarid and subhumid regions of northern China, *Quaternary Sciences*, 2, 136-144, 1998.
- Eryuan, L., Shao, X.M., and Qin, N.S.: Tree-ring based summer temperature reconstruction for the source region of the Yangtze River on the Tibetan Plateau, *Global and Planetary Change*, 61, 313-320, 2008.

- Esper, J., Cook, E.D., and Schweingruber, F.H.: Test of the RCS method for preserving low-frequency variability in long tree-ring chronologies, *Tree-Ring Research*, 59(2), 81-98, 2003.
- Fang, X.Q., Ren, L.L., and Li, Q.F.: Variations of hydrological elements in Laohahe River Basin, *Journal of Hohai University(Natural Sciences)*, 37(6), 620-624, 2009.
- Fritts, H.C.: *Tree ring and climate*, London: Academic Press., 6-8, 1976.
- Giovanna, B., David, F., and Ulf, B.: Five centuries of Central European temperature extremes reconstructed from tree-ring density and documentary evidence, *Global and Planetary Change*, 72, 182-191, 2010.
- Gou, X.H., Yang, M.X., Peng, J.F., et al.: Maximum temperature reconstruction for Animaqing mountain over past 830 years based on tree-ring records, *Quaternary Sciences*, 26(6), 991-998, 2006.
- Gu, Y., Wang, X.L., and Lin, J.: Analysis on change and trend of dry season runoff for major rivers in China since past 60 years, *Water Resources and Hydropower Engineering*, 4, 131-137, 2011.
- Hao, L., Wang, J.A., Gao, L., et al.: On the runoff in Laohahe river basin in the last 40 years, *Journal of Beijing Normal University(Natural Science)*, 44(6): 629-633, 2008.
- Hughes, M.K., Leggett, P., Gray, B., et al.: Climatic signals in British Isles tree-ring chronologies, *Nature*, 72, 605-606, 1978.
- Joseph, R., Mc, A., Louis, A., et al.: Tree-ring record of hillslope erosion and valley floor dynamics: Landscape responses to climate variation during the last 400yr in the Colorado Plateau, northeastern Arizona, *Global and Planetary Change*, 50, 184-201, 2006.

- Jiang, S.H., Ren, L.L., Yong, B., et al.:The effects of climate variability and human activities on runoff from the Laohahe River Basin in Northern China,Water Resources Protection.,26(6),1-15,2010.
- Kang, L.L., Wang, Y.Z., Ma, Y., et al.:Reconstruction of the natural runoff Series in recent 523 years at Huayankou station in Yellow River,Journal of Water Resources and Water Engineering.,19(6), 10-13,2008.
- Kang, X.C., Chen, G.D., Kang, E.S., et al.:Reconstruction the export runoff in the Black River using tree rings in nearly thansoud years,Science in China,Ser.D.,32(8),675-685,2002.
- Law, B.E., Turner,D., Lefsky,M., et al.:Carbon fluxes across regions:Ob-servational constraints at multiple scales. In: Wu J, Jones B, Li H, et al, eds. Scaling and Uncertainty Analysis in Ecology: Methods and Appli-cations,New York: Columbia University Press.,167-190,2006.
- Li, A.M., Han, Z.W., Huang, C.H., et al.:Remote sensing monitoring on dynamic of sandy desertification degree in Horqin Sandy Land at the beginning of 21st century,Journal of Desert Research.,27(4),546-551,2006.
- Li, J.F., Yuan, Y.J., You, X.R., et al.: A-360 years' runoff reconstruction in the RMQI river basin using tree rings,China Quaternary Sciences.,2,131-137,1997.
- Li, J.F., Yuan, Y.J., You, X.R.,et al.:The research and application in tree-ring hydrology,Beijing,science press.,106-107,2000.
- Li, Q., Feng, Q.,- and Zhai, L.X.:Study of the height growth dynamic based on tree-ring data in Populus euphratica from the lower reach of the Heihe River,China,Dendrochronologia.,28,49-64,2010.
- Liu, P.X., Chen, F.H., Jin, L.Y., et al.: About 100-year reconstruction of spring streamflow based

- on tree rings in the lower reaches of Heihe River, *Arid Land Geography*, 30(5), 696-700, 2007.
- Liu, X.H., Qin, D.H., Shao, X.M., et al.: Temperature changes in the middle Qilian Mountain over the last millennium using tree-rings record. *Science in China, Ser.D.*, 34(1), 89-95, 2004.
- Liu, Y., Sun, J.Y., Song, H.M., et al.: Tree-ring hydrologic reconstructions for the Heihe River watershed, western China since AD 1430, *Water Research*, 44, 2781-2792, 2010.
- Liu, Y., Tian, H., Song, H.M., et al.: Tree ring based reconstruction of the May-June mean temperature since A.D. 1884 in WeiChang, HeBei province, *China Quaternary Sciences*, 29(5), 896-904, 2009.
- Liu, Y., Yang, Y.K., Cai, Q.F., et al.: Tree ring based reconstruction of the June-July runoff in HuangShui since past 248 years, *Journal of Arid Land Resources and Environment*, 20(6), 69-73, 2009.
- Ma, L., and Liu, T.X.: X. Relationship of Elm growth and hydrometeorological factors of Horqin Sandy Land and climatic series reconstruction, *Journal of Glaciology and Geocryology*, 29(5), 802-807, 2009.
- Ma, L., Liu, T.X., Jin, H.L., et al.: Reconstruction in elm tree-ring width chronologies and precipitation in Horqin sandy Since 1826, *China Quaternary Sciences*, 31(2), 360-369, 2011.
- Magda, V.N., Zelenova, A.V., and Andreev, S.G.: A 280-year reconstruction of Baikal Lake water level from tree rings, *Conference "Tree Rings and People"*, 22-26, 2001.
- Meko, D.M., Therrell, M.D., Baisan, C.H., et al.: Sacramento River flow reconstructed to A.D. 869 from tree rings, *Journal of the American Water Resources Association*, 37(4), 1029-1040, 2001.
- Michael, G., Loso, R.S., Anderson, S.P., et al.: A 1500-year record of temperature and glacial

- response inferred from varved Iceberg Lake, southcentral Alaska, *Quaternary Research*, 66, 12-24, 2006.
- Nicoletta, M.: Climate from dendrochronology: Latest developments and results, *Glob Planet Change*, 40, 129-139, 2004.
- Qin, N.S., Jin, L.Y., Shi, X.H., et al.: A 518-year runoff reconstruction of Tongtian river basin using tree-ring width chronologies, *Acta Geographica Sinica*, 59(4), 550-556, 2004.
- Samuli, H., Heikki, S.H., John, B., et al.: Reconciling pollen-stratigraphical and tree-ring evidence for high- and low-frequency temperature variability in the past millennium, *Quaternary Science Reviews*, 29, 3905-3918, 2010.
- Shao, X.M.: Some development of chronology round the tree, *Quaternary Sciences*, 3, 265-271, 1997.
- Shao, X.M., Huang, L., Liu, H.B., et al.: Precipitation in Qinghai Millennium Delhi region using tree-ring record, *Science in China, Ser.D*, 34(2), 145-153, 1997.
- Shao, X.M., Wang, S.Z., Xu, Y., et al.: A 3500-year master tree-ring dating chronology from the northeast part of the Qaidam Basin, *Quaternary Sciences*, 27(4), 477-485, 1997..
- Sophie, R., Heike, L., Harald, B., et al.: Understanding the low-temperature limitations to forest growth through calibration of a forest dynamics model with tree-ring data, *Forest Ecology and Management*, 246, 251-263, 1997.
- Stockton, C .W., and Meko, D.M.: A long-term history of drought occurrence in western United States as inferred from tree rings, *Weatherwise*, 28(6), 245-249, 1975.
- Sun, J.Y., Liu, Y., Cai, Q.F., et al.: Tree-ring based reconstruction of June-July for the DaTong river watershed runoff, western china since AD1525, *Marine Geology & Quaternary*

- Geology.,31(3): 109-115,2011.
- Van, R.B., Stahle, D.W., and Cleaveland, M.K.:Earthquake signals in tree-ring data from the New Madrid Seismic zone and implications for paleoseismicity, Geology.,26(6),515-518,1998.
- Wang, G.Q., Jin, J.L., Wang, J.X., et al.:Study on hydrological characteristics of Liaohe River basin in response to climate change,Advances in Earth Science.,26(4),433-440,2011.
- Wang, T., Chen, G.T., Zhao, H.L., et al.:Research progress on aeolian desertification process and controlling in north of china,Journal of Desert Research.,22(4),507-516,2011.
- Wang, T., Wu, W., Xue, X., et al.:Spatial-temporal changes of sandy desertified Land during Last five decades in northern china,Acta Geographica Sinica.,59(2),203-212, 2004.
- Wang, T., Wu, W., Zhao, H.L., et al.:Analyses on driving factors to sandy desertification process in Horqin region, China Journal of Desert Research.,24(5), 519-528, 2004.
- Wang, T., and Zhu, Z.D.:The desertification problems of northern China,Quaternary Sciences., 21(1), 56-65, 2004.
- Wang, T., Zhu, Z.D., and Wu, W.:Sandy desertification in the north China,Science in China:Series D., 45(Supp),23-34,2002.
- Wang, Y.J., Chen, F.H., Gou, X.H., et al.:Change in runoff. During March to June in Heihe since 230-year ago, Journal.,26(2), 202-206,2004.
- Woodhouse, C.A.: A tree-ring reconstruction of streamflow for the Colorado FrontRange,Journal of the American Water Resources Association., 37 (3),561-570,2001.
- Wu, X.D., Sun, L., and Zhan, X.Z.:The initial attempt on past climate in central Tibet use of tree-ring reconstruction,Acta Geographica Sinica., 44 (3), 334-341,2001.
- Yang, H.S., Liu, J., and Liang, H.Y.:Change characteristics of climate and water resources in west

- Liaohe River Plain, Chinese Journal of Applied Ecology., 20(1), 84-90, 2009.
- Zhao, H.L., Zhou, R.L., et al.: Effects of human activity and climate changes on vegetation in Horqin Sandy grassland, Inner Mongolia, Advances in Earth Science., 23(4), 408-414, 2008.
- Zhang, J., and He, J.S.: Analysis of runoff series characteristics in the Liaohe Basin, China Rural Water and Hydropower., 4, 10-13, 2011.
- Zhang, J.Y., Zhang, S.L., Wang, J.X., et al.: Study on runoff trends of the six larger basins in China over the past 50 years, Advances in Water Science., 18(2): 230-234, 2011.
- Zhu, H.F., Zheng, Y.H., Shao, X.M., et al.: The temperature changes on Qing Hai Wu Lan based on tree-ring record nearly thousand years, Chinese Science Bulletin., 53(15), 1835-1841, 2008.

Table C1. Correlation coefficient of standard chronology and runoff.

	Runoff in March	Runoff in April	Runoff in August	Runoff in September	Runoff in October	Runoff in November	◀ Annual Runoff
Standard chronology	0.365**	0.296*	0.455**		0.512**	0.549**	0.281*
Standard chronology in year $t + 1$	0.384**	0.491**	0.298*		0.483**	0.545**	0.277*
Standard chronology in year $t + 2$	0.301*	0.374**	0.449**		0.520**	0.584**	0.454*
Standard chronology in year $t + 3$	0.444**	0.558**	0.386**	0.368**	0.566**	0.626**	0.455*

Note: ** is 99% confidence level, * is 95% confidence level.

- 带格式的: 居中
- 带格式的: 英语(美国)
- 带格式的: 英语(美国)
- 带格式的: 英语(美国)
- 带格式的: 英语(美国)
- 带格式的: 字体: 倾斜
- 带格式的: 字体: 倾斜
- 带格式的: 字体: 倾斜

Table C2. Correlation coefficients of runoff and precipitation.

	Runoff in March	Runoff in April	Runoff in July	Runoff in August	Runoff in September	Runoff in October	Runoff in November	Annual Runoff
Precipitation in July	0.395**	0.293*	0.469**	0.288*	0.398**	0.366**	0.445**	0.519**
Precipitation in August				0.572**	0.414**	0.329*		0.283*
Precipitation in September					0.574**	0.291*	0.285*	
Annual Precipitation	0.276*	0.316*	0.443**	0.359**	0.417**	0.415**	0.430**	0.381**

Note: ** is 99% confidence level, * is 95% confidence level.

带格式的: 居中

带格式的: 制表位: 不在 9.11 字符 + 20.12 字符

Table C3. Mean runoff changes in the reconstruction sequence.

Years [†] mes	Mean runoff (× 10,000 m ³)	Times	Mean runoff (× 10,000 m ³)
1820s	1415.61	1920s	911.33
1830s	1017.40	1930s	1019.15
1840s	975.48	1940s	1119.74
1850s	1225.76	1950s	1410.31
1860s	961.44	1960s	972.87
1870s	1088.97	1970s	887.31
1880s	719.28	1980s	824.57
1890s	971.15	1990s	635.25
1900s	1127.83	2000s	344.35
1910s	862.07		

带格式的：居中

带格式的：居中

带格式的：上标

带格式的：上标




Fig. B2. Regional elm STD chronologies and sample size.




Fig. B3. Contrast curve among the annual runoff, runoff in October, runoff in November, as well as the added runoffs of October and November.




Fig. B4. Correlation among precipitation, runoff, and chronology.




Fig. B5. Contrast curve of the runoff in measured and reconstructed values in ~~Oct-October~~ and ~~Nov-November~~.




Fig. B6. Changes in the runoff reconstruction sequence K_p as well as the distribution in continuous wet and dry season times.




Fig. B7. Sliding changes and trends in the runoff reconstruction sequence and the measured precipitation for ten years.