

Interactive comment on “Impacts of forest changes on hydrology: a case study of large watersheds in the upper reach of Yangtze River Basin” by X. Cui et al.

Anonymous Referee #1

Received and published: 30 July 2012

1. As a reviewing article, the valid references cited in this paper are too less to reaching an effective results. Most viewpoints are depended on only three references (i.e. Liu S, Xu Q, and Zhang Y), which prevented it to discuss deeply on a wide-attention theoretical issue. The climate change, as example, conduced from references before 2002, most data used from 1960s to 2000. How about was it last 10 years? Whether the climate change could be compared with forest change? 2. In section 4, author cited some research results of canopy interception and listed in Table 1. Different people provided dominate different interception ratio, from 17% to 50%, indicate that the variation and indeterminacy are very large. Thus, how did the author introduced the conclusion that

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



the canopy interception of forest in Minjiang River is higher than in most studies forest ecosystems in China and elsewhere? In Dadu River, close to the Minjiang River, some research results shown that the interception percentage for the young and middle-aged *Abies fabri* stand was 23% and 21% of gross rainfall, respectively. 3. The isotope composition $\delta^{18}\text{O}$ and $\delta^2\text{H}$ are usually used to determine the underground water source, sub-surface water movement and its relationship with underground water, et al. In this article, I think the relationships among precipitation, ET, surface runoff and groundwater runoff in different type forests, and the variation of water yield in watershed scale are the key issues should be discussed. In the upper reach of Yangtze River, I knew there were a few documents on those topics. However, author cited only two references from same author on soil water movement and water use pattern of three species of sub alpine forest. 4. It is unclear in determining the differences of water cycle for natural forests and plantation forests. We knew that the interception, water-holding capacity in litter layers and ET differed dominantly between the two typical forests. But in sub-alpine forest land, when natural forests turned to plantation forests, the changes in water yield and the reason are not clear. 5. In section 5, Author pointed out that the contribution of glaciers and snow melt water to base flow was dominant, ranging from 63.8% to 92.6 %, while rain contributions vary from 7.4% to 36.2 %. Although this data were related to a major tributary to the Minjiang River watershed, I suggested that the author must reconsider those statement, the data of the contribution of glaciers and snow melt water were too larger for a tributary river basin in Minjiang river. In source region of Yangtze river, the Tongtian river watershed, the contribution of glaciers and snow melt water to the total runoff is about 5.2%-9.2%, and to the base flow is lower than 50%. 6. At watershed scale, author concluded that the large-scale reforestation or conversion programs consistent with the wide-held “trade-off” relationship between forest and water yield in Minjiang River. But it is lack of sufficient proofs to support this conclusion. It is better to add some effective data illustrating the “trade-off” relationship in watershed scale.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

