

***Interactive comment on “How will climate change modify river flow regimes in Europe?” by C. Schneider et al.***

**MS Richter (Referee)**

brichter@tnc.org

Received and published: 21 August 2012

This is well-executed study with conclusions that should be highly relevant to water management and policy-making in Europe. The methods applied are scientifically sound and the design of this study reflects the authors' strong familiarity with biodiversity conservation, ecological flow management and aquatic ecological theory, ecosystem services, global hydrology modeling, and climate change. This paper also very well demonstrates the potency and flexibility of WaterGAP3 for addressing the implications of changes in hydrological driving forces, e.g., the capability of generating both natural and climate-modified flow regimes.

This paper is very well written and enjoyable to read. The logical structure of the C3929

analysis is well described and easy to follow. In setting up the context for the study, the authors have done an excellent job of summarizing the state of our understanding of river flow influences on biota and ecosystem functions, and appropriately emphasized the economic and other social values tied to ecosystem services that could be lost with changes in flow regimes.

The conclusions of the study are easily understood and supported by modeling results. The only substantive shortcoming is that I wanted to see some comparisons between climate change-induced flow changes and what has already taken place due to anthropogenic water uses. I realize that a comparison with current conditions is somewhat outside the scope of the study's intent, but I know that WaterGAP has been used to describe anthropogenic impacts on flow regimes previously (i.e., Doll et al 2009 in HESS). I wonder if these authors might consider the possibility of adding some results from anthropogenically-altered model runs, as a point of comparison with natural and climate-changed flow regimes? It would be really interesting to at least plot the monthly flow regimes for natural, existing, and future climate conditions in Figures 8-13, for example. Without such a comparison, the reader is left wondering whether future climate effects are greater or less than existing anthropogenic impacts on flow regimes, and whether climate effects will be additive or offsetting with anthropogenic influences.

The authors need to be careful about the asymmetry of percentage changes when comparing low versus high flows. A relatively small change in low flows can produce a large percentage change, with the opposite result for high flows. This does not affect most of the conclusions, tables, or figures in this paper, but it does have some implication for Figure 3, in which changes in individual flow characteristics were added together. Caution should also be exercised in comparing smaller vs. larger rivers for this same reason – for example, the authors might want to run a quality-control check for any significant differences in their results when rivers are parsed into different size classes.

Finally, a small but important suggestion: try to use the same vertical scaling in Figures

8-13 to the extent possible. When you use different vertical scales it can give misleading impression of magnitude of differences between natural and climate-affected flow regimes.

All in all, very nice work!

---

Interactive comment on *Hydrol. Earth Syst. Sci. Discuss.*, 9, 9193, 2012.

C3931