Hydrol. Earth Syst. Sci. Discuss., 9, C5626–C5627, 2012

www.hydrol-earth-syst-sci-discuss.net/9/C5626/2012/ © Author(s) 2012. This work is distributed under the Creative Commons Attribute 3.0 License.



## Interactive comment on "The potential for material processing in hydrological systems – a novel classification approach" by C. E. Oldham et al.

## A. Lorke

lorke@uni-landau.de

Received and published: 2 December 2012

I came across this discussion paper while looking for concepts for characterizing pelagic redoxclines in lakes (my background is environmental fluid dynamics; and I mainly work in surface waters). I found the concepts discussed in the manuscript very useful for our systems and was a little bit surprised to see the rather critical reviews. I agree that the manuscript would benefit from a clear definition of the system and corresponding boundary conditions in both - the conceptual discussion and the examples. However, the limitations of the proposed scaling also apply to the redoxlines in our studies, and also here the (turbulent) diffusion coefficient is scale-dependent, non-local, and varies in space and time... But still I can use the concept (e.g. the

C5626

non-dimensional number Ne) to identify scales at which this heterogeneity will affect turnover rates of redox-reaction and the conditions, e.g., enhanced transport along preferential transport paths, under which hot-spots are potentially important. The approach further allows to estimate the relative importance of transport limitation for different redox-sensitive elements, and I can apply it to sinking particles to evaluate to importance of transformations during sinking... The point I want to make is that, by taking all limitations into account, the concepts described in the manuscript provide a very useful framework for our current study, and potentially many others more.

Andreas Lorke

Interactive comment on Hydrol. Earth Syst. Sci. Discuss., 9, 10487, 2012.