

***Interactive comment on “Global multi-scale  
segmentation of continental and coastal waters  
from the watersheds to the continental margins”  
by G. G. Laruelle et al.***

**G. G. Laruelle et al.**

[goulven@geo.uu.nl](mailto:goulven@geo.uu.nl)

Received and published: 23 January 2013

**[Authors reply to Interactive comment on “Global multi-scale segmentation of continental and coastal waters from the watersheds to the continental margins” by G. G. Laruelle et al.](#)**

*[Anonymous Referee #2 Received and published: 13 November 2012](#)*

The authors present a global multi-scale segmentation of continental and coastal waters from the watersheds to the continental margins.

C6440

The paper is generally well written and follows a clear structure, although it is quite lengthy in many places and I would suggest more concise sections, especially section 3 is very long compared to the rest of the paper. Maybe have results and discussion as separate sections.

[1] We believe that separating the results and discussion would not make the manuscript easier to read. We did, however, take the reviewer’s recommendation into account and condensed our text in several places (in particular sections 3.2 and 3.4, see below). The introduction and conclusion have also been largely re-written (see Comment #1, Reviewer #1).

When I read the abstract I got the impression that the authors were proposing a ‘new’ global scale segmentation composed of three different layers that “includes the wholeaquatic continuum with its riverine, estuarine and shelf sea components.” However, when reading the paper more carefully it seems that the authors have used existing layer created in separate form previously and aggregated those to one single database.

This should be stated in the abstract.

[2] Our work largely uses already existing layers and the abstract has been modified to better reflect this. “*Our work delineates comprehensive ensembles by harmonizing previous segmentations and typologies in order to retain the most important physical characteristics of both the land and shelf areas.*”

Also it should be made clearer in the Introduction why each of the three levels are needed?

[3] The end of the 3rd paragraph now lists the databases that are required for budget and flux calculations. The 4th paragraph describes their integration into all 3 levels.

C6441

*"[...]Moreover, environmental databases gathering monitoring data, climatological forcings and average earth surface properties are available under various forms and at different spatial resolution. Some consists in gridded maps, files or model outputs at 0.5 or 1 degree resolution (World Ocean Atlas, DaSilva et al., 1994, Levitus et al., 1998, GLOBALNEWS, Mayorga et al., 2010, Seitzinger et al., 2005) while others are databases containing measurements from millions (SOCAT, Pfeil et al., 2012) to thousands (GLORICH) or just several dozen (Lonborg and Alvarez-Salgado, 2012, Laruelle et al., 2010, Seiter et al., 2005) of unevenly distributed sampling points. Thus, for environmental budgeting purposes, a multi-scale approach is required to integrate and combine this variety of databases.*

*In this study, we present a harmonized multi-scale segmentation for the land-ocean continuum, from the watershed to the outer limit of the continental shelf. It is based on three increasing levels of aggregation and the inter compatibility of these levels not only allows to integrate a wide variety of databases compiled at various spatial resolutions, but also to compare and combine them with one another. The first level, at the finer resolution of 0.5 degrees, is based on the work of Vorosmarty et al., (2002) and resolves the watersheds and river routing. It also attributes an estuarine type to each watershed following the typology of Durr et al. (2011) which includes small deltas, tidal systems, lagoons and fjords. This spatial resolution allows for a realistic representation of the global river network and is compatible with many global databases (World Ocean Atlas, LOICZ, Buddemeier et al, 2008, Crossland, 2005). This level is also suitable for detailed regional analyses of coastal regions and their corresponding watersheds (Regnier et al., submitted). The second level is built on an updated version of the COSCAT segmentation (Meybeck et al., 2006) which distinguishes different segments of the global coastline based on a combination of terrestrial watershed characteristics and coastal geomorphologic features. It is extended here to include the relevant portions of the adjacent continental shelves. The highest level in the hierarchy is termed MARCATS (for MARGins and CATchment Segmentation) and consists of aggregated COSCAT units according to the main*

C6442

*climatological, morphological and oceanographic characteristics of the coastal zone. It is based on the recent synthesis by Liu et al. (2010) and allows for coarser regional analysis and upscaling calculations when datasets are limited. It nevertheless retains the major physical features of many different coastal regions and identifies a number of widely studied systems such as the main regional seas and some major coastal currents. It can be viewed as an analogue to the coarse segmentation of Takahashi et al. (2009) for the estimation of CO2 fluxes in the open ocean."*

Why is the MARCATS level needed? It is essentially an upscaling of the COSCAT units if I understand correctly.

[4] Correct. The MARCATS are an aggregation of COSCATs, in the same way that COSCATs aggregate single watersheds. The purpose of MARCATS is to design larger scale units accounting for coastal currents, regional seas and large climatic zones that are broader than the COSCATs. It provides a suitable scale at which meaningful regional analysis or data aggregation can be performed for given variables, processes or fluxes when data availability is scarce (Seiter et al., 2005, Laruelle et al., 2010, Lonborg and Alvarez Salgado, 2012). An example of such analysis based on the MARCATS units is given in the new section 3.5, which provides regionalized air-water CO2 fluxes estimates in estuarine environments. In this example, the MARCATS segmentation allows to calculate fluxes from pCO2 data for about a third of the segments. The calculations also require values for estuarine surface areas, which are only available at that resolution (see table 3).

Was any of the listed parameter values in tables 1 to 3 validated or compared to other existing estimates, models or indeed observations, other than the relatively crude global estimates already stated in the text? Or is this the first time such numbers (such as freshwater residence times, watershed surfaces, freshwater discharge, etc) are presented? I imagine some alternative estimates or observations would exist for some of those parameters at least for some locations around the globe?

C6443

[5] In a few of the best studied regions of the world (e.g. Western Europe, Eastern US), some data are already available but none of the parameter values in Table 1-3 have been synthesized at the global-scale. The purpose of our work precisely consisted in a systematic evaluation of these parameters for the entire world and this has been reiterated in the introduction. *"The COSCAT and MARCATS segmentation were used to calculate, for different isobaths, the surface area and volume of each segment of the coastal ocean. The surface areas of watersheds and estuaries are also reported at the same levels. These segmentations can be used in conjunction with biogeochemical databases (e.g. World Ocean Atlas, LOICZ, Hexacoral, GLORICH, SOCAT, and so forth) to establish regional budgets and, eventually, refine global assessments of the carbon and nutrient cycles."*

Note that inconsistent geographic delimitations are often used among different studies, in particular for limit of the outer edge of the shelf. As a result, it remains difficult to relate the surface area of a coastal zone segment to its corresponding watershed. We believe that the data compiled in table 2 and 3 and supplementary material is useful in this respect as they provide comprehensive and consistent synthesis. We agree, however, that some validation for well surveyed regions is useful and we thus compared our results with published estimates for the North Sea, Baltic Sea, Hudson Bay and Persian Gulf (see new table and [7] below).

System	North Sea COSCAT 403	Baltic Sea MARCATS 18	Hudson Bay MARCATS 12	Persian Gulf MARCATS 29
Watershed Surface Area (103 km <sup>2</sup> )	<b>850</b> (1) 870	<b>1650</b> (1) 1619	<b>3700</b> (2) 3601	n.a.
Shelf Surface Area (103 km <sup>2</sup> )	<b>573.3</b> (3) 592	<b>374.6</b> (4) 383	<b>1040</b> (5) 1064	<b>239</b> (6) 233
Shelf Volume (103 km <sup>3</sup> )	<b>42.3</b> (3) 36.3	<b>20.5</b> (4) 20.1	<b>100</b> (7) 105	<b>8.8</b> (6) 8.3

C6444

(1) OSPAR, 2010 (2) Dery et al., 2005 (3) Thomas et al., 2005 (4) Wulff et al., 2001 (5) Macdonald and Kuzyk, 2011 (6) Pous et al., 2012 (7) Saucier et al., 2004

The authors should highlight some of the important implications their data could highlight (for the first time maybe?). For instance, with their data, can they allude to what implications there are on a global level for example of the different freshwater residence times?

Section 3.2 gives a little bit of assessment and compares some of the numbers obtained to previous estimations, which is very useful, but I feel the assessment of the proposed database is not as complete and insightful as it should be.

[6] Following Reviewer #2's suggestion, we have condensed the descriptive part of section 3.2 and elaborate further on the implications of our work. In particular, the second paragraph of the section now explicitly discusses the value of the newly calculated, better constrained, surface areas of continental shelf seas.

*"Most coastal ocean surface area evaluations yield values in the range 25-30 106 km<sup>2</sup> (Laruelle et al., 2010; Walsh et al., 1988; Cai et al., 2006; Chen and Borges, 2005), which corresponds to 8% of the world's ocean. These values have been largely used in the literature to constrain global budgets and box models (Borges et al., 2005, Laruelle et al., 2010, Mackenzie et al., 1993, Rabouille et al., 2001, Ver, 1998, Wanninkhof et al., 2012) but the use of surface areas varying by 20% from one study to another has implications regarding the accuracy of the budgets calculated. The common definition of a single proper limit for the outer edge of the continental shelf remains a matter of debate in the literature (Borges et al., 2005, Laruelle et al., 2010, Liu, 2010) and the choice of this limit depends on various sedimentological and morphological criteria but also, to some degree, on convenience of use. Convenience is the main reason why the 200 m isobath has often been selected as it provides a consistent limit which is easy to manipulate and allows for inter-comparability between studies. However, the morphological heterogeneity of the coastal zone cannot be accounted for by such a*

C6445

simple boundary. Liu et al. (2010) proposed a definition based on the increase in slope of the continental shelf as an alternative which is also used here (see section 2.1) and, although our estimate of 30 106 km<sup>2</sup> falls within the range of previously reported values, the method allows for a more rigorous regional analysis of the shelf area distribution around the globe. The shelf break depths for each COSCAT are provided in table 2. Furthermore, the surface areas and volumes between the calculated isobaths for all COSCAT segments are available as supplementary material as well as GIS files providing the exact geographic extend of each unit. This allows for comparisons between studies relying on different definitions for the boundary between the open and the coastal ocean. It also provides a clear boundary for oceanic studies which either exclude the coastal zone (Takahashi et al., 2009) or treat it differently from the open ocean (Wanninkhof et al., 2012).

The last paragraph of section 3.4 concerning the fresh water residence times has also been modified. It is now more concise and highlights the wide spatial heterogeneity of fresh water residence times in continental shelves and their potential use as indicator of strong riverine influence on coastal regions.

"For each COSCAT and MARCATS, the ratio between the shelf volume and the corresponding riverine discharge has been calculated (Fig 5a-5f). Globally, the comparison between the volume of continental shelf seas (3860 103 km<sup>3</sup>) and the annual fresh water input into the ocean (39 103 km<sup>3</sup> yr<sup>-1</sup>) yields an average value of ~100. However, this "fresh water residence time" is somewhat skewed by the very large contribution of Antarctic shelves to the total (figure 6). If they are excluded from the calculation, the fresh water residence time drops to ~55 years which remains significantly higher than the average residence time of ~8-10 years calculated on the basis of the exchange with the open ocean through upwelling fluxes (Brink et al., 1995; Rabouille et al., 2001; Ver, 1998). Therefore, our results reveal that the renewal of continental shelf waters by fresh water inputs is 5-7 times slower than through upwelling fluxes on average. It should however be noted that the globally averaged

C6446

box-model calculations for upwellings fluxes do not account for the significant spatial and temporal variability in intensity of upwelling processes which can locally renew coastal waters in just weeks (Gruber et al., 2011). Furthermore, neither the box model nor our calculations resolve the lateral transport by along-shore coastal currents. The ratio of fresh water discharge to continental shelf volume varies significantly from one region to the other, from 2 years (for COSCAT 1103 where the Amazon flows) to several thousands of years in many arid regions. Only 17 of the 149 COSCATs have fresh water residence times shorter than 10 years and the cumulative annual fresh water input of these 17 COSCAT segment amounts to 16 103 km<sup>3</sup>, which corresponds to 41% of the global water flux. These regions can be identified as coastal waters under strong riverine influence and bear resemblance, in that respect, to the RiOMAR which are defined as continental margins over which biogeochemical processes are dominated by riverine influences (McKee et al., 2004)."

Without any sort of validation/error characteristics, these numbers do not say much and actually the suggested value of the created database as suggested by the authors might not be entirely 'realistic' (such as regional analyses and for upscaling and biogeochemical budgets and application in Earth System analysis).

[7] We understand the need for some sort of validation. The new table 4 (see [3]) and the accompanying text at the beginning of the updated section 3.3 provides further insights on the robustness of our estimates:

"Table 3 summarizes the surface areas of watersheds, estuaries and continental shelves for every MARCATS. The surface areas of watersheds and continental shelves are also compared with published values for the North Sea, Baltic Sea, Hudson Bay and Persian Gulf (Table 4). The consistency between our estimates and literature data is fairly good and the discrepancy never exceeds 3%. Table 4 also provides comparison between reported values and our estimates for the continental shelf volumes. Only 2%, 5% and 6% deviation are obtained for BAL, HUD and PER,

C6447

respectively. The discrepancy reaches 15% in the case of the North Sea (COSCAT 403) but this is likely due to the use of a slightly different geographic definition of the extent of the North Sea in Thomas et al., (2005), which includes a very deep trench located on the Eastern side.”

On a related note, which regional analysis do the authors mean (see abstract)?

[8] Basically, we aim at regionalized budget and flux analyses for biogeochemical variables. This has now been clarified in the abstract, introduction and conclusion. The new section 3.5 in which MARCATS units are used to calculate regionalized estuarine CO<sub>2</sub> fluxes while still relying on the level 1 segmentation for the allocation of estuarine types within each segment should clarify our objectives.

The results and discussion section is very descriptive in many places and there is not much detail on the actual value these data might provide.

[9] In the revised manuscript, sections 3.2 and 3.4 are now better balanced between description and analysis (see [6]). The new section 3.5 also focuses more on the implications of our multi-scale segmentation, being also now relevant for the study of biogeochemical processes (i.e. CO<sub>2</sub> fluxes).

In the conclusion, the authors need to clarify what they mean by:

- “The 0.5 degree resolution of our level I compares to the highest resolution globally available”. This is interesting and useful but 50 km resolution is still very coarse and I’m unsure whether stating that at this resolution, “the majority of river networks are properly represented” is adequate?

[10] This resolution was used because it is the resolution at which numerous global and regional databases are available and this allows a coupling between these databases and our segmentation (LOICZ, hexacoral, World Ocean Atlas...). It is that compatibility which allows combining the global estuarine typology of Durr et al. (2011) with the

C6448

global fresh water discharge of Fekete et al., (2002) and the COSCAT and MARCATS units. “the majority of river networks are properly represented” means that most watersheds produced at a 0.5 degree display a proper river routing and the water discharge calculated by terrestrial GIS models are realistic for watersheds larger than ten 0.5 degree grid cells. This has been checked by previous studies (Vorosmarty and al., 2000, Beusen et al., 2005) and by calculations of the authors. Our statement has been made more explicit now in the conclusion section:

*“At this resolution, the routing amongst the vast majority of river networks is properly represented and terrestrial GIS models are able to produce reliable riverine discharge estimates for large and medium sized rivers (watersheds > ten terrestrial cells, Beusen et al., 2005).”*

- “The multi-scale segmentation of the Land-Ocean-continuum provides an appropriate support for the progressive integration of global databases for carbon, nutrients and green house gas characteristics into lateral land-ocean matter flux budgets”; How?

[11] Again, our new section 3.5 should now illustrate how our segmentation can be applied to establish budgets of biogeochemically relevant variables such as CO<sub>2</sub> fluxes in spite of the limited available measurements. The purpose of our segmentation is to integrate, combine and compare various databases by, for example, calculating average properties for any given segment (watershed, COSCAT or MARCATS) depending on the data density and availability. This is, for instance, what is being performed in section 3.4 in which fresh water discharges extracted from the Global NEWS database are compared to the estuarine typology of Durr et al. (2011), each at 0.5 degree resolution, and fresh water residence times are calculated for each COSCAT segment using continental shelf volumes. Based on these average properties per COSCAT, extrapolation to other regions or upscaling to the entire world is possible as illustrated in the new section 3.5, and allows for a new set of studies on biogeochemical element

C6449

cycling based on a harmonized database, allowing to compare uncertainties due the approaches chosen while minimizing the effects of the application of different geodata-products.

- "robust regional and global budgets of relevance to environmental and climatic research"; how would research communities use these?

[12] Constraining regional and global air-water CO<sub>2</sub> flux for estuaries has been the object of intense research over the past decade (Abril and Borges, 2004, Borges 2005, Borges et al., 2005, Chen and Borges, 2009, Laruelle et al., 2010, Borges and Abril, 2011, Chen et al., 2012, Cai, 2011). Similar applications can be ambitioned for the other compartments of the Land-Ocean continuum such as inland waters and the coastal ocean. Other biogeochemically relevant gas or chemical components could be covered too (e.g. the fate of organic carbon or silica impacting the biological carbon pump). This work fits within the broad target of constraining better the natural and anthropogenic carbon cycle and thus complements the work currently carried out on the source and sinks for terrestrial ecosystems and the open ocean (see e.g. RECCAP analysis in Special issue of Biogeosciences Discussion 2012: REgional Carbon Cycle Assessment and Processes (RECCAP), Editor(s): J. Canadell, P. Ciais, C. Sabine, and F. Joos). In this context, the contribution of the aquatic continuum remains poorly constrained, especially its role in the global anthropogenic CO<sub>2</sub> budget and climate perturbation. Thus the dataset can be used to improve the knowledge about the carbon cycle, but also about other relevant elements influencing feedback mechanisms of the Earth's system like N, P or Si.

**NB: The updated sections of the manuscript (introduction, section 3.5, conclusions and outlook) are available in a PDF as supplementary material to this letter.**

#### References:

C6450

Abril G, Borges AV (2004) Carbon dioxide and methane emissions from estuaries. Greenhouse gases emissions from natural environments and hydroelectric reservoirs: fluxes and processes, Chapter 7: 187-207. Tremblay A., Varfalvy L., Roehm C. and Garneau M. (eds.). Environmental Science Series, Springer, Berlin, Heidelberg, New York.

Beusen, A. H. W., Dekkers, A. L. M., Bouwman, A. F., Ludwig, W., and Harrison, J. Estimation of global river transport of sediments and associated particulate C, N, and P. *Global Biogeochemical Cycles*, 19 (4), doi: 10.1029/2005GB002453, 2005.

Borges A.V. and G. Abril (2011) Carbon Dioxide and Methane Dynamics in Estuaries, In: Editors-in-Chief: Eric Wolanski and Donald McLusky, *Treatise on Estuarine and Coastal Science - Volume 5: Biogeochemistry*, Academic Press, Waltham, 2011, Pages 119-161, ISBN 9780080878850, 10.1016/B978-0-12-374711-2.00504-0.

Borges AV, Delille B, Frankignoulle M (2005) Budgeting sinks and sources of CO<sub>2</sub> in the coastal ocean: Diversity of ecosystems counts. *Geophys Res Lett* 32: L14601.

Brink, K. H., Abrantes, F. F. G., Bernal, P. A., Dugdale, R. C., Estrada, M., Hutchings, L., Jahnke, R. A., Muller, P. J. and Smith, R. L.: Group report: How do coastal upwelling systems operate as integrated physical, chemical, and biological systems and influence the geological record?, in *Upwelling in the Ocean: Modern Processes and Ancient Records* edited by C. P. Summerhayes, K.-C. Emeis, M. V. Angel, R. L. Smith and B. Zeitzschel, 103-124, 1995

Buddemeier, R.W., Smith, S.V., Swaney, D.P., Crossland, C.J., Maxwell, B.A., 2008. Coastal typology: an integrative "neutral" technique for coastal zone characterization and analysis. *Estuarine, Coastal and Shelf Science* 77 (2008) 197-205.

Cai WJ (2011) Estuarine and coastal ocean carbon paradox: CO<sub>2</sub> sinks or sites of terrestrial carbon incineration? *Annu Rev Mar Sci* 3: 123-145.

Chen CTA, Borges AV (2009) Reconciling opposing views on carbon cycling in the

C6451

coastal ocean: continental shelves as sinks and near-shore ecosystems as sources of atmospheric CO<sub>2</sub>. *Deep-Sea Res PT II* 56 : 578-590 Chen CTA, Huang TH, Fu YH, Bai Y, He X (2012) Strong sources of CO<sub>2</sub> in upper estuaries become sinks of CO<sub>2</sub> in large river plumes. *Curr Opin Environ Sustain* 4: 179-185.

Crossland, C. J., Kremer, H. H., Lindeboom, H. J., Marshall Crossland, J. I., and LeTissier, M. D. A.: Coastal Fluxes in the Anthropocene. *Global Change - The IGBP Series*: Berlin, Heidelberg, Springer, 2005.

DaSilva A., A. C. Young, and S. Levitus, "Atlas of surface marine data 1994, volume 1.: Algorithms and procedures.," Tech. Rep. 6, U.S. Department of Commerce, NOAA, NESDIS, 1994.

Dery, S.J., Wood, E.F., 2005. Decreasing river discharge in northern Canada. *Geophysical Research Letters* 32, L10401. doi:10.1029/2005GL022845.

Durr, H. H., Laruelle, G. G., van Kempen, C. M., Slomp, C. P., Meybeck, M., and Middelkoop, H.: Worldwide Typology of Nearshore Coastal Systems: Defining the Estuarine Filter of River Inputs to the Oceans. *Estuaries and Coasts*, 34(3), 441-458, doi:10.1007/s12237-011-9381-y, 2011.

Fekete, B. M., Vorosmarty, C. J., and Grabs W.: High-resolution fields of global runoff combining observed river discharge and simulated water balances. *Global Biogeochemical Cycles* 16: 1042. doi:10.1029/1999GB001254, 2002.

Gruber, N., Lachkar, Z., Frenzel, H., Marchesiello, P., Munnich, M., McWilliams, J. C., Nagai, T. and Plattner, G.-K.: Mesoscale eddy-induced reduction in eastern boundary upwelling systems. *Nature Geosciences*, 4, 787-792, 2011.

Laruelle, G. G., Durr, H. H., Slomp, C. P., and Borges, A. V.: Evaluation of sinks and sources of CO<sub>2</sub> in the global coastal ocean using a spatially-explicit typology of estuaries and continental shelves, *Geophys. Res. Lett.*, 37, L15607, doi:10.1029/2010GL043691, 2010.

C6452

Levitus, S., T.P. Boyer, M.E. Conkright, T. O' Brien, J. Antonov, C. Stephens, L. Stathopoulos, D. Johnson, R. Gelfeld, 1998: NOAA Atlas NESDIS 18, World Ocean Database 1998: VOLUME 1: INTRODUCTION, U.S. Gov. Printing Office, Wash., D.C., 346pp.

Lonborg C., Alvarez-Salgado X.A. 2012. Recycling versus export of bioavailable dissolved organic matter in the coastal ocean and efficiency of the continental shelf pump. *Global biogeochemical cycles* 26, GB3018, <http://dx.doi.org/10.1029/2012GB004353>.

Liu K.-K. et al. (eds.), *Carbon and Nutrient Fluxes in Continental Margins*, *Global Change - The IGBP Series*, 3 doi 10.1007/978-3-540-92735-8 1, Springer-Verlag Berlin Heidelberg, 2010.

Macdonald R. W. and Z. Z. A. Kuzyk. The Hudson Bay System. Edited by Robie W. Macdonald and Zou Zou A. *Kuzyk Journal of Marine Systems*, Volume 88, Issue 3, Pages 337-488 (2011)

Mayorga, E., Seitzinger, S. P., Harrison, J. A., Dumont, E., Beusen, A. H. W., Bouwman, A. F., Fekete, B. M., Kroeze, C. and Van Drecht, G.: Global Nutrient Export from WaterSheds 2 (NEWS 2): Model development and implementation. *Environmental Modelling and Software* 25(7):837-853, doi:10.1016/j.envsoft.2010.01.007, 2010.

Mackenzie F. T., Ver L. M., Sabine C., Lane M., and Lerman A. (1993) C, N, P, S global biogeochemical cycles and modeling of global change. In *Interactions of C, N, P and S Biogeochemical Cycles and Global Change* (eds. R. Wollast, F. T. Mackenzie, and L. Chou), pp. 1-62. Springer-Verlag.

McKee, B. A., Aller, R. C., Allison, M. A., Bianchi, T. S. and Kineke, G. C.: Transport and transformation of dissolved and particulate materials on continental margins influenced by major rivers: Benthic boundary layer and seabed processes. *Continental Shelf Research* 24: 899-926, 2004.

Meybeck, M., Durr, H. H., and Vorosmarty, C. J.: Global coastal segmentation and its

C6453

river catchment contributors: A new look at land-ocean linkage. *Global Biogeochemical Cycles*, 20, GB1S90:1-15, doi: 10.1029/2005GB002540, 2006.

OSPAR. 2010. The Quality Status Report 2010.

Pous, S., Carton, X. and P. Lazure. 2012. A Process Study of the Tidal Circulation in the Persian Gulf. *Open Journal of Marine Science*, 2012, 2, 131-140, doi:10.4236/ojms.2012.24016

Rabouille, C., Mackenzie, F. T., and Ver, L. M.: Influence of the human perturbation on carbon, nitrogen, and oxygen biogeochemical cycles in the global coastal ocean. *Geochimica et Cosmochimica Acta* 65: 3615-3641, 2001.

Regnier P., Arndt S., Goossens N., Volta C., Laruelle G. G., Lauerwald R., Hartmann J. Modeling estuarine biogeochemical dynamics: from the local to the global scale, Submitted to *Aquatic Geochemistry*

Saucier, F.J., Senneville S., Prinsenbergh S., Roy F., Smith G., Gachon P., Caya D., Laprise R. 2004. Modeling the ice-ocean seasonal cycle in Hudson Bay, Foxe Basin and Hudson Strait, Canada. *Clim. Dyn.* 23: 303-326.

Seiter, K., Hensen, C., and Zabel, M.: Benthic carbon mineralization on a global scale. *Global Biogeochemical Cycles*, 19, GB1010, doi:10.1029/2004GB002225, 2005.

Seitzinger, S. P., Harrison, J. A., Dumont, E., Beusen, A. H. W., and Bouwman, F. B. A.: Sources and delivery of carbon, nitrogen, and phosphorus to the coastal zone: An overview of global Nutrient Export from Watersheds (NEWS) models and their application. *Global Biogeochemical Cycles* 19: GB4S01. doi:10.1029/2005GB002606, 2005.

Takahashi, T., S. C. Sutherland, R. Wanninkhof, C. Sweeney, R. A. Feely, D. W. Chipman, B. Hales, G. Friederich, F. Chavez, A. Watson, D. C. E. Bakker, U. Schuster, N. Metzl, H. Yoshikawa-Inoue, M. Ishii, T. Midorikawa, Y. Nojiri, C. Sabine, J. Olafsson, Th. S. Arnarson, B. Tilbrook, T. Johannessen, A. Olsen, Richard Bellerby, A. Kortzinger, T.

C6454

Steinhoff, M. Hoppema, H. J. W. de Baar, C. S. Wong, Bruno Delille and N. R. Bates (2009). Climatological mean and decadal changes in surface ocean pCO<sub>2</sub>, and net sea-air CO<sub>2</sub> flux over the global oceans. *Deep-Sea Res. II*, 56, 554-577

Thomas, H., Bozec, Y., de Baar, H. J. W., Elkalay, K., Frankignoulle, M., Schiettecatte, L.-S., Kattner, G., and Borges, A. V.: The Carbon budget of the North Sea, *Biogeosciences*, 2, 87-96, 2005.

Ver L. M. (1998) Global kinetic models of the coupled C, N, P, and S biogeochemical cycles: Implications for global environmental change. Ph.D. dissertation, University of Hawaii. Vorosmarty, C. J., Fekete, B.M., Meybeck, M., and Lammers, R. B.: Geomorphometric attributes of the global system of rivers at 30-minute spatial resolution. *Journal of Hydrology* 237: 17-39, 2000.

Walsh, J. J.: On the nature of continental shelves. Academic Press, San Diego, New York, Berkeley, Boston, London, Sydney, Tokyo, Toronto, 1988.

Wanninkhof R, Park GH, Takahashi T, Sweeney C, Feely R, Nojiri Y, Gruber N, Doney SC, McKinley GA, Lenton A, Le Quere C, Heinze C, Schwinger J, Graven H, Khatiwala S (2012) Global ocean carbon uptake: magnitude, variability and trends. *Biogeosci Discuss* 9.

Wulff, F., E. Bonsdorff, I-M. Gren, S. Johansson and A. Stigebrandt, 2001. Giving advice on cost effective measures for a cleaner Baltic Sea: a challenge to science. *Ambio* 30: 254-259.

C6455