

# Geochemical inverse modeling of chemical and isotopic data from groundwaters in Sahara (Ouargla basin, Algeria)

R. Slimani<sup>a</sup>, A. Guendouz<sup>b</sup>, F. Trolard<sup>c</sup>, A.S. Moulla<sup>d</sup>, B. Hamdi-Aïssa<sup>a</sup>, G. Bourrié<sup>c</sup>

<sup>a</sup>Univ Ouargla, Fac. des sciences de la nature et de la vie, Lab. Biochimie des milieux désertiques, Ouargla 30000, Algeria

<sup>b</sup>Blida University, Science and Engineering Faculty, P.O.Box 270 Soumaa, Blida, Algeria

<sup>c</sup>INRA, UMR 1114 Emmah, Avignon, France

<sup>d</sup>Nuclear Research Centre, 16000 Algiers, Algeria

---

## Abstract

~~New samples were collected in the three major Saharan aquifers namely, the “Continental Intercalaire” (CI), the “Complexe Terminal” (CT) and the Phreatic aquifer (Phr) and completed with unpublished more ancient chemical and isotopic data.~~ Instead of classical Debye-Hückel extended law, Specific Interaction Theory (SIT) model, recently incorporated in Phreeqc 3.0 was used. Inverse modeling of hydrochemical data constrained by isotopic data was used here to quantitatively assess the influence of geochemical processes: at depth, the dissolution of salts from the geological formations during upward leakage *without evaporation* explains the transitions from CI to CT and to a first pole of Phr (pole I); near the surface, the dissolution of salts from sebkhas by rainwater explains another pole of Phr (pole II). In every case, secondary precipitation of calcite occurs during dissolution. All Phr waters result from the mixing of these two poles together with calcite precipitation and ion exchange processes. These processes are quantitatively assessed by Phreeqc model. Globally, gypsum dissolution and calcite precipitation were found to act as a carbon sink.

*Keywords:* hydrochemistry, stable isotopes, Sahara, Algeria

---

## 1. INTRODUCTION

A scientific study published in 2008 showed that 85% of the world population lives in the driest half of the Earth. More than 1 billion people residing in arid and semi-arid areas of the world have only access to little or no renewable water resources (OECD, 2008). In many arid regions such as Sahara, groundwater is the only source of water supply for domestic, agricultural or industrial purposes, causing most of the time overuse and / or degradation of water quality.

The groundwater resources of Ouargla basin (Lower-Sahara, Algerian) (Fig. 1) are contained in three main reservoirs (UNESCO, 1972; Eckstein and Eckstein, 2003; OSS, 2003, 2008):

- at the top, the phreatic aquifer (Phr), located in sandy gypsum permeable formations of Quaternary, is almost unexploited (~~only~~ north of Ouargla) due to its salinity (50 g/L);

---

Email address: [slm\\_rabia@yahoo.fr](mailto:slm_rabia@yahoo.fr) (R. Slimani)

- 34 • in the middle, the “Complexe Terminal” (CT) aquifer, (Cornet and Gouscov, 1952; UN-  
35 ESCO, 1972) which is the most exploited, and includes several aquifers in different geo-  
36 logical formations. It circulates in one or two lithostratigraphic formations of the Eocene  
37 and Senonian carbonates or Miopliocene sands;
- 38 • at the bottom, the “Continental Intercalaire” (CI) aquifer, where water is contained in the  
39 lower Cretaceous continental formations (Barremian and Albian), mainly composed of  
40 sandstones, sands and clays. It is only partially exploited because of its significant depth.

41 After use, waters are discharged in a closed system (endorheic basin) and constitute a poten-  
42 tial hazard to the environment, to public health and may jeopardize the sustainability of agricul-  
43 ture (rising of the phreatic aquifer watertable, extension of soil salinization and so on) (Hamdi-  
44 Aïssa et al., 2004; Slimani, 2006). Several previous studies (Guendouz, 1985; Fontes et al., 1986;  
45 Guendouz and Moulla, 1996; Edmunds et al., 2003; Guendouz et al., 2003; Hamdi-Aïssa et al.,  
46 2004; Foster et al., 2006; OSS, 2008; Al-Gamal, 2011) tried, starting from chemical and isotopic  
47 information ( $^2\text{H}$ ,  $^{18}\text{O}$ ,  $^{234}\text{U}$ ,  $^{238}\text{U}$ ,  $^{36}\text{Cl}$ ) to best characterize the relationships between aquifers.  
48 They were more specifically tackling the issue of the Continental Intercalaire recharge. These  
49 investigations dealt particularly with water chemical facies, mapped isocontents of various pa-  
50 rameters, and reported typical geochemical ratios ( $[\text{SO}_4^{2-}]/[\text{Cl}^-]$ ,  $[\text{Mg}^{2+}]/[\text{Ca}^{2+}]$ ) as well as other  
51 correlations. Minerals / solutions equilibria were checked by computing saturation indices with  
52 respect to calcite, gypsum, anhydrite and halite, but processes were only qualitatively assessed.

53 In the present study, new data were collected in order to characterize the hydrochemical and  
54 the isotopic composition of the major aquifers in Ouargla’s region. They also aimed at identify-  
55 ing the origin of the mineralization and water-rock interactions that occur along the flow. New  
56 possibilities offered by progress in geochemical simulations, were used. More specifically, the  
57 inverse modeling of chemical reactions allows us to select the best conceptual model for the in-  
58 terpretation of the geochemical evolution of the Ouargla aquifer. The stepwise inversion strategy  
59 involves designing a list of the scenarios that includes the most plausible combinations of geo-  
60 chemical processes, solving scenarios in a stepwise manner, and selecting the scenario that pro-  
61 vides the best conceptual geochemical model (Dai et al., 2006). Inverse modeling with Phreeqc  
62 3.0 was used to quantitatively assess the influence of the processes that explain the acquisition  
63 of solutes for the different aquifers: dissolution, precipitation, mixing and ion exchange. This  
64 results in constraints on mass balances as well as on the exchange of matter between aquifers.

## 65 2. METHODOLOGY

### 66 2.1. Presentation of the study area

67 The study area is located in the northeastern desert of Algeria “Lower-Sahara” (Le Houérou,  
68 2009) near the city of Ouargla (Fig. 1),  $31^\circ 54'$  to  $32^\circ 1'$  N and  $5^\circ 15'$  to  $5^\circ 27'$  E, with a mean eleva-  
69 tion of 134 (masl). It is located in the quaternary fossil valley of Oued Mya basin. Present climate  
70 belongs to the arid Mediterranean-type (Dubief, 1963; Le Houérou, 2009; ONM, 1975/2013).  
71 This climate is characterized by a mean annual temperature of  $22.5^\circ\text{C}$ , a yearly rainfall of  
72  $43.6\text{ mm/yr}$  and a very high evaporation rate of  $2,138\text{ mm/yr}$ .

73  
74 Ouargla’s region and the entire Lower Sahara has experienced during its long geological  
75 history alternating marine and continental sedimentation phases. During Secondary era, vertical

76 movements affected the Precambrian basement and Primary causing particularly progressive col-  
77 lapse of its central part, along an axis passing substantially through the Oued Righ valley and the  
78 upper portion of the valley oued Mya. According to (Furon, 1960), a epicontinental sea spread  
79 to the Lower Eocene of northern Sahara. After the Oligocene, the sea gradually withdrew. It is  
80 estimated at present that this sea did not reach Ouargla and transgression stopped at the edge of  
81 the bowl (Lelièvre, 1969). The basin is carved into Miopliocene (MP) deposits, which alternate  
82 with red sands, clays and sometimes marls; gypsum is not abundant and dated from Pontian (MP)  
83 (Cornet and Gouscov, 1952; Dubief, 1953; Ould Baba Sy and Besbes, 2006). The continental  
84 Pliocene consists of a local limestone crust with puddingstone or lacustrine limestone (Fig. 2),  
85 shaped by eolian erosion into flat areas (regs). The Quaternary formations are lithologically com-  
86 posed of alternating layers of permeable sand and relatively impermeable marl (Aumassip et al.,  
87 1972; Chellat et al., 2014).

88 The exploitation of Miopliocene aquifer is ancient and at the origin of the creation of the  
89 oasis (Lelièvre, 1969; Moulias, 1927). The piezometric level was higher (145 m a.s.l.) but over-  
90 exploitation at the end of the XIXth century led to a catastrophic decrease of the resource, with  
91 presently more than 900 boreholes (ANRH, 2011).

92 The exploitation of Senonian aquifer dates back to 1953 at a depth 140 m to 200 m depth,  
93 with a small initial rate *ca.* 540 L mn<sup>-1</sup>; two boreholes have been exploited since 1965 and 1969,  
94 with a total flowrate *ca.* 2,500 L mn<sup>-1</sup>, for drinking water and irrigation.

95 The exploitation of Albian aquifer dates back to 1956, with a piezometric level 405 m and a  
96 pressure 22 kg cm<sup>-2</sup>. Presently, two boreholes are exploited:

- 97 • El Hedeb I, 1,335 m depth, with a flowrate 141 L s<sup>-1</sup>;
- 98 • El Hedeb II, 1,400 m depth, with a flowrate 68 L s<sup>-1</sup>.

## 99 2.2. *Sampling and analytical methods*

100 The sampling scheme complies with the flow directions of the two formations (Phr and CT  
101 aquifers); for the CI aquifer only five points are available, so it is impossible to choose a transect  
102 (Fig. 3). Groundwater samples ( $n = 107$ ) were collected during a field campaign in 2013, along  
103 the main flow line of Oued Mya, 67 piezometers tap the phreatic aquifer, 32 wells tap the CT  
104 aquifer and 8 boreholes tap the CI aquifer (Fig. 3). Analyses of Na<sup>+</sup>, K<sup>+</sup>, Ca<sup>2+</sup>, Mg<sup>2+</sup>, Cl<sup>-</sup>,  
105 SO<sub>4</sub><sup>2-</sup> and HCO<sub>3</sub><sup>-</sup> were performed by ion chromatography at Algiers Nuclear Research Center  
106 (CRNA). Previous and yet unpublished data (Guendouz and Moulla, 1996) sampled in 1996 are  
107 used here too: 59 samples for Phr aquifer, 15 samples for CT aquifer and 3 samples for the CI  
108 aquifer for chemical analyses, data <sup>18</sup>O and <sup>3</sup>H (Guendouz and Moulla, 1996).

## 109 2.3. *Geochemical method*

110 Phreeqc (Parkhurst and Appelo, 2013) was used to check minerals / solution equilibria us-  
111 ing the specific interaction theory (SIT), *i.e.* the extension of Debye-Hückel law by Scatchard  
112 and Guggenheim incorporated recently in Phreeqc 3.0. Inverse modeling was used to calculate  
113 the number of minerals and gases' moles that must respectively dissolve or precipitate/degas to  
114 account for the difference in composition between initial and final water end members (Plum-  
115 mer and Back, 1980; Kenoyer and Bowser, 1992; Deutsch, 1997; Plummer and Sprinkle, 2001;  
116 Güler and Thyne, 2004; Parkhurst and Appelo, 2013). This mass balance technique has been  
117 used to quantify reactions controlling water chemistry along flow paths (Thomas et al., 1989).  
118 It is also used to quantify the mixing proportions of end-member components in a flow system

119 (Kuells et al., 2000; Belkhiri et al., 2010, 2012). The Inverse modeling involves designing a list  
120 of the scenarios that includes the most plausible combinations of geochemical processes. For  
121 example, the way to identify whether calcite dissolution/precipitation is relevant or not consists  
122 of solving the inverse problem under two alternate scenarios: (1) considering a geochemical sys-  
123 tem in which calcite is present, and (2) considering a geochemical system without calcite. After  
124 solving the two scenarios, it is usually possible to select the better result as the solution of the  
125 inverse problems and conclude whether calcite dissolution/precipitation is relevant or not. This  
126 stepwise strategy allows us to identify the relevance of a given chemical process by solving the  
127 inverse problem under alternative scenarios in which the process is either occurring or not.

### 128 3. RESULTS AND DISCUSSION

129 Tables 1 to 5 illustrate the results of the chemical and the isotopic analyses. Samples are  
130 ordered according to an increasing ~~salt content that was estimated from their specific~~ electric  
131 conductivity (EC). In both phreatic and CT aquifers, temperature is close to 25 °C, while for  
132 CI aquifer, temperature is close to 50 °C. The results presented in ~~those~~ tables are raw ana-  
133 lytical data that were corrected for defects of charge balance before computing activities with  
134 Phreeqc. As analytical errors could not be ascribed to a specific analyte, the correction was  
135 made proportionally. The corrections do not affect the anions to anions mole ratios such as for  
136  $[\text{HCO}_3^-]/([\text{Cl}^-] + 2[\text{SO}_4^{2-}])$  or  $[\text{SO}_4^{2-}]/[\text{Cl}^-]$ , whereas they affect the cation to anion ratio such  
137 as for  $[\text{Na}^+]/[\text{Cl}^-]$ .

#### 138 3.1. Characterization of chemical facies of the groundwater

139 Piper diagrams drawn for the studied groundwaters (Fig. 4) broadly show a scatter plot  
140 dominated by a Chloride-Sodium facies. However, when going into small details, the widespread  
141 chemical facies of the Phr aquifer is closer to the NaCl pole than those of CI and CT aquifers. The  
142 facies of the Phreatic aquifer most concentrated samples are in the following order: Ca-sulfate  $\leq$   
143 Na-sulfate  $\equiv$  Mg-sulfate  $\leq$  Na-chloride. This sequential order of solutes is comparable to that of  
144 other groundwater occurring in North Africa, and especially in the neighboring area of the chotts  
145 (depressions where salts concentrate by evaporation) Merouane and Melrhir (Vallès et al., 1997;  
146 Hamdi-Aïssa et al., 2004).

#### 147 3.2. Spatial distribution of the mineralization

148 The salinity of the phreatic aquifer varies considerably depending on the location (~~near~~ wells  
149 or drains) and time (influence of irrigation) (Fig. 5a).

150 Its salinity is low around irrigated and fairly well-drained areas, such as the palm groves of  
151 Hassi Miloud, just north of Ouargla (Fig. 3) that benefit from freshwater and are drained to the  
152 sebkha Oum el Raneb. However, the three lowest salinity values are observed in the wells of  
153 Ouargla palm-grove itself, where the Phr aquifer watertable is deeper than 2 m.

154 Conversely, the highest salinity waters are found in wells drilled in the chotts and sebkhas (a  
155 sebkha is the central part of a chott where salinity is the largest) (Safioune and Oum er Raneb)  
156 where the aquifer is often shallower than 50 cm.

157 The salinity of the ~~Complexe Terminal~~ (Miopliocene) aquifer (Fig. 5b) is much lower than  
158 that of the Phr aquifer, and ranges from 1 to 2 g/L; however, its hardness is larger and it contains  
159 more sulfate, chloride and sodium than the waters of the Senonian formations and those of the

160 CI aquifer. The salinity of the Senonian aquifer ranges from 1.1 to 1.7 g/L , while the average  
161 salinity of the ~~Continental Intercalaire~~ is 0.7 g/L (Fig. 5c).

162 A likely contamination of the Miopliocene aquifer by phreatic groundwaters through casing  
163 leakage in an area where water is heavily loaded with salt and therefore particularly aggressive  
164 cannot be excluded.

### 165 3.3. Saturation Indices

166 The calculated saturation indices reveal that waters from CI at 50 °C are close to equilibrium  
167 with respect to calcite, except for 3 samples that are slightly oversaturated. They are however all  
168 undersaturated with respect to gypsum (Fig. 6 ).

169 Moreover, they are oversaturated with respect to dolomite and undersaturated with respect to  
170 anhydrite and halite (Fig. 7).

171 Waters from CT and phreatic aquifers show the same pattern, but some of them are more  
172 largely oversaturated with respect to calcite, at 25 °C.

173 However, several phreatic waters (P031, P566, PLX4, PL18, P002, P023, P116, P066, P162  
174 and P036) that are located in the sebkhas of Sefioune, Oum-er-Raneb, Bamendil and Ain el  
175 Beida's chott are saturated with gypsum and anhydrite. This is in accordance with high evapora-  
176 tive environments found elsewhere (UNESCO, 1972; Hamdi-Aïssa et al., 2004; Slimani, 2006).

177 No significant ~~saturation indices' evolution~~ from the south to the north upstream and down-  
178 stream of Oued Mya (Fig. 7) is observed. This suggests that the acquisition of mineralization  
179 is due to geochemical processes that have already reached equilibrium or steady state in the  
180 upstream areas of Ouargla.

### 181 3.4. Change of facies from the carbonated pole to the evaporites' pole

182 The facies shifts progressively from the carbonated (CI and CT aquifers) to the evaporites'  
183 one (Phr aquifer) with an increase in sulfates and chlorides at the expense of carbonates (SI  
184 of gypsum, anhydrite and halite). This is illustrated by a decrease of ~~the following two ratios:~~  
185  $[\text{HCO}_3^-]/([\text{Cl}^-] + 2[\text{SO}_4^{2-}])$  (Fig. 8) from 0.2 to 0 and of ~~the ratio~~  $[\text{SO}_4^{2-}]/[\text{Cl}^-]$  from 0.8 to  
186 values ranging from 0.3 and 0 (Fig. 9) while salinity increases. Carbonate concentrations tend  
187 towards very small values, while it is not the case for sulfates. This is due to both gypsum  
188 dissolution and calcite precipitation.

189 Chlorides in groundwater may come from three different sources: (i) ancient sea water en-  
190 trapped in sediments; (ii) dissolution of halite and related minerals that are present in evaporite  
191 deposits and (iii) dissolution of dry fallout from the atmosphere, particularly in these arid regions  
192 (Matiatos et al., 2014; Hadj-Ammar et al., 2014).

193 For most of the sampled points the  $[\text{Na}^+]/[\text{Cl}^-]$  ratio remains close to 1, but significant ranges  
194 are observed: from 0.85 to 1.26 for CI aquifer, from 0.40 to 1.02 for the CT aquifer and from 0.13  
195 to 2.15 for the Phr aquifer. All the measured points from the three considered aquifers are more  
196 or less linearly scattered around the unity slope straight line that stands for halite dissolution  
197 (Fig. 10). The latter appears as the most dominant reaction occurring in the medium. However,  
198 at very high salinity,  $\text{Na}^+$  seems to swerve from the straight line, towards smaller values.

199 A further scrutiny of (Fig. 10) shows that CI waters are very close to the 1:1 line. CT  
200 waters are enriched in both  $\text{Na}^+$  and  $\text{Cl}^-$  but slightly lower than the 1:1 line while phreatic waters  
201 are largely enriched and much more scattered. CT waters are closer to the seawater mole ratio  
202 (0.858), but some lower values imply a contribution from another source of chloride than halite  
203 or from entrapped seawater. Conversely, a  $[\text{Na}^+]/[\text{Cl}^-]$  ratio larger than 1 is observed for phreatic

204 waters, which implies the contribution of another source of sodium, most likely sodium sulfate,  
205 that is present as mirabilite or thenardite in the chotts and the sebkhas areas.

206  $[\text{Br}^-]/[\text{Cl}^-]$  ratio ranges from  $2 \times 10^{-3}$  to  $3 \times 10^{-3}$ . The value of this molar ratio for halite is  
207 around  $2.5 \times 10^{-3}$ , which matches the aforementioned range and confirms that halite dissolution  
208 is the most dominant reaction taking place in the studied medium.

209 In these aquifers, calcium originates both from carbonate and sulfate (Fig. 11 and 12). Three  
210 samples from CI aquifer are close to the  $[\text{Ca}^{2+}]/[\text{HCO}_3^-]$  1:2 line, while calcium sulfate disso-  
211 lution explains the excess of calcium. However, a small but significant number of samples (9)  
212 from phreatic aquifer are depleted in calcium, and plot under the  $[\text{Ca}^{2+}]/[\text{HCO}_3^-]$  1:2 line. This  
213 cannot be explained by precipitation of calcite, as some are undersaturated with respect to that  
214 mineral, while others are oversaturated.

215 In this case, a cation exchange process seems to occur leading to a preferential adsorption  
216 of divalent cations, with a release of  $\text{Na}^+$ . This is confirmed by the inverse modeling that is  
217 developed below and which implies  $\text{Mg}^{2+}$  fixation and  $\text{Na}^+$  and  $\text{K}^+$  releases.

218 Larger sulfate values observed in the phreatic aquifer (Fig. 12) with  $[\text{Ca}^{2+}]/[\text{SO}_4^{2-}] < 1$  can  
219 be attributed to a sodium-magnesium sulfate dissolution from a mineral bearing such elements.  
220 This is for instance the case of bloedite.

221

### 222 3.5. Isotope geochemistry

223 CT and CI aquifer exhibit depleted and homogeneous  $^{18}\text{O}$  contents, ranging from  $-8.32 \text{‰}$   
224 to  $-7.85 \text{‰}$ . This was already previously reported by many authors (Edmunds et al., 2003;  
225 Guendouz et al., 2003; Moulla et al., 2012). On the other hand,  $^{18}\text{O}$  values for the phreatic  
226 aquifer are widely dispersed and vary between  $-8.84 \text{‰}$  to  $3.42 \text{‰}$  (Table 6).

227 Waters located north of the Hassi Miloud to Sebkheth Safioune axis are more enriched in  
228 heavy isotopes and therefore more evaporated. In that area, water table is close to the surface  
229 and mixing of both CI and CT groundwaters with phreatic ones through irrigation is nonexistent.  
230 Conversely, waters located south of Hassi Miloud up to Ouargla city show depleted values. This  
231 is the clear fingerprint of a contribution to the Phr waters from the underlying CI and CT aquifers  
232 (Gonfiantini et al., 1975; Guendouz, 1985; Fontes et al., 1986; Guendouz and Moulla, 1996).

233 Phreatic waters result from a mixing of two end-members. An evidence for this is given  
234 by considering the ( $[\text{Cl}^-]$ ,  $^{18}\text{O}$ ) relationship (Fig. 13). The two poles are: i) a first pole of  $^{18}\text{O}$   
235 depleted groundwater (Fig. 14), and ii) another pole of  $^{18}\text{O}$  enriched groundwater with positive  
236 values and a high salinity. The latter is composed of phreatic waters occurring in the northern  
237 part of the study region.

238 Pole I represents the waters from CI and CT whose isotopic composition is depleted in  $^{18}\text{O}$   
239 (average value around  $-8.2 \text{‰}$ ) (Fig. 13). They correspond to an old water recharge (palæorecharge);  
240 whose age estimated by means of  $^{14}\text{C}$ , exceeds 15.000 years BP (Guendouz, 1985; Guendouz and  
241 Michelot, 2006). So, it is not a water body that is recharged by recent precipitation. It consists of  
242 CI and CT groundwaters and partly of phreatic waters, and can be ascribed to an upward leakage  
243 favored by the extension of faults near Amguid El-Biod dorsal.

244 Pole II, observed in Sebkheth Safioune, can be ascribed to the direct dissolution of surficial  
245 evaporitic deposits conveyed by evaporated rainwater.

246 Evaporation alone cannot explain the distribution of data that is observed (Fig. 13). An  
247 evidence for this is given in a semi-logarithmic plot (Fig. 14), as classically obtained according  
248 to the simple approximation of Rayleigh equation (cf. Appendix):

$$\delta^{18}\text{O} \approx 1000 \times (1 - \alpha) \log[\text{Cl}^-] + cte, \quad (1)$$


$$\approx -\epsilon \log[\text{Cl}^-] + cte, \quad (2)$$

249 where  $\alpha$  is the fractionation factor during evaporation, and  $\epsilon \equiv -1000 \times (1 - \alpha)$  is the enrich-  
250 ment factor (Ma et al., 2010; Chkir et al., 2009).

251 CI and CT waters are better separated in the semi-logarithmic plot because they are differen-  
252 tiated by their chloride content. According to equation (1), simple evaporation gives a straight  
253 line (solid line in Fig. 14). The value of  $\epsilon$  used is the value at 25 °C, which is equal to  $-73.5$ .  
254 There is only one sample (P115) on the evaporation straight line, which could be considered as  
255 an outlier in Fig. 13 ( $[\text{Cl}^-] \approx 0$ ). All other samples fit on the logarithmic curve derived from the  
256 mixing line illustrated by Fig. 13.

257 The phreatic waters that are close to pole I (Fig. 13) correspond to groundwaters occurring in  
258 the edges of the basin (Hassi Miloud, piezometer P433) (Fig. 14). They are low-mineralized and  
259 acquire their salinity via two processes namely: dissolution of evaporites along their underground  
260 transit up to Sebket Safioune and dilution through upward leakage by the less-mineralized wa-  
261 ters of CI and CT aquifers (for example Hedeb I for CI and D7F4 for CT) (Fig. 14) (Guendouz,  
262 1985; Guendouz and Moulla, 1996).

263 The rates of the mixing that are due to upward leakage from CI to CT towards the phreatic  
264 aquifer can be calculated by means of a mass balance equation. It only requires knowing the  $\delta$   
265 values of each fraction that is involved in the mixing process.

266 The  $\delta$  value of the mixture is given by:  $\delta_{\text{mix}} = f_1 \times \delta_1 + f_2 \times \delta_2$   (3)

267 where  $f_1$  is the fraction of CI aquifer,  $f_2$  the fraction of the CT and  $\delta_1$ ,  $\delta_2$  are the respective  
268 isotope contents.

269 Average values of mixing fractions from each aquifer to the phreatic waters computed by  
270 means of equation (3) gave the rates of 65 % for CI aquifer and 35% for CT aquifer.

271 A mixture of a phreatic water component that is close to pole I (*i.e.* P433) with another ~~one~~  
272 which is rather close to pole II (*i.e.* P039) (Fig. 13 and 14), for an intermediate water with a  $\delta^{18}\text{O}$   
273 signature ranging from  $-5\text{‰}$  to  $-2\text{‰}$  gives mixture fraction values of 52 % for pole I and 48 %  
274 for pole II. Isotope results will be used to independently cross-check the validity of the mixing  
275 fractions derived from an inverse modeling involving chemical data (*cf. infra*).

276 Turonian evaporites are found to lie in between CI deep aquifer, and the Senonian and  
277 Miocene formations bearing CT aquifer. CT waters can thus simply originate from ascend-  
278 ing CI waters that dissolve Turonian evaporites, a process which does involve any change in  $^{18}\text{O}$   
279 content. Conversely, phreatic waters result to a minor degree from evaporation, and mostly from  
280 dissolution of sebkhas evaporites by  $^{18}\text{O}$  enriched rainwater and mixing with CI-CT waters.

### 281 3.5.1. Tritium content of water

282 Tritium contents of Phr aquifer are relatively small (Table 6), they vary between 0 and 8 TU.  
283 Piezometers PZ12, P036 and P068 show values close to 8 TU, piezometers P018, P019, P416,  
284 P034, P042 and P093 exhibit values ranging between 5 and 6 TU, and the rest of the samples'  
285 concentrations are lower than 2 TU.

286 ~~This~~ values are dated back to November 1992 so they are old values. ~~This is the main reason~~  
287 ~~why~~ they are considered high comparatively to what is expected to be found nowadays. In fact,

288 at present times, tritium figures have fallen lower than 5 TU in precipitation measured in the  
289 northern part of the country.

290 The comparison of these results with that of precipitation which was 16 TU in 1992 was  
291 collected from the National Agency for Water Resources station from Ouargla).

292 This value seems to be high but we can note that we are in an arid area (desert) where precip-  
293 itation is very scarce and irregular. Precipitation takes place in the form of sudden thunderstorms  
294 in an unsaturated atmosphere and a great part of this precipitation evaporates back into the mois-  
295 ture unsaturated atmosphere sometimes during many cycles. Consequently, an enrichment in  
296 tritium happens because when water evaporates back, the lightest fractions (isotopes) are the  
297 ones that evaporate first causing an enrichment in Tritium in the remaining fraction. The 16 TU  
298 value would thus correspond then to a rainy event that had happened during the same sampling  
299 period (Nov. 1992). It's the only available value and it is not a weighted mean for a long period  
300 of time. It is the most representative value for that region and for that time. Unfortunately, all the  
301 other stations (Algiers, Ankara, and Tenerife) (Martinelli et al., 2014) are subject to a completely  
302 different climatic regime and besides the fact that they have more recent values, can absolutely  
303 not be used for our case. Therefore all the assumptions based on recent tritium rain values do not  
304 apply to this study.

305 Depleted contents in  $^{18}\text{O}$  and low tritium concentrations for phreatic waters fit well the mix-  
306 ing scheme and confirm the contribution from the older and deeper CI/CT groundwaters. The  
307 affected areas were clearly identified in the field and correspond to locations that are subject to  
308 a recycling and a return of irrigation waters whose origin are CI/CT boreholes. Moreover, the  
309 mixing that is clearly brought to light by the  $\text{Cl}^-$  vs.  $^{18}\text{O}$  diagrams (Fig. 13 and 14) could partly  
310 derive from an ascending drainage from the deep and confined CI aquifer (exhibiting depleted ho-  
311 mogenous  $^{18}\text{O}$  contents and very low tritium), a vertical leakage that is favoured by the Amguid  
312 El-biod highly faulted area (Guendouz and Moulla, 1996; Edmunds et al., 2003; Guendouz et al.,  
313 2003; Moulla et al., 2012).

### 314 3.6. Inverse modeling

315 We assume that the relationship between  $^{18}\text{O}$  and  $\text{Cl}^-$  data obtained in 1996 is stable with  
316 time, which is a logical assumption as times of transfer from CI to both CT and Phr are very long.  
317 Considering both  $^{18}\text{O}$  and  $\text{Cl}^-$  data, CI, CT and Phr data populations can be categorized. The CI  
318 and CT do not show appreciable  $^{18}\text{O}$  variations, and can be considered as a single population.  
319 The Phr samples consist however of different populations: Pole I, with  $\delta^{18}\text{O}$  values close to -8,  
320 and small  $\text{Cl}^-$  concentrations, more specifically less than  $35 \text{ mmol L}^{-1}$ ; Pole II, with  $\delta^{18}\text{O}$  values  
321 larger than 3, and very large  $\text{Cl}^-$  concentrations, more specifically larger than  $4,000 \text{ mmol L}^{-1}$   
322 (Table 7); intermediate Phr samples result from mixing between poles I and II (mixing line in  
323 Fig. 13, mixing curve in Fig. 14) and from evaporation of pole I (evaporation line in Fig. 14).

324 The mass-balance modeling has shown that relatively few phases are required to derive ob-  
325 served changes in water chemistry and to account for the hydrochemical evolution in Ouargla's  
326 region. The mineral phases' selection is based upon geological descriptions and analysis of rocks  
327 and sediments from the area (OSS, 2003; Hamdi-Aïssa et al., 2004).

328 The inverse model was constrained so that mineral phases from evaporites including gypsum,  
329 halite, mirabilite, glauberite, sylvite and bloedite were set to dissolve until they reach saturation,  
330 and calcite, dolomite were set to precipitate once they reached saturation. Cation exchange reac-  
331 tions of  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{K}^+$  and  $\text{Na}^+$  on exchange sites were included in the model to check which  
332 cations are adsorbed or desorbed during the process. Dissolution and desorption contribute as



333 positive terms in the mass balance, as elements are released in solution. On the other hand,  
334 precipitation and adsorption contribute as negative terms, while elements removed from the so-  
335 lution.  $\text{CO}_{2(g)}$  dissolution is considered by Phreeqc as a dissolution of a mineral, whereas  $\text{CO}_{2(g)}$   
336 degassing is dealt with as if it were a mineral precipitation.

337 Inverse modelling leads to a quantitative assessment of the different solutes' acquisition pro-  
338 cesses and a mass balance for the salts that are dissolved or precipitated from CI, CT and Phr  
339 groundwaters (Fig. 14, Table 8), as follows:

- 340 • transition from CI to CT involves gypsum, halite and sylvite dissolution, and some ion  
341 exchange namely calcium and potassium fixation on exchange sites against magnesium  
342 release, with a very small and quite negligible amount of  $\text{CO}_{2(g)}$  degassing. The maximum  
343 elemental concentration fractional error equals 1%. The model consists of a minimum  
344 number of phases (*i.e.* 6 solid phases and  $\text{CO}_{2(g)}$ ); Another model implies as well dolomite  
345 precipitation with the same fractional error;
- 346 • transition from CT to an average water component of pole I involves dissolution of halite,  
347 sylvite, and bloedite from Turonian evaporites, with a very tiny calcite precipitation. The  
348 maximum fractional error in elemental concentration is 4%. Another model implies  $\text{CO}_{2(g)}$   
349 escape from the solution, with the same fractional error. Large amounts of  $\text{Mg}^{2+}$  and  $\text{SO}_4^{2-}$   
350 are released within the solution (Sharif et al., 2008; Li et al., 2010; Carucci et al., 2012);
- 351 • the formation of Phr pole II can be modeled as being a direct dissolution of salts from the  
352 sebkha by rainwater with positive  $\delta^{18}\text{O}$ ; the most concentrated water (P036 from Sebkhet  
353 Safioune) is taken here for pole II, and pure water as rainwater. In a decreasing order  
354 of amounts respectively involved in that process, halite, sylvite, gypsum and huntite dis-  
355 solve, and little calcite precipitates while some  $\text{Mg}^{2+}$  are released versus  $\text{K}^+$  fixation on  
356 exchange sites. The maximum elemental fractional error in the concentration is equal to  
357 0.004%. Another model implies dolomite precipitation with some more huntite dissolv-  
358 ing, instead of calcite precipitation, but salt dissolution and ion exchange are the same.  
359 Huntite, dolomite and calcite stoichiometries are linearly related, so both models can fit  
360 field data, but calcite precipitation is preferred compared to dolomite precipitation at low  
361 temperature;
- 362 • the origin of all phreatic waters can be explained by a mixing in variable proportions of  
363 pole I and pole II. For instance, waters from pole I and pole II can easily be separated by  
364 their  $\delta^{18}\text{O}$  respectively close to  $-8\text{‰}$  and  $3.5\text{‰}$  (Fig. 13 and 14). Mixing the two poles  
365 is of course not an inert reaction, but rather results in the dissolution and the precipitation  
366 of minerals. Inverse modeling is then used to compute both mixing rates and the extent  
367 of matter exchange between soil and solution. For example, a phreatic water (piezometer  
368 P068) with intermediate values ( $\delta^{18}\text{O} = -3$  and  $[\text{Cl}^-] \approx 2\text{M}$ ) is explained by the mixing  
369 of 58% water from pole I and 42% from pole II. In addition, calcite precipitates,  $\text{Mg}^{2+}$   
370 fixes on exchange sites, against  $\text{Na}^+$  and  $\text{K}^+$ , gypsum dissolves as well as a minor amount  
371 of huntite (Table 8). The maximum elemental concentration fractional error is 2.5% and  
372 the mixing fractions' weighted the  $\delta^{18}\text{O}$  is  $-3.17\text{‰}$ , which is very close to the measured  
373 value ( $-3.04\text{‰}$ ). All the other models, making use of a minimum number of phases, and  
374 not taking into consideration ion exchange reactions are not found compatible with isotope  
375 data. Mixing rates obtained with such models are for example 98% of pole I and 0.9% of

376 pole II, which leads to a  $\delta^{18}O = (-7.80\text{‰})$  which is quite far for the real measured value  
377  $(-3.04\text{‰})$ .

378 The main types of groundwaters occurring in Ouargla basin are thus explained and could  
379 quantitatively be reconstructed. An exception is however sample P115, which is located exactly  
380 on the evaporation line of Phr pole I. Despite numerous attempts, it could not be quantitatively  
381 rebuilt. Its  $^3H$  value (6.8) indicates that it is derived from a more or less recent water component  
382 with very small salt content, most possibly affected by rainwater and some preferential flow  
383 within the piezometer. As this is the only sample on this evaporation line, there remains a doubt  
384 on its significance.

385 Globally, the summary of mass transfer reactions occurring in the studied system (Table 8)  
386 shows that gypsum dissolution results in calcite precipitation and  $CO_{2(g)}$  dissolution, thus acting  
387 as an inorganic carbon sink.

#### 388 4. CONCLUSIONS

389 Groundwater hydrochemistry is a good record indicator for the water-rock interactions that  
390 occur along the groundwater flowpath. The mineral load reflects well the complex processes  
391 taking place while water circulates underground since its point of infiltration.

392 The hydrochemical study of the aquifer system occurring in Ouargla's basin allowed us to  
393 identify the origin of its mineralization. Waters exhibit two different facies: sodium chloride and  
394 sodium sulfate for the phreatic aquifer (Phr), sodium sulfate for the Complexe Terminal (CT)  
395 aquifer and sodium chloride for the Continental Intercalaire (CI) aquifer. Calcium carbonate pre-  
396 cipitation and evaporite dissolution explain the facies change from carbonate to sodium chloride  
397 or sodium sulfate. However reactions imply many minerals with common ions, deep reactions  
398 without evaporation as well as shallow processes affected by both evaporation and mixing. Those  
399 processes are separated by considering both chemical and isotopic data, and quantitatively ex-  
400 plained making use of an inverse geochemical modeling. The main result is that Phr waters do  
401 not originate simply from infiltration of rainwater and dissolution of salts from the sebkhas. Con-  
402 versely, Phr waters are largely influenced by the upwardly mobile deep CT and CI groundwaters,  
403 fractions of the latter interacting with evaporites from Turonian formations. Phreatic waters oc-  
404 currence is explained as a mixing of two end-member components: pole I, which is very close to  
405 CI and CT, and pole II, which is highly mineralized and results from the dissolution by rainwater  
406 of salts from the sebkhas.

407 At depth, CI leaks upwardly and dissolves gypsum, halite and sylvite, with some ion ex-  
408 change, to give waters of CT aquifer composition.

409 CT transformation into Phr pole I waters involves the dissolution of Turonian evaporites  
410 (halite, sylvite and bloedite) with minor calcite precipitation.

411 At the surface, direct dissolution by rainwater of salts from sebkhas (halite, sylvite, gypsum  
412 and some huntite) with precipitation of calcite and  $Mg^{2+}/K^+$  ion exchange results in pole II Phr  
413 composition.

414 All phreatic groundwaters result from a mixing of pole I and pole II water that is accompanied  
415 by calcite precipitation, fixation of  $Mg^{2+}$  on ion exchange sites against the release of  $K^+$  and  $Na^+$ .

416 Moreover, some  $CO_{2(g)}$  escapes from the solution at depth, but dissolves much more at the  
417 surface. The most complex phenomena occur during the dissolution of Turonian evaporites while  
418 CI leaks upwardly towards CT, and from Phr I to Phr II, while the transition from CT to Phr I

419 implies a very limited number of phases. Globally, gypsum dissolution and calcite precipitation  
 420 processes both act as an inorganic carbon sink.

#### 421 **ACKNOWLEDGEMENTS**

422 The authors wish to thank the staff members of the National Agency for Water Resources in  
 423 Ouargla (ANRH) and the Laboratory of Algerian waters (ADE) for the support provided to the  
 424 Technical Cooperation within which this work was carried out. Analyses of  $^{18}\text{O}$  were funded by  
 425 the project CDTN / DDHI (Guendouz and Moulla, 1996). The supports of University of Ouargla  
 426 and of INRA for travel grants of R. Slimani and G. Bourrié are gratefully acknowledged too.

#### 427 **APPENDIX**

428 According to a simple Rayleigh equation, the evolution of the heavy isotope ratio in the  
 429 remaining liquid  $R_l$  is given by:

$$R_l \approx R_{l,0} \times f_l^{\alpha-1}, \quad (4)$$

430 where  $f_l$  is the fraction remaining liquid and  $\alpha$  the fractionation factor.

431 The fraction remaining liquid is derived from chloride concentration, as chloride can be con-  
 432 sidered as conservative during evaporation: all phreatic waters are undersaturated with respect to  
 433 halite, that precipitates only in the last stage. Hence, the following equation holds:

$$f_l \equiv \frac{n_{w,1}}{n_{w,0}} = \frac{[\text{Cl}^-]_0}{[\text{Cl}^-]_1}. \quad (5)$$

434 By taking natural logarithms, one obtains:

$$\ln R_l \approx (1 - \alpha) \times \ln[\text{Cl}^-] + cte, \quad (6)$$

435 As, by definition,

$$R_l \equiv R_{std.} \times \left(1 + \frac{\delta^{18}\text{O}}{1000}\right), \quad (7)$$

436 one has:

$$\ln R_l \equiv \ln R_{std.} + \ln\left(1 + \frac{\delta^{18}\text{O}}{1000}\right), \quad (8)$$

$$\approx \ln R_{std.} + \frac{\delta^{18}\text{O}}{1000}, \quad (9)$$

437 hence, with base 10 logarithms:

$$\delta^{18}\text{O} \approx 1000(1 - \alpha) \log[\text{Cl}^-] + cte, \quad (10)$$

$$\approx -\epsilon \log[\text{Cl}^-] + cte, \quad (11)$$

438 where as classically defined  $\epsilon = 100(\alpha - 1)$  is the enrichment factor.

439 **References**

- 440 Al-Gamal, S.A., 2011. An assessment of recharge possibility to North-Western Sahara Aquifer System (NWSAS) using  
441 environmental isotopes. *Journal of Hydrology* 398, 184 – 190.
- 442 ANRH, 2011. Inventaire des forages de la Wilaya de Ouargla. Rapport technique. Agence Nationale des Ressources  
443 Hydrauliques.
- 444 Aumassip, G., Dagorne, A., Estorges, P., Lefèvre-Witier, P., Mahrour, F., Nesson, C., Rouvillois-Brigol, M., Trecolle, G.,  
445 1972. Aperçus sur l'évolution du paysage quaternaire et le peuplement de la région de Ouargla. *Libyca*, 205–257.
- 446 Belkhir, L., Boudoukha, A., Mouni, L., Baouz, T., 2010. Application of multivariate statistical methods and inverse  
447 geochemical modeling for characterization of groundwater — A case study: Ain Azel plain (Algeria). *Geoderma*  
448 159, 390 – 398.
- 449 Belkhir, L., Mouni, L., Boudoukha, A., 2012. Geochemical evolution of groundwater in an alluvial aquifer: Case of El  
450 Eulma aquifer, East Algeria. *Journal of African Earth Sciences* 66–67, 46 – 55.
- 451 Carucci, V., Petitta, M., Aravena, R., 2012. Interaction between shallow and deep aquifers in the Tivoli Plain (Central  
452 Italy) enhanced by groundwater extraction: A multi-isotope approach and geochemical modeling. *Applied Geochem-*  
453 *istry* 27, 266 – 280. URL: <http://www.sciencedirect.com/science/article/pii/S0883292711004628>,  
454 doi:<http://dx.doi.org/10.1016/j.apgeochem.2011.11.007>.
- 455 Chellat, S., Bourefis, A., Hamdi-Aïss, a.B., Djerrab, A., 2014. Paléoenvironnementale reconstitution of Mio-pliocenes  
456 sandstones of the lower-Sahara at the base of exoscopic and sequential analysis. *Pensee Journal* 76, 34 – 51.
- 457 Chkir, N., Guendouz, A., Zouari, K., Hadj Ammar, F., Moulla, A., 2009. Uranium isotopes in groundwater from the  
458 continental intercalaire aquifer in Algerian Tunisian Sahara (Northern Africa). *Journal of Environmental Radioac-*  
459 *tivity* 100, 649 – 656. URL: <http://www.sciencedirect.com/science/article/pii/S0265931X09001143>,  
460 doi:<http://dx.doi.org/10.1016/j.jenvrad.2009.05.009>.
- 461 Cornet, A., Gouscov, N., 1952. Les eaux du Crétacé inférieur continental dans le Sahara algérien: nappe dite "Albien",  
462 in: *Congrès géologique international, Alger* p. 30.
- 463 Dai, Z., Samper, J., Ritzi, R., 2006. Identifying geochemical processes by inverse modeling of multicomponent reactive  
464 transport in the aquia aquifer. *Geosphere* 2, 210–219.
- 465 Deutsch, W., 1997. *Groundwater Chemistry-Fundamentals and Applications to Contamination*. New York.
- 466 Dubief, J., 1953. *Essai sur l'hydrologie superficielle au Sahara*. Direction du service de la colonisation et de  
467 l'hydraulique, Service des études scientifiques.
- 468 Dubief, J., 1963. *Le climat du Sahara*. Hors-série, Institut de recherches sahariennes.
- 469 Eckstein, G., Eckstein, Y., 2003. A hydrogeological approach to transboundary ground water resources and international  
470 law. *American University International Law Review* 19, 201–258.
- 471 Edmunds, W., Guendouz, A., Mamou, A., Moulla, A., Shand, P., Zouari, K., 2003. Groundwater evolution in the  
472 continental intercalaire aquifer of southern Algeria and Tunisia: trace element and isotopic indicators. *Applied*  
473 *Geochemistry* 18, 805–822.
- 474 Fontes, J., Yousfi, M., Allison, G., 1986. Estimation of long-term, diffuse groundwater discharge in the northern sahara  
475 using stable isotope profiles in soil water. *Journal of Hydrology* 86, 315 – 327.
- 476 Foster, S., Margat, J., Droubi, A., 2006. Concept and importance of nonrenewable resources. Number 10 in IHP-VI  
477 *Series on Groundwater, UNESCO*.
- 478 Furon, R., 1960. *Géologie de l'Afrique*. 2eme édition, Payot.
- 479 Güler, C., Thyne, G., 2004. Hydrologic and geologic factors controlling surface and groundwater chemistry in Indian  
480 wells–Owens valley area, southeastern California, USA. *Journal of Hydrology* 285, 177–198.
- 481 Gonfiantini, R., Conrad, G., Fontes, J.C., Sauzay, G., Payne, B., 1975. Étude isotopique de la nappe du Continental  
482 Intercalaire et de ses relations avec les autres nappes du Sahara septentrional. *Isotope Techniques in Groundwater*  
483 *Hydrology* 1, 227–241.
- 484 Guendouz, A., 1985. Contribution à l'étude hydrochimique et isotopique des nappes profondes du Sahara nord-est  
485 septentrional, Algérie. Phd thesis. Université d'Orsay, France.
- 486 Guendouz, A., Michelot, J., 2006. Chlorine-36 dating of deep groundwater from northern Sahara. *Journal of Hydrology*  
487 328, 572–580.
- 488 Guendouz, A., Moulla, A., 1996. Étude hydrochimique et isotopique des eaux souterraines de la cuvette de Ouargla,  
489 Algérie. Rapport technique. CDTN/DDHI.
- 490 Guendouz, A., Moulla, A., Edmunds, W., Zouari, K., Shands, P., Mamou, A., 2003. Hydrogeochemical and isotopic  
491 evolution of water in the complex terminal aquifer in Algerian Sahara. *Hydrogeology Journal* 11, 483–495.
- 492 Hadj-Ammar, F., Chkir, N., Zouari, K., Hamelin, B., Deschamps, P., Aigoun, A., 2014. Hydro-  
493 geochemical processes in the Complexe Terminal aquifer of southern Tunisia: An integrated investi-  
494 gation based on geochemical and multivariate statistical methods. *Journal of African Earth Sciences*  
495 100, 81 – 95. URL: <http://www.sciencedirect.com/science/article/pii/S1464343X14001940>,  
496 doi:<http://dx.doi.org/10.1016/j.jafrearsci.2014.06.015>.

- 497 Hamdi-Aïssa, B., Vallès, V., Aventurier, A., Ribolzi, O., 2004. Soils and brines geochemistry and mineralogy of hyper  
498 arid desert playa, Ouargla basin, Algerian Sahara. *Arid Land Research and Management* 18, 103–126.
- 499 Kenoyer, G., Bowser, C., 1992. Groundwater chemical evolution in a sandy aquifer in northern Wisconsin. *Water*  
500 *Resources Research* 28, 591–600.
- 501 Kuells, C., Adar, E., Udluft, P., 2000. Resolving patterns of ground water flow by inverse hydrochemical modeling in a  
502 semiarid Kalahari basin. *Tracers and Modelling in Hydrogeology* 262, 447–451.
- 503 Le Houérou, H., 2009. *Bioclimatology and biogeography of Africa*. Springer Verlag.
- 504 Lelièvre, R., 1969. Assainissement de la cuvette de Ouargla. rapports Géohydraulique n° 2. Ministère des Travaux  
505 Publiques et de la construction.
- 506 Li, P., Qian, H., Wu, J., Ding, J., 2010. Geochemical modeling of groundwater in southern plain area of Pengyang  
507 County, Ningxia, China. *Water Science and Engineering* 3, 282–291.
- 508 Ma, J., Pan, F., Chen, L., Edmunds, W., Ding, Z., Zhou, K., He, J., Zhou, K., Huang, T., 2010. Isotopic and geochemical  
509 evidence of recharge sources and water quality in the Quaternary aquifer beneath Jinchang city, NW China. *Applied*  
510 *Geochemistry* 25, 996–1007.
- 511 Martinelli, G., Chahoud, A., Dadomo, A., Fava, A., 2014. Isotopic features of emilia-romagna region  
512 (north italy) groundwaters: Environmental and climatological implications. *Journal of Hydrology* 519, Part  
513 B, 1928 – 1938. URL: <http://www.sciencedirect.com/science/article/pii/S0022169414007690>,  
514 doi:<http://dx.doi.org/10.1016/j.jhydrol.2014.09.077>.
- 515 Matiatos, I., Alexopoulos, A., Godelitsas, A., 2014. Multivariate statistical analysis of the hydrogeochemical and isotopic  
516 composition of the groundwater resources in northeastern Peloponnesus (Greece). *Science of The Total Environment*  
517 476–477, 577 – 590. URL: <http://www.sciencedirect.com/science/article/pii/S0048969714000515>,  
518 doi:<http://dx.doi.org/10.1016/j.scitotenv.2014.01.042>.
- 519 Moulias, D., 1927. L'eau dans les oasis sahariennes, organisation hydraulique, régime juridique. Phd thesis. Alger.
- 520 Moulla, A., Guendouz, A., Cherchali, M.H., Chaid, Z., Ouarezki, S., 2012. Updated geochemical and isotopic data  
521 from the Continental Intercalaire aquifer in the Great Occidental Erg sub-basin (south-western Algeria). *Quaternary*  
522 *International* 257, 64–73.
- 523 OECD, 2008. *OECD Environmental Outlook to 2030. Technical Report 1*. Organisation for Economic Cooperation and  
524 Development.
- 525 ONM, 1975/2013. *Bulletins mensuels de relevé des paramètres climatologiques en Algérie*. Office national  
526 météorologique.
- 527 OSS, 2003. *Système aquifère du Sahara septentrional*. Technical Report. Observatoire du Sahara et du Sahel.
- 528 OSS, 2008. *Système aquifère du Sahara septentrional (Algérie, Tunisie, Libye): gestion concertée d'un bassin trans-*  
529 *frontalier*. Technical Report 1. Observatoire du Sahara et du Sahel.
- 530 Ould Baba Sy, M., Besbes, M., 2006. Holocene recharge and present recharge of the Saharan aquifers — a study by  
531 numerical modeling, in: *International symposium - Management of major aquifers*.
- 532 Parkhurst, D., Appelo, C., 2013. Description of Input and Examples for PHREEQC (Version 3) — A computer program  
533 for speciation, batch-reaction, one-dimensional transport, and inverse geochemical calculations. Technical Report 6.  
534 U.S. Department of the Interior, U.S. Geological Survey. URL: <http://pubs.usgs.gov/tm/06/a43>.
- 535 Plummer, L., Back, M., 1980. The mass balance approach: application to interpreting the chemical evolution of hydro-  
536 logical systems. *American Journal of Science* 280, 130–142.
- 537 Plummer, L., Sprinkle, C., 2001. Radiocarbon dating of dissolved inorganic carbon in groundwater from confined parts  
538 of the upper Floridan aquifer, Florida, USA. *Journal of Hydrology* 9, 127–150.
- 539 Sharif, M., Davis, R., Steele, K., Kim, B., Kresse, T., Fazio, J., 2008. Inverse geochemical modeling of groundwater  
540 evolution with emphasis on arsenic in the Mississippi River Valley alluvial aquifer, Arkansas (USA). *Journal of Hy-*  
541 *drology* 350, 41 – 55. URL: <http://www.sciencedirect.com/science/article/pii/S0022169407007093>,  
542 doi:<http://dx.doi.org/10.1016/j.jhydrol.2007.11.027>.
- 543 Slimani, R., 2006. Contribution à l'évaluation d'indicateurs de pollution environnementaux dans la région de Ouargla:  
544 cas des eaux de rejets agricoles et urbaines. Master's thesis. Université de Ouargla.
- 545 Stumm, W., Morgan, J., 1999. *Aquatic Chemistry: Chemical Equilibria and Rates in Natural Waters*. John Wiley and  
546 Sons.
- 547 Thomas, J., Welch, A., Preissler, A., 1989. Geochemical evolution of ground water in smith creek valley - a hydrologi-  
548 cally closed basin in central Nevada, USA. *Applied Geochemistry* 4, 493–510.
- 549 UNESCO, 1972. *Projet ERESS, Étude des ressources en eau du Sahara septentrional*. Technical Report 10. UNESCO.
- 550 Vallès, V., Rezagui, M., Auque, L., Semadi, A., Roger, L., Zouggari, H., 1997. Geochemistry of saline soils in two arid  
551 zones of the Mediterranean basin. I. Geochemistry of the Chott Melghir-Mehrouane watershed in Algeria. *Arid Soil*  
552 *Research and Rehabilitation* 11, 71–84.



Table 1: Field and analytical data for the Continental Intercalaire aquifer.

Locality	Lat.	Long.	Elev.	Date	EC	T	pH	/mmolL <sup>-1</sup>									
								Alk	Cl <sup>-</sup>	SO <sub>4</sub> <sup>2-</sup>	Na <sup>+</sup>	K <sup>+</sup>	Mg <sup>2+</sup>	Ca <sup>2+</sup>	Br <sup>-</sup>		
Hebbs I	3,534,750	723,986	134.8	09/11/2012	2.01	46.5	7.65	3.5	5.8	6.79	10.7	0.63	2.49	3.3	0.034		
Hebbs I	3,534,750	723,986	134.8	1996	1.9	49.3	7.35	0.42	5.81	1.07	5.71	0.18	0.77	0.48			
Hebbs II	3,534,310	724,290	146.2	1996	2.02	47.4	7.64	0.58	6.19	1.22	5.06	0.2	1.28	0.82			
Aouinet Mousa	3,548,896	721,076	132.6	22/02/2013	2.2	48.9	7.55	1.28	6.49	1.28	5.65	0.16	1.14	1.17			
Aouinet Mousa	3,548,896	723,986	134.8	11/12/2010	2.2	48.9	7.55	3.19	9.8	3.89	6.3	0.69	5.71	1.27			
Hebbs I	3,534,310	724,290	146.2	24/02/2013	2.19	49.3	7.35	1.91	12.4	4.58	10.7	0.7	3.77	2.35			
Hebbs II	3,534,750	724,290	146.2	27/02/2013	2.43	47.4	7.64	2.11	13.1	5.24	13.9	0.53	4.53	1.41	0.033		
Hassi Khfif	3,591,659	721,636	110	09/11/2012	2.01	50.5	6.83	2.98	14.3	5.24	10.8	0.84	3.44	4.63			
Hassi Khfif	3,534,750	723,986	134.8	22/02/2013	2	46.5	7.65	3.46	15.1	7.67	11.8	0.51	5.57	5.16			
Hassi Khfif	3,591,659	721,636	110	09/11/2012	2.96	50.1	7.56	3.31	15.3	7.77	12.2	0.59	5.77	4.95			
El-Bour	3,560,264	720,366	160	22/02/2013	2.96	54.5	7.34	2.88	18.6	6.21	20.6	0.66	4.79	1.38			

Table 2: Field and analytical data for the Complex Terminal aquifer.

Locality	Site	Aquifer	Lat.	Long.	Elev.	Date	EC	T	pH	Alk.	Cl <sup>-</sup>	SO <sub>4</sub> <sup>2-</sup>	Na <sup>+</sup>	K <sup>+</sup>	Mg <sup>2+</sup>	Ca <sup>2+</sup>	Br <sup>-</sup>
				/m			/mS cm <sup>-1</sup>	/°C					/mmol L <sup>-1</sup>				
Ramerdill	D2F4	M	3,560,759.6	720,586.2	206	20/01/2013	2.02	20.1	7.86	1.63	10.1	5.79	9.88	0.68	3.92	2.51	
Ramerdill	D2F5	M	3,567,891.7	720,586.2	206	19/06	2.02	21.1	8.2	1.96	10.75	3.54	10.98	0.99	2.32	2.12	
Ifr	D1F151	S	3,538,891.7	721,060.5	204	1996	2.67	23.5	7	1.26	10.75	2.71	7.99	0.73	2.32	2.12	
Saïd Ouba	D2F66	M	3,540,257.3	720,085.4	216	1996	2.31	18	8	1.43	11.02	4.73	11.47	0.16	2.07	3.33	
Oglat Larbaâ	D6F64	M	3,566,201.4	729,369.3	177	27/01/2013	2.31	24	7.9	1.41	11.36	6.85	11.59	2.31	1.96	4.58	
El-Bour	D4F94	M	3,536,245.2	722,641.7	100.6	1996	3.05	26.2	7.37	1.61	12.8	6.79	5.15	1.94	1.65	9.13	
Saïd Ouba I	D2F71	S	3,557,412.4	718,272.8	211.9	1996	2.27	24.2	8.2	1.54	13.53	5.72	14.99	0.33	3.28	2.57	
Deïche	D6F61	M	3,547,557.1	717,067.1	173.5	26/01/2013	2.22	22.9	7.74	1.78	14.22	8.41	12.6	0.66	3.38	4.43	
El-Bour	D2F71	S	3,557,412.4	718,272.8	211.9	1996	2.22	22.9	7.74	1.78	14.22	8.41	12.6	0.66	3.38	4.43	
Saïd Ouba I	D2F71	S	3,557,412.4	718,272.8	211.9	1996	2.22	22.9	7.74	1.78	14.22	8.41	12.6	0.66	3.38	4.43	
Rouissat III	D3F10	S	3,535,068.1	722,552.1	248	26/01/2013	5.63	25.1	7.34	2.38	14.3	6.86	13.1	0.4	3.36	5.43	0.034
Ifr	D1F151	S	3,538,891.7	721,060.5	204	20/01/2013	2.37	22.9	7.79	1.75	15.4	8.31	13.7	0.22	5.17	4.75	
Saïd Ouba	D2F66	M	3,540,257.3	720,085.4	216	31/01/2013	2.37	22.9	7.79	1.75	15.4	8.31	13.7	0.22	5.17	4.75	
Oglat Larbaâ	D6F64	M	3,566,201.4	729,369.3	177	31/01/2013	2.38	24.9	7.91	2.19	16.1	16.3	8.65	16.5	0.74	4.29	
SAR Mekhadma	D1F91	S	3,536,257.7	717,822.3	221	03/02/2013	2.43	23.7	7.62	2.3	16.3	8.53	16.1	0.68	5.27	4.92	
Saïd Koutid	D6F73	S	3,549,386.5	729,828.4	199	25/01/2013	2.47	25.8	7.75	3.43	16.3	8.53	16.1	0.68	5.27	4.92	
Am N'sara	D6F51	S	3,559,333.1	729,828.4	225.5	25/01/2013	3.36	24.7	7.95	1.98	16.8	9.21	15.9	0.35	3.50	7.97	0.033
Al Louisa	D4F73	S	3,537,523.4	721,904.6	310	26/01/2013	2.57	24	7.49	1.98	17.4	9.04	13.9	1.99	5.78	5.05	
Charzalet A.H	D6F79	M	3,598,750.2	720,356.8	119	02/02/2013	2.84	22.5	7.55	3.47	17.4	9.35	16.6	0.62	6.24	4.96	
Am moussa II	D9F30	S	3,537,814.1	719,665.1	220.6	02/02/2013	7.52	23.9	7.52	2.37	17.5	8.24	17.3	0.39	3.1	6.46	0.033
Am N'sara	D6F50	S	3,559,333.6	716,868.4	255	02/02/2013	2.62	23.8	7.65	2.11	17.7	9.19	15.5	1.13	6.11	4.73	
H.Miloud	D1F135	M	3,547,557.1	717,067.1	173	03/02/2013	2.76	21.6	7.55	3.32	17.9	9.22	16.5	1.01	6.17	4.91	
El Bour	D6F61	M	3,540,257.1	715,816.0	169	25/01/2013	2.65	19.9	8.02	2.14	17.9	5.28	15.8	1.6	3.84	4.73	
H.Miloud	D1F135	M	3,547,557.1	717,067.1	173	03/02/2013	2.76	21.6	7.55	3.32	17.9	9.22	16.5	1.01	6.17	4.91	
N'youssa El Hou	D6F51	S	3,556,256.7	718,976.5	198	31/01/2013	2.97	22.9	7.52	2.03	18.4	9.71	17.9	0.32	6.49	5.14	
El Koum	D6F67	S	3,573,694.1	721,639.7	143	1996	3.07	22.9	8.09	3.52	18.4	9.71	17.9	0.32	6.49	5.14	
El Koum	D6F67	S	3,573,694.1	721,639.7	143	1996	2.5	25	7.6	1.5	18.79	7.17	10.18	3.43	4.97	5.81	
ITAS	D1F150	M	3,536,186.6	717,046.1	93.1	21/01/2013	3.66	23.9	7.54	1.48	18.8	7.07	10.1	3.41	4.94	5.77	
Am moussa V	D9F13	M	3,538,409.2	718,680.2	210.2	08/02/2013	2.39	25.3	7.22	2.28	19.4	9.45	18.8	0.39	3.31	7.61	
El-Bour	D4F94	M	3,536,245.2	722,641.7	100.6	1996	2.3	21.2	7.9	1.58	20.05	7.21	12.09	2.62	5.76	5.17	
Rouissat I	D3F18	M	3,535,564.2	722,498.9	80.4	26/07/2013	2.13	21	7.84	1.85	21.26	11.26	17.2	0.17	5.98	6.01	
St. poupage ebott	D5F80	S	3,541,656.9	723,521.9	224.1	04/02/2013	3.28	24.5	8.23	3.91	22.16	11.9	19.9	2.13	7.64	6.28	
Chart Palmerate	D5F77	S	3,538,219.3	725,541.3	242.8	05/02/2013	3.37	24.6	7.53	3.26	22.3	12.1	20.9	1.15	8.25	6.78	
Bour El Hakeha	D1F134	M	3,545,333.1	720,391.7	86	06/02/2013	3.4	22.2	7.34	4.13	23.2	12.2	21.2	1.49	8.61	6.01	
Guet Chemia	D2F69	M	3,552,504.9	712,786.3	137.1	03/02/2013	3.54	24.6	7.61	2.24	24.7	12.7	21.1	1.65	8.45	6.47	
Abzatt	D2F69	M	3,552,504.9	712,786.3	137.1	03/02/2013	4.05	28	7.3	2.21	25.9	9.47	25.4	0.57	3.64	7.17	
Frame	D6F62	M	3,570,175.8	717,133.8	167.5	27/01/2013	3.79	24.2	7.95	2.27	25.9	13.5	22.6	0.64	8.91	7.16	
Am N'sara	D6F62	M	3,570,175.8	717,133.8	167.5	27/01/2013	3.79	24.2	7.95	2.27	25.9	13.5	22.6	0.64	8.91	7.16	
N'youssa El Hou	D6F51	S	3,556,256.7	718,976.5	198	25/01/2013	3.15	23.2	8.05	2.59	28.59	8.61	23.14	0.62	4.42	8.01	0.035
H.Miloud Benyaza	D1F138	M	3,551,192.5	717,042.1	88.9	28/01/2013	3.85	25.2	7.61	2.44	28.4	14.2	28.4	1.66	10.01	7.12	
Am L'arab	D6F49	M	3,558,822.6	716,799.1	156.5	28/01/2013	3.97	23.7	7.33	2.16	28.9	9.01	23.9	0.53	5	7.72	0.037
H.Miloud Benyaza	D1F138	M	3,551,192.5	717,042.1	88.9	1996	2.9	22.8	7.5	2.16	28.92	9.03	23.87	0.52	4.99	7.7	
Rouissat	D3F8	M	3,545,470.7	732,837.6	332.4	03/02/2013	4.38	25.4	7.51	1.71	29.8	8.33	22.8	1.23	6.23	6.08	
Rouissat	D3F8	M	3,545,470.7	732,837.6	332.4	1996	6.16	25.3	7.22	1.71	29.81	8.33	22.86	1.23	6.23	6.08	
Am El'Arch	D3F8	M	3,544,683.8	733,316	333.0	1996	6.16	25.3	7.22	1.71	29.81	8.33	22.86	1.23	6.23	6.08	
St. poupage ebott	D3F80	S	3,541,666.9	723,521.9	224.1	1996	3.69	25.4	7.67	2.28	42.22	13.53	36.77	1.12	7.43	9.73	

M = Mioplisene aquifer; S = Senonian aquifer

Table 3: Field and analytical data for the Phreatic aquifer.

Locality	Site	Lat.	Long.	Elev.	Date	EC		T	pH	Alk.	Cl <sup>-</sup>	SO <sub>4</sub> <sup>2-</sup>	Na <sup>+</sup>	K <sup>+</sup>	Mg <sup>2+</sup>	Ca <sup>2+</sup>	Br <sup>-</sup>
						mS cm <sup>-1</sup>	°C										
				/m								/mmolL <sup>-1</sup>					
Khezama	P433	3,597,046	719,626	118	20/01/2013	2,09	22,7	9,18	1,56	12,02	7,3	13	0,99	4,34	2,8		
Khezama	P433	3,597,046	719,626	118	1996	2	22,1	8,86	1,46	12	6,87	11,57	0,93	4,4	2,9		
Hassi Mhould	R059	3,547,216	718,358	124	27/01/2013	2,1	23,9	8,15	1,86	13	7,3	12,6	1,25	4,43	3,43	0,024	
Ain Kheir	P106				1996	4,01	23,79	7,52	1,86	14,15	17,89	15,89	0,61	10,61	7,5		
Hassi Nagra	PLX3	3,584,761,4	717,604,5	125	20/01/2013	2,93	23	8,09	2,04	17,77	9,4	16,6	0,93	5,75		0,031	
	LTP 30				1996	4,08	23,73	7,12	5,25	18,21	9,97	24,29	0,41	1,43	8,13		
Maison de culture	PL31	3,537,988	720,114	124	1996	2,31	23,83	8,08	1,46	18,91	7,8	26,05	0,62	2,13			
El Bour	R006	3,547,216	719,421	161	1996	2,96	23,45	7,88	1,27	18,98	7,74	12,41	4,25	5,32	5,31		
Hassi Mhould	R059	3,547,216	718,358	124	1996	2,77	23,45	7,83	3,29	22,1	12,4	21,8	2,21	8,61	5,47		
Oglet Larba	P430	3,567,287,5	750,058,8	139	24/01/2013	4,5	27,5	8,29	4,22	22,6	8,6	28,4	2,21	4,01	3,17		
Maison de culture	PL31	3,537,988	720,114	124	28/01/2013	3,7	22,2	8,23	4,22	22,6	13,4	21,8	2,21	1,86	8,25		
France El Koum	P401	3,572,830,2	719,211,4	112	20/01/2013	3,44	27,5	7,52	2,21	23,3	13,4	50,56	2,82	9,33	6,28		
Gherbouz	PL15	3,537,962	718,744	134	1996	2,47	23,47	7,72	2,21	23,3	13,4	21,8	2,21	1,77	4,18		
Bour El Hachia	P408	3,544,999,3	719,930,6	110	1996	2,43	23,46	7,75	2,99	24,16	13,23	41,89	0,88	2,34	2,23	0,025	
Station d'epuration	PL30	3,538,398	721,404	130	20/01/2013	5,31	23,80	7,39	3,01	24,32	21,22	24,26	1,77	9,09	7,91		
France Ank Djemel	P422	3,575,339	724,063,3	127	1996	4,7	23,61	7,22	4,39	25,3	9,5	23,7	2,32	4,96	7,46		
Kouate Ain Mousssa	PLX2	3,577,944,8	714,428,5	111	20/01/2013	4,1	25,2	7,61	3,03	26,2	10,36	14,83	0,24	2,32	7,46	0,033	
H Chegga	R058	3,547,329,7	716,520,7	129	27/01/2013	3,66	24,6	8,1	3	27,7	10,6	24	2,29	4,96	6,55		
Hassi Mhould	R057	3,548,943	717,353	133	1996	5,3	23,44	7,69	1,34	28,21	11,48	17,38	2,03	11,48	8,57		
Kouate Ain Mousssa	PL15	3,533,586	714,060	141,6	1996	2,62	23,87	7,76	2,84	30,87	16,66	58,74	0,03	0,83	0,73		
Kouate Ain Mousssa	PL15	3,537,109,4	718,419,1	137	1996	4,67	23,87	7,76	1,75	30,87	16,66	58,74	0,03	0,83	0,73		
Mekranah	PL18	3,537,270	721,119	119	31/01/2013	4,67	22,2	7,89	1,75	31,2	15,4	21,3	2,03	11,17	8,57		
H Chegga	PLX4	3,577,944,8	714,428,5	111	1996	4,49	23,67	7,58	1,5	31,2	10,08	20,05	0,8	7,53	6,5		
Kouate El Gokla	PL16	3,532,463	713,715	117	1996	4,49	23,69	7,62	1,45	31,94	12,83	22,23	0,8	7,53	7,89		
Gherbouz	PL15	3,537,962	718,744	134	21/01/2013	4,65	23,3	8,16	1,78	32,4	14,6	10,83	0,8	6,76	10,55		
Kouate El Gokla	PL17	3,531,435	713,298	111	1996	4,77	23,70	7,70	1,55	32,81	12,85	27,7	0,96	6,19	7,57		
Kouate Ain Mousssa	R057	3,548,943	717,353	133	26/01/2013	5,7	26,2	7,64	2,48	33,5	12,1	30,18	0,96	5,98	5,74		
Ecole primaire/collège	PL52	3,538,478	720,170	131	21/01/2013	5,72	22,9	8,21	1,96	33,6	12,1	29,2	3,33	5,93	8,17		
DSA	PL10	3,537,035	719,746	114	1996	6,08	23,71	7,69	1,32	35,01	13,52	37,1	1,92	6,36	8,17		
Kouate El Gokla	PL17	3,531,435	713,298	111	03/02/2013	5,5	22,5	7,72	1,66	35,4	13,8	37,1	1,92	6,36	5,68		
Kouate El Gokla	PL16	3,532,463	713,715	117	03/02/2013	5,8	22,5	8,04	1,66	36,3	11,6	28,5	3,21	6,75	6,75		
Station d'epuration	PL30	3,538,398	719,746	130	31/01/2013	5,29	25,1	7,84	1,66	38,4	14,6	31,9	3,21	6,75	8,37		
Hassi Debeh	P416	3,537,035	719,746	106	24/01/2013	5,5	24,6	8,86	2,37	38,6	18	22,3	0,89	4,45	11,62		
DSA	PL10	3,538,292,9	720,429	114	24/01/2013	5,51	24,6	8,44	2,37	38,6	16,9	22,3	0,89	4,45	11,62		
Hospital	LTPSN2	3,536,077	720,429	132	27/01/2013	6,09	24,5	7,78	2,57	39,7	11,7	36	1,93	9,03	9,21		
PARC SONNACOM	PL28	3,549,999,3	719,558	134	21/01/2013	6,08	24,5	8,13	1,82	42	10,72	30,6	1,86	5,97	5,97		
Bour El Hachia	P408	3,544,999,3	719,930,6	110	27/01/2013	6,22	23,1	8,07	1,82	42,14	19,1	18,57	1,86	5,31	8,46		
Kouate Ain Mousssa	R056	3,549,933	717,022	128	1996	7,62	23,65	7,93	0,36	42,5	10,72	18,57	1,86	5,31	8,46		
Kouate Ain Mousssa	R056	3,549,933	717,022	128	1996	5,98	24,6	7,63	2,16	42,5	10,72	18,57	1,86	5,31	8,46		
Ecole Oksa B. Nafaa	PL41	3,538,660	719,831	127	31/01/2013	6,36	24,1	7,68	2,11	44,9	13,2	36,2	1,18	6,32	6,68		



Table 4: Field and analytical data for the Phreatic aquifer (continued).

Locality	Site	Lat.	Long.	Elev.	Date	EC /mS.cm <sup>-1</sup>	T /°C	pH	Alk.	Cl <sup>-</sup>	SO <sub>4</sub> <sup>2-</sup>	Na <sup>+</sup> /mmol.L <sup>-1</sup>	K <sup>+</sup>	Mg <sup>2+</sup>	Ca <sup>2+</sup>
PARC HYDRAULIQUE	P419	3,539,494	725,605	132	31/01/2013	7,03	26,4	7,84	2,05	45,1	14,4	41,4	10,78	5,95	6,91
Parc hydraulique	PL13	3,536,550	720,200	123	21/01/2013	7,22	24,5	7,51	3,24	47,8	14,5	44,4	10,55	6,35	6,59
Mekhadma	P506	3,536,230	718,708	129	21/01/2013	7,64	27,1	7,94	1,78	48	14,5	42,9	6,56	7,4	7,61
Said Oba	P506	3,535,238,1	725,075,1	126	04/02/2013	8,32	24,3	8,12	1,71	52,6	14,6	42,8	10,97	7,51	7,83
Mekhadma	P566	3,535,238,1	725,075,1	115	19/06	6,7	23,28	7,46	1,8	54,39	17,58	33,32	4,11	22,16	5,17
Mekhadma	PL17	3,540,433,1	719,661,3	115	27/01/2013	9	24,6	7,64	1,72	62,5	15,2	71,6	3,03	4,61	6,06
Mekhadma	PL13	3,536,908	718,511	130	21/01/2013	9,4	24,5	8,06	3,39	63,2	15,6	77,2	2,51	4,08	5,11
Mekhadma	PL25	3,530,116,2	722,775,1	130	04/02/2013	10,09	30,2	7,91	1,63	63,6	21,5	88,3	4,08	4,21	4,65
Mekhadma	PL25	3,536,230	718,708	129	19/06	9,5	23,72	7,96	0,63	75,57	10,62	10,22	2,64	32,94	9,54
Mekhadma (Bab-sha)	P066	3,542,636,5	718,957,4	126	19/06	7,75	23,48	7,62	1,51	80,23	12,45	45,87	2,46	23,59	5,91
CEM Malek B. Nabi	PL03	3,540,010,9	725,738,1	130	19/06	7,34	23,86	7,60	3,04	84,14	30,58	108,55	2,23	10,17	8,99
ENTV	PL21	3,536,074	721,268	128	19/06	9,73	23,82	7,25	4,46	84,26	23,68	61,62	3,75	33,53	1,88
Hôtel Transat	PL23	3,538,419	720,950	126	19/06	1,5	24,2	8,2	4,53	86,6	16,7	79,9	3,21	14,54	6,85
Mekhadma	PL21	3,536,074	721,268	128	28/01/2013	16,41	25,7	7,45	1,97	99,9	17,4	85,5	5,7	15,66	7,6
Mekhadma	PL05	3,537,109,4	718,419,1	137	21/01/2013	16,8	24,8	7,64	2,02	101,3	17,7	85,9	5,85	16,69	7,59
Beni Thour	PL44	3,536,039,3	721,673,9	134	19/06	4,68	23,85	7,19	2,74	109,75	67,21	134,67	5,71	42,02	8,77
Tazegrant	PLSNI	3,537,675	719,416	125	22/01/2013	17,08	24,9	8	3,41	114,2	18,1	92,9	12,8	16,85	7,24
CEM Malek B. Nabi	PL03	3,540,010,9	725,738,1	130	27/01/2013	10,84	23,1	7,54	3,29	117,3	14,7	116,4	2,06	8,99	7,24
El Bour	P006	3,564,272	719,421	161	03/02/2013	18,31	23,6	7,71	2,38	131,9	18,1	96,3	8,61	52,44	6,25
Ain Moussa	P015	3,551,711	720,591	103	19/06	12,42	26,4	7,85	4,03	138	16,7	108,8	13,06	19,51	7,99
Station de pompage	PL04	3,541,410,1	723,501,1	138	27/01/2013	19,01	26,4	7,67	2,68	142,22	24,5	125,9	3,1	27,11	8,72
Drain Chott Ouargla	D.Ch				19/06		23,88	7,8	4,96	153	17,7	96,31	3,16	44,22	3,02
Beni Thour	PL44	3,536,039,3	721,673,9	134	28/01/2013	20,18	25,8	7,8	4,96	153	17,7	96,31	3,16	44,22	3,02
CNMC	PL27	3,535,474	718,407	126	21/01/2013	21,23	24,8	8,11	1,7	169,4	18,4	130,3	6,29	22,83	8,08
Bamerdil	P076	3,540,137	716,721	118	26/01/2013	22,31	27,2	7,57	4,33	171,5	17,1	130,8	4,89	27,81	8,63
N'Goussa	P041	3,559,563	716,543	135	25/01/2013	25,94	24,5	8,18	7,95	208,6	13,4	198,9	3,61	18,1	8,83
N'Goussa	P009	3,559,388	717,707	123	26/01/2013	27,51	28,4	8,39	11,45	208,8	15,8	195,1	2,65	18,7	9,01
LTP16					19/06	11,53	23,78	7,48	3,84	213,35	48,63	147,9	7,46	75,31	4,25
Chert Adjadja Aven	PLX1	3,540,758,8	726,115,6	132	28/01/2013	17,18	23,64	7,59	3,37	235,01	46,44	264,84	4,74	25,57	5,56
Route Frane	P003	3,569,043	721,496	134	02/02/2013	32,93	23,4	7,95	4,44	245,6	20,9	141,4	26,88	44,56	17,66
El Bour-N'geuca	P007	3,562,236	718,651	129	26/01/2013	31,03	23,5	8,01	6,91	252,7	17,9	208,2	9,41	29,99	10,03
Route Ain Bida	PLX2	3,537,323,9	724,063,3	127	21/01/2013	30,07	28,4	7,76	5,42	254,7	15,5	270,4	10,43	28,82	7,51
Ain Moussa	P015	3,551,711	720,591	103	25/01/2013	43,25	25,7	8,07	5,15	262,2	9,3	206,9	15,5	62,77	21,46
Ain Moussa	P402	3,549,503	721,514	138	25/01/2013	32,02	22,7	8,03	2,95	263	15,4	206,9	6,56	32,12	9,95
Route Frane	P001	3,572,148	722,336	127	19/06	60	28,7	8,6	7,69	313,2	93,9	442,8	23,26	12,56	10,17
Ain Moussa	P014	3,551,466	719,339	131	02/02/2013	60,58	23,63	8,37	4	323,62	58,13	331,43	5,01	49,77	3,97
N'Goussa	P019	3,562,060	717,719	113	26/01/2013	61,06	23,40	7,31	3,98	336,96	64,29	336,96	5,53	62,37	5,45
N'Goussa	P018	3,562,122	716,590	110	26/01/2013	61,06	27,8	7,65	6,02	356,2	96	432,5	29,77	21,02	26,23
Ain Moussa	P014	3,551,466	719,339	131	25/01/2013	49,04	26,2	8,42	6,46	372,4	82,3	347,1	22,64	60,71	26,63
Route Sedrata	PL13	3,535,586	714,576	105	03/02/2013	62,24	24,8	7,89	5,96	399,7	21,1	389,3	2,41	18,97	7,39
N'Goussa	P009	3,559,388	717,707	123	19/06	62,24	23,27	7,84	2,4	426,85	57,81	393,83	9,13	59,13	12,02

Table 5: Field and analytical data for the Phreatic aquifer (continued).

Locality	Site	Lat.	Long.	Elev.	Date	EC	T	pH	Alk.	Cl <sup>-</sup>	SO <sub>4</sub> <sup>2-</sup>	NH <sub>4</sub> <sup>+</sup>	K <sup>+</sup>	Mg <sup>2+</sup>	Ca <sup>2+</sup>
			/m			/mS cm <sup>-1</sup>	/°C					/mmol/L			
Roume France	P001	3.572,148	723,366	127	02/02/2013	66,16	28,3	7,24	6,49	468,7	101,5	350,3	25,96	116,21	35,31
Sebkhet Safoume	P031	3.577,804	720,172	120	1996		23,75	7,31	6,32	481,83	43,35	326,82	12,61	94,15	23,56
Sebkhet Safoume	P031	3.577,804	720,172	120	02/02/2013	75,96	27,9	8,06	5,85	500,3	110,3	470,5	28,67	79,12	35,47
Roume France	P002	3.570,523	722,028	108	1996		23,81	7,76	6,29	522,39	182,95	653,78	9,97	104,7	10,99
Sebkhet Safoume	P030	3.577,253	721,936	130	1996		23,52	7,72	4,43	527,7	123,48	533,79	11,59	106,21	10,65
Oumt Karabeh	P012	3.584,089	718,612	114	25/01/2013	64,05	30,3	7,83	7,77	534,3	20,9	529,6	6,41	19,73	4,73
Oumt Karabeh	P012	3.584,089	718,612	114	1996		23,41	7,46	2,72	539,35	60,64	413,55	5,55	112,77	9,42
ANK Djemel	P423	3.540,881	723,178	102	31/01/2013	90,8	23,5	7,48	6,19	636,5	101,3	495,5	8,31	125,81	30,32
Saïd Othob-Chout	P006	3.540,265	724,729	111	1996		23,59	7,71	3,69	645,07	78,46	357,28	5,89	208,4	12,86
ANK Djemel	P030	3.577,253	721,936	130	03/02/2013		23,1	7,83	3,71	671,8	90,3	742,9	15,97	41,46	7,65
Sebkhet Safoume	P006	3.560,256	715,781	130	26/01/2013	100,1	31	7,43	3,78	679,3	114,1	597,8	10,71	125,85	26,29
N'Crassa	P021	3.573,943	723,161	105	1996		23,55	7,43	4,24	700,77	154,45	608,68	53,6	85,85	163,08
ANK Djemel	P017	3.573,943	723,161	105	1996		23,30	7,42	2,37	716,27	14,1	560,07	7,04	77,72	11,04
Station de pompage	P004	3.541,410,1	717,719	138	02/02/2013	63,82	25,9	7,57	1,65	748,5	34,75	651,5	14,72	77,72	27,29
Roume France	P002	3.570,523	722,028	108	03/02/2013	68,31	25,9	7,72	2,37	771	62,6	615,9	23,46	77,72	30,39
Saïd Othob-Chout	P006	3.540,265	724,729	111	1996		23,30	7,18	1,24	779,13	77,13	711,46	9,23	69,64	18,06
N'Crassa	P019	3.562,960	717,719	113	1996		26,2	8,7	1,54	800,4	824	1,240,7	18,95	37,63	12,05
Saïd Othob(Bib-shou)	P006	3.542,636,5	718,957,4	126	25/01/2013	82,28	29,6	7,64	2,29	842,8	283	824	49,54	319,35	25,59
ANK Djemel	P021	3.573,943	723,161	105	03/02/2013	150,6	30,7	7,15	2,43	848,8	81	244,21	49,54	33,47	17,74
N'Crassa	P018	3.562,122	716,590	110	1996		23,29	7,46	1,24	886,7	94,4	1,309,9	13,3	33,47	11,67
Oumt Karabeh	P162	3.535,133	725,129	98	25/01/2013	160	27,4	7,70	2,81	954,89	289,9	997,52	7,53	86,69	14,24
Roume Sedrara	P113	3.547,234	714,576	110	1996		23,49	7,37	2,88	980,1	15,5	930,8	19,14	270,91	13,3
Hotel Transit	PZ12	3.577,198	722,931	105	05/02/2013	114,9	23,49	7,42	2,25	1,103,31	91,14	1,058,21	11,72	133,47	12,41
Sebkhet Safoume	P023	3.577,198	725,726	126	1996		23,32	7,42	1,76	1,176,99	14,7	1,051,1	18,27	56,37	17,38
Sebkhet Safoume	P023	3.579,698	725,633	97	05/02/2013	130	29,4	7,64	1,85	1,379,35	139,61	1,257,42	41,55	190,51	10,03
Sebkhet Safoume	P034	3.577,198	725,726	99	1996		23,60	7,64	1,94	1,638,66	143,36	1,321,87	26,85	331,38	12,26
Chout Adiafja	PLXI	3.540,758,8	726,115,6	132	05/02/2013	117,9	23,77	7,71	1,72	1,860,53	91,55	1,434,73	26,2	278,77	13,25
Sebkhet Safoume	P063	3.545,586,8	725,667,4	99	1996		23,26	7,67	1,43	1,887,9	92,9	1,455,8	26,66	282,88	13,44
Bamerndi	LTT06	3.540,137	716,721	118	1996		23,53	7,71	1,41	1,860,53	143,36	1,434,73	27,33	171,23	6,54
El Boue-N'poca	P076	3.562,236	718,651	129	1996		23,26	7,79	1,49	2,106,07	18,27	1,765,47	29,49	278,18	10,44
Sebkhet Safoume	P063	3.545,586,8	725,667,4	99	05/02/2013	178,9	23,39	7,49	1,49	1,820,8	182,08	1,957,53	29,49	278,18	10,44
Oumt Karabeh	P068	3.547,234	722,931	110	1996		23,42	7,59	1,1	2,330,85	101,22	1,963,71	52,19	248,1	11,24
Hassi Dabch	PZ12	3.581,097	730,922	106	1996		23,31	7,84	3,35	2,335,67	222,08	2,302,25	26,84	219,9	7,19
N'Crassa	P041	3.559,563	716,543	135	1996		23,38	7,94	4,33	2,405,55	109,92	2,178,55	25,23	199,35	12,65
Sebkhet Safoume	P034	3.579,698	725,633	97	1996		23,37	7,85	1,95	2,433,73	178,87	2,361,09	24,34	196,07	9,2
Sebkhet Safoume	P039				1996		23,34	7,85	1,94	2,599,74	324,58	2,878,99	44,57	152,83	10,97
Sebkhet Safoume	P074				1996		23,54	6,87	4,17	2,752	134,14	2,616,77	24,42	180,14	9,23
Sebkhet Safoume	P074				1996		23,36	6,92	1,52	4,356,48	201,44	4,042,62	27,9	257,81	22,63
Sebkhet Safoume	P036				1996		23,35	7,54	1,4	4,953,84	184,54	4,759,9	2,9	347,57	7,86
Sebkhet Safoume	P036				1996		23,35	7,54	1,4	4,972,75	108,12	4,692,23	36,84	221,13	9,63

Table 6: Isotopic data  $^{18}\text{O}$  and  $^3\text{H}$  and chloride concentration in Continental Intercalaire, Complexe Terminal and Phreatic aquifers (sampling campaign in 1992).

Phreatic aquifer											
Piezometer	$\text{Cl}^-$ /mmolL $^{-1}$	$\delta^{18}\text{O}$ /‰	$^3\text{H}$ /UT	Piezometer	$\text{Cl}^-$ /mmolL $^{-1}$	$\delta^{18}\text{O}$ /‰	$^3\text{H}$ /UT	Piezometer	$\text{Cl}^-$ /mmolL $^{-1}$	$\delta^{18}\text{O}$ /‰	$^3\text{H}$ /UT
P007	1.860.5	-2.49	0	PL15	23.54	-7.85	0.6(1)	P074	4.356.4	3.42	6.8(8)
P009	426.85	-6.6	1.2(3)	P066	80.23	-8.14	0.8(1)	PL06	14.15	-8.13	1.0(2)
P506	54.39	-6.83	1.6(3)	PL23	1,103.32	-6.1	0	PL30	24.32	-7.48	2.4(4)
P018	818.67	-2.95	6.2(11)	P063	1,379.3	-3.4	8.7(15)	P002	522.39	-5.71	0.6(1)
P019	779.13	-4.67	5.6(9)	P068	2,335.6	-3.04	8.8(14)	PL21	84.26	-7.65	1.2(2)
PZ12	2,405.5	-2.31	8.1(13)	P030	527.7	-6.57	2.4(4)	PL31	18.91	-7.38	1.6(3)
P023	1,176.9	-2.62	0.2(1)	P076	1,743.5	-5.56	2.8(5)	P433	12	-8.84	0
P416	2,433.7	-7.88	5.9(9)	P021	700.7	-5.16	2.6(4)	PL03	84.14	-7.35	1.7(3)
P034	2,752	-1.77	5.7(9)	PL04	716.27	-2.89		PL44	109.75	-8.82	1.0(2)
P036	4,972.7	3.33	2.1(4)	P093	2,198.5	-2.64	5.1(8)	PL05	30.87	-7.44	1.9(3)
P037	4,953.8	3.12	1.8(3)	P096	645.07	-6.13	4.8(8)	P408	24.16	-7.92	0
P039	4,189.5	0.97	2.2(4)	PLX1	1,296.6	-5.6	1.1(2)	PL16	31.94	-7.18	1.1(2)
P041	2,599.7	-0.58	7.3(13)	PLX2	25.68	-7.6	1.3(2)	LTP16	213.35	-7.48	1.6(3)
P044	2,106.1	-4.46	2.7(5)	P015	134.68	-6.77	3.0(5)	PL17	32.81	-6.92	0.1
P014	336.96	-6.9	2.8(5)	P001	323.62	-4.66	2.5(4)	PL10	35.01	-7.31	0.2(1)
P012	539.3	-6.41	2.2(4)	P100	235.01	-5.81	0	PL25	75.57	-7.41	0.9(2)
P042	2,330.8	2.05	6.0(10)	P056	42.14	-7.03	2.9(5)	LTP30	18.21	-7.5	1.1(2)
P006	18.98	-6.64	0.5(1)	P113	954.89	-4.75	0.8(2)	LTP06	1,638.6	-1.97	2.8(5)
P057	28.21	-7.33	1.1(2)	PLX4	31.52	-7.1	0.3(1)	P031	481.83	-6.06	3.0(5)
P059	20.83	-7.81	0	P115	28.77	-2.54	6.8(12)				

Complexe Terminal aquifer											
Borehole	$\text{Cl}^-$ /mmolL $^{-1}$	$\delta^{18}\text{O}$ /‰	$^3\text{H}$ /UT	Borehole	$\text{Cl}^-$ /mmolL $^{-1}$	$\delta^{18}\text{O}$ /‰	$^3\text{H}$ /UT	Borehole	$\text{Cl}^-$ /mmolL $^{-1}$	$\delta^{18}\text{O}$ /‰	$^3\text{H}$ /UT
D5F80	42.22	-7.85		D1F138	28.92	-8.13	0.7(1)	D2F71	13.53	-8.23	0.6(1)
D3F8	29.81	-8.14	1.4(2)	D3F18	21.66	-8.23	0.2(1)	D7F4	10.6	-8.27	0.1(1)
D3F26	34.68	-7.97	0.8(1)	D3F10	14.27	-7.88	1.5(2)	D2F66	11.02	-8.3	
D4F94	20.05	-8.18	0.6(1)	D6F51	28.39	-7.9	0.7(1)	D1F151	10.75	-8.32	0.4(1)
D6F67	18.79	-8.23	3.7(6)	D1F135	18.08	-7.97	1.1(2)	D6F64	11.36	-8.28	4.3(7)

Continental Intercalaire aquifer											
Borehole	$\text{Cl}^-$ /mmolL $^{-1}$	$\delta^{18}\text{O}$ /‰	$^3\text{H}$ /UT	Borehole	$\text{Cl}^-$ /mmolL $^{-1}$	$\delta^{18}\text{O}$ /‰	$^3\text{H}$ /UT	Borehole	$\text{Cl}^-$ /mmolL $^{-1}$	$\delta^{18}\text{O}$ /‰	$^3\text{H}$ /UT
Hadeb I	5.8	-8.02	0	Hadeb II	6.19	-7.93	0.1(1)	Aouinet Moussa	6.49	-7.88	1.1(2)

Table 7: Statistical parameters for Continental Intercalaire (CI), Complexe Terminal (CT) and Phreatic (Phr) aquifers samples selected on the basis of  $\delta^{18}\text{O}$  and  $\text{Cl}^-$  data (see text).

Aquifer	Size	Parameter	EC /mS cm $^{-1}$	T /°C	pH	Alk.	$\text{Cl}^-$	$\text{SO}_4^{2-}$	$\text{Na}^+$	$\text{K}^+$	$\text{Mg}^{2+}$	$\text{Ca}^{2+}$
CI	11	Average	2.2	49.0	7.5	2.3	11.0	4.7	10.3	0.51	3.6	2.4
CI	11	Stdd. dev.	0.3	2.0	0.2	1.0	4.6	2.5	4.6	0.23	2.0	1.8
CT	50	Average	3.2	23.0	7.8	2.3	20.0	8.9	17.0	1.0	5.5	5.6
CT	50	Stdd. dev.	1.1	2.4	0.4	0.8	7.0	2.6	6.0	0.8	2.2	1.7
Phr pole I	30	Average	3.9	24.0	7.9	2.3	24.7	11.8	24.2	2.1	7.2	5.3
Phr pole I	30	Stdd. dev.	1.3	1.3	0.4	1.0	6.9	3.4	11.0	1.7	5.0	2.7
Phr pole II	3	Average		23.4	7.0	2.4	4,761.0	158.0	4,021.0	32.4	500.0	13.0
Phr pole II	3	Stdd. dev.		0.1	0.5	1.6	350.0	43.0	1,093.0	28.0	378.0	8.0

Table 8: Summary of mass transfer for geochemical inverse modeling. Phases and thermodynamic database are from Phreeqc 3.0 (Parkhurst and Appelo, 2013).

Phases	Stoichiometry	CI/CT	CT/Phr I	Rainwater/P036	PhrI/PhrII 60%/40%
Calcite	CaCO <sub>3</sub>	—	$-6.62 \times 10^{-6}$	$-1.88 \times 10^{-1}$	$-2.26 \times 10^{-1}$
CO <sub>2</sub> (g)	CO <sub>2</sub>	$-6.88 \times 10^{-5}$	—	$8.42 \times 10^{-4}$	$5.77 \times 10^{-4}$
Gypsum	CaSO <sub>4</sub> · 2 H <sub>2</sub> O	$4.33 \times 10^{-3}$	—	$1.55 \times 10^{-1}$	$1.67 \times 10^{-1}$
Halite	NaCl	$7.05 \times 10^{-3}$	$3.76 \times 10^{-3}$	6.72	1.28
Sylvite	KCl	$2.18 \times 10^{-3}$	$1.08 \times 10^{-3}$	$4.02 \times 10^{-1}$	—
Bloedite	Na <sub>2</sub> Mg(SO <sub>4</sub> ) <sub>2</sub> · 4 H <sub>2</sub> O	—	$1.44 \times 10^{-3}$	—	—
Huntite	CaMg <sub>3</sub> (CO <sub>3</sub> ) <sub>4</sub>	—	—	$4.74 \times 10^{-2}$	$5.65 \times 10^{-2}$
Ca ion exchange	CaX <sub>2</sub>	$-1.11 \times 10^{-3}$	—	—	—
Mg ion exchange	MgX <sub>2</sub>	$1.96 \times 10^{-3}$	—	$1.75 \times 10^{-1}$	$-2.02 \times 10^{-1}$
Na ion exchange	NaX	—	—	—	$3.92 \times 10^{-1}$
K ion exchange	KX	$-1.69 \times 10^{-3}$	—	$-3.49 \times 10^{-1}$	$1.20 \times 10^{-2}$

Values are in mol/kg (H<sub>2</sub>O). Positive (mass entering solution) and negative (mass leaving solution) phase mole transfers indicate dissolution and precipitation, respectively; — indicates no mass transfer.




Figure 1: Localisation and schematic relations of aquifers in Ouargla. *Blue lines represent limits between aquifers, and the names of aquifers are given in bold letters; as the limit between Senonian and Miopliocene aquifers is not well defined, a dashed blue line is used. Names of villages and cities are given in roman (Bamendil, Ouargla, Sidi Khouiled), while geological/geomorphological features are in italic (Glacis, Sebkhah, Chott, Dunes). Depths are relative to the ground surface. Letters a and b refer to the cross section (fig. 2) and to the localisation map (fig. 3).*




Figure 2: Geologic cross section in the region of Ouargla.  
*The blue pattern used for Chott and Sebka correspond to the limit of the saturated zone.*




Figure 3: Localisation map of sampling point





Figure 4: Piper diagram for Continental Intercalaire (filled squares), Complexe Terminal (open circles) and Phreatic aquifer (open triangles).





5

Figure 5: Contour maps of the salinity (expressed as global mineralization) in the aquifer system, (a) Phreatic aquifer; (b) and (c) Complexe Terminal [(b) Miopliocene and (c) Senonian]; figures are isovalues of global mineralization (values in g/L).




Figure 6: Equilibrium diagrams of calcite (top) and gypsum (bottom) for Continental Intercalaire (filled squares), Complexe Terminal (open circles) and Phreatic aquifer (open triangles). Equilibrium lines are defined as:  $\log[\text{Ca}^{2+}] + \log[\text{CO}_3^{2-}] = \log K_{sp}$  for calcite, and  $\log[\text{Ca}^{2+}] + 2 \log[\text{H}_2\text{O}] + \log[\text{SO}_4^{2-}] = \log K_{sp}$  for gypsum.




Figure 7: Variation of saturation indices with distance from south to north in the region of Ouargla.




Figure 8: Change from carbonate facies to evaporite from Continental Intercalaire (filled squares), Complexe Terminal (open circles) and Phreatic aquifer (open triangles).




Figure 9: Change from sulfate facies to chloride from Continental Intercalaire (filled squares), Complexe Terminal (open circles) and Phreatic aquifer (open triangles).




Figure 10: Correlation between  $\text{Na}^+$  and  $\text{Cl}^-$  concentrations in Continental Intercalaire (filled squares), Complexe Terminal (open circles) and Phreatic aquifer (open triangles). Seawater composition (star) is  $[\text{Na}^+] = 459.3 \text{ mmol L}^{-1}$  and  $[\text{Cl}^-] = 535.3 \text{ mmol L}^{-1}$  (Stumm and Morgan, 1999, p.899).




Figure 11: Calcium vs.  $\text{HCO}_3^-$  diagram in Continental Intercalaire (filled squares), Complex Terminal (open circles), Phreatic aquifer (open triangles) and Seawater composition (star) is  $[\text{Ca}^{2+}] = 10.2 \text{ mmol L}^{-1}$  and  $[\text{HCO}_3^-] = 2.38 \text{ mmol L}^{-1}$  (Stumm and Morgan, 1999, p.899).




Figure 12: Calcium vs.  $\text{SO}_4^{2-}$  diagram in Continental Intercalaire (filled squares), Complexe Terminal (open circles), Phreatic aquifer (open triangles) and Seawater composition (star) is  $[\text{Ca}^{2+}] = 10.2 \text{ mmol L}^{-1}$  and  $[\text{SO}_4^{2-}] = 28.2 \text{ mmol L}^{-1}$  (Stumm and Morgan, 1999, p.899).




Figure 13: Chloride concentration versus  $\delta^{18}\text{O}$  in Continental Intercalaire (filled squares), Complexe Terminal (open circles) and Phreatic aquifer (open triangles) from Ouargla.




Figure 14: Log  $[\text{Cl}^-]$  concentration versus  $\delta^{18}\text{O}$  in Continental Intercalaire (filled squares), Complexe Terminal (open circles) and Phreatic aquifer (open triangles) from Ouargla.