

Response to Rereree #1

We would like to much thank Prof. Peng Gao (Referee #1) for his detailed comments and suggestions on the original manuscript. These comments and suggestions have been used to greatly improve the manuscript during the following revision. A point-by-point response to your comments is addressed below.

In this study, the authors attempted to explain the different channel planforms of four reaches in the source area of Yellow River, China using (partially) measured water discharge, stage, and cross section data, as well as qualitative description. First, I think this area is unique regarding to the world large rivers and thus is worth studying. Second, the river dynamics in this area is very complex and hence is very hard to capture. Therefore, I think this study is significant and potentially very useful for understanding the river environment in source areas for large rivers in general. However, I think the authors need to fix a series of problems in the current manuscript before it reaches the level for publication.

Response: Many thanks for your positive comments and pointing out the weakness about the manuscript. We have updated the manuscript according to your suggestions.

I describe them in details: Introduction: the authors spend too many sentences describe the progress in channel planforms, in particular braided and anabranching rivers (pages 2 to 5). Instead, I think they should reduce these part.

Response: We have shorten the introduction and focus on the knowledge gap from the existing studies. We feel it is important to contextualize this study, highlighting similarities and differences with conventional literatures. We paid very careful attention to this issue.

In the meantime, they should expand the studies on river diversity in the source area of Yellow River (lines 17-31, page 5) and explain what we need to explore further in details, which could lead to the objectives of this study. There are quite a few English problems in this section.

1 **Response: Good suggestions. We enhanced this part so as to make our studies more**
2 **sense, emphasizing how and why this intriguing yet understudied part of the work**
3 **relates to other areas. Meanwhile, the English has been polished and double-checked.**

4

5 Sections 2 and 3: these two should be combined into one section. Also, Figs 1 and 2 should be
6 combined.

7 **Response: Section 2 and 3 have been combined into one section. Figs 1 and 2 have also**
8 **been combined.**

9

10 Section 4: it is very important that the authors specify what value of Manning's n are used for
11 each of the four reaches when they introduce their model because comparison of their
12 difference would provide a quantitative means of showing the impact of vegetation on river
13 morphology. The sentence in lines 4-5 on page 9 does not make sense.

14 **Response: We have specified Manning's n , and discuss variability in roughness. We**
15 **reconsider the sentences.**

16

17 Section 5.1.1: Page 10, Lines 14-15: what is the difference between middle and high flood
18 stages? It might be more informative if this description is tied to Figure 8. For example, can
19 one say that middle flood stage may be represented by the high discharges in September and
20 high flood stage may be reflected by the high discharges in July?

21 **Response: Yes, the difference between middle and high flood stages is not very distinct.**
22 **Here the middle stage means that the channel flow partly submerges the bar surface,**
23 **but the stage does not completely inundate the vegetation. Therefore, we need to add**
24 **more explanation on the middle and high flood stages in the section.**

25

26 Page 10, Line 18: if a water depth of 2.0 m represents the bankfull discharge, then what does
27 the water depth of 3.0 m represent? Can I say that $h = 2.0$ m is the height at the top of the
28 stable bars in middle channels?

1 **Response: Sure. If a water depth of 2.0 m represents the bankfull discharge, the water**
2 **depth of 3.0m represent that 1m is inundation water depth. We can think $h=2.0m$ is the**
3 **height at the top of the stable bars surface, but does not submerge the vegetation (i.e.,**
4 **trees)**

5

6 Page 10, lines 23-32: the message delivered by this paragraph is very vague. It seems to me
7 that the data in October in both 1968 and 1984 follow the curve formed by the data in June
8 and July. If the authors believe there are significant difference between June and July, and
9 August and September, why not use the data in the two periods to run non-linear regression
10 (power function) and see if the exponents of the two are significantly different?

11 **Response: Using non-linear regression for the data in different months in both 1968 and**
12 **1984 is very good choice. We have done this job to obtain the exponents so as to**
13 **quantitatively explain the difference.**

14

15 The authors should explain quantitatively the geomorphological significance of the two
16 different trends in Fig. 8a and 8b (i.e., the trend formed by the data in March and April against
17 the trend formed by the remaining data). Also, the difference between the high scatter trend
18 for low discharges (probably low flood stages) and regular trends for high discharges
19 (probably high flood stages) should be elaborated. The key is to explain why channels in this
20 reach is semi-braided and semi-anabranching. My guess is vegetation on bars assures that
21 during low and middle flood stages, bars and islands are relatively stable, while during high
22 flood stage, they are unstable. Figure 8 should be used to make this point clear.

23 **Response: We agree with this suggestion regarding analysis on Fig.8a and Fig. 8b and**
24 **their underlying meaning. Accordingly, we have explained quantitatively the**
25 **geomorphological significance of the two different trends and why channels in this reach**
26 **are semi-braided and semi-anabranching, further, emphasizing the role of vegetation.**

27

28 Page 11, Lines 1-11: this paragraph is about Fig. 9. I think the figure shows a completely
29 different aspect of stream channels in this reach: channel morphology before 1976 is different
30 from that after 1976. This difference is represented by the two different trends of the data. The
31 authors should run non-linear regression to establish power functions for the two different

1 trends in each listed month and then link this difference to the possible difference of
2 vegetation cover in the two different time periods. This would strengthen the analysis a lot.
3 Fig. 10 is not well tied to the data shown in Figs.8 and 9. It is nice, but there lacks evidence to
4 support it.

5 **Response: OK, it is very good suggestion. We adopt non-linear regression to build power**
6 **functions in each listed month and link these differences to the possible difference of**
7 **vegetation cover. Moreover, we rethought Fig.10 and augment the analysis on Fig.9.**

8

9 Section 5.1.2: First, the authors should mentioned Fig. 11 first and then Fig. 12.

10 **Response: OK. We have corrected this.**

11

12 Second, the big problem here is that the postulation raised here (lines 15-19 on page 11) is not
13 fully supported by the only data shown in Fig. 11. The stage data in Fig. 11 are not sufficient
14 to argue the change of flow regime exactly because the channels here are anabranching
15 channels. This means that the same flow stage in different seasons might be associated with
16 different water discharges. Maybe there are no water discharge data available in this reach. If
17 this is the case, the authors should re-think their arguments: the fact that these channels are
18 stable means that sediment (bedload) supplied from upstream (i.e., the Dari reach) is balanced
19 by the sediment transport capacity in this reach. One way might be useful is to compare the
20 supplied bed load based on the prediction made for the Dari reach with the transport capacity
21 predicted in this reach. The authors should expect that they are similar or very close to each
22 other. Then, the impact of vegetation on the hydraulics might be reflected in Manning's n
23 used in the bedload model. Comparing this value with the one used in the Dari reach may
24 show the impact of vegetation on the stable status of this reach.

25 **Response: We agree with the detailed analysis above. Since Fig.11 did not fully support**
26 **our analysis, we continue to collect the data of monthly-channel discharge and monthly-**
27 **sediment transport rate in four hydrological stations (Huangheyuan, Dari, Maqu,**
28 **Tangnaihai). New data and analyses can strengthen this section, in particular, the**
29 **impact of vegetation more distinct.**

30

1 Section 5.1.3: This reach is a tributary. If the authors have no water discharge data in this
2 tributary, I suggest to delete this part completely from the current manuscript. This is because
3 only showing a postulated diagram (i.e. Fig. 13) is insufficient to convince the readers about
4 the status of this reach.

5 **Response: OK. In section 5.1.3, the Lanmucuo River is a small meandering river which**
6 **has no hydrological data, but we conducted field investigations during 2011-2015.**
7 **Especially, in 2015 we measured the cross-section and mean velocity in the middle reach.**
8 **Perhaps we delete the Fig. 13, but add other data or figure so as to keep the integrity of**
9 **this study.**

10

11 Section 5.1.4: This reach is unstable. Again, just using the temporal changes of channel
12 sections between three years (i.e., Fig. 15) is not enough to explain how vegetation affects
13 them. Again, I think it is very important for the authors to predict bedload transport rates and
14 then use them to calculate the mean sediment load in this reach. By comparing this (or these)
15 mean value(s), the authors may argue that why the reach is not stable. In the meantime,
16 comparing the value of manning's n used in this reach with those used in the first and second
17 reaches along the main river would provide evidence of the impact of vegetation on river
18 morphology.

19 **Response: Yes, the braided reach of Daheba River is quite unstable and the vegetation**
20 **effect can be ignored here. Actually, the authors have predicted bedload transport rates**
21 **in this reach. Unfortunately, there are no measured data of bedload transport rates for**
22 **comparison.**

23

24 Minor points:

25 Lines 5-8 on page 6: this description is very confusing;

26 **Response: OK. We have revised it.**

27 Lines 17-18 on page 7: why should the stable reach have high bedload transport capacity?

28 **Response: The reach is very stable because the dense trees develop on bars/islands as**
29 **well as river banks. If over-capacity bed load is incoming, the reach is very stable**
30 **because trees densely develop on bars/islands. If over-capacity bed load is incoming, the**

1 **stable anabranching channel can not be widened and keep high velocity within the**
2 **channel so as to efficiently transport bedload relative to unstable braided channel.**

3 Lines 4-5 on page 9: what does the rivers in an arid area have anything to do with rivers in the
4 study area?

5 **Response: Here we cited the references for arid area to justify the correctness of using**
6 **the Manning formula as flow resistance.**

7 Figure 1: Please mark R1, R2, R3, and R4. Also, only use the arrow to show the direction of
8 flow.

9 **Response: OK, we add R1,R2, R3, R4 in Fig.1 and correctly use the arrow.**

10 In the legend, 'Tributary' and 'Trunk stream' should be reversed. Please use 'Main stream'
11 rather than 'Trunk stream';

12 **Response: Yes, I have changed it immediately.**

13 Figures 3-6: these figures should be combined into one figure;

14 **Response: No problems. We are able to combine them into one figure.**

15 Figure8: please use the same legend for the two figures;

16 **Response: OK, I have revised it quickly.**

17 Figure 14: it does not help much in understanding the difference between the regular and
18 flood conditions;

19 **Response: We have chosed this figure and better images.**

20

21

Response to Rereree #2

We appreciate Prof. Coenders-Gerrits M. (Referee #2)'s comments and suggestions. These comments are very used to enhance the manuscript during the following revision. A point-by-point responses to each comment are addressed below.

The authors present a relevant study on the effect of vegetation on bedload transport capacity and channel stability. Therefore, they study 4 reaches of the upper Yellow River, China. The 4 reaches differ in planform. Despite the potential interest, the paper is highly descriptive and hypothetical. Barely any data is collected to justify the conclusions. This leads to the question what we can learn from this study. The river planform is not really something we can easily adjust and the role of vegetation is more a result of the planform, than a cause. Maybe this also relates to the fact that there is no study objective given.

Response: Many thanks for your objective remarks about the manuscript. We confess that the interesting phenomenon in the Yellow River source needs more data to verify our conclusion. This study about river planform of the Yellow River source is an intriguing but understudied part of the world – altitude, plateau landscapes, and its global significance, so we need strong foundation studies to set up further analyses-given data limitations, these will be inherently descriptive in the first instance, but it is important to get this right. We still believe the role of vegetation plays a great role on the planform in this region, though there is a lack of direct evidence. Perhaps we need to go further in making relations to other parts of the world, in terms of the influence of landscape and environmental setting upon river diversity that these relations are the same here, or there are some notable differences.

Abstract:

The abstract starts immediately with describing what the study entails, but the existing knowledge gap is missing. As well as the 'reason for this study'.

Response: It is a very good suggestion. We add 1-2 sentence to explain the existing knowledge gap missed and the reason of this study.

Introduction:

The introduction is really long and very general. It seems like a 'lecture' on river planforms in relation to bars. I would advise to shorten the introduction and focus on what is currently missing (knowledge

1 gap) and why this study is relevant (what will it bring). Furthermore, I would also explain how the
2 existing study differ from exiting studies.

3 **Response: OK, we are pleased to accept this valuable advice to compress the introduction. The**
4 **knowledge gap has been seriously considered and answer why this study is relevant and differs**
5 **from existing studies.**

6

7 P9L1-25:

8 Add dimensions or units to symbols

9 **Response: OK, I can do it.**

10 Equation 3-5:

11 Why do you need Eq. 4 if you can also derive it from Eq. 3 and 5?

12 **Response: Definitely Eq.4 is derived from Eq.3 and Eq.5. Eq. 3 gives us the dimensionless**
13 **bedload transport rate per channel width, but we want to obtain the dimensional bedload**
14 **transport rate per unit channel width. So keeping Eq.4 in text is reasonable.**

15 Section 5:

16 Based on what can the authors conclude how the bars are developed/eroded? (fig 10, 12,13). Can this
17 not better be answered with satellite images over several years?

18 **Response: Figure 10, 12, 13 are simple sketches of the bars in braided, anabranching and**
19 **meandering channel based on our field investigation and satellite images. Adopting the satellite**
20 **images is a good option, but the difference of water depth in the different satellite images so that**
21 **the submerging range in channel varies. After discussing with other authors, we w seriously**
22 **considered the availability of satellite images in this study.**

23 Figure 1:

24 Naming R1, R2, R3, and R4 are not visible in the figure

25 **Response: OK, I can do it.**

26 Figure 8:

27 What's happening during the low flows? This seems to weird behaviour. How can the stage drop when
28 Q increases? That is remains constant is possible if the river width increase after a certain threshold,
29 but this seems unrealistic. Please elaborate/explain.

30 **Response: Your questions make sense. We believe the data is correct. During the low flows, the**
31 **channel partly is frozen in December, January, February, March, and April. Because the water**

1 in lower layer is frozen, the stage of incoming flow increases but the discharge still very lower or
2 keep constant. Therefore, in the low flows, the stage increases when Q is nearly constant.

3

4 Figure 9:

5 Please be consistent. The upper graphs are Qh-plots, while the lower two are hQ-plots. Furthermore,
6 the coloring is not that clear, which makes the plot difficult to interpret.

7 **Response: Many thanks for pointing out this mistake. The upper and lower graphs are Q-h plots,**
8 **but the coordinate texts of the lower graphs are wrong. Meanwhile, we adopt Adobe Photoshop**
9 **CS to processing the coloring image by increasing the resolution.**

10

11 Figure 11:

12 Is the stage unit correct? What is the datum of this stage?

13 **Response: The stage value is correct. The datum of this stage is the elevation of water surface.**
14 **We will double-check the data and add the explanation in data source avoiding the**
15 **misunderstanding.**

16

17 Throughout the entire manuscript:

18 • Textual: after "i.e." and "e.g." a comma should be placed

19 **Response: Good point! I have added a comma for all "i.e." and "e.g.".**

20

21 • Order appearance figures in text, is order figure numbers (e.g., figure 11 and 12).

22 Please check

23 **Response: OK. I have updated the figures order in text.**

24

25

Reply to the editor

We deeply appreciate the constructive comments of the reviewers (Prof. Peng Gao and Prof. Coenders-Gerrits M.) and the major revisions from the Editor (Prof. Günter Blöschl) on our manuscript of 'hess-2015-526'. These suggestions are quite helpful for us and we have incorporated all comments into the revised manuscript. During the last two month, we have revised the paper so as to greatly improve the quality. We have replied all comments point-by-point. The revised words, sentences and references of the manuscript were highlighted in red color in the marked manuscript. Meanwhile, the reason and explanation will be addressed below one by one to the major revisions. The authors would like to continue to polish the paper until reaching the level of publication.

The major revisions are addressed below.

1. The authors add two sentences to address the existing knowledge gap in the Abstract and Introduction. Meanwhile, we shorten the Introduction about the bar, following the suggestion of the reviewers.
2. We have run non-linear regression (power function) for the data of the water stage and discharge in June, July, August, and September in 1966 to 1984, and run the regression analysis in July from 1964-1984, and accordingly analyze the difference of the coefficients and exponents in different periods. R2 of all regression analysis is up to 0.98.
3. We slightly adjusted the parameters of Maqu and Daheba reaches in Table 3 after discussing with co-authors.
4. Revised Figure 1 and combined the previous Figures 1 and 2 into Figure 1.
5. Figures 3-6 are combined into Figure 3.
6. Revised the legend of Figure 8.
7. Add a comma after "i.e." and "e.g."
8. Double-check the list and citation of references.

The minor revisions are addressed below.

1. Page 1, add a new affiliate in Line 6-7, and add a sentence in the Abstract in Line 23-25.
2. Page 2, revised the sentence in Line 2-3, and add a sentence in 22-23.
3. Page 5, add a citation in Line 14, and "previous" in Line 18.

- 1 4. Page 6, revise the sentence in Line 3-6, and add two citation in Line 9 and 13.
- 2 5. Page 7, revise the number of figures in Line 6-7, 14-15, and 31. And add “1.5-3.0 km
- 3 wide” in Line 13.
- 4 6. Page 8, revise the number of figures. Add “median size” in Line 13, “ the” in Line 23.
- 5 7. Page 9, add the unit for all parameters. Add “Daheba reach”, “Lanmucuo River”, “Dari
- 6 River”, and “ Maqu reach” in Line 9-11.
- 7 8. Page 10, revise the number of figures in Line 9 and 19. Add the regression analysis in
- 8 Line 25-31.
- 9 9. Page 11, revise the number of figures in Line 4-5, 23-24. Add the regression analysis in
- 10 Line 9-14 and 26-32. “occurred” was replaced by “changed” in Line 17.
- 11 10. Page 12-13, revise the number of figures. And revise two sentences in Line 14-18.
- 12 11. Page 15, revise the acknowledgements.
- 13

1 **Vegetative impacts upon bedload transport capacity and** 2 **channel stability for differing alluvial planforms in the** 3 **Yellow River Source Zone**

4
5 **Z. W. Li¹, G. A. Yu², G. Brierley³, and Z.Y. Wang⁴**

6 [1]{School of Hydraulic Engineering, **Key Laboratory of Water Sediment Sciences and Water**
7 **Disaster Prevention of Hunan Province**, Changsha University of Science & Technology,
8 Changsha, China}

9 [2]{Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of
10 Sciences, Beijing, China}

11 [3]{School of Environment, University of Auckland, Auckland, New Zealand}

12 [4]{State Key Laboratory of Hydrosience and Engineering, Tsinghua University, Beijing,
13 China}

14 Correspondence to: G. A. Yu (yuga@igsnr.ac.cn)

15 16 **Abstract**

17 The influence of vegetation upon bedload transport and channel morphodynamics is
18 examined along a channel stability gradient ranging from meandering through anabranching
19 through anabranching-braided to fully braided planform conditions along trunk and tributary
20 reaches of the Upper Yellow River in western China. Although the regional geology and
21 climate are relatively consistent across the study area, there is a distinct gradient in the
22 presence and abundance of riparian vegetation for these reaches atop the Qinghai-Tibet
23 Plateau (elevations in the study area range from 2800-3400 m a.s.l.). **To date, the influence of**
24 **vegetative impacts upon channel planform and bedload transport capacity of alluvial reaches**
25 **of the Upper Yellow River remains unclear because of a lack of hydrological and field data.**
26 **In this region**, the types and pattern of riparian vegetation vary with planform type as follows:
27 trees exert the strongest influence in the anabranching reach, the meandering reach flows
28 through meadow vegetation, the anabranching-braided reach has a grass, herb, and sparse
29 shrub cover, and the braided reach has no riparian vegetation. A non-linear relation between

1 vegetative cover on the valley floor and bedload transport capacity is evident, wherein
2 bedload transport capacity is highest for the anabranching reach, **roughly** followed by **the**
3 **braided, meandering, and anabranching-braided reaches** respectively. The relationship
4 between the bedload transport capacity of a reach and sediment supply from upstream exerts a
5 significant influence upon channel stability. Bedload transport capacity during the flood
6 season (June-September) in the braided reach is much less than the rate of sediment supply,
7 inducing bed aggradation and dynamic channel adjustments. Rates of channel adjustment are
8 less pronounced for the anabranching-braided and anabranching reaches, while the
9 meandering reach is relatively stable (*i.e.*, this is a passive meandering reach).

10

11 **1 Introduction**

12 Transitions in river character and behaviour are a key focal point of enquiry in fields such as
13 geomorphology, hydrology, and sedimentology. Such concerns have significant management
14 applications, especially relating to issues such as management of flood risk and sedimentation
15 hazards. These issues are likely to become even more pronounced in the future, as rivers
16 adjust in response to climate and land use changes, and management actions. Putting aside
17 concerns for terminological issues associated with differentiation of river types and their
18 morphological attributes (see Lewin and Ashworth, 2014; Carling et al., 2014; Tadaki et al.,
19 2014), process-based understandings of morphodynamic adjustments are required to address
20 concerns for prospective future river changes (Beechie et al., 2010). Here we evaluate the
21 influence of riparian vegetation upon process interactions along relatively understudied
22 reaches of the Upper Yellow River atop the Qinghai-Tibet Plateau in western China.
23 **Qualitative description and analysis of this complex influence on bedload transport capacity**
24 **remains unclear to our knowledge.**

25 Channel bars are products of instream deposition of bedload materials, whether at the channel
26 margin (bank-attached forms) or mid-channel bars (Brierley and Fryirs, 2005). Typically, bars
27 mutually adjust with channel geometry, such that they scale to the size of the channel in
28 which they form (Task Force on Bed Forms in Alluvial Channels, 1966; Nicholas et al., 2013).
29 If these features become vegetated and stabilized, they are referred to as islands (Fryirs and
30 Brierley, 2012). Unit bars (migrating lobate bed forms with heights and lengths that scale with
31 channel depth and width) are differentiated from larger, more complex compound bars (*e.g.*,
32 Bridge, 1993; Brierley, 1989, 1991; Smith, 1974). Compound bars are products of multiple

1 phases of accretion and reworking, with stacked sequences of unit bar, dune, and smaller bed
2 form deposits that are often trimmed at their margins by bank erosion processes or dissected
3 by chute channels (Ashworth et al., 2011; Best et al., 2003; Bridge, 2003; Brierley and Fryirs,
4 2005; McGowen and Garner, 1970; Reesink et al., 2014; Sambrook Smith et al., 2009).
5 Various studies have characterized the main morphological elements of large bars and islands,
6 while other studies have developed conceptual models of bar evolution (e.g., Ashworth et al.,
7 2000; Gurnell et al., 2001; Latrubesse and Franzinelli, 2002; Mertes et al., 1996).

8 There is notable variability in the presence, form and hydraulic/sedimentologic
9 (morphodynamic) role of bars along the continuum of channel planform (Bridge, 1993;
10 Brierley, 1996). By definition, as suspended-load rivers have limited bedload-calibre
11 materials, they have very few, if any, bars. The prominence of fine-grained (silt-clay) deposits
12 under low energy conditions (often very low channel gradient) promotes passive channel
13 behaviour, typically with a low sinuosity, passive meandering or anabranching (anastomosing)
14 planform (Eaton et al., 2010; Fryirs and Brierley, 2012; Makaske, 2001; Wang et al., 2005).
15 Patterns of bar formation in mixed- and bedload-dominated rivers reflect the flow-sediment
16 balance along any given reach, with a spectrum of planform types ranging from active
17 meandering and wandering variants through to fully braided rivers (see Ashworth, 1996;
18 Ashworth and Lewin, 2012; Burge, 2006; Church and Rice, 2009). Braiding results from the
19 inability of flow to transport all sediments that are made available to the channel, such that
20 mid-channel sedimentation occurs (i.e., competence and/or capacity limits are exceeded).
21 Recurrent reworking of bedload materials via thalweg shift during flood events alters the
22 number, shape, and location of bars. Bar dissection and avulsion create multi-thread channel
23 systems with a disorderly river planform, extremely unstable bars, and inconstant flow paths
24 (Ashmore, 1991; Ashworth et al., 2000; Jerolmack and Mohrig, 2007).

25 If channel boundary conditions induce sufficient bank strength, and flows are able to transport
26 available bedload sediments, the river adopts a configuration with better-defined, less mobile
27 channels with a much lower width-depth ratio, whether within a single-channel configuration
28 (typically passive meandering) or a multi-channel anabranching configuration (Eaton et al.,
29 2010; Song and Bai, 2015). Controversy abounds in our theoretical understanding of process
30 controls upon anabranching river behaviour (see Carling et al., 2014; Nicholas et al., 2013).
31 While Huang and Nanson (2007) and Jansen and Nanson (2004, 2010) attribute an
32 anabranching configuration to the least action principle, wherein channels adjust their form to

1 transport available sediment in the most hydraulically efficient manner, Eaton and co-workers
2 postulate an alternative theoretical framing in which anabranching channels adjust to
3 minimize their capacity to transport materials (Eaton and Church, 2004, 2007; Eaton et al.,
4 2010). It is not our concern here to address this issue directly. Rather, our focus lies with
5 analysis of relationships between bedload transport capacity and channel morphodynamics
6 along a continuum of channel planform types that is coincident with a gradation in riparian
7 vegetation cover along the Upper Yellow River (Yu et al., 2014).

8 Bar stability is the key distinguishing attribute of braided and anabranching rivers. In some
9 instances, vegetation may support the long-term development of stable sandbars within a
10 stable multi-channel system – a variant of an anabranching river (Latrubesse, 2008; Nanson
11 and Knighton, 1996; Murray and Paola, 2003; Tal and Paola, 2010). Vegetation increases
12 flow resistance and stabilizes the channel bed and bank along anabranching rivers, thereby
13 altering channel geometry, bedload transport rates, and the resulting rates and patterns of bed
14 deposition or erosion. Once a particular morphology has been formed, the configuration of
15 channels and associated distribution of bars and roughness elements fashions process
16 responses to subsequent flood events (Hooke, 1986, 2015; Hooke and Yorke, 2011; Luchi et
17 al., 2010). If critical threshold conditions are exceeded, alterations to the balance and patterns
18 of erosion and deposition processes may bring about transitions to different planform types.

19 Mutual adjustments between patterns of vegetation types (size, spacing, and density) and
20 flow-sediment dynamics (patterns and rates of erosion and deposition) vary at different
21 positions on the valley floor. Vegetation encroachment by pioneer species and successional
22 processes induce abiotic and biotic transitions in geomorphic processes from the unvegetated
23 channel bed and bar surfaces to grassland, shrubs, and treed areas at the margins of
24 bars/islands and on floodplains (Corenblit et al., 2007, 2011; Gurnell, 2014; Hickin, 1984;
25 Hupp and Osterkamp, 1996; Millar, 2000; Tooth and Nanson, 2000). Vegetation attributes
26 influence the pattern of roughness elements and the associated distribution of flow energy,
27 thereby affecting the distribution of erosional and depositional processes, and resulting
28 morphological attributes (including the grain size distribution of bed/bar materials). Hence,
29 vegetative controls influence the stability and behaviour of alluvial bed and bars, and the
30 influence of vegetation upon flow-sediment interactions, vary for differing planform types
31 (Gran and Paola, 2001; Gradzinski et al., 2003; Jang and Shimizu, 2007; McBride et al.,
32 2007).

1 Although the prominence of seasonal low flow stages and nutrient-rich fine sands may
2 support the growth of annual or perennial herbs and shrubs on mid-channel and transverse
3 bars in braided rivers, this sparse vegetation cover has negligible impact upon sediment
4 deposition and erosion patterns, and is removed easily at flood stage (Coulthard, 2005). This
5 mutual interaction between vegetation and erosion-deposition can be viewed as a threshold
6 condition: if sufficient vegetation establishment occurs, resistance may exceed the erosion-
7 deposition capability of a normal flood such that stabilization ensues, prospectively altering
8 sedimentation patterns, increasing bank strength, and reducing channel width-depth ratio
9 (Gran and Paola, 2001; Coulthard, 2005; Eaton *et al.*, 2010). In anabranching channels the
10 vegetation cover on mid-channel bars inhibits lateral migration, inducing a stable branching
11 channel condition. During lower frequency floods, when bars are partially or completely
12 submerged by flow, vegetation increases flow resistance, traps sediment, and inhibits erosion.

13 This study builds upon previously-reported exploratory analyses of river diversity in the
14 source zone of the Yellow River (Blue *et al.*, 2013; Brierley *et al.*, 2016; Li *et al.*, 2013; Yu *et al.*,
15 2014). In this region, herbs and sparse shrubs that establish on the sand/gravel bars of
16 braided rivers have a trivial influence upon channel morphodynamics, while establishment of
17 dense shrubs and sparse trees on sand/gravel bars promotes the emergence of anabranching
18 channel configurations. Building on these *previous* observations, here we appraise process
19 interactions along a vegetative gradient of river morphologic adjustments for four reaches:
20 Dari and Maqu reaches of the Yellow River main stream, and Daheba and Lanmucuo River
21 tributaries of the Upper Yellow River (Table 1). Dari reach has a semi-stable braided channel,
22 where sandbars are covered by herbaceous vegetation and sparse shrubs. Maqu reach has a
23 very stable anabranching channel with dense willows (*Salix atopantha*) on sandbars. The
24 study reach along Lanmucuo River has a stable gravel meandering river with herb coverage.
25 The study reach along Daheba River has a very unstable gravel braided channel without
26 vegetation cover. We develop and apply a simplified model to explain the interaction between
27 sediment transport capacity and river bed deposition in these reaches, examining the effect of
28 vegetation resistance and adjustment of fluvial hydraulic geometry. From this, we
29 quantitatively analyse the stability and evolution of braided, anabranching, and meandering
30 reaches during flood events.

31

1 2 Study area and methods

2 Upstream of Tangnaihai hydrological station the source zone of the Yellow River drains an
3 area of 132,000 km² (see Fig. 1(a)). In the 1950s the Yellow River Conservancy Commission
4 established four hydrological stations along the Yellow River in this area, namely (from
5 upstream to downstream), Huangheyan station in Maduo County, Jimai station in Dari County,
6 Maqu station in Maqu County, and Tangnaihai station in Xinhai County. The reach from
7 Huangheyan to Jimai station is 325 km long and drains an area of 24,089 km². In this reach
8 the valley is quite wide, with semi-braided and semi-anabranching planform morphologies
9 characterized by disordered channels with many bars (Brierley et al., 2016). The reach from
10 Jimai to Maqu is 585 km long and drains an area of 41,029 km². The upper section of this
11 reach has a deeply incised (confined), sinuous valley between the Anyemaqen and Bayan Har
12 Mountains. Flowing into the Ruoergai alluvial basin, there is a diverse array of planform
13 types (Blue et al., 2013; Brierley et al., 2016; Li et al., 2013). The reach from Maqu to
14 Tangnaihai station is 373 km long and drains an area of 35,924 km². Most of this reach
15 comprises a steep and incised canyon, with many deeply carved meander bends.

16 For this study, field investigations of vegetative influences upon bed/bar geomorphic
17 processes were conducted four times in the summers of 2011-2014. Particle size distributions
18 of bed and bank materials size were analyzed using a laser particle size analyzer (Mastersizer
19 2000) and field sieves were used to test ten samples of river bed and bank materials in each
20 reach. As a supplement, photographs of gravel and cobbles on the bed/bar surface were taken
21 to visually estimate bed particle size. To estimate bedload transport capacity, channel
22 geometry was assessed in the field and from remote sensing images of the branching channel
23 network (Google Earth images from 2005-2014, with a resolution of about 0.24 m).

24 The best available hydrological data that could be accessed for this study were daily stage-
25 discharge data from Jimai (1964-1985), monthly stage-discharge data from Maqu (1959-
26 1970), monthly cross-section elevation change data from Shangcun station along the Daheba
27 River (1.8 km upstream from its confluence with the Yellow River, 2009-2011), and 2011-
28 2014 field data for the Lanmucuo River (a tributary of the Yellow River in Maqu-Tangnaihai
29 section, at an elevation of 3400-4200 m a.s.l., for which upstream and mid-catchment reaches
30 have a typically meandering channel, while the downstream reach has a confined bedrock
31 channel). There are no intensive human activities in this area of the Yellow River Source
32 Zone.

1

2 **3 Basic characteristics of four alluvial reaches**

3 **Fig.1 (b)** shows the planform morphology of the four channel reaches. Figure 2 shows the
4 channel morphology, pattern of bar types, and bed sediment. Basic channel characteristics of
5 the study reaches are summarized in Table 2.

6 Dari reach has a semi-braided and semi-anabranching channel in a wide valley (**Fig.1 (R1)**
7 **and Fig. 2(R1)**). This braided-anabranching transition zone is considered to be semi-stable,
8 with an active channel zone that is around 1 km wide. The braided part of the channel is
9 made up of many small longitudinal and transverse bars, with multiple connected branching
10 channels. In the anabranching part, the large bars/islands have a dense grassland vegetation.
11 Given the extensive width of the active channel zone, annual floods during June-September
12 exert negligible impacts upon these relatively stable surfaces.

13 The Maqu reach is located in a wide alluvial valley (**1.5-3.0 km wide**). The dense tree cover
14 of the vegetated islands is indicative of a stable channel configuration (see **Fig. 1(R2) and Fig.**
15 **2(R2)**). During the flood season, tree trunks are partly submerged into water, but the trees are
16 sufficiently dense to limit erosion. As a result, the anabranching system as a whole is quite
17 stable with high bedload transport capacity.

18 Lanmucuo River is a meadow meandering river with nearly 100% vegetation cover (see **Fig.**
19 **1(R3) and Fig. 2(R3)**). The root system of riparian grasses induces considerable protection
20 from near-bank erosion. Field investigations from 2011-2014 indicate that the lateral
21 migration induced by cantilever bank failure occurred at a rate of less than 0.2 m/yr. The
22 gravel-bed channel has a low bedload transport rate in the flood season. In some local sections,
23 mid-channel bars with dense grass coverage have developed at the apex of bends. The whole
24 channel is quite stable, in spite of short-term outer bank failures and long-term meander neck
25 cutoffs.

26 Daheba River has incised into the Gonghe-Xinhai sedimentary basin. Severe gully erosion has
27 incised river-lacustrine sediments to a depth of 50-100 m, supplying large volumes of
28 gravel/cobble to the middle and lower Daheba channel, inducing significant bed aggradation
29 and the formation of a braided planform. Alluvial fans at gully outlets not only supply
30 additional sediment, but also push the channel to the opposite side of the valley floor (a big
31 fan is shown near D point in **Fig. 1(R4) and Fig. 2(R4)**). As a result, the main branching
32 channels are subjected to frequent and recurrent avulsion. Flows erode new small branching

1 channels during the flood season, but a main channel coexists with several branching channels
2 in the non-flood season. Unstable mid-channel bars are unvegetated other than sparse
3 vegetation coverage (grass and shrubs) on riparian banks. The gravel-cobble bed and high
4 bedload transport rate restrict vegetation establishment and growth, resulting in a typically
5 unstable braided river.

6 Bank strength induced by sediment material mix and vegetation root networks exerts a critical
7 influence upon the stability of alluvial channels (Eaton and Giles, 2009). Reinforcement of
8 bank strength reinforced by grass, shrub, and tree roots is related to the density, depth, and
9 spatial structure of the root network (Abernethy and Rutherford, 2001). Fig.3 shows
10 representative photographs of river banks in the four study reaches. The diverse bank material
11 composition and vegetation cover affect the relative strength of banks and their capacity to
12 resist nearbank flow scour. The river bank in Dari reach has a two-layer structure, with a 20-
13 30 cm deep soil-root layer (d_{50} is median size, $d_{50} = 0.02$ mm) lying atop a gravel-sand layer
14 ($d_{50} = 6.0$ mm) (Fig. 3(a)). The river bank in Maqu reach has a dense grass, shrub, and tree
15 cover (Figure 3(b)), with no indication of flow scour in the flood season. The study reach
16 along the Lanmucuo River has a typical composite bank sedimentology of a mixed load river
17 (Fig. 3(c)). An upward-fining sequence is characterized by a basal gravel unit ($d_{50} = 5.5$ mm)
18 extending to a 10-30 cm thick silt/sand layer ($d_{50} = 0.03$ mm) that is capped by a 10-50 cm
19 thick fine-grained soil-root complex ($d_{50} = 0.02$ mm). Conversely, the bank of the middle
20 Daheba River has characteristic deposits of a bedload-dominated river, with non-cohesive
21 gravels and a sparse grass cover (Fig. 3(d)). Adjacent terraces that are more than 10m high
22 limit the capacity for channel widening, while actively supplying gravels. Mobile gravel
23 banks influence the braided characteristics of the Daheba River. In summary, bank strength of
24 the four study reaches varies from high to low as follows: Maqu reach, Lanmucuo River, Dari
25 reach, and Daheba River.

26

27 **4 Estimation of bedload transport capacity**

28 Given the lack of observed data of bed load transport rate, bedload transport capacity has been
29 estimated for a rectangular cross-section using the theoretical bed load formulae outlined
30 below. Channel flow follows the laws of flow continuity, flow resistance and sediment
31 transport with flow continuity law taking the form:

$$1 \quad Q = AV \quad (1)$$

2 where Q , A , and V are flow discharge (m^3/s), channel cross-sectional area (m^2), and average
3 flow velocity (m/s), respectively, $A=WH$, W is channel width (m), H is water depth (m).

4 This study adopts the Manning formula to embody the law of flow resistance for uniform
5 alluvial channel flow:

$$6 \quad V = \frac{1}{n} R^{2/3} S^{1/2} \quad (2)$$

7 where R is hydraulic radius (m), $R=WH/(2H+W)$, S is flow energy slope, n is Manning's
8 roughness coefficient. In this study, following Chow (1959), $n = 0.050$ if no vegetation in
9 gravel-bed channels at high stages (i.e., **Daheba reach**), $n = 0.030$ in floodplain with short
10 grass (i.e., **Lanmucuo River**), $n = 0.050$ in floodplain with scattered brush and heavy weeds
11 (i.e., **Dari River**), and $n = 0.150$ in floodplain with dense willows at flood stage (i.e., **Maqu**
12 **reach**).

13 Among numerous bedload formulae, the Meyer-Peter and Muller equation has been
14 extensively and successfully applied (Meyer-Peter and Müller, 1948). The modification
15 developed by Wong and Parker (2006) has been used in this study:

$$16 \quad \Phi = 3.97(\Psi - 0.0495)^{3/2}. \quad (3)$$

17 where Φ and Ψ are the dimensionless bedload transport rate per unit channel width and the
18 dimensionless flow shear stress, respectively, that are defined as

$$19 \quad \Phi = \frac{q_b}{\sqrt{(\rho_s / \rho - 1) g d_{50}^3}}. \quad (4)$$

$$20 \quad \Psi = \frac{RS}{(\rho_s / \rho - 1) d_{50}}. \quad (5)$$

21 where q_b is the dimensional bedload transport rate per unit channel width ($\text{kg}/\text{s}/\text{m}$), ρ_s is the
22 density of sediments transported (kg/m^3), ρ is the density of water (kg/m^3), g is the
23 acceleration of gravity (m/s^2), and d_{50} is the median sediment size (mm).

24 Cross-section and water depth were measured based on field survey and remote sensing
25 images (see Table 2). Estimated hydraulic parameters and bedload transport capacity for the
26 four reaches, derived using Eq.(1)-(5), are summarised in Table 3. Note that channel width is

1 effective bankfull width in the flood season, not valley width. The adopted mean grain size is
2 lower than bed sediment size. Results shown here are considered to be approximations, and
3 are analysed solely in relational rather than absolute terms. Results show that the bedload
4 transport capacity of the four reaches from high to low is as follows: Maqu, Daheba ,
5 Lanmucuo, Dari reaches.

6

7 **5 Effect of vegetation and bedload capacity on channel stability**

8 **5.1.1 Dari reach (braided/anabranching river with grass and shrub cover)**

9 Dari reach is a wide semi-braided and semi-anabranching channel, where the channel width is
10 up to 1600 m (Fig.2 R1(a)). Some large stable gravel bar or islands have a dense grass and
11 sparse shrub cover. Many unstable bars with low vegetation cover are subjected to recurring
12 erosion and channel adjustment. Vegetation may inhibit erosion and enhance bar stability at
13 middle flood stage, but the resistance effect of vegetation at high flood stage is very limited.
14 As a result, the whole channel may be eroded at high flow stage, resulting in disordered
15 patterns of mid-channel gravel bars. The estimated bedload transport capacity per unit channel
16 width is 1.77 kg/m/s for 2.0 m water depth (see Table 2). If the water depth increases to 3.0 m
17 in the flood season, bedload transport capacity per unit width significantly increases up to
18 14.93 kg/m/s. It is likely that these flow depths cause intense erosion that divides the stable
19 bars into many unstable bars.

20 Fig. 4(a) and (b) show monthly stage-discharge relationships for 1968 and 1984, respectively.
21 Since Dari reach is a multi-thread channel system, the stage-discharge relationship is not a
22 single function relationship. In non-flood months (December, January, February, March, and
23 April) the river bed is frozen. May and November are pivotal times in the stage-discharge
24 relationship (the former reflects ice melt, the latter freezing). In flood months (June, July,
25 August, and September) the stage-discharge relationship adjusts due to strongly erosion and
26 deposition within the channel. The stage-discharge data (June, July, August, and September)
27 in the 1968 and 1984 are to run non-linear regression (power function, $Z = a * Q^b$, Z is water
28 stage (m), Q is discharge (m^3/s), a is the coefficient and b is the exponent). Two coefficients of
29 a (b) in 1968 are 4.7(0.09), 4.8(0.09), 5.4(0.07), and 4.5(0.10) for June, July, August, and
30 September, respectively; and accordingly, 2.9(0.12), 2.8(0.13), 3.0(0.12), and 3.5(0.09) in
31 1984. The results of the regression show that a obviously decreased and b was almost

1 unchanged, indicating the increase of water depth slowed down with the incoming discharge
2 increasing from 1968 to 1984 because the sediment deposition led to the wider channel
3 year by year.

4 For instance, different discharges for the same flow stage in June and July 1968 are
5 considered to reflect erosion of the channel (Fig. 4(a)). In the other instance shown here, the
6 maximum discharge in 1984 occurred in July (Fig.4(b)), probably marking the transition from
7 erosion to deposition phases. The geomorphological significance of the two different trends is
8 showed in Fig. 4 (a) and (b) (i.e., the trend formed by the data in March and April against the
9 trend formed). Also, there is a difference between the high scatter trend for low discharges
10 (probably low flood stages) and regular trends for high discharges (probably high flood stages)
11 because the water submerged the bars in high discharge and the multi-thread channels
12 appears in low discharges. The Dari reach is defined in this study as semi-braided and semi-
13 anabranching since vegetation (grass and some shrubs) partially develops on channel bars,
14 and bars are relatively stable during low and middle flood stages, while are prone to change
15 during high flood stages.

16 Fig. 5 shows the stage-discharge relationships of the Upper Yellow River at Dari from June to
17 September in 1964-1984. Apparently, the stages of 1975 are out of line with 1978, perhaps
18 indicating that the elevation benchmark of the station changed in 1976 or 1977. In the same
19 month of different years, the stage-discharge relationship does not have a simple
20 corresponding relation, especially in August and September. This may reflect: 1) responses of
21 the channel bed to strong deposition in June and July, and thereafter the high stage
22 corresponds to low discharge such as August in 1978-1984 and September in 1964-1975; 2)
23 the channel bed strongly erodes in June and July, and thereafter the high stage corresponds to
24 high discharge such as August in 1964-1975 and September in 1978-1984. Overall, Figure 2
25 R1(a) and (b) indicate that the channel of Dari reach is quite unstable during the flood season,
26 with erosion and deposition changing the stage-discharge relationship. A sketch showing how
27 flow erosion divides bars and deposits to form new bars is shown in Fig. 6. The stage-
28 discharge data in July from 1964-1984 are to run power function regression ($Z = a * Q^b$). Two
29 coefficients of a (b) are 5.3(0.08) in 1964, 4.7(0.10) in 1965, 5.4(0.07) in 1966, 4.8(0.09) in
30 1968, 5.2(0.07) in 1969, 5.3(0.07) in 1970, 5.3(0.06) in 1973, 4.6(0.10) in 1975, 2.9(0.11) in
31 1978, 2.7(0.13) in 1979, 2.7(0.13) in 1980, 2.4(0.15) in 1981, and 2.8(0.13) in 1984.

1 Obviously, this difference of a (b) is represented by the two different trends of the data before
2 and after 1976, i.e, a decreased and b increased.

3

4 **5.1.2 Maqu reach (anabranching river with tree cover)**

5 Maqu reach in wide Ruorgai basin is covered by dense tress (*Salix atopantha*) and has a
6 stable anabranching channel planform (Fig. 2 R2(a)). It is postulated that a herb and shrub
7 cover gradually supports the stabilization of new bars, facilitating sediment deposition on the
8 body of the bar during low and middle flood stages, and protecting the bar from erosion at
9 high flood stages. Subsequent development of trees presents a tall green barrier in the flood
10 period. Although the water floods trees, their density induces sufficient resistance to decrease
11 the flow velocity and trap fine sand and gravel on the body of the bar. Therefore, this
12 anabranching channel system is basically stable over a decadal timescale.

13 Water stage change at Maqu station from 1959-1970 is shown in Fig. 7. The stage peak
14 occurs in July and September. The maximum difference of 2.43 m occurred between June and
15 September in 1963. If the water depth increases to 8.0 m from 4.0 m, bedload transport
16 capacity increases to 18.52 kg/s/m from 2.75 kg/s/m. As a result, the branching channel bed
17 may erode if the transport capacity exceeds upstream sediment supply. However, protection
18 by trees is strong enough to inhibit erosion of bars. In contrary, if the upstream sediment
19 supply surpasses the transport capacity, increasing bed deposition with flow stage further
20 increases the transport capacity of the reach. This agrees with analyses by Huang and Nanson
21 (2007) who stated that anabranching channels can achieve the optimal transport efficient
22 without increasing bed gradient. Even though these reaches may appear to promote deposition
23 on the channel bed during extreme floods (see Fig. 8), the flow erodes the bed later in the
24 flood season, thereby maintaining an equilibrium cross-section. As a result, the anabranching
25 channel of Maqu reach maintains a long-term stable situation.

26 **5.1.3 Lanmucuo River (passive meandering river with meadow cover)**

27 Lanmucuo River is a typical meandering river covered by dense meadow. Although typically
28 characterized by large bends in a flat valley, mid-channel gravel bar covered by herbs
29 sometimes form at the apex of bends (Fig. 2 R3(a)). The meandering channel and bars are
30 very stable because of low sediment supply in the flood season and good vegetation coverage.

1 The tight root-soil complex on concave banks inhibits flow scour. When cantilevered bank
2 failures do occur, slump blocks restrict further erosion of the bank. Grass develops on the
3 point bars of convex banks. If the overbank flow submerges the point bar, the herbaceous
4 vegetation can increase flow resistance and promote fine sand deposition (Fig. 9), thereby
5 maintaining channel geometry with a relatively low migration rate. Growth of herbs on mid-
6 channel bars and apices (Fig. 2 R3(b)) helps to increase the flow resistance and trap fine
7 sediment, facilitating channel stability.

8

9 **5.1.4 Daheba River (unvegetated braided river)**

10 The gravel bed of Daheba River is characterized by deposition in the flood season and erosion
11 in the non-flood season. This makes it difficult for vegetation to develop on bars and banks of
12 the braided channels. Fig. 10 shows morphological changes of the riverbed before and after
13 the flood season in 2005. The main branching and sub-branching channels of the channel
14 completely changed, with an initial phase of sediment deposition followed by flood-induced
15 division of bars and the re-emergence of a multi-thread braided system. Table 3 shows
16 derived estimates of the bed load transport capacity per width, $q_b=2.25$ kg/s/m. This capacity
17 is seemingly unable to efficiently transport the excess sediment supply from upstream. As a
18 result, serious deposition occurs along Daheba River in the flood season.

19 Adjustments to channel geometry as a result of erosion and deposition processes before,
20 during and after the flood season are shown in Fig. 11. The elevation of the riverbed on July
21 29, 2009 was 0.27 m higher than on April 1, 2009. Other than slight erosion of the left bank,
22 the subsequent phase was depositional, with up to 1.59 m of aggradation occurring by
23 October 23, 2009. The elevation of riverbed was increased by 0.27 m after the flood season in
24 2010. The elevation of the riverbed in July 1, 2011 was 0.26 m higher than on April 29, 2011.
25 Trivial deposition occurred from July 1 to July 8, but 0.24 m of erosion occurred by July 23,
26 with subsequent deposition of 0.27 m by October 23. As a result, the riverbed elevation was
27 0.24 m higher after flood season in 2011, but multiple phases of deposition and erosion has
28 occurred. The deposition-erosion-deposition phases may reflect lower bedload transport
29 capacity relative to sediment supply in the early flood season, but widespread deposition
30 increases the local bed slope, thereby increasing bedload transport capacity. According to Eq.
31 (3), a 10% increase in bed slope increases the transport capacity by 85% in Daheba reach, so

1 bed erosion occurs again. Bed erosion decreases the bed slope until the transport capacity has
2 adjusted to reduced sediment supply, thereby inducing riverbed deposition once more.
3 Consequently, alternative deposition and erosion leads to the extreme instability in the middle
4 and lower Daheba River.

5

6 **6 Discussion and Conclusions**

7 This study has outlined the complex interplay between flow and sediment supply in the flood
8 season, and the geomorphic/hydrodynamic role of vegetation cover on the valley floor, as
9 determinants of channel morphodynamics/stability and bedload transport capacity for four
10 alluvial reaches of the Yellow River source zone. Although the elevation of four reaches is
11 different (Dari = 3960 m, Maqu = 3465 m, Lanmucuo River = 3604 m, and Daheba = 2832
12 m), the precipitation, temperature, and bed sediment size are basically similar. Nevertheless,
13 vegetation coverage in the four reaches is quite different. The Dari reach (anabranching-
14 braided) has a herb and shrub cover, Maqu (anabranching) reach has trees, Lanmucuo River
15 (meandering) has meadow, and Daheba River (braided) has no vegetation cover. As shown
16 elsewhere, bar morphodynamics vary markedly for differing planform types, with key
17 differences outlined here for braided, anabranching, and meandering channels (cf., Hooke,
18 1986; Kleinhans, 2010; Kleinhans and van den Berg, 2010; Church and Ferguson, 2015). Bar
19 development and stability reflect the ability of vegetation to trap sediments and stabilize
20 banks, which in turn is directly influenced by flow energy relationships (*i.e.*, these are mutual
21 adjustments; Corenblit et al., 2007; Gurnell et al., 2012; Gurnell, 2014; Osterkamp and Hupp,
22 2010; Pietsch and Nanson, 2011). In this study, riparian vegetation and its root network are
23 considered to restrict channel width and increase hydraulic efficiency, inducing greater
24 bedload transport capacity in multi-thread channels (Allmendinger et al., 2005; Huang and
25 Nanson, 2007). Islands and floodplains are able to trap more fine-grained sediment in the
26 flood season, enhancing the longer-term (decadal) stability of anabranching channels, as
27 shown by the stable islands of Maqu reach.

28 Relative to the passive (resisting) role of vegetation, bedload transport actively affects short-
29 term patterns and rates of bed erosion and deposition. This, in turn, is affected by
30 relationships between the flow regime (especially flood events and formative flows) and the
31 influence of sediment supply upon bedload transport for differing river types (Church and
32 Ferguson, 2015; Dunne et al., 2010). The supply of bed material sediment to an alluvial

1 channel accelerates the growth of longitudinal, transverse, and point bars, thereby enhancing
2 thalweg development and locally increasing flow velocity. Non-equilibrium between
3 sediment supply and transport induces local channel instability, accentuating either bed
4 erosion or deposition (Jansen and Nanson, 2010; Nanson and Huang, 2008). In this study, a
5 channel stability gradient accords with both sediment movement and vegetation cover,
6 wherein bedload transport capacity (a function of bed slope, hydraulic geometry, and
7 sediment particle size) is related to the influence of riparian vegetation upon channel
8 geometry/planform. We contend that the differing vegetation cover and planform response
9 reflects the delicate balance between erosion and deposition on the channel bed and bank as
10 influenced by bedload sediment supply in the flood season. Only when the bedload transport
11 capacity is equivalent or greater than sediment supply, does vegetation act as a key
12 determinant of channel stability.

13

14 **Author contributions.** Z.W. Li and G.A. Yu designed and conducted the field
15 investigations. G. Brierley and Z.Y. Wang supervised the research and helped to contextualize
16 the findings. Z.W. Li prepared the manuscript with contributions from all co-authors.

17

18 **Acknowledgements**

19 This study is funded by [the CRSRI Open Research Program \(Program CKWV2016369/KY\)](#),
20 Alexander von Humboldt Foundation, the Natural Science Foundation of China (NSFC,
21 [41571009; 41330751; 91547112; 91547113](#)), and the [Key Laboratory of Water Sediment](#)
22 [Sciences and Water Disaster Prevention of Hunan Province \(2016SS04\)](#). X.Z. Wang, C.D.
23 Zhang, and X.D. Zhou are acknowledged for field assistance (2011-2014). We deeply
24 appreciate the careful comments and constructive suggestions from two reviewers (Prof. Peng
25 Gao and Prof. Coenders-Gerrits M.).

26

27 **References**

28 Abernethy, B. and Rutherford, I. D.: The distribution and strength of riparian tree roots in
29 relation to riverbank reinforcement. *Hydrol. Process.*, 15(1), 63–79, 2001.

- 1 Allmendinger, N.E., Pizzuto, J.E., Potter, N. Jr., Johnson, T.E., and Hession, W.C.: The
2 influence of riparian vegetation on stream width, eastern Pennsylvania, USA. *GSA Bulletin*,
3 117, (1-2), 229-243, 2005.
- 4 Ashmore, P.E.: How do gravel-bed river braid? *Canadian J. of Earth Sci.*, 28, 326-341, 1991.
- 5 Ashworth, P.J.: Mid-channel bar growth and its relationship to local flow strength and
6 direction. *Earth Surf. Proc. and Landforms*, 21, 103-123, 1996.
- 7 Ashworth, P.J., Best, J.L., Roden, J.E., Bristow, C.S., and Klaassen, G.J.: Morphological
8 evolution and dynamics of a large, sand braid-bar, Jamuna River, Bangladesh. *Sedimentology*,
9 47, 533-555, 2000.
- 10 Ashworth, P.J. and Lewin, J.: How do big rivers come to be different? *Earth-Science Reviews*,
11 114, 84-107, 2012.
- 12 Ashworth, P.J., Sambrook Smith, G.H., Best, J.L., Bridge, J.S., Lane, S.N., Lunt, I.A.,
13 Reesink, A.J.H., Simpson, C.J., and Thomas, R.E.: Evolution and sedimentology of a channel
14 fill in the sandy braided South Saskatchewan River and its comparison to the deposits of an
15 adjacent compound bar. *Sedimentology*, 58, 1860-1883, 2011.
- 16 Beechie, T.J., Sear, D.A., Olden, J.D., Pess, G.R., Buffington, J.M., Moir, H., Roni, P., and
17 Pollock, M.M.: Process-based principles for restoring river ecosystems. *BioScience*, 60(3),
18 209-222, 2010.
- 19 Best, J.L., Ashworth, P.J., Bristow, C. S., and Roden, J. E.: Three-dimensional sedimentary
20 architecture of a large, mid-channel sand braid bar Jamuna River, Bangladesh. *J. of*
21 *Sedimentary Res.*, 73, 516-530, 2003.
- 22 Blue, B., Brierley, G. J., and Yu, G.A.: Geodiversity in the Yellow River source zone. *J. of*
23 *Geographical Sciences*, 23(5), 775-792, 2013.
- 24 Bridge, J.S.: Description and interpretation of fluvial deposits: a critical perspective.
25 *Sedimentology*, 40(4), 801-810, 1993.
- 26 Bridge, J.S.: *Rivers and Floodplains: Forms Processes and Sedimentary Record*, Blackwell
27 Publishing, London, 2003.
- 28 Brierley, G.J.: Bar sedimentology of the Squamish River, British Columbia: Definition and
29 application of morphostratigraphic units. *J. of Sedimentary Petrology*, 61, 211-225, 1989.

- 1 Brierley, G.J.: Floodplain sedimentology of the Squamish River, British Columbia: relevance
2 of element analysis. *Sedimentology*, 38, 735–750, 1991.
- 3 Brierley, G.J.: Channel morphology and element assemblages: A constructivist approach to
4 facies modelling. In: Dawson, M. and Carling, P.A. (Eds.). *Recent advances in fluid
5 mechanics and alluvial stratigraphy*. Chichester, Wiley Interscience, 1996.
- 6 Brierley, G.J. and Fryirs, K. A.: *Geomorphology and River Management: Applications of the
7 River Styles Framework*. Blackwell Publications, Oxford, UK, 2005.
- 8 Brierley, G.J. and Hickin, E.J.: Channel planform as a non-controlling factor in fluvial
9 sedimentology: The case of the Squamish River floodplain, British Columbia. *Sedimentary
10 Geology*, 75, 67–83, 1991.
- 11 Brierley, G.J., Li, X.L., Cullum, C., and Gao, J.: *Landscape and ecosystem diversity,
12 dynamics and management in the Yellow River source zone*. Springer, Switzerland, 2016.
- 13 Burge, L.M.: Stability, morphology and surface grain size patterns of channel bifurcation in
14 gravel-cobble bedded anabranching. *Earth Surf. Proc. and Landforms*, 31, 1211–1226, 2006.
- 15 Carling, P., Jansen, J., and Meshkova, L.: Multichannel rivers: their definition and
16 classification. *Earth Surf. Proc. and Landforms*, 39, 26–37, 2014.
- 17 Chow, V.T.: *Open Channel Hydraulics*, McGraw-Hill, New York, 1959.
- 18 Church, M. and Ferguson, R.I.: Morphodynamics: River beyond steady state. *Water Resour.
19 Res.*, 51: 1883–1897, doi: 10.1002/2014WR016862, 2015.
- 20 Church, M., Rice, S.P.: Form and growth of bars in a wandering gravel-bed river. *Earth Surf.
21 Proc. and Landforms*, 34, 1422–1432, 2009.
- 22 Corenblit, D., Baas, A.C.W., Bornette, G., Bornette, G., Darrozes, J., Delmotte, S., Francis,
23 R.A., Gurnell, A.M., Julien, F., Naiman, R.J., and Steiger, J.: Feedbacks between
24 geomorphology and biota controlling earth surface processes and landforms: A review of
25 foundation concepts and current understandings. *Earth-Science Reviews*, 106(3–4), 307–331,
26 2011.
- 27 Corenblit, D., Tabacchi, E., Steiger, J., and Gurnell, A.M.: Reciprocal interactions and
28 adjustments between fluvial landforms and vegetation dynamics in river corridors: a review of
29 complementary approaches. *Earth-Science Reviews*, 84(1–2), 56–86, 2007.

- 1 Coulthard, T.J.: Effects of vegetation on braided stream patterns and dynamics. *Water Resour.*
2 *Res.*, 41, W04003, doi:10.1029/2004WR003201, 2005.
- 3 Dunne, T., Constantine, J.A, and Singer, M.B.: The role of sediment transport and sediment
4 supply in the evolution of river channel and floodplain complexity. *Transactions, Japanese*
5 *Geomorphological Union*, 31–32, 155–170, 2010.
- 6 Eaton, B.C., Church, M., and Millar, R.G.: Rational regime model of alluvial channel
7 morphology and response. *Earth Surf. Proc. and Landforms*, 29(4), 511–529, 2004.
- 8 Eaton, B.C. and Church, M.: Predicting downstream hydraulic geometry: A test of rational
9 regime theory. *J.of Geophys. Res.*, 112, F03025, doi: 10.1029/2006JF000734, 2007.
- 10 Eaton, B.C. and Giles, T.R.: Assessing the effect of vegetation-related bank strength on
11 channel morphology and stability in gravel-bed streams using numerical models. *Earth Surf.*
12 *Proc. and Landforms*, 34(5), 712–724, 2009.
- 13 Eaton, B.C., Millar, R.G., and Davidson, S.: Channel patterns: braided, anabranching, and
14 single-thread. *Geomorphology*, 120, 353–364, 2010.
- 15 Fryirs, K.A. and Brierley, G.J.: *Geomorphic analysis of river systems: an approach to reading*
16 *the landscape*. Wiley-Blackwell, Chichester, UK, 2012.
- 17 Gradzinski, R., Baryla, J., Doktor, M., Gmur, D., Gradzinski, M., Kedzior, A., Paszkowski,
18 M., Soja, R., Zielinski, T., and Zurek, S.: Vegetation-controlled modern anastomosing system
19 of the upper Narew River (NE Poland) and its sediments. *Sedimentary Geology*, 157, 253–276,
20 2003.
- 21 Gran, K. and Paola, C.: Riparian vegetation controls on braided stream dynamics. *Water*
22 *Resour. Res.*, 37(12), 3275–3283, 2001.
- 23 Gurnell, A.: Plants as river system engineers. *Earth Surf. Proc. and Landforms*, 39(1), 4–25,
24 2014.
- 25 Gurnell, A. M., Bertoldi, W., and Corenblit, D.: Changing river channels: The roles of
26 hydrological processes, plants and pioneer fluvial landforms in humid temperate, mixed load,
27 gravel bed rivers. *Earth-Science Reviews*, 111,129–141, 2012.
- 28 Gurnell, A.M., Petts, G.E, Hannah, D.M., Smith, B.P., Edwards, P.J., Kollmann, J., Ward,
29 J.V., and Tockner, K.: Riparian vegetation and island formation along the gravel-bed Fiume
30 Tagliamento, Italy. *Earth Surf. Proc. and Landforms*, 26, 31–62, 2001.

- 1 Hickin, E.J.: Vegetation and river channel dynamics. *Canadian Geographer*, 28(2), 111–126,
2 1984.
- 3 Hooke, J.M.: The significance of mid-channel bars in an active meandering river.
4 *Sedimentology*, 33, 839–850, 1986.
- 5 Hooke, J.M.: Variations in flood magnitude-effect relations and the implications for flood risk
6 assessment and river management. *Geomorphology*, 251, 91–107, 2015.
- 7 Hooke, J.M. and Yorke, L.: Channel bar dynamics on multi-decadal timescales in an active
8 meandering river. *Earth Surf. Proc. and Landforms*, 36, 1910–1928, 2011.
- 9 Huang, H.Q. and Nanson, G.C.: Why some alluvial rivers develop an anabranching pattern.
10 *Water Resour. Res.*, 43(7), doi: 10.1029/2006WR005223, 2007.
- 11 Hupp, C.R. and Osterkamp, W.R.: Riparian vegetation and fluvial geomorphic processes.
12 *Geomorphology*, 14, 277–295, 1996.
- 13 Jang, C.L. and Shimizu, Y.: Vegetation effects on the morphological behavior of alluvial
14 channels. *J. of Hydraul. Res.*, 45(6), 763–772, 2007.
- 15 Jansen, J.D. and Nanson, G.C.: Anabranching and maximum flow efficiency in Magela Creek,
16 northern Australia. *Water Resour. Res.*, 40(4), doi: 10.1029/2003WR002408, 2004.
- 17 Jansen, J.D. and Nanson, G.C.: Functional relationships between vegetation, channel
18 morphology, and flow efficiency in an alluvial (anabranching) river. *J. of Geophys. Res.*, 115,
19 F04030, doi: 10.1029/2010JF001657, 2010.
- 20 Jerolmack, D.J. and Mohrig, D.: Conditions for branching in depositional rivers. *Geology*,
21 35(5), 463–466, 2007.
- 22 Kleinhans, M.G.: Sorting out river channel patterns. *Progress in Physical Geography*, 34(3),
23 287–326, 2010.
- 24 Kleinhans, M.G. and van den Berg, J.H.: River channel and bar patterns explained and
25 predicted by an empirical and a physics-based method. *Earth Surf. Proc. and Landforms*,
26 36(6), 721–738, 2010.
- 27 Latrubesse, E.M.: Patterns of anabranching channels: The ultimate end-member adjustment of
28 mega rivers. *Geomorphology*, 101, 130–145, 2008.

- 1 Latrubesse, E.M. and Franzinelli, E.: The Holocene alluvial plain of the middle Amazon
2 River, Brazil. *Geomorphology*, 44, 241–257, 2002.
- 3 Lewin, J. and Ashworth, P.J.: Defining large river channel patterns: Alluvial exchange and
4 plurality. *Geomorphology*, 215, 83–98, 2014.
- 5 Li, Z.W., Wang, Z.Y., Pan, B.Z., Du, J., Brierley, G., Yu, G.A., and Blue, B.: Analysis of
6 controls upon channel planform at the First Great Bend of the Upper Yellow River, Qinghai-
7 Tibet Plateau. *J.of Geographi. Sci.*, 23(5), 833–848, 2013.
- 8 Luchi, R., Zolezzi, G., and Tubino, M.: Modelling mid-channel bars in meandering channels.
9 *Earth Surf. Proc. and Landforms*, 35, 902–917, 2010.
- 10 Makaske, B.: Anastomosing rivers: a review of their classification, origin and sedimentary
11 products. *Earth-Science Reviews*, 53, 149–196, 2001.
- 12 McBride, M., Hession, W.C., Rizzo, D.M., and Thompson, D.M.: The influence of riparian
13 vegetation on near-bank turbulence: a flume experiment. *Earth Surf. Proc. Landforms*, 32,
14 2019–2037, 2007.
- 15 McGowen, J.H. and Garner, L.: Physiographic features and stratification types of coarse-
16 grained point bars: Modern and ancient examples. *Sedimentology*, 14, 77–111, 1970.
- 17 Mertes, L., Dunne, T., and Martinelli, L.: Channel floodplain geomorphology along the
18 Solimões–Amazon River, Brazil. *Geological Society of America Bulletin*, 108, 1089–1107,
19 1996.
- 20 Meyer-Peter, E. and Müller, R.: Formulas for bed load transport, paper presented at 2nd
21 Meeting, International Association for Hydraulic Environmental Engineering and Research,
22 Madrid, 1948.
- 23 Millar, R.G.: Influence of bank vegetation on alluvial channel patterns. *Water Resour. Res.*,
24 36(4), 1109–1118, 2000.
- 25 Murray, A.B. and Paola, C.: Modelling the effect of vegetation on channel pattern in bedload
26 rivers. *Earth Surf. Proc. and Landforms*, 28,131–143, 2003.
- 27 Nanson, G.C. and Huang, H.Q.: Least action principle, equilibrium states, iterative adjustment,
28 and the stability of alluvial channels, *Earth Surf. Proc. Landforms*, 33, 923–942, 2008.

- 1 Nanson, G.C. and Knighton, A.D.: Anabranching rivers: their cause, character and
2 classification. *Earth Surf.Proc. and Landforms* 21, 217–239, 1996.
- 3 Nicholas, A.P., Ashworth, P.J., Smith, G.H.S., and Sandbach, S. D.: Numerical simulation of
4 bar and island morphodynamics in anabranching megarivers. *J. of Geophys.Res.: Earth Surf.*,
5 118, 2019–2044, 2013.
- 6 Osterkamp, W.R. and Hupp, C.R.: Fluvial processes and vegetation-Glimpses of the past, the
7 present, and perhaps the future. *Geomorphology*, 116, 274–285, 2010.
- 8 Pietsch, T.J. and Nanson, G.C.: Bankfull hydraulic geometry: the role of in-channel
9 vegetation and downstream declining discharges in the anabranching and distributary
10 channels of the Gwydir distributive system, southeastern Australia. *Geomorphology*, 129,
11 152–165, 2011.
- 12 Reesink, A.J., Ashworth, P.J., Sambrook Smith, G.H., Best, J.L., Parsons, D.R., Amsler, M.L.,
13 Hardy, R.J., Lane, S.N., Nicholas, A.P., Orfeo, O., Sandbach, S.D., Simpson, C.J., and
14 Szupiany, R.N.: Scales and causes of heterogeneity in bars in a large multi-channel river: R ó
15 Paran á Argentina. *Sedimentology*, 61, 1055–1085, 2014.
- 16 Sambrook, S.G.H., Ashworth, P.J., Best, J. L., Lunt, I.A., Orfeo, O., and Parsons, D.R.: The
17 sedimentology and alluvial architecture of a large braid bar R ó Paran á Argentina, *J. of*
18 *Sedimentary Res.*, 79, 629–642, 2009.
- 19 Smith, N.D.: Sedimentology and bar formation in the Upper Kicking Horse River, a braided
20 outwash stream. *J. of Geology*, 82, 205–223, 1974.
- 21 Song, X.L. and Bai, Y.C.: A new empirical river pattern discriminant method based on flow
22 resistance characteristics. *Catena*, 135, 163–172, 2015.
- 23 Tadaki, M., Brierley, G., and Cullum, C.: River classification: theory, practice, politics. *Wiley*
24 *Interdisciplinary Reviews: Water*, 1(4), 349–367, 2014.
- 25 Tal, M. and Paola, C.: Effects of vegetation on channel morphodynamics: results and insights
26 from laboratory experiments. *Earth Surf.Proc. and Landforms*, 35, 1014–1028, 2010.
- 27 Task Force on Bed Forms in Alluvial Channels: Nomenclature for Bed Forms in Alluvial
28 Channels. *J. of the Hydraul. Division*, 92(3), 51–64, 1966.

- 1 Tooth, S. and Nanson, G.C.: The role of vegetation in the formation of anabranching channels
2 in an ephemeral river, Northern plains, arid central Australia. *Hydrol. Process.*, 14(16-17),
3 3099-3117, 2000.
- 4 Wang, S.J., Chen, Z.Y., and Smith, D.G.: Anastomosing river system along the subsiding
5 middle Yangtze River basin, southern China. *Catena*, 60,147-163,2005.
- 6 Wong, M. and Parker, G.: Reanalysis and correction of bed load relation of Meyer-Peter and
7 Müller using their own database, *J. of Hydraul. Engineering*, 132(11), 1159-1168, 2006.
- 8 Yu, G.A., Brierley, G., Huang, H.Q., Wang, Z., Blue, B., and Ma, Y.X.: An environmental
9 gradient of vegetative controls upon channel planform in the source region of the Yangtze and
10 Yellow Rivers. *Catena*, 119, 14-153, 2014.
- 11

1 Table 1. Characteristics of the four study reaches (Flood season = June-September)

Alluvial reach	Planform type	Catchment area (km ²)	Flood-season mean discharge (m ³ /s)	Channel gradient	Vegetation cover
Dari	braided-anabranching	45020	270	0.00120	dense grasses/ sparse brush
Maqu	anabranching	86000	920	0.00050	dense trees
Lanmucuo	meandering	660	15	0.00150	dense grass
Daheba	braided	5200	70	0.00144	non-vegetation

2

1 Table 2. Characteristics and bed material of alluvial channels in the four study reaches

Alluvial reach	Channel width (m)	Water depth (m)	Bed material d_{50} (m)	Branching channels	Stability
Dari	450-1600	1.0-3.0	0.025	>5	semi-stable
Maqu	300-1000	2.0-5.0	0.015	>3	very stable
Lanmucuo	10-20	0.3-1.0	0.030	≤ 2	very stable
Daheba	150-500	0.5-2.0	0.060	>3	unstable


2

3

1 Table 3. Estimation of hydraulic coefficients and bedload transport rates


River reach	Bankfull channel width (m)	Bankfull water depth (m)	Channel gradient	Median grain size (m)	Manning coefficient	Average velocity (m/s)	Channel discharge (m ³ /s)	q_b (kg/s/m)
Dari	200	2.0	0.00120	0.015	0.05	0.90	269.67	1.77
Maqu	400	4.0	0.00050	0.015	0.15	0.37	593.14	2.75
Lanmucuo	20	0.8	0.00150	0.010	0.03	1.06	16.91	2.35
Daheba	50	1.5	0.0018	0.016	0.05	0.96	71.75	2.25

2



1
2
3
4
5
6
7

Figure 1. (a) The course of the Upper Yellow River. R1 is Dari reach, R2 is Maqu reach, R3 is Lanmucuo River, and R4 is Daheba River, (b) Planform morphology of the study reaches (R1 is Dari reach, R2 is Maqu reach, R3 is Lanmucuo River reach, and R4 is Daheba River reach). R1, R2, and R4 are Google Earth images and R3 is a photograph taken from nearby hills. Points A, B, C, and D are the location of photographs shown in [Figure 2](#).




1
 2 Figure 2. (R1) Channel morphology and gravel bed of Dari reach (photographs taken on 2
 3 July, 2012, 33.7553 N, 99.6414 E, 3960 m elevation), (R2) Channel morphology and gravel
 4 bed of Maqu reach (photographs taken on 8 July, 2012, 33.3594 N, 102.0553 E, 3465 m
 5 elevation), (R3) Channel morphology and gravel bed of a grass covered bar in middle
 6 Lanmucuo River (to photographs taken on 5 July, 2012, 34.4287 N, 101.4663 E, 3604 m
 7 elevation), (R4) Channel morphology and gravel bed of middle Daheba River (photographs
 8 taken on 6 August, 2011, 35.5169 N, 100.0183 E, 2832 m elevation).

9




1
 2 Figure 3. River bank of the study reaches (a) Dari reach, (b) Maqu reach, (c) Lanmucuo River
 3 reach, and (d) Daheba River reach.

4



1




2

3 Figure 4. Monthly stage-discharge relationships for Jimai hydrological station in Dari reach (a)

4 1968 (b) 1984 (Note: number refers to month, e.g., 1 for January and 12 for December).

5




1

2

3 Figure 5. Annual stage-discharge relationship (1964-1984) of Dari reach in Jimai hydrological
 4 station.


5



1

2 Figure 6. Sketch of channel bed deposition and erosion in flood season in Dari reach.


3



1

2 Figure 7. Monthly stage change of Maqu hydrological station (1959-1970).

3




1

2 Figure 8. Sketch of branching channel deposition and stage increasing in flood season in

3 Maqu reach


4



1

2 Figure 9. Sketch of submerged bend apex with a mid-channel bar in the Lanmucuo River


3



1

2 Figure 10. Braided channels evolution of the middle Daheba River in 2005 (a) in non-flood
3 season, (b) in flood season

4



1
 2 Figure 11. Elevation change of cross-section in Shangcun hydrological station (2009-2011)
 3 (left for left bank, right for right bank)