1 The importance of snowmelt spatio-temporal variability for

2 isotope-based hydrograph separation in a high-elevation 3 catchment

Jan Schmieder¹, Florian Hanzer¹, Thomas Marke¹, Jakob Garvelmann², Michael Warscher²,
 Harald Kunstmann² and Ulrich Strasser¹

⁶ ¹Institute of Geography, University of Innsbruck, Innsbruck, 6020, Austria

7 ²Institute of Meteorology and Climate Research - Atmospheric Environmental Research, Karlsruhe Institute of

8 Technology, Garmisch-Partenkirchen, 82467, Germany

9 Correspondence to: Jan Schmieder (Jan.Schmieder@uibk.ac.at)

10 Abstract. Seasonal snow cover is an important temporary water storage in high-elevation regions. Especially in 11 remote areas, the available data is often insufficient to explicitly quantify snowmelt contributions to streamflow. 12 The limited knowledge about the spatio-temporal variability of the snowmelt isotopic composition, as well as 13 pronounced spatial variation in snowmelt rates lead to high uncertainties in applying the isotope-based 14 hydrograph separation method. The stable isotopic signatures of snowmelt water samples collected during two 15 spring 2014 snowmelt events at a north- and a south-facing slope were volume-weighted with snowmelt rates derived from a distributed physics-based snow model in order to transfer the measured plot-scale isotopic 16 composition of snowmelt to the catchment scale. The observed $\delta^{18}O$ values and modelled snowmelt rates showed 17 18 distinct inter- and intra-event variations, as well as marked differences between north- and south-facing slopes. 19 Accounting for these differences, two-component isotopic hydrograph separation revealed snowmelt 20 contributions to streamflow of 35 ± 3 % and 75 ± 14 % for the early and peak melt season, respectively. These 21 values differed from those determined by formerly used weighting methods (e.g. using observed plot-scale melt 22 rates) or considering either the north- or south-facing slope by up to 5 and 15 %, respectively.

23 **1 Introduction**

24 In many headwater catchments, seasonal water availability is strongly dependent on cryospheric processes and

understanding these processes becomes even more relevant in a changing climate (APCC, 2014; IPPC, 2013;

26 Weingartner and Aschwanden, 1992). The seasonal snow cover is an important temporary water storage in alpine

27 regions. The timing and amount of water released from this storage is important to know for water resources

28 management, especially in downstream regions where the water is needed (drinking water, snow making,

29 hydropower, irrigation water) or where it represents a potential risk (flood, drought). Environmental tracers are a

30 common tool to investigate the hydrological processes, but scientific studies are still rare for high-elevation

31 regions because of the restricted access and high risk for field measurements in these challenging conditions.

32 Two-component isotopic hydrograph separation (IHS) is a technique to separate streamflow into different time

33 source components (event water, pre-event water) (Sklash et al., 1976). The event component depicts water that

34 enters the catchment during an event (e.g. snowmelt) and is characterized by a distinct isotopic signature,

35 whereas pre-event water is stored in the catchment prior to the onset of the event (i.e. groundwater and soil water

36 which form baseflow) and is characterized by a different isotopic signature (Sklash and Farvolden, 1979; Sklash

37 et al., 1976). The technique dates back to the late 1960s (Pinder and Jones, 1969) and was initially used for

1 separating storm hydrographs in humid catchments. The first snowmelt-based studies were conducted in the

2 1970s by Dincer et al. (1970) and Martinec et al. (1974). These studies showed a large pre-event water fraction

3 (>50 %) of streamflow that changed the understanding of the processes in catchment hydrology fundamentally

4 (Klaus and McDonnell, 2013; Sklash and Farvolden, 1979) and forced a paradigm shift, especially for humid

5 temperate catchments. However, other snowmelt-based studies (Huth et al., 2004; Liu et al., 2004; Williams et

6 al., 2009) reveal a large contribution of event water (>70 %), i.e. in permafrost or high-elevation catchments,

depending on the system state (e.g. frost layer thickness and snow depth), catchment characteristics and runoff
 generation mechanisms.

9 Klaus and McDonnell (2013) highlighted the need to quantify and account for the spatial variability of the 10 isotope signal of event water, which is still a vast uncertainty in snowmelt-based IHS. In the literature 11 inconclusive results prevail with respect to the variation of the isotopic signal of snowmelt. Spatial variability of 12 snowmelt isotopic composition was statistically significant related to elevation (Beaulieu et al., 2012) in a 13 catchment in British Columbia, Canada with 500 m relief. Moore (1989) and Laudon et al. (2007) found no 14 statistical significant variation in their snowmelt δ^{18} O data, due to the low gradient and small elevation range (approximately 30 m and 290 m) in their catchments, which favours an isotopically more homogenous snow 15 16 cover. The effect of the aspect of the hillslopes on isotopic variability and IHS results in topographically 17 complex terrain has-also been rarely investigated. Dahlke and Lyon (2013) and Dietermann and Weiler (2013) 18 surveyed the snowpack isotopic composition and showed a notable spatial variability in their data, particularly 19 between north- and south-facing slopes. They conclude that the spatial variability of snowmelt could be high and 20 that the timing of meltwater varies with the morphology of the catchment. Dietermann and Weiler (2013) also 21 concluded that an elevation effect (decrease of snowpack isotopic signature with elevation), if observed, is 22 disturbed by fractionation due to melt/refreeze processes during the ablation period. These effects most likely superimpose the altitudinal gradient. Aspect and slope are therefore important factors that affect the isotopic 23 24 evolution of the snow cover and its melt (Cooper, 2006). In contrast, there have been various studies that have 25 investigated the temporal variability of the snowmelt isotopic signal, e.g. by the use of snow lysimeters (Hooper 26 and Shoemaker, 1986; Laudon et al., 2002; Liu et al., 2004; Maulé and Stein, 1990; Moore, 1989; Williams et 27 al., 2009). During the ablation season the isotopic evolution of the snowpack progresses due to percolating rain water and fractionation caused by melting, refreezing and sublimation (Dietermann and Weiler, 2013; Lee et al., 28 29 2010; Unnikrishna et al., 2002; Zhou et al., 2008), which leads to a homogenization of the isotopic profile of the 30 snowpack (Árnason et al., 1973; Dincer et al., 1970; Stichler, 1987) and an increase in heavy isotopes of 31 meltwater throughout the freshet period (Laudon et al., 2007; Taylor et al., 2001; Taylor et al., 2002; 32 Unnikrishna et al., 2002). Therefore, the characterization and the use of the evolving isotopic signal of snowmelt 33 water instead of single snow cores is crucial for applying IHS (Taylor et al., 2001; 2002). 34 There have been various approaches to cope with the temporal variability of the input signal. If one uses more 35 than one δ^{18} O snowmelt value for applying the IHS method, it is important to weight the values with appropriate

36 melt rates, e.g. measured from the outflow of a snow lysimeter. Common weighting methods are the volume-

37 weighted average approach (VWA), as used by Mast et al. (1995), and the current meltwater approach (CMW),

38 applied by Hooper and Shoemaker (1986). Laudon et al. (2002) developed the runoff-corrected event water

39 approach (runCE), which accounts for both, the temporal isotopic evolution and temporary storage of meltwater

40 in the catchment and overcomes the shortcoming of the exclusion of residence times by VWA and CMW. This

1 method was also deployed in several other snowmelt-based IHS (Beaulieu et al., 2012; Carey and Quinton,

2 2004; Laudon et al., 2004; Laudon et al., 2007).

3 Tracers have successfully been used in modelling studies to provide empirical insights into runoff generation 4 processes and catchment functioning (Birkel and Soulsby, 2015; Birkel et al., 2011; Capell et al., 2012; 5 Uhlenbrook and Leibundgut, 2002), but the combined use of distributed modelling and isotope tracers in snow-6 dominated environments is rare. Ahluwalia et al. (2013) used an isotope and a-modelling approach to derive 7 snowmelt contributions to streamflow and determined differences between the two techniques of 2 %. 8 Distributed modelling can provide areal melt rates that can be used for weighting the measured isotopic 9 composition of meltwater. Pomeroy et al. (2003) described the differences of insolation between north- and 10 south-facing slopes in complex terrain that lead to spatial varying melt rates of the snowpack throughout the 11 freshet period. The use of the areal snowmelt data from models will likely reduce the uncertainty that arises from 12 the representativeness of measured melt rates at the plot-scale.

13 The overall goal of our study was to quantify the contribution of snowmelt to streamflogd hence to improve

14 the knowledge of hydroclimatological processes in high-elevation catchments. This study aims to test a

- 15 (technique to enhance the reliability of isotope-based hydrograph separation, and thus the estimation of snowmelt
- contributions to streamflow by considering the distinct spatio-temporal variability of snowmelt and its isotopic
 signature in a high-elevation study region. This study has the following three objectives: 1) the estimation of the
- 18 spatio-temporal variability of snowmelt and its isotopic composition, 2) the quantification of the impact of the
- 19 spatial variability in snowmelt rates and its isotopic composition on isotope-based hydrograph separation (IHS)
- and 3) to assess the combined use of a physically-based snowmelt model and traditional IHS. Distributed melt
- 21 rates provided by a surface energy balance model were used to weight the measured isotopic composition of
- snowmelt in order to characterize the event water isotopic composition. Traditional weighting methods (e.g.
- 23 using plot-scale observed melt rates) were compared with the model approach. This study provides an integrated
- 24 approach to understand the streamflow components based on experimental field work (data collection) and
- 25 modelling, as requested in previous studies (e.g. Seibert and McDonnell, 2002).

26 2 Study area

- The 98 km² high-elevation catchment of the stream Rofenache is located in the Central Eastern Alps (Oetztal Alps, Austria), close to the main Alpine ridge. The study area has a dry inner-alpine climate. Mean annual precipitation is 800 mm yr⁻¹, of which 44 % falls as snow. The mean annual temperature at the gauging station in Vent (1890 m.a.s.l., reference period: 1982-2003) is 2°C. Seasonal snow cover typically lasts from October to the end of June at the highest regions of the valley. The basin ranges in elevation from approximately 1900 m.a.s.l. to 3770 m.a.s.l.. Average slope is 25° and average elevation is 2930 m.a.s.l. (calculated from a 50 m digital elevation model). A narrow riparian zone (<100 m width) is located in the valley floor. The
- 34 predominantly south- (SE) and north-facing (NNW) slopes form the main valley (cf. Fig. 1a), which trends
- 35 roughly from southwest to northeast (cf. Fig. 1b),
- 36 The bedrock consists of mainly paragneiss and mica schist and is overlain by a mantle of glacial deposits and
- 37 thin soils (< 1 m). The bedrock outcrops and unconsolidated bare rocks cover the largest part (42 %) of the
- 38 catchment (CLC, 2006). Glaciers cover approximately a third of the Rofenache catchment (35 %), while pastures
- 39 and coniferous forests are located in the lowest parts of the catchment and cover less than 0.5 % (CLC, 2006).

- 1 Sparsely vegetated areas and natural grassland cover 15 and 7.5 %, respectively (CLC, 2006). Besides seasonally
- 2 frozen ground at slopes of various expositions, permafrost is likely to occur at an altitude over 2600 m.a.sl. at the
- 3 north-facing slopes (Haeberli, 1975). The annual hydrograph reveals a highly seasonal flow regime. The mean
- 4 annual discharge is 4.5 $\text{m}^3 \text{s}^{-1}$ (reference period: 1971-2009) and is dominated by snow and glacier melt during
- 5 the ablation season, which typically lasts from May to September. The onset of the early snowmelt season in the
- 6 lower part of the basin is typically in April.

7 **3 Methods**

8 **3.1 Field sampling, measurements and laboratory analysis**

9 The field work was conducted during the 2014 snowmelt season between the beginning of April and the end of 10 June. Two short-term melt events (3 days) were investigated to illustrate the difference between early spring 11 season melt and peak melt. The events were defined as warm and precipitation-free spells, with clear sky and dry 12 antecedent conditions (i.e. no precipitation was observed 48 h prior to the event). Low discharge and air 13 temperatures with a small diurnal variation and low melt rates, as well as a snow-covered area (SCA) of about 14 90 % in the basin (Fig. 2a) characterize the conditions of the early melt event at the end of April (cf. Fig. 3b). In 15 contrast, the peak melt period at the end of June is characterized by high discharge and melt rates, a flashy 16 hydrograph, high air temperatures with remarkable diurnal variations (Fig. 3c) and a strongly retreated snowline 17 (SCA: 66 %; cf. Fig. 2c). Discharge data are available at an hourly resolution for the gauging station in Vent and meteorological data are obtained by 2 tomatic weather stations (hourly resolution) located in and around the 18 19 basin (Fig. 1).

20 The stream water sampling for stable isotope analysis consisted of pre-freshet baseflow samples at the beginning 21 of March, sub-daily samples (temporal resolution ranges between 1 and 4 hours) during the two studied events 22 and a post-event sample in July as indicated in Fig. 3a (grey-shaded area). Samples of snowmelt, snowpack and 23 surface overland flow (if observed) were collected at the south- (S1, S2) and north-facing slope (N1, N2), as well 24 as on a wind-exposed ridge shown in Fig. 1b using a snowmelt collector. At each test site a snow pit was dug to 25 install a 0.1 m² polyethylene snowmelt collector at the ground-snowpack interface. The snowmelt collector 26 consists of a pipe that drains the percolating meltwater into a fixed plastic bag. Tests yield a preclusion of 27 evaporation for this sampling method. Composite daily snowmelt water samples (bulk sample) were collected in 28 these bags and transferred to polyethylene bottles in the field before the onset of the diurnal melt cycle. 29 Furthermore, sub-daily grab melt samples were collected at S1 (on 23 April) and at N2 (on 07 June) to define the 30 diurnal variability of the respective melt event. Unfortunately further sub-daily snowmelt sampling was not 31 feasible. The pit face was covered with white styrofoam to protect it from direct sunlight. Stream, surface 32 overland flow and grab snowmelt water samples were collected in 20 mL polyethylene bottles. Snow samples 33 from snow pit layers were filled in airtight plastic bags and melted below room temperature before transferring 34 them in bottles. Overall, 144 samples were taken during the study period. Snow water equivalent (SWE), snow 35 height (HS), snow density (SD), and various snowpack observations (wetness and hand hardness index) were 36 observed before the onset of the diurnal melt cycle at the study plots (Fig. 1). Mean SWE was determined by averaging five snow tube measurements within an area of 20 m² at each site. Daily melt rates were calculated by 37 38 subtracting succeeding SWE values. Sublimation was neglected, as it contributes only to a small percentage (~10 39 %) to the seasonal water balance in high altitude catchments in the Alps (Strasser et al., 2008).

All samples were treated by the guidelines as-proposed by Clark and Fritz (1997) and were stored dark and cold until analysis. The isotopic composition of the samples ($\delta^{18}O$, δD) was measured with cavity ring-down spectroscopy (Picarro L1102-i). Results are expressed in the delta notation as parts per thousand relative to the Vienna Standard Mean Ocean Water (VSMOW2).The mean laboratory precision (replication of 8 measurements) for all measured samples was 0.06 ‰ for $\delta^{18}O$. Due to the covariance of $\delta^{2}H$ (δD) and $\delta^{18}O$ (Fig. 5), all analyses were done with oxygen-18 values.

7 **3.2 Model description**

8 For the simulation of the daily melt rates, the non-calibrated, distributed, and physically-based 9 hydroclimatological model AMUNDSEN (Strasser, 2008) was applied. Model features include interpolation of 10 meteorological fields from point measurements (Marke, 2008; Strasser, 2008); simulation of short- and 11 longwave radiation, including topographic and cloud effects (Corripio, 2003; Greuell et al., 1997); 12 parameterization of snow albedo depending on snow age and temperature (Rohrer, 1992); modelling of forest 13 snow and meteorological processes (Liston and Elder, 2006; Strasser et al., 2011); lateral redistribution of snow 14 due to gravitational (Gruber, 2007) and wind-induced (Helfricht, 2014; Warscher et al., 2013) processes; and 15 determination of snowmelt using an energy balance approach (Strasser, 2008). Besides having been applied for 16 various other Alpine sites in the past (Hanzer et al., 2014; Marke et al., 2015; Pellicciotti et al., 2005; Strasser, 17 2008; Strasser et al., 2008; Strasser et al., 2004), AMUNDSEN has recently been set up and extensively 18 validated for the Oetztal Alps region (Hanzer et al., 2016). This setup was also used to run the model in this 19 study for the period 2013–2014 using a temporal resolution of 1 hour and a spatial resolution of 50 meters. In 20 order to determine the model performance during the study period, catchment-scale snow distribution by 21 satellite-derived binary snow cover maps and plot-scale observed SWE data were used for the validation (cf. 22 Section 4.2). Therefore the spatial snow distribution as simulated by AMUNDSEN was compared with a set of 23 MODIS (500 m spatial resolution) and Landsat (30 m resolution, subsequently resampled to the 50 m model 24 resolution) snow maps with less than 10 % cloud coverage over the study area using the methodology described 25 in Hanzer et al. (2016). Model results were evaluated using the performance measures BIAS, accuracy (ACC) 26 and critical success index (CSI) (Zappa, 2008). ACC represents the fraction of correctly classified pixels (either 27 snow-covered or snow-free both in the observation and the simulation). CSI describes the number of correctly 28 predicted snow-covered pixels divided by the number of times where snow is predicted in the model and/or 29 observed, and BIAS corresponds to the number of snow-covered pixels in the simulation divided by the 30 respective number in the observation. ACC and CSI values range from 0 to 1 (where 1 is a perfect match), while 31 for BIAS values below 1 indicate underestimations of the simulated snow cover, and values above 1 indicate 32 overestimations. At the plot-scale, observed SWE values were compared with AMUNDSEN SWE values 33 represented by the underlying pixel at the location of the snow course. Catchment-scale melt rates are calculated 34 by subtracting two consecutive daily SWE grids, neglecting sublimation losses, as also done to achieve observed 35 melt rates at the plot-scale. Subsequently, the DEM was used to calculate an aspect grid and further to divide the catchment into two parts: grid cells with aspects ranging from $\geq 270^{\circ}$ to $\leq 90^{\circ}$ were classified as 'north-facing'. 36 37 while the remaining cells were attributed to the class 'south-facing'. Finally, these two grids were combined to derive melt rates for the south-facing $(melt_s)$ and for the north-facing slope $(melt_n)$. 38

1 3.3 Isotopic hydrograph separation, weighting approaches and uncertainty analysis

- IHS is a steady-state tracer mass balance approach and several assumptions underlie this simple principle, which
 are described and reviewed in Buttle (1994) and Klaus and McDonnell (2013):
- 4 (1) The isotopic compositions of event and pre-event water are significantly different.
- 5 (2) The event water isotopic signature has no spatio-temporal variability, or variations can be accounted 6 for.
- 7 (3) The pre-event water isotopic signature has no spatio-temporal variability, or variations can be8 accounted for.
- 9 (4) Contributions from the vadose zone must be negligible or soil water should be isotopically similar to10 groundwater.
- 11 (5) There is no or minimal discharge contribution from surface storage.
- 12 The focus of this study is on one of the assumptions: the spatio-temporal variability of event water isotopic 13 signature is absent or can be accounted for. The fraction of event water (f_e) contributing to streamflow was
- 14 calculated from Eq. (1).

15
$$f_e = \frac{(c_p - c_s)}{(c_p - c_e)}$$
 (1)

- 16 The tracer concentration of the pre-event component (C_p) is the δ^{18} O composition of baseflow prior to the onset
- 17 of the freshet period, constituted mainly by groundwater and potentially by soil water which was assumed to
- 18 have the same isotopic signal as groundwater. Tracer concentration C_s is the isotopic composition of stream
- 19 water for each sampling time. The isotopic compositions of snowmelt samples were weighted differently to
- 20 compose the event water tracer concentration (C_e) using the following five weighting approaches:
- 21 (1) volume-weighted with observed plot-scale melt rates (VWO)
- 22 (2) equally weighted, assuming an equal melt rate on north- and south-facing slopes (VWE)
- 23 (3) no weighting, only south-facing slopes considered (SOUTH)
- 24 (4) no weighting, only north-facing slopes considered (NORTH)
- 25 (5) volume-weighted with simulated catchment-scale melt rates (VWS)

Equation (2) is the VWS approach with simulated melt rates for north- and south-facing slopes as described in Section 3.2, where *M* is the simulated melt rate (in mm d⁻¹), $\delta^{18}O$ is the isotopic composition of sampled snowmelt and subscripts *s* and *n* indicate north and south, respectively. For obtaining the value of C_e a daily timestep (*t*) is used, considering daily melt rates and the isotopic composition of the daily bulk snowmelt samples.

31
$$C_e(t) = \frac{M_s(t)\delta^{18}O_s(t) + M_n(t)\delta^{18}O_n(t)}{M_s(t) + M_n(t)}$$
(2)

An uncertainty analysis (Eq. (3)) was performed according to the Gaussian standard error method proposed by
 Genereux (1998):

$$34 \qquad W_{f_e} = \left\{ \left[\frac{c_p - c_s}{(c_p - c_e)^2} W_{C_e} \right]^2 + \left[\frac{c_s - c_e}{(c_p - c_e)^2} W_{C_p} \right]^2 + \left[\frac{-1}{(c_p - c_e)^2} W_{C_s} \right]^2 \right\}^{1/2}, \tag{3}$$

35 where *W* is the uncertainty, *C* is the isotopic composition, *f* is the fraction and the subscripts *p*, *s* and *e* refer to 36 the pre-event, stream and event component. This assumes negligible errors in the discharge measurements and

- the melt rates (modelled and observed). The uncertainty of streamflow (W_{C_s}) is assumed to be equal to the laboratory precision (0.06 %). For the uncertainty of the event component (W_{C_e}), the diurnal temporal variability 2 (standard deviation) of the snowmelt isotopic signal (from one site and one day) was multiplied by the 3 4 appropriate value of the two-tailed t-table (dependent on sample number) and used for the event, as proposed by Genereux (1998). Different uncertainty values were applied for the early melt event ($W_{C_e} = 0.2 \%$) and the peak 5 melt event ($W_{C_e} = 0.5$ ‰). An error of 0.04 ‰ was assumed for the pre-event component (W_{C_p}), which reflects 6 7 the standard deviation of two baseflow samples. A 95 % confidence level was used. Spatial variation in 8 snowmelt and its isotopic composition were not considered in this error calculation method as they represent the 9 hydrologic signal of interest.
- 4 Results

10

32

4.1. Spatio-temporal variability of streamflow and stable isotopic signature of sampled of water sources 11

12 Two major snowmelt pulses (Mid-May and beginning of June) and four less pronounced ones between mid-13 March to early May occurred during the snowmelt season (Fig. 3a). Peak melt occurred at the beginning of June with maximum daily temperatures and runoff of 15 °C and 18 mm d⁻¹, respectively. The following high-flows 14 15 were affected by rain (Fig. 3a) and glacier melt due to the strongly retreated snow line and snow-free ablation 16 area of the glaciers in July. Diurnal variations in discharge were strongly correlated with diurnal variations of air 17 temperature (Fig. 3b and c) with a time lag of 3-5 hours for the early melt event and 2-3 hours for the peak melt event. An inverse relationship between streamflow δ^{18} O and discharge was found for the early melt event (Fig. 18 4a and c). Small diurnal responses of streamflow δ^{18} O were identified for both events, but were masked due to 19 20 missing data during the recession of the hydrograph.

21 The quality control of the isotopic data was performed by the $\delta^2 H - \delta^{18} O$ plot (Fig. 5), which indicated no shift in 22 the linear regression line and thus no secondary fractionation effects (evaporation) during storage and transport 23 of the samples. The slope of the linear regression (slope=8.5, n=144, R²=0.93) of the measurement data slightly 24 deviates from that of the global meteoric (slope=8) and local meteoric water line (slope=8.1) based on monthly 25 data from the ANIP (Austrian Network of Isotopes in Precipitation) sampling site in Obergurgl, which is located 26 in an adjacent valley (reference period: 1991-2014). The small deviation (visible in Fig. 5) of the sampled water 27 (i.e. snowpack and snowmelt) could indicate fractionation effects induced by phase transition (i.e. melt/refreeze 28 and sublimation). The significant differences between the isotopic signatures of pre-event streamflow and

- 29 snowmelt water enabled the IHS.
- Overall, the complete δ^{18} O values ranged from -21.5 to -15.0 ‰ while snowpack samples were characterized by 30
- 31 the most negative and pre-event baseflow samples by the least negative values. Snowpack samples showed a
- signal mixing of the water components. Figure 6 shows the δ^{18} O data of the water samples grouped into different 33

wide isotopic range, while streamflow samples revealed the narrowest spread, reflecting a composite isotopic

- categories and split into early and peak melt data. It shows the different δ^{18} O ranges and medians of the sampled 34
- 35 water sources (Fig. 6a), as well as marked spatio-temporal variations in the isotopic signal (Fig. 6b and c). It is
- apparent that the snowpack δ^{18} O values have a larger variation compared to the snowmelt data due to 36
- homogenization effects (Fig. 6a), as was also shown by Árnason et al. (1973), Dincer et al. (1970) and Stichler 37
- (1987). The median of the δ^{18} O of snowmelt was higher than that of the snowpack indicating fractionation. The 38
- median δ^{18} O of surface overland flow was higher than that of snowmelt (Fig. 6a) for the early and peak melt 39

- 1 period. Overall, the peak melt δ^{18} O values (Fig. 6b) were less variable and had a higher median than the early
- 2 melt values, because fractionation effects (due to melt/refreeze and sublimation) most likely altered the isotopic
- 3 composition of the snowpack over time (cf. Taylor et al., 2001, 2002). One major finding was that the north-
- 4 facing slope δ^{18} O values had a larger range and a lower median compared to the opposing slope (Fig. 6c).
- 5 Samples from the wind drift influenced site (also south-exposed) were more depleted in heavy isotopes
- 6 compared to the south-facing slope samples (Fig. 6c).
- 7 In general, the average snowmelt and snowpack isotopic composition was more depleted for the early melt
- 8 period (Table 1) and changed over time because fractionation likely altered the snowpack and its melt. It is 9 obvious that the isotopic evolution (gradually enrichment) on the south-facing slope took place earlier in the
- annual melting cycle of the snow, and indicates a premature snowpack concerning the enrichment of isotopes
- 11 and earlier ripening compared to the north-facing slope.
- 12 Table 1 shows that meltwater sampling throughout the entire snowmelt period is required to account for the
- temporal variation in the isotopic composition of the snowpack (cf. Taylor et al., 2001, 2002). In detail, the snowpack and snowmelt δ^{18} O data highlighted a marked spatial inhomogeneity between north- and south-facing
- 15 slopes throughout the study period. The snowpack isotopic composition from both sampled slopes was
- 16 statistically different for the early melt, but not for the peak melt (with Kruskal-Wallis test at 0.05 significance
- 17 level), whereas the snowmelt δ^{18} O showed a significant difference throughout the study period (Fig. 7).
- 18 Sub-daily snowmelt samples (n=5) at S1 (23 April 2014) had a range of 0.1 % in δ^{18} O, and the bulk sample
- 19 (integrating the entire diurnal melt cycle) was within the scatter of those values (Fig. 8). The intra-daily
- 20 variability of snowmelt (n=3) at N2 (07 June 2014) was relatively higher with values ranging from -17.9 to -
- 21 18.1 ‰. The bulk sample (-17.9 ‰) was at the upper end of those values (Fig. 8).
- 22 Stream water isotopic composition was more enriched in heavy isotopes during the early melt period and
- 23 successively became more depleted throughout the freshet period, resulting in more negative values during peak
- melt (Table 2). The standard deviation and range of stream water δ^{18} O during early melt was higher and could be
- 25 related to an increasing snowmelt contribution throughout the event and larger diurnal amplitudes of snowmelt
- 26 contribution compared to peak melt (Table 2).

27 **4.2 Snow model validation and snowmelt variability**

- Figure 9 shows the values for the selected performance measures based on the available MODIS and Landsat
- 29 scenes during the period March–July 2014. The results indicate a reasonable model performance with a tendency
- 30 to slightly overestimate the snow cover during the peak melt season (BIAS >1). In general the CSI does not drop
- 31 below 0.7 and 80 % of the pixels are correctly classified (ACC) throughout the study period. Fig. 2 shows the
- 32 observed and simulated spatial snow distribution around the time of the two events. Despite a higher SCA during
- 33 the early melt season (Fig. 2a and b) compared to the peak melt season (Fig. 2c and d) one can see the
- 34 overestimation of the simulated SCA compared to the observed (MODIS/Landsat) SCA. Table 3 shows the
- 35 observed and simulated SWE values at the plot-scale. The model slightly underestimated SWE during peak melt,
- 36 but generally appears to be in quite good agreement, suggesting well simulated snowpack processes. Throughout
- 37 the study period the model deviates by 13 % from the observed SWE values, but the representativeness (small-
- 38 scale effects) of SWE values represented by the respective 50 m pixel should be considered.
- 39 Snowmelt (observed and simulated daily losses of SWE) showed a distinct spatial variation between the north-
- 40 facing and the south-facing slope for the early melt (23/24 April), but less marked variations for the peak melt

1 (07/08 June) period (Fig. 10). Relative day-to-day differences are more pronounced for the early melt season. 2 Both simulated and observed melt rates are higher for the peak melt event on the south-facing slope, but not for

- 3 the north-facing slope. Simulated melt intensity on the south-facing slope at the end of April was twice the rate
- 4 on the north-facing slope, while simulated melt rates were approximately the same for the opposing slopes
- 5 during peak melt. Simulated (catchment scale) snowmelt rates were markedly higher during the early melt (23
- 6 and 24 April) on the north-facing slope compared to the observed (plot scale) melt rates (Fig. 10a), but
- 7 differences between them were small during peak melt for both slopes (07 and 08 June; Fig. 10).

8 4.3 Weighting techniques and isotope-based hydrograph separation

- 9 Differences between the applied snowmelt weighting techniques, induced by the high spatial variability of 10 snowmelt (Section 4.2), led to different event water isotopic compositions (C_e) for the IHS analyses (Table 4). The event water component is depleted in δ^{18} O by roughly 0.3 % for the second day (24 April) of the early melt 11 12 event compared to the preceding day, but inter-daily variation during the peak melt is almost absent. Especially 13 during early melt (23/04 to 24/04) strong deviations between observed plot-scale melt rates and distributed 14 (areal) melt rates obtained by AMUNDSEN occurred (Fig. 11), and led to more different event water isotopic 15 compositions between the VWS and the VWO approach (Table 4).
- 16 The hydrograph and the results of the IHS applied with the VWS method for the early and peak melt event are 17 presented in Fig. 4 and highlighted lower flow rates and higher pre-event fractions during early melt (Fig. 4c) 18 and vice versa for the peak melt period (Fig. 4d). The total runoff volume during the peak melt period was 19 approximately six times higher than in the early melt period. The fraction of snowmelt (volume) estimated with 20 the VWS approach was 35 and 75 % with calculated uncertainties (95 % confidence level) of ± 3 and ± 14 % for 21 the early and peak melt event, respectively. The uncertainty calculated from Eq. (3) of the IHS applied with the 22 VWS method was higher (14%) for the peak melt event than for the early melt event because the difference 23 between isotopic composition of pre-event water and event water was smaller than for the early melt event
- 24 (uncertainty: 3%) (cf. Table 2 and 4).
- 25 Throughout the early melt event, the snowmelt fraction increased from 25 to 44 % (Fig. 4c; Table 5). This trend mirrors the stream isotopic composition, which became more depleted (Fig. 4a). Event water contributions 26 27 during peak melt were generally higher but had a smaller range (70 to 78 %; Fig. 4d). Diurnal isotopic variations
- 28 of stream water were small for both events (Fig. 4a and b), and could not clearly be obtained due to missing data
- 29 at the falling limb of the hydrographs.

- 30 The use of the different weighting approaches led to strongly varying estimated snowmelt fractions of
- streamflow (Fig. 12). Especially the differences between the SOUTH and the NORTH approach during both 32 investigated events (up to 24 %), and the differences between the VWS and the VWO approach (5 %) during
- 33 early melt (Fig. 12a) are notable. Event water contributions estimated by the different weighting methods ranged
- 34 from 21-28 % at the beginning of the early melt event up to 31-55 % at the end of the event (cf. Fig. 12a, Table
- 35 6). Minimum event water contributions during the peak melt were estimated at 60-84 % and maxima ranged
- 36 between 67-94 % for the different weighting methods (Table 6, Fig. 12b). Beside these intra-event variations in
- 37 snowmelt contribution, the volumetric variations at the event-scale were smaller and ranged between 28 to 40 %
- 38 and 66 to 90 %, for the early and peak melt event, respectively (Table 6).
- 39 Considering only spatial variation of snowmelt isotopic signatures (i.e. comparing the NORTH/SOUTH
- 40 approach with the VWE approach) for IHS led to differences in estimated event water fractions up to 7 and 14 %

- for the early and peak melt period, respectively (Table 6). However, considering only spatial variation in snowmelt rates (i.e. comparing the VWS/VWO approach with the VWE approach) led to differences in event water fraction up to 3 and 2 % for the early and peak melt period, respectively (Table 6).
- 4 Surface overland flow was not considered in the IHS analyses, but if applied, it would most likely increase the
- 5 calculated snowmelt fraction slightly. Furthermore, snowmelt samples from the wind-exposed site were not used
- 6 in the IHS analyses because this site was only sampled on the south-facing slope during early melt and is not
- 7 representative for the catchment due to its limited coverage. However, an-incorporation of this data would
- 8 decrease the calculated snowmelt fraction by approximately 2 %.

9 5 Discussion

10 **5.1 Temporal variation in streamflow during the melting season**

Snowmelt is a major contributor to streamflow during the spring freshet period in alpine regions and large 11 12 amounts of snowmelt water infiltrate into the soil and recharge groundwater (Penna et al., 2014). The 13 hydrological response of the stream followed the variations of air temperature, as already observed by 14 Braithwaite and Olesen (1989) (Fig. 3a). The observed time lags (Fig. 3b and c) between maximum daily air temperature and daily peak flow are common in mountain catchments (Engel et al., 2016; Schuler, 2002). During 15 16 peak melt, the flashy hydrograph revealed less variation in the timing of peak discharge of 7 day data 17 (cf. Fig. 3c) compared to the early melt, as well reported by Lundquist and Cayan (2002). The increase in discharge coincides with decreasing streamflow δ^{18} O during the early melt event (Fig. 4a and c) and confirms 18 19 earlier findings of Engel et al. (2016) who identified inverse relationships between streamflow δ^{18} O and 20 discharge during several 24-hour events in an adjacent valley on the southern side of the main Alpine ridge, 21 although their findings rely on streamflow contributions from snow and glacier melt. The lower stream water 22 isotopic composition during peak melt suggests a remarkable contribution of more depleted snowmelt to 23 streamflow and therefore confirms the results of the IHS.

24 5.2 Spatio-temporal variability of snowmelt and its isotopic signature

25 The rate of snowmelt varies spatially in catchments with complex topography (Carey and Quinton, 2004; Dahlke 26 and Lyon, 2013; Pomeroy et al., 2003). This was also demonstrated for the Rofen valley in this study (Fig. 10, 27 Table 3). Snowmelt results from a series of processes (e.g. energy exchange snow-atmosphere) that are spatially 28 variable - especially in complex terrain. This also becomes obvious when comparing the snowmelt rates on 23 29 April 2014 in Fig. 10a. Differences of observed and simulated snowmelt rates might result from the non-30 representativeness of point measurements for catchment averages and refer to the scale issue of data collection. 31 The peak melt period was characterised by less spatial and day-to-day variation in observed melt rates (Fig. 10). 32 The modelled daily snowmelt during this period was similar for north- and south-facing slopes, likely because of 33 higher melt rates but also a smaller snow-covered area of the south-facing slope in contrast to the north-facing 34 slope during peak melt (Fig. 11). The model performance was good for SWE (Table 3) and snow cover extent 35 (Fig. 2 and 9). The spatial variation of snowpack isotopic composition are significant, as can be seen in the 36 differences for north- and south-facing slopes, as also shown by Carey and Quinton (2004), Dahlke and Lyon 37 (2013) and Dietermann and Weiler (2013) in their high-gradient catchments, whereas there are unclear

38 differences for the spatial variation of the snowmelt isotopic signal in the literature. It is not clear to which extent

1 altitude is important, as Dietermann and Weiler (2013) stated that a potential elevation effect (decrease of snowmelt δ^{18} O with elevation) is likely to be disturbed by melting processes (isotopic enrichment) depending on 2 3 catchment morphology (aspect, slope) during the ablation period. Beaulieu et al. (2012) detected elevation as a predictor, which explained most of the variance they observed in snowmelt $\delta^{18}O$ from four distributed snow 4 lysimeters. Moore (1989) and Laudon et al. (2007) found no significant difference of δ^{18} O in their lysimeter 5 outflows, likely due to the small elevation gradient of their catchments which favours an isotopically 6 7 homogenous snowpack, whereas Unnikrishna et al. (2002) found a remarkable small-scale spatial variability. An 8 altitudinal gradient was not considered in this study, but possible effects on IHS are discussed in Section 5.6. 9 The difference of snowmelt (not snowpack) isotopic signature between north- and south-facing slopes was 10 clearly shown in this study. The dataset is small, but reveals clear differences induced by varying magnitudes 11 and timing of melt due to different intensities in solar radiation on the opposing slopes (cf. Fig. 7). Temporal 12 variability in snowmelt isotopic composition is greater for the north-facing slope compared to the south-facing 13 slope (Fig. 7), which was also pointed out by Carey and Quinton (2004) in their subarctic catchment. Earlier 14 homogenization in the isotopic profile of the snowpack and earlier melt out are responsible for this phenomenon 15 (cf. Dincer et al., 1970; Unnikrishna et al., 2002). Fractionation processes likely controlled this homogenization of the snowpack between the two investigated melt events. The isotopic homogenization of the snowpack on the 16 17 south-facing slope started earlier in the melting period and caused a smaller spatial and temporal variation 18 compared the north-facing snowpack, as also reported by Unnikrishna et al. (2002) and Dincer et al. (1970). The 19 differences between these investigated snowpacks in the early melt season were larger than in the peak melt 20 season. This affects the IHS results, especially because the snowmelt contributions from the south- and north-21 facing slope - with marked isotopic differences - were distinct. Due to melt, fractionation processes proceeded 22 and the snowpack became more homogenous throughout the snowmelt season. However, inter-daily variations 23 of snowpack isotopic composition, especially for the north-facing slope, were still observable during the peak 24 melt period. The gradual isotopic enrichment of the snowpack was also observed for snowmelt, as described by 25 many others (Feng et al., 2002; Shanley et al., 2002; Taylor et al., 2001; Taylor et al., 2002; Unnikrishna et al., 26 2002).

Intra-daily variations of snowmelt δ^{18} O could be quantified for two sites (Fig. 8). At S1 on the south-facing slope 27 during the early melt event, the 0.1 % range in δ^{18} O (n=5) was smaller than the range at N2 on the north-facing 28 29 slope during the peak melt event (n=3, range=0.2 ‰). This sub-daily variability is markedly smaller than the differences between the investigated slopes (cf. Table 1), which ranged from 0.8 ‰ (peak melt) to 1.4 ‰ (early 30 melt). Unnikrishna et al. (2002) described significant temporal variations of snowmelt δ^{18} O during large 31 32 snowmelt events (peak melt). However, these findings could not be confirmed within in this study, probably due 33 to the temporally limited data and should be tested with a larger dataset. The bulk sample at S1 (23 April 2014) 34 was isotopically closer to the sub-daily values compared to the bulk sample at N2 (07 June 2014) that was at the 35 upper range of the appropriate sub-daily samples (Fig. 7). Therefore one could argue that for the south-facing 36 slope there is a negligible uncertainty if one uses a single snowmelt value (at one time) for IHS instead of using a 37 bulk sample, but this is not the case for the north-facing slope (cf. Fig. 8, site N2). Unfortunately the sample 38 numbers are small, because more frequent and more distributed sampling (at different sites) was not feasible due 39 to logistical issues. Hence these results should be used with caution and should be investigated in further studies. 40 If the focus and the scale of the study is not on the sub-daily variability, the authors recommend to use bulk 41 samples, because these integrate (automatically weighed with snowmelt rate) the diurnal variations.

2 **5.3 Validity of isotopic hydrograph separation**

3 The validity of IHS relies on several assumptions (cf. Section 3.3; Buttle, 1994; Klaus and McDonnell, 2013). 4 The assumption that the isotopic composition of event and pre-event water differ significantly (Assumption 1) 5 was successfully proven, because the snowmelt isotopic values were markedly lower than pre-event baseflow 6 values (cf. Table 2 and 4, Fig. 5). Spatio-temporal variations of event water isotopic composition (Assumption 7 2) were accounted for by collecting daily and sub-daily samples during both events throughout the freshet period 8 and meltwater sampling at a north- and south-facing slope, respectively. The spatially variable input of event 9 water was considered by dividing the catchment into two parts -a north- and a south-facing slope. This study 10 supports the findings of Dahlke and Lyon (2013) and Carey and Quinton (2004), emphasizing the highly 11 variable snowpack/snowmelt isotopic composition due to enrichment in complex topography catchments. The 12 temporal variability of event water isotopic composition was considered by using bulk daily samples, which 13 integrate snowmelt from the entire diurnal melting cycle, but smooth out a sub-daily signal. Because the focus of 14 this study was more on the inter-event than the intra-daily scale, this approach seemed reliable. The spatio-15 temporal variability of the isotopic composition of pre-event water (Assumption 3) is a major limitation and could not be clearly identified due to a lack of data and was therefore assumed to be constant. Small differences 16 between the pre-event samples (-15.00 % and -15.05 % for δ^{18} O) and post-event stream water isotopic 17 18 composition support this assumption (Table 2). The assumption of soil water having the same isotopic 19 composition as groundwater in time and space (Assumption 4) is quite critical. Some studies reveal no 20 significant differences (e.g. Laudon et al., 2007), whereas others do (e.g. Sklash and Farvolden 1979). Isotopic 21 differences between groundwater and soil water were not considered due to a lack of data. Furthermore, it is not 22 known to which degree the vadose zone contributes to baseflow in the study area. Winter baseflow used in the 23 analyses is assumed to integrate mainly groundwater and partly soil water. Soil water could be hypothesized to have a negligible contribution to baseflow during winter due to the recession of the soil storage in autumn and 24 frozen soils in whether the assumption that no or minimal surface storage occurs (Assumption 5) is plausible 25 26 because water bodies like lakes or wetlands do not exist in the study catchment and due to the steep topography 27 detention storage is likely limited. The transit time of snowmelt was assumed to be less than 24 h. This short 28 travel time is characteristic for headwater catchments (Lundquist et al., 2005) with high in-channel flow 29 velocities, steep hillslopes, a high drainage density with snow-fed tributaries, thin soils, most snowmelt 30 originating from the edge of the snow-line (small average travel distances), partly frozen soil, and observed 31 surface overland flow. The state-of-the-art method (runCE) to include residence times of snowmelt in the event 32 water reservoir proposed by Laudon et al. (2002) was applied in several IHS studies (Beaulieu et al., 2012; Carey 33 and Quinton, 2004; Petrone et al., 2007), but was not feasible due to the short-term character and temporally 34 limited data of the experimental design.

35 5.4 Hydrograph separation results and inferred runoff generation processes

36 High contributions from snowmelt to streamflow are common in high-elevation catchments. Daily contributions

- 37 between 35 and 75 % in the Rofen valley are comparable to the results of studies conducted in other
- 38 mountainous regions, mostly outside the European Alps. Beaulieu et al. (2012) estimated snowmelt contributions
- ranging from 7 to 66 % at the seasonal scale for their 2.4 km² catchment and reported contributions of 34 and

1 62 %, for the early melt and peak melt, respectively. The hydrograph is dominated by pre-event water during early melt in April (Fig. 4c), which is in accordance with the results obtained by other IHS studies (Beaulieu et 2 3 al., 2012; Laudon et al., 2004; Laudon et al., 2007; Moore, 1989). The snowmelt contribution increased as the 4 freshet period progressed and peaked with high contributions at the beginning of June. Beaulieu et al. (2012) and 5 Sueker et al. (2000) reported comparable results for their physically similar catchments during peak melt with 62 6 and up to 76 % event water contributions to streamflow, respectively. At the event-scale comparable studies are 7 rare. Engel et al. (2016) report a maximum daily snowmelt contribution estimated with a three-component 8 hydrograph separation of 33 % for an 11 km² catchment southwest of the Rofen valley with similar 9 physiographic characteristics, but on the southern side of the main Alpine ridge. It should be mentioned that in 10 their study, runoff was fed by three components (snowmelt, glacier melt and groundwater) and lower snowmelt 11 contributions were prevalent because most of the catchment area (69 %) was snow-free.

12 Initial snowmelt events flush the pre-event water reservoir as snowmelt infiltrates into the soil and causes the 13 pre-event water to exfiltrate and contribute to the streamflow. As the soil and groundwater reservoir becomes 14 gradually filled with new water (snowmelt), the event water fraction in the stream increases. The system is also 15 wetter during peak melt. The dominance of event water in the hydrograph is interpreted as an outflow of pre-16 event water stored in the subsurface and the gradual replenishment of the soil and groundwater reservoirs by 17 event water. The higher water table – compared to the early melt period – could cause a transmissivity feedback 18 mechanism (Bishop, 1991). This is a common mechanism in catchments with glacial till (Bishop et al., 2011) 19 characterised by higher transmissivities and hence increased lateral flow velocities towards to the surface. 20 Runoff generation is spatially very variable in the study area. There are areas (meadow patches between rock 21 fields) where saturation excess overland flow is dominant (observed mainly at plots S1, S2 and Wind) and areas 22 (with larger rocks and debris) where rapid shallow subsurface flow can be assumed (plot N2). Catchment 23 morphology controls various hydrologic processes and hence the shape of the hydrograph. Upslope residence 24 times of snowmelt are usually smaller due to thinner-soils (observed during the field work), steeper slopes 25 (Sueker et al., 2000) and higher contributing areas of glaciers with impermeable ice (Behrens, 1978) and would 26 be indicators for the more flashy hydrograph during the peak melt season.

5.5 Impact of spatial varying snowmelt and its isotopic composition on IHS (Assessment of weighting approaches)

29 Klaus and McDonnell (2013) stress in their review paper the need for investigating the effects of the spatially 30 varying snowmelt and its isotopic composition on IHS. This study quantified the impact of the spatially varying 31 isotopic composition of snowmelt between north- and south facing slopes on IHS results for the first time. The 32 difference in volumetric snowmelt contribution to streamflow at the event-scale determined using the five 33 different weighting methods for IHS is maximal 24 % (NORTH approach vs. SOUTH approach). The data show 34 that the variations between the weighting approaches (VWS, VWO and VWE) are higher throughout the early 35 melt season (Table 6), because small-scale variability of snowmelt and its isotopic composition are more 36 pronounced in the early melt season. Thus the influence of spatial variability of snowmelt and its isotopic 37 composition on the event water fraction calculated with IHS is larger during this time. Melt rates strongly differ 38 between the south- and the north-facing slope (Fig. 11), which was deceptively gathered by manually measured SWE, likely due to micro-topographic effects. As the contributions from both slopes are used in Eq. (3), they 39 strongly influence the applied weighting tect the weighting method SOUTH (or NORTH) represents the 40

hypothetical and most extreme scenario in which only one sampling site is used for the IHS analysis. Because 1 snowmelt is more depleted in δ^{18} O and closer to pre-event water isotopic composition on the south-facing slope 2 3 during peak melt, this scenario has the greatest effect on IHS and leads to the strongest deviation in estimated 4 snowmelt fractions (up to 15 % overestimation compared to the VWS approach). These scenarios 5 (NORTH/SOUTH) are theoretical and it is obvious that it is not recommended to conduct a IHS analysis by 6 using only samples from either north- or south-facing slopes in catchments with complex terrain. Similar to the 7 VWE method, snowmelt isotopic data was not volume-weighted in other studies (e.g. Engel et al., 2016) where 8 snowmelt data was not available. This has a more distinct effect on IHS during the early melt season because of 9 the higher spatio-temporal variability in snowmelt (and its isotopic composition) compared to the peak melt 10 season and led to a deviation in the snowmelt fraction in streamflow of 2 % and 3 % compared to the VWS and 11 VWO approach, respectively. These differences are small, because the differing snowmelt and isotopic values 12 offset each other in this particular case (Table 6). Nevertheless the results of VWS are more correct for the right 13 reason, because single observed plot-scale melt rates do not represent distributed snowmelt contribution at the 14 catchment-scale. Therefore one can hypothesize that distributed simulated melt rates enhance the reliability and 15 feasibility of IHS, whereas plot-scale weighting introduces a very large error caused by the difficulty in finding 16 locations that represent the melt rate in complex terrain. The IHS resul 17 variability of snowmelt δ^{18} O, than spatial variations of snowmelt rates (Table 6). This is even more pronounced for the peak melt period, because snowmelt rates were similar for the north- and south-facing slope, probably 18 19 due to a ripe snow cover throughout the catchment.

20 **5.6 Limitations of the study**

21 Collecting water samples in high-elevation terrain is challenging due to limited access and high risk (e.g. 22 avalanches), limiting high-frequency sampling. Hence some limitations are inherent in this study. Potential 23 elevation effects on snowmelt isotopic composition were not tested. The opposing sampling sites (S1-N1 and S2-24 N2) were at the same elevation (Fig. 1). It was assumed that the differences of north- and south-facing slopes 25 were much greater than a possible altitudinal gradient in snowmelt isotopic composition. This hypothesis was 26 not tested, but based on the results of other studies (Dietermann and Weiler, 2013). However, accounting for a potential altitudinal gradient (decrease of snowmelt δ^{18} O with elevation) would lead to more depleted isotopic 27 signatures of event water and hence to lower event water fractions. Ther disadvantage is that no snow survey 28 29 was conducted prior to the onset of snowmelt (peak accumulation) to estimate spatial variability in bulk snow 30 δ^{18} O. Because snowmelt is used for applying IHS, it is not clear to which degree the spatial variability of the 31 snowpack isotopic composition is important. Two-component isotopic hydrograph separation was successfully 32 applied using the end-members snowmelt and baseflow, but potential contributions of glacier melt were 33 neglected (here defined as ice/firn melt). Because glaciers in the catchment were still covered by snow during the 34 peak melt season, a significant contribution from ice/firn melt was assumed to be unlikely. Nevertheless 35 negligible amounts of basal (ice) meltwater could originate from temperate glaciers. No samples could be 36 collected during the recession of the hydrograph (at night). Even though the spatial variability of the event water 37 signal was the focus of the study, only temporal variability was considered in the Genereux-based uncertainty 38 analyses. Although the temporal variability of winter baseflow isotopic composition seems to be insignificant, 39 the sample number (n=2) could be too small to characterize the pre-event component and should be clearly 40 investigated in future work. Penna et al. (2016) used two approaches to determine the isotopic composition of

1 pre-event water and described differences in the estimated event water contributions during snowmelt events. 2 They advise to take pre-event samples prior to the onset of the melt season, because using pre-event samples 3 taken prior to the onset of the diurnal melt cycle could be affected by snowmelt water from the previous melt 4 pulses and therefore could lead to underestimated snowmelt fractions and high uncertainties. Furthermore, model 5 results and observed discharges were assumed to be free of error in the analyses. As pointed out, instrumentation 6 and accessibility are major problems for high-elevation studies and their sampling strategies. For this study it 7 turned out that composite snowmelt samples were easier to collect, representing the day-integrated melt signal. A 8 denser network of melt collectors would be desirable, as well as a snow lysimeter to gain high-frequency data 9 automatically. Representative samples of the elevation zones and different vegetation belts could be important 10 too, especially in partly forested catchments with a distinct relief (cf. Unnikrishna et al., 2002).

11 **6** Conclusions

12 This study provides new insights into the variability of the isotopic composition in snowmelt and highlights its 13 impact on IHS results in a high-elevation environment. The spatial variability in snowmelt isotopic signature was 14 considered by experimental investigations on south- and north-facing slopes to define the isotopic composition 15 of the snowmelt end-member with greater accuracy. This study clearly shows that distributed snowmelt rates 16 simulated by a model, fed with meteorological data from local automatic weather stations, affect the weighting 17 of the event water isotopic signal, and hence the estimation of the snowmelt fraction in the stream by IHS. The 18 study provides a variety of relevant findings that are important for hydrologic research in high-alpine 19 environments. There was a distinct spatial variability in snowmelt between north- and south-facing slopes shown 20 for this complex terrain, especially during the early melt season. The isotopic composition of snowmelt water 21 was significantly different between north-facing and south-facing slopes, which resulted in a pronounced effect 22 on the estimated snowmelt contributions to streamflow with IHS. The differences in the estimated snowmelt 23 fraction due to the weighting methods used for IHS were as large as 24 %. This study also shows that it is hardly 24 possible to characterize the event water signature of larger slopes based on plot-scale snowmelt measurements. 25 Applying a distributed model reduced the uncertainty of the spatial snowmelt variability inherent in point-scale 26 observations. Hence, applying the VWS method provided more reasonable results than the VWO method. This 27 study highlighted that sampling north- and south-facing slopes is of major importance in conducting snowmelt-28 based IHS in mountainous catchments with complex topography in which a non-uniform input of snowmelt can 29 be expected, and that the selection of sampling sites has a major effect on IHS results. Sampling at least north-30 facing and south-facing slopes in complex terrain and using distributed melt rates to weight the snowmelt 31 isotopic composition of the differing exposures is therefore highly recommended for applying snowmelt-based 32 IHS.

33

34 Acknowledgments. The authors wish to thank the Institute of Atmospheric and Cryospheric Sciences of the University of 35 Innsbruck, the Zentralanstalt für Meteorologie and Geodynamik, the Hydrographic Service of Tyrol and the TIWAG-Tiroler 36 Wasserkraft AG for providing hydrological and meteorological data, the Amt der Tiroler Landesregierung for providing the 37 DEM, the Center of Stable Isotopes (CSI) for laboratory support, as well as many individuals who have helped to collect data 38 in the field. We also thank the reviewers for their valuable suggestions that have improved the manuscript a lot and the editor 39 for the careful handling of the manuscript.

1 References

- 2 Ahluwalia, R. S., Rai, S. P., Jain, S. K., Kumar, B., and Dobhal, D. P.: Assessment of snowmelt runoff
- 3 modelling and isotope analysis: a case study from the western Himalaya, India, Annals of Glaciology, 54, 299-
- 4 304, doi:10.3189/2013AoG62A133, 2013.
- 5 APCC: Austrian Assessment Report (AAR14). Summary for Policymakers (SPM), Austrian Panel on Climate
- 6 Change, Vienna, Austria, 2014.
- 7 Árnason, B., Buason, T., Martinec, J., and Theodorson, P.: Movement of water through snowpack traced by
- deuterium and tritium. In: The role of snow and ice in hydrology. Proc. Banff Symp., UNESCO-WMO-IAHS
 (Ed.), IAHS Publ. No., 107, 1973.
- 10 Beaulieu, M., Schreier, H., and Jost, G.: A shifting hydrological regime: a field investigation of snowmelt runoff
- 11 processes and their connection to summer base flow, Sunshine Coast, British Columbia, Hydrological Processes,
- 12 26, 2672-2682, doi:10.1002/hyp.9404, 2012.

13 Behrens, H., Moser, H., Oerter, H., Rauert, W., Stichler, W. and Ambach, W.: Models for the runoff from a

- 14 glaciated catchment area using measurements of environmental isotope contents, 1978, 829-846.
- 15 Birkel, C. and Soulsby, C.: Advancing tracer-aided rainfall-runoff modelling: a review of progress, problems and
- 16 unrealised potential, Hydrological Processes, 29, 5227-5240, doi:10.1002/hyp.10594, 2015.
- 17 Birkel, C., Tetzlaff, D., Dunn, S. M., and Soulsby, C.: Using time domain and geographic source tracers to
- 18 conceptualize streamflow generation processes in lumped rainfall-runoff models, Water Resources Research, 47,
- 19 W02515, doi:10.1029/2010WR009547, 2011.
- 20 Bishop, K.: Episodic increase in stream acidity, catchment flow pathways and hydrograph separation, 1991.
- 21 Bishop, K., Seibert, J., Nyberg, L., and Rodhe, A.: Water storage in a till catchment. II: Implications of
- 22 transmissivity feedback for flow paths and turnover times, Hydrological Processes, 25, 3950-3959,
- 23 doi:10.1002/hyp.8355, 2011.
- 24 Braithwaite, R. J. and Olesen, O. B.: Calculation of glacier ablation from air temperature, West Greenland. In:
- 25 Glacier Fluctuations and Climatic Change, Glaciology and Quaternary Geology, Oerlemans, J. (Ed.), Kluwer
- 26 Academic Publisher, Dordrecht, 1989.
- 27 Buttle, J. M.: Isotope hydrograph separations and rapid delivery of pre-event water from drainage basins,
- 28 Progress in Physical Geography, 18, 16-41, doi:10.1177/030913339401800102, 1994.
- 29 Capell, R., Tetzlaff, D., and Soulsby, C.: Can time domain and source area tracers reduce uncertainty in rainfall-
- 30 runoff models in larger heterogeneous catchments?, Water Resources Research, 48, W09544,
- doi:10.1029/2011WR011543, 2012.
- 32 Carey, S. K. and Quinton, W. L.: Evaluating snowmelt rnoff generation in a discontinuous permafrost catchment
- using stable isotope, hydrochemical and hydrometric data, Nordic Hydrology, 35, 309-324, 2004.
- 34 Clark, I. D. and Fritz, P.: Environmental Isotopes in Hydrogeology, Lewis Publishers, Ney York, 1997.
- 35 CLC: Corine Land Cover 2006 raster data. European Environment Agency. The European Topic Centre on Land
- 36 Use and Spatial Information. 2006.
- 37 Cooper, L. W.: Isotopic fractionation in snow cover. In: Isotope tracers in catchment hydrology, Kendall, C. and
- 38 McDonnell, J. J. (Eds.), Elsevier Science, Amsterdam, Netherlands, 2006.
- 39 Corripio, J. G.: Vectorial algebra algorithms for calculating terrain parameters from DEMs and the position of
- 40 the sun for solar radiation modelling in mountainous terrain, International Journal of Geographical Information
- 41 Science, 17, 1-23, 2003.

- 1 Dahlke, H. E. and Lyon, S. W.: Early melt season snowpack isotopic evolution in the Tarfala valley, northern
- 2 Sweden, Annals of Glaciology, 54, 149-156, doi:10.3189/2013AoG62A232, 2013.
- 3 Dietermann, N. and Weiler, M.: Spatial distribution of stable water isotopes in alpine snow cover, Hydrology
- 4 and Earth System Sciences, 17, 2657-2668, doi:10.5194/hess-17-2657-2013, 2013.
- 5 Dinçer, T., Payne, B. R., Florkowski, T., Martinec, J., and Tongiorgi, E.: Snowmelt runoff from measurements 6 of tritium and oxygen-18, Water Resources Research, 6, 110-124, doi:10.1029/WR006i001p00110, 1970.
- 7 Engel, M., Penna, D., Bertoldi, G., Dell'Agnese, A., Soulsby, C., and Comiti, F.: Identifying run-off
- 8 contributions during melt-induced run-off events in a glacierized alpine catchment, Hydrological Processes, 30,
- 9 343-364, doi:10.1002/hyp.10577, 2016.
- 10 Feng, X., Taylor, S., Renshaw, C. E., and Kirchner, J. W.: Isotopic evolution of snowmelt 1. A physically based
- 11 one-dimensional model, Water Resources Research, 38, 35-31-35-38, doi:10.1029/2001WR000814, 2002.
- 12 Genereux, D.: Quantifying uncertainty in tracer-based hydrograph separations, Water Resources Research, 34,
- 13 915-919, doi:10.1029/98WR00010, 1998.
- 14 Greuell, W., Knap, W. H., and Smeets, P. C.: Elevational changes in meteorological variables along a
- 15 midlatitude glacier during summer, Journal of Geophysical Research: Atmospheres, 102, 25941-25954,
- 16 doi:10.1029/97JD02083, 1997.
- 17 Gruber, S.: A mass-conserving fast algorithm to parameterize gravitational transport and deposition using digital
- 18 elevation models, Water Resources Research, 43, W06412, doi:10.1029/2006WR004868, 2007.
- Haeberli, W.: Untersuchungen zur Verbreitung von Permafrost zwischen Flüelapass und Piz Grialetsch(Graubünden), 1975.
- 21 Hanzer, F., Helfricht, K., Marke, T., and Strasser, U.: Multilevel spatiotemporal validation of snow/ice mass
- balance and runoff modeling in glacierized catchments, The Cryosphere, 10, 1859-1881, doi:10.5194/tc-10-
- 23 1859-2016, 2016..
- 24 Hanzer, F., Marke, T., and Strasser, U.: Distributed, explicit modeling of technical snow production for a ski
- area in the Schladming region (Austrian Alps), Cold Regions Science and Technology, 108, 113-124,
 doi:http://dx.doi.org/10.1016/j.coldregions.2014.08.003, 2014.
- 27 Helfricht, K.: Analysis of the spatial and temporal variation of seasonal snow accumulation in Alpine catchments
- 28 using airborne laser scanning. Basic research for the adaptation of spatially distributed hydrological models to
- 29 mountain regions, PhD, University of Innsbruck, Innsbruck, 134 pp., 2014.
- 30 Hock, R.: Temperature index melt modelling in mountain areas, Journal of Hydrology, 282, 104-115,
- 31 doi:10.1016/S0022-1694(03)00257-9, 2003.
- 32 Hooper, R. P. and Shoemaker, C. A.: A Comparison of Chemical and Isotopic Hydrograph Separation, Water
- 33 Resources Research, 22, 1444-1454, doi:10.1029/WR022i010p01444, 1986.
- 34 Huth, A. K., Leydecker, A., Sickman, J. O., and Bales, R. C.: A two-component hydrograph separation for three
- high-elevation catchments in the Sierra Nevada, California, Hydrological Processes, 18, 1721-1733,
 doi:10.1002/hyp.1414, 2004.
- 20 doi:10.1002/hjp.111,2001.
- 37 IPPC: Summary for Policymakers. Climate Change 2013: The Physical Science Basis. Contribution of Working
- 38 Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge, United
- 39 Kindom and New York, NY,USA., 2013.
- 40 Klaus, J. and McDonnell, J. J.: Hydrograph separation using stable isotopes: Review and evaluation, Journal of
- 41 Hydrology, 505, 47-64, doi:10.1016/j.jhydrol.2013.09.006, 2013.

- 1 Laudon, H., Hemond, H. F., Krouse, R., and Bishop, K. H.: Oxygen 18 fractionation during snowmelt:
- 2 Implications for spring flood hydrograph separation, Water Resources Research, 38, 40-41-40-10,
- 3 doi:10.1029/2002WR001510, 2002.
- 4 Laudon, H., Seibert, J., Köhler, S., and Bishop, K.: Hydrological flow paths during snowmelt: Congruence
- 5 between hydrometric measurements and oxygen 18 in meltwater, soil water, and runoff, Water Resources
- 6 Research, 40, W03102, doi:10.1029/2003WR002455, 2004.
- 7 Laudon, H., Sjöblom, V., Buffam, I., Seibert, J., and Mörth, M.: The role of catchment scale and landscape
- 8 characteristics for runoff generation of boreal streams, Journal of Hydrology, 344, 198-209, 9 doi:10.1016/j.jhydrol.2007.07.010, 2007.
- 10 Lee, J., Feng, X., Faiia, A. M., Posmentier, E. S., Kirchner, J. W., Osterhuber, R., and Taylor, S.: Isotopic
- 11 evolution of a seasonal snowcover and its melt by isotopic exchange between liquid water and ice, Chemical
- 12 Geology, 270, 126-134, doi:10.1016/j.chemgeo.2009.11.011, 2010.
- 13 Liston, G. E. and Elder, K.: A Distributed Snow-Evolution Modeling System (SnowModel), Journal of
- 14 Hydrometeorology, 7, 1259-1276, doi:10.1175/JHM548.1, 2006.
- 15 Liu, F., Williams, M. W., and Caine, N.: Source waters and flow paths in an alpine catchment, Colorado Front
- Range, United States, Water Resources Research, 40, 1-16, doi:10.1029/2004WR003076, 2004. 16
- 17 Lundquist, J. D. and Cayan, D. R.: Seasonal and Spatial Patterns in Diurnal Cycles in Streamflow in the Western
- 18 United States, 3. 591-603. Journal of Hydrometeorology, doi:10.1175/1525-19 7541(2002)003<0591:SASPID>2.0.CO;2, 2002.
- 20 Lundquist, J. D., Dettinger, M. D., and Cayan, D. R.: Snow-fed streamflow timing at different basin scales: Case
- 21 study of the Tuolumne River above Hetch Hetchy, Yosemite, California, Water Resources Research, 41,
- 22 W07005, doi:10.1029/2004WR003933, 2005.
- 23 Marke, T.: Development and Application of a Model Interface to couple Regional Climate Models with Land
- 24 Surface Models for Climate Change Risk Assessment in the Upper Danube Watershed, Dissertation der Fakultät
- 25 für Geowissenschaften, Digitale Hochschulschriften der LMU München, München, 2008.
- 26 Marke, T., Strasser, U., Hanzer, F., Stötter, J., Wilcke, R. A. I., and Gobiet, A.: Scenarios of Future Snow
- 27 Conditions in Styria (Austrian Alps), Journal of Hydrometeorology, 16, 261-277, doi:10.1175/JHM-D-14-
- 28 0035.1, 2015.
- 29 Martinec, J., Siegenthaler, U., Oeschger, H., and Tongiorgi, E.: New insights into the run-off mechanism by
- environmental isotopes. In: Isotope techniques in groundwater hydrology, Symposium, P. o. a. I. (Ed.), IAEA, 30
- 31 Vienna, Austria, 1974.
- 32 Mast, A. M., Kendall, K., Campbell, D. H., Clow, D. W., and Back, J.: Determination of hydrologic pathways in
- 33 an alpine-subalpine basin using isotopic and chemical tracers, Loch Vale Watershed, Colorado, Usa. In: IAHS 34
- Publ. Ser. Proc. Rep.Int. Assoc. Hydrol. SCI., 1995.
- 35 Maulé, C. P. and Stein, J.: Hydrologic Flow Path Definition and Partitioning of Spring Meltwater, Water 36 Resources Research, 26, 2959-2970, doi:10.1029/WR026i012p02959, 1990.
- 37 Moore, R. D.: Tracing runoff sources with deuterium and oxygen-88 during spring melt in a headwater
- 38 southern Journal Hydrology, catchment, Laurentians, Quebec, of 112, 135-148,
- 39 doi:http://dx.doi.org/10.1016/0022-1694(89)90185-6, 1989.

- 1 Pellicciotti, F., Brock, B., Strasser, U., Burlando, P., Funk, M., and Corripio, J.: An enhanced temperature-index
- 2 glacier melt model including the shortwave radiation balance: development and testing for Haut Glacier d'Arolla,
- 3 Switzerland, Journal of Glaciology, 51, 573-587, 2005.
- 4 Penna, D., van Meerveld, H. J., Zuecco, G., Dalla Fontana, G., and Borga, M.: Hydrological response of an
- 5 Alpine catchment to rainfall and snowmelt events, Journal of Hydrology, 537, 382-397,
 6 doi:http://dx.doi.org/10.1016/j.jhydrol.2016.03.040, 2016.
- 7 Penna, D., Engel, M., Mao, L., Dell'Agnese, A., Bertoldi, G., and Comiti, F.: Tracer-based analysis of spatial and
- 8 temporal variations of water sources in a glacierized catchment, Hydrology and Earth System Sciences, 18,
- 9 5271-5288, doi:10.5194/hess-18-5271-2014, 2014.
- 10 Petrone, K., Buffam, I., and Laudon, H.: Hydrologic and biotic control of nitrogen export during snowmelt: A
- combined conservative and reactive tracer approach, Water Resources Research, 43, 1-13,
 doi:10.1029/2006WR005286, 2007.
- 13 Pinder, G. F. and Jones, J. F.: Determination of the ground-water component of peak discharge from the
- 14 chemistry of total runoff, Water Resources Research, 5, 438-445, doi:10.1029/WR005i002p00438, 1969.
- 15 Pomeroy, J. W., Toth, B., Granger, R. J., Hedstrom, N. R., and Essery, R. L. H.: Variation in Surface Energetics
- during Snowmelt in a Subarctic Mountain Catchment, Journal of Hydrometeorology, 4, 702-719,
 doi:10.1175/1525-7541(2003)004<0702:VISEDS>2.0.CO;2, 2003.
- 18 Rohrer, M. B.: Die Schneedecke im Schweizer Alpenraum und ihre Modellierung, ETH, Zürich, 1992.
- Schuler, T.: Investigation of water drainage through an alpine glacier by tracer experiments and numericalmodeling., 2002.
- 21 Seibert, J. and McDonnell, J. J.: On the dialog between experimentalist and modeler in catchment hydrology:
- 22 Use of soft data for multicriteria model calibration, Water Resources Research, 38, 23-21-23-14,
- 23 doi:10.1029/2001WR000978, 2002.
- 24 Shanley, J. B., Kendall, C., Smith, T. E., Wolock, D. M., and McDonnell, J. J.: Controls on old and new water
- contributions to stream flow at some nested catchments in Vermont, USA, Hydrological Processes, 16, 589-609,
- 26 doi:10.1002/hyp.312, 2002.
- Sklash, M. G. and Farvolden, R. N.: The role of groundwater in storm runoff, Journal of Hydrology, 43, 45-65,
 doi:10.1016/0022-1694(79)90164-1, 1979.
- 29 Sklash, M. G., Farvolden, R. N., and Fritz, P.: A conceptual model of watershed response to rainfall, developed
- 30 through the use of oxygen-18 as a natural tracer, Canadian Journal of Earth Sciences, 13, 271-283,
- doi:10.1139/e76-029, 1976.
- 32 Stichler, W.: Snowcover and Snowmelt Processes Studied by Means of Environmental Isotopes. In: Seasonal
- 33 Snowcovers: Physics, Chemistry, Hydrology, Jones, H. G. and Orville-Thomas, W. J. (Eds.), D. Reidel
- 34 Publishing Company, Dordrecht, Holland, 1987.
- 35 Strasser, U.: Modelling of the mountain snow cover in the Berchtesgaden National Park, Research Rep., 2008.
- 36 Strasser, U., Bernhardt, M., Weber, M., Liston, G. E., and Mauser, W.: Is snow sublimation important in the 37 alpine water balance?, The Cryosphere, 2, 53-66, doi:10.5194/tc-2-53-2008, 2008.
- 38 Strasser, U., Corripio, J., Pellicciotti, F., Burlando, P., Brock, B., and Funk, M.: Spatial and temporal variability
- 39 of meteorological variables at Haut Glacier d'Arolla (Switzerland) during the ablation season 2001:
- 40 Measurements and simulations, Journal of Geophysical Research: Atmospheres, 109, D03103,
- 41 doi:10.1029/2003JD003973, 2004.

- 1 Strasser, U., Warscher, M., and Liston, G. E.: Modeling Snow-Canopy Processes on an Idealized Mountain,
- 2 Journal of Hydrometeorology, 12, 663-677, doi:10.1175/2011JHM1344.1, 2011.
- 3 Sueker, J. K., Ryan, J. N., Kendall, C., and Jarrett, R. D.: Determination of hydrologic pathways during
- 4 snowmelt for alpine/subalpine basins, Rocky Mountain National Park, Colorado, Water Resources Research, 36,
- 5 63-75, doi:10.1029/1999WR900296, 2000.
- 6 Taylor, S., Feng, X., Kirchner, J. W., Osterhuber, R., Klaue, B., and Renshaw, C. E.: Isotopic evolution of a
- 7 seasonal snowpack and its melt, Water Resources Research, 37, 759-769, doi:10.1029/2000WR900341, 2001.
- 8 Taylor, S., Feng, X., Williams, M., and McNamara, J.: How isotopic fractionation of snowmelt affects
- 9 hydrograph separation, Hydrological Processes, 16, 3683-3690, doi:10.1002/hyp.1232, 2002.
- 10 Uhlenbrook, S. and Leibundgut, C.: Process-oriented catchment modelling and multiple-response validation,
- 11 Hydrological Processes, 16, 423-440, doi:10.1002/hyp.330, 2002.
- 12 Unnikrishna, P. V., McDonnell, J. J., and Kendall, C.: Isotope variations ni a Sierra Nevada snowpack and their
- 13 relation to meltwater, 260, 38-57, 2002.
- 14 Warscher, M., Strasser, U., Kraller, G., Marke, T., Franz, H., and Kunstmann, H.: Performance of complex snow
- 15 cover descriptions in a distributed hydrological model system: A case study for the high Alpine terrain of the
- 16 Berchtesgaden Alps, Water Resources Research, 49, 2619-2637, doi:10.1002/wrcr.20219, 2013.
- 17 Weingartner, R. and Aschwanden, H.: Discharge regime the basis for the estimation of average flows. In:
- 18 Hydrological Atlas of Switzerland, 1992.
- 19 Williams, M. W., Seibold, C., and Chowanski, K.: Storage and release of solutes from a subalpine seasonal
- 20 snowpack: soil and stream water response, Niwot Ridge, Colorado, Biogeochemistry, 95, 77-94,
- 21 doi:10.1007/s10533-009-9288-x, 2009.
- 22 Zappa, M.: Objective quantitative spatial verification of distributed snow cover simulations—an experiment for
- the whole of Switzerland, Hydrological Sciences Journal, 53, 179-191, doi:10.1623/hysj.53.1.179, 2008.
- 24 Zhou, S., Nakawo, M., Hashimoto, S., and Sakai, A.: The effect of refreezing on the isotopic composition of
- 25 melting snowpack, Hydrological Processes, 22, 873-882, doi:10.1002/hyp.6662, 2008.
- 26



Figure 1: (a) Distribution of slope aspects in the study area; (b) Study area (Rofen valley) with underlying orthophoto, sampling and measurement locations.



Figure 2: Comparison of observed and simulated snow distributions for (a, b) May 5 (MODIS scene) and (c, d)
 June 10, 2014 (Landsat scene).



1

Figure 3: (a) Daily precipitation, air temperature, and discharge during the complete study period; Hourly hydroclimatologic data of a 7-day period around the (b) early melt and (c) peak melt event. Data was measured at the outlet of the catchment. Grey-shaded areas indicate the investigated events.



Figure 4: Linearly interpolated stream isotopic content of Rofenache for (a) the early melt and (b) the peak melt event. Dots indicate measurements. Event and pre-event water contributions during (c) the early melt and (d) the peak melt event calculated with the VWS approach.





Figure 5: Relationship between δ²H and δ¹⁸O values of water sources sampled during the snowmelt season 2014 in the
 Rofen valley, Austrian Alps.



Figure 6: Jittered dot plots for δ^{18} O of collected water samples split into (a) water sources, (b) stage of snowmelt and (c) spatial origin. Grey circles indicate early melt samples and black circles are for peak melt samples. The grey and

1 2 black line represents the median of early and peak melt data, respectively. Ng is the number of early melt samples and n_p is the number of peak melt samples.



3

- Figure 7: Jittered dot plots for δ^{18} O of (a) snowpack and (b) snowmelt of north- and south-facing slopes. Grey circles
- 4 5 6 7 indicate early melt samples and black circles are for peak melt samples. The grey and black line indicates the median of the early and peak melt data, respectively. N_p is the number of early melt samples and n_p is the number of peak melt samples.



8

- 9 Figure 8: Comparison of snowmelt $\delta^{18}O$ between bulk sample (dashed line) and sub-daily samples (circles) for two 10 sites (S1, N2).
- 11





Figure 9: Performance measures (a) Accuracy (ACC), (b) Critical Success Index (CSI), and (c) BIAS as calculated by comparing AMUNDSEN simulation results with satellite-derived (MODIS/Landsat) snow maps.





Figure 10: Differences between the observed (plot scale) and simulated (catchment scale) daily snowmelt on (a) the north-facing and (b) the south-facing slope for the early melt (23/24 April) and peak melt (07/08 June).



Figure 11: Comparison of the spatial contribution of weighting apperbase. VWS: volume-weighted with simulated (areal) melt rates. VWO: volume-weighted with observed (plot-scale) melt rates.





Figure 12: Comparison of weighting techniques used for estimating snowmelt fraction with IHS during (a) early melt
 and (b) peak melt. Scale of Y-axis in b) differs from that in a).

10Table 1: Average isotopic content of snowpack and snowmelt with standard deviation for north- and south-facing11slopes during the early and the peak melt event. Values are averages of three consecutive days.

North-fac	ing slope	South-facing slope		
Snowpack $\delta^{18}O$ (‰)	Snowmelt $\delta^{18}O(\%)$	Snowpack $\delta^{18}O(\%)$	Snowmelt $\delta^{18}O(\%)$	

Early melt event	-19.7±0.6	-18.8±0.2	-17.3±0.3	-17.4±0.2
	(n=12)	(n=3)	(n=4)	(n=8)
Peak melt event	-17.6±0.4	-17.9±0.1	-17.9±0.1	-17.1±0.0
	(n=18)	(n=3)	(n=15)	(n=2)

Table 2: Descriptive statistics of streamflow isotopic content (Rofenache) during events of the snowmelt season 2014.
 Data is sampled at the outlet of the basin.

	Pre-event	Early melt	Peak melt	Post-event
Date	07/03	23/04 - 25/04	07/06 - 09/06	11/07
Average (δ^{18} O ‰)	-15.02	-15.97	-16.87	-15.09
Standard deviation ($\delta^{18}O$ ‰)	0.04	0.16	0.05	n/a
Range (δ^{18} O ‰)	0.05	0.50	0.20	n/a
Number of samples	2	17	30	1

7 Table 3: Comparison of observed and simulated (represented by the underlying pixel) SWE values at the plot-scale.

Site	Date	Stage of snowmelt	SWE [mm]		Difference between
		season	Observed	Simulated	simulated SWE [%]
S1	2014-04-23	Early melt	141	151	7
N1	2014-04-23	Early melt	351	356	1
Wind	2014-04-24	Early melt	201	229	14
S 1	2014-04-25	Early melt	113	78	-31
N1	2014-04-25	Early melt	270	293	9
N2	2014-06-07	Peak melt	594	477	-20
N2	2014-06-08	Peak melt	568	435	-23
N2	2014-06-09	Peak melt	537	390	-27
Mean deviation between observed and simulated SWE					13

10 Table 4: Isotopic characterization of the event water component by the applied weighting techniques

	Event	Event water isotopic composition $(\delta^{18}O \%)$			
	23/04	24/04	07/06	08/06	
VWS	-17.9	-18.2	-17.5	-17.5	
VWO	-18.3	-18.6	-17.4	-17.5	
VWE	-18.1	-18.3	-17.5	-17.5	

NORTH	-18.6	-18.8	-17.9	-17.9
SOUTH	-17.6	-17.9	-17.1	-17.1

Table 5: Discharge quantities of the Rofenache for the early and peak melt event at the outlet of the basin.

	Event		
	Early Melt	Peak Melt	
Date	23/04 - 25/04	07/06 - 09/06	
Mean discharge	$1.5 \text{ m}^3 \text{ s}^{-1}$	$11.5 \text{ m}^3 \text{ s}^{-1}$	
Peak discharge	$1.9 \text{ m}^3 \text{ s}^{-1}$	$17.4 \text{ m}^3 \text{ s}^{-1}$	
Volume runoff	3.3 mm	20.7 mm	
Mean event water fraction	35±3 %	75±14 %	
Peak event water fraction	44±4 %	78±15 %	

Table 6: Event water contribution to streamflow <u>estimated with different</u> weighting techniques. The error indicates
 the variability (standard deviation) and the <u>brackets</u> depict the range.

	Event water contribution (%)				
	VWS	VWO	VWE	NORTH	SOUTH
Early melt event	35±6	30±4	33±5	28±3	40±9
	(25-44)	(22-35)	(24-39)	(21-31)	(28-55)
Peak melt event	75±2	78±3	76±2	66±2	90±3
	(70-78)	(71-82)	(70-78)	(60-67)	(84-94)