

Interactive comment on “Bridging glacier and river catchment scales: an efficient representation of glacier dynamics in a hydrological model” by Michel Wortmann et al.

Anonymous Referee #3

Received and published: 14 July 2016

I enjoyed working through this manuscript. The authors have introduced lots of new ideas and sensible approaches to this kind of modeling. I agree that they are addressing an important gap between hydrological and glaciological models, at the catchment scale, and this effort is a genuine bridge between those worlds. There are new ideas and the model is applied in two interesting and well-selected locations. The paper is generally well-written, with strong conclusions that are well-supported by the model results. I think this work will find a receptive audience and it is likely that others will build on the model presented here. For these reasons, I recommend publication, with a number of minor points and perhaps one or two larger concerns to be addressed.

Main concerns

[Printer-friendly version](#)

[Discussion paper](#)



1. I have one specific concern about the model, which might just need more explanation or might suggest a 'push-back' towards more substantial revisions. On p.5, l.17 and Eq. (1), the critical snow thickness to be ice does not make sense to me. H_c . For typical slopes, like 10 degrees, $\tan\alpha \sim 0.1$ and $H_c = 100$ m. Does this mean that snow that survives the summer does not turn into ice unless there is at least 100 m of snow/ice accumulated? What about the snow thermal and albedo properties, etc., those should match ice after one year. Also, ice that is thinner than this creeps and slides. It does not wait until it reaches a critical shear stress before it starts moving. That is a misinterpretation of τ_s , which is more a 'balance' value where steady-state fluxes allow an equilibrium ice thickness. If I understand correctly, this seems odd an arbitrary for a glacier not to exist, thermodynamically and mechanically, until this much ice has accumulated. It would preclude many of today's present glaciers (and parts of them), which are thinner than this.

I also did not understand Eq. 3 or the area treatment on the next page, this could be explained more clearly I think. As I read it, H_c is maintained while area decreases in the lowest elevation band? I like this general idea, it is a nice new idea, but it seems unrealistic to maintain a steep and non-thinning layer of glacier ice while the area retreats. Shouldn't both decline at once, following a realistic volume-area relationship or what one would expect for a 'wedge-shaped' terminus?

2. As the title of my review suggests, I did struggle with whether this model is sufficiently physically-based and state-of-the-art to actually be useful. That sounds harsh, perhaps, but there are more complex and realistic models out there (many of them cited by the authors), and the model proposed here has several free parameters of an empirical nature, e.g., degree day factors, which are not actual physical variables. These are tuned to observed discharge in the specific basins, and results are reasonable, but how portable are they in space and time? The authors do nicely balance complexity with pragmatism, with a relatively simple treatment of a lot of the processes, but perhaps appropriate for the large-scale objectives. I do agree with the authors that forcing

[Printer-friendly version](#)

[Discussion paper](#)



data are not commonly available for more complex models (e.g., energy balance melt models, or more detailed glacier processes such as sliding). I just have concerns that this model is heavily parameterized and tuned in ways that are not representing the actual physics, which make it unclear how broadly useful it is. For instance, no attention is paid to conservation of energy or mass at the catchment scale; precipitation and mass balance are scaled as necessary, and melt rates are turned up or down ad hoc in order to match discharge observations. In the final lines, the authors note an intention to use this modeling approach in future projections, but is it reasonable to take climate model precipitation and energy fields and manipulate them in this way, paying no heed to basic conservation? I have concerns that this is too far from constrained reality. Still, this model presents an initial step into coupled glacio-hydrological modeling that has yet to be done well at large scales. For this reason I don't think my concern here is fatal.

Some specific questions on this point:

How do the final parameter sets vary for the different catchments/sub-catchments? Are there generally sensible, robust, and repeatable parameter values, that you would feel comfortable to use in other environments or in future projections?

Is the glacier melt model the same as that for the SWIM snow model? I did not realize that degree day melt models were still in broad use, scientifically. The range of melt factors explored here (Table 1) looks like values that are common literature values used for snow and ice melt, without incorporating the effects of shading, aspect, debris cover, etc. I have trouble to imagine that this fully represents the range of potential values.

Daily mean temperature is also simplistic when it comes to estimating PDD for snow and ice melt. e.g., a mean temperature of -1 C means no melt, although much of the day will be above 0 C. Minimum and maximum temperature are widely available and can be used to generate a daily temperature cycle - can this be considered?

In general the modeling approach emphasizes the model parsimony, that it does not

[Printer-friendly version](#)

[Discussion paper](#)



need many input variables, and only things like temperature that are 'more known'. But it does need regional mass balances, across individual glaciers: both in balance, for the spinup, and maybe also in time, for the model calibration - I was less sure here. Are these widely known, or known well enough to be able to use this model in much of the world? It seems like this is 'higher order' knowledge than some of the basic meteorological variables you would need for an energy balance model.

Back to degree day factors.. I don't understand why these are based on length of day. Why not actually calculate potential shortwave radiation as a function of latitude, day of year, slope, aspect and shading? This is a more direct and realistic way to include this effect, and can be pre-processed easily. Do the length of day calculations include slope and aspect effects, and shading?

And some minor points:

p.1, l.20, "strongly heterogeneous processes like glacier dynamics" - is that really true? glacier dynamics work roughly the same way everywhere. But ice thickness and slope vary strongly in space.

p.3, l.30, "is described"

p.5, l.3, there is some jargon throughout, like 'cleaning area' - please define. Also 'hydrotope', referenced below on this page. How does a hydrotope compare with an HRU, or are they equivalent?

p.5, l.12, 30 cm of soil cover actually seems like a lot for steep alpine terrain, which is more likely bare rock. Am I mid-understanding here?

p.7, Eq. 10. I like this in general, the approach to separate melt and sublimation. But note that in fact M in field-based PDD calibration studies includes sublimation, i.e. it is actually observed ablation, M+S. But generally $S \ll M$ where PDD factors are being calibrated, so this is maybe OK. This is not true everywhere though.

p.9, debris discussion. I was not sure that these are the main processes involved. Also

[Printer-friendly version](#)

[Discussion paper](#)



consider landslide/rock avalanche debris? Also aerosol deposition, which can be a blanket or it can also be concentrated by ablation and streams, e.g. where streams intersect crevasses. These processes should at least be noted, as they can be locally or regionally important.

p.12, l.28, "complemented". Next line down, "its"

p.13, l.19, implemented

Figure 4, outlet of the Rhone. It looks like there are diurnal cycles here - is this really daily discharge, or is it hourly? If the former, what are the oscillations?

p.15, l.13, comparison "is" shown

p.16, l.8, principal

p.16, l.9, I don't think one year can have a climate. The year's weather?

p.16, l.14, "varies"

p.18, l.16, I don't think it is wise to say that it covers "all major glacier processes". As the authors point out, things like glacier sliding, surging, and calving are not included, and these could be considered major processes. Also seasonal albedo evolution and many other details of glacier ablation.

Interactive comment on Hydrol. Earth Syst. Sci. Discuss., doi:10.5194/hess-2016-272, 2016.

Printer-friendly version

Discussion paper

