

1 **River water quality changes in New Zealand over 26 years (1989 – 2014): Response to land**
2 **use intensity**

3
4 Jason P. Julian*^{1,5}, Kirsten M. de Beurs^{2,5}, Braden Owsley^{2,5}, Robert J. Davies-Colley³, Anne-
5 Gaelle E. Ausseil⁴

6 ¹Department of Geography, Texas State University, San Marcos, TX, USA

7 ²Department of Geography and Environmental Sustainability, The University of Oklahoma,
8 Norman, OK, USA

9 ³National Institute of Water and Atmospheric Research Ltd (NIWA), Hamilton, New Zealand

10 ⁴Landcare Research, Palmerston North, New Zealand

11 ⁵Landscape & Land Use Change Institute (LLUCI), <http://tethys.dges.ou.edu/main/>, USA

12 *Corresponding author: Jason.Julian@txstate.edu

13

14 **Abstract**

15 Land use-water quality relationships are complex with interdependencies, feedbacks, and legacy
16 effects. Most river water quality studies have assessed catchment land use as areal coverage, but
17 here, we hypothesize and test whether land use *intensity* – the inputs (fertilizer, livestock) and
18 activities (vegetation removal) of land use – is a better predictor of environmental impact. We
19 use New Zealand as a case study because it has had one of the highest rates of agricultural land
20 intensification globally over recent decades. We interpreted water quality state and trends for the
21 26 years from 1989 to 2014 in the National Rivers Water Quality Network (NRWQN) –
22 consisting of 77 sites on 35 mostly large river systems with an aggregate catchment amounting to
23 half of NZ's land area. To characterize land use intensity, we analyzed spatial and temporal

24 changes in livestock density and land disturbance (i.e. bare soil resulting from vegetation loss by
25 either grazing or forest harvesting) at the catchment-scale, as well as fertilizer inputs at the
26 national scale. Using simple multivariate statistical analyses across the 77 catchments, we found
27 that visual water clarity was best predicted by areal coverage of high-producing pastures. The
28 primary predictor for all four nutrient variables, however, was cattle density, with plantation
29 forest coverage as the secondary predictor variable. While land disturbance was not itself a
30 strong predictor of water quality, it did help explain outliers of land use-water quality
31 relationships. From 1990 to 2014, visual clarity significantly improved in 34/77 catchments,
32 which we attribute mainly to increased dairy cattle exclusion from rivers (despite dairy
33 expansion) and the considerable decrease in sheep numbers across the NZ landscape, from 58
34 million sheep in 1990 to 31 million in 2012. Nutrient concentrations increased in many of NZ's
35 rivers with dissolved oxidized nitrogen significantly increasing in 27/77 catchments, which we
36 largely attribute to increased cattle density and legacy nutrients that have built up on high-
37 producing grasslands and plantation forests since the 1950s and are slowly leaking to the rivers.
38 Despite recent improvements in water quality for some NZ rivers, these legacy nutrients and
39 continued agricultural intensification are expected to pose broad-scale environmental problems
40 for decades to come.

41

42 1. Introduction

43 River water quality reflects multiple activities and processes within its catchment,
44 including geomorphic processes, vegetation characteristics, climate, and anthropogenic land uses
45 (Brierley, 2010). Relationships between water quality and these catchment characteristics are not
46 straightforward because all of these factors interact over both space and time. For example, if
47 intensive livestock grazing occurs on steep slopes, surface runoff and consequently river

48 turbidity is expected to be greater than if grazing occurs on flatter areas. Or if fertilizers are
49 heavily applied to sandy soils with high drainage density, rivers will likely become eutrophied
50 over a period of decades due to legacy nutrients slowly leaking to the rivers through groundwater
51 (McDowell et al., 2008). The influence of land use on water quality has also been shown to vary
52 among different climates (Larned et al., 2004). With all of the various types of intensive land
53 uses that have occurred across diverse landscapes over hundreds of years, rivers with degraded
54 water quality are now widespread.

55 Historically, water quality in rivers was managed to meet minimally acceptable standards
56 or maximum pollutant load limits (Baron et al., 2002; Boesch, 2002; Howard-Williams et al.,
57 2010). However, in the last decade, a greater emphasis has been placed on maximizing the
58 ecosystem services provided by healthy rivers, which is driving efforts to further improve water
59 quality (Brauman et al., 2007; Davies-Colley, 2013). Early efforts in developed countries to
60 improve water quality focused on point-source pollution, particularly wastewater discharges
61 from factories and treatment plants (Campbell et al., 2004). While the broad-scale reduction in
62 point-source pollution elevated many water quality variables above minimal standards, most
63 rivers globally still have water quality impairments due to diffuse pollution – fine sediments,
64 nutrients, pathogens, toxicants, salts, and other contaminants that are delivered from unknown or
65 many indistinguishable sources across the catchment (Vorosmarty et al., 2010). Although
66 considerable effort has been directed at monitoring and reducing diffuse pollution with some
67 success, the legacy of pollutants from various land uses remains (Boesch, 2002; Kronvang et al.,
68 2008; Zobrist and Reichert, 2006). Agricultural land uses are by far the greatest contributors of
69 diffuse pollution, globally (Foley et al., 2005; Vitousek et al., 1997); however, the ‘intangible’

70 sources of diffuse pollution make it difficult to assign cause-and-effect relationships (Campbell
71 et al. 2004).

72 Many studies have used theoretical or numerical models to examine relationships
73 between land use and water quality because of the lack of consistent water quality monitoring
74 over long periods (bracketing land use change). While modelling approaches can be useful for
75 small catchments where much is known about its landscape, modelling may not work well for
76 larger catchments because land-water relationships are complex with interdependencies,
77 feedbacks, and legacy effects. Empirical studies can shed light on some of these complexities,
78 but they are only useful for their particular catchments and may have limited generality or
79 transferability. Comparisons of many diverse catchments is probably most useful to advance
80 understanding of broad-scale land-water relationships (Zobrist and Reichert, 2006).

81 One of the most comprehensive empirical multi-catchment studies to date on land use-
82 water quality relationships has been Varanka and Luoto's (2012) study of 32 boreal rivers in
83 Finland. They analyzed five water quality variables over ten years as a function of a suite of
84 physiographic, climate, and land use variables. A similar study was conducted on many of the
85 same rivers in Finland, but with a more sophisticated temporal analysis (Ekholm et al., 2015).
86 And several other studies have used this same river water quality dataset to investigate
87 environmental drivers. In a study of 11 Swiss watersheds, Zobrist and Reichert (2006) analyzed
88 export coefficients of six water quality variables from biweekly, flow proportional, composite
89 samples over a 24-year period within the context of land use.

90 All of these studies, and most catchment land use studies, assessed land use (or land use
91 change) as areal coverage. However, land use *intensity* – the inputs (e.g. fertilizer, livestock) and
92 activities (e.g. vegetation removal) of land use – could be a better predictor of environmental

93 impact for being a more direct measure of impact than areal coverage (Blüthgen et al., 2012;
94 Ramankutty et al., 2006). Unfortunately, our understanding of the patterns, processes, and
95 impacts of land use intensity is inadequate because of (1) its complex, multidimensional
96 interactions with other landscape variables, and (2) the lack of appropriate datasets across broad
97 spatiotemporal scales (Kuemmerle et al., 2013; Erb et al., 2016). New Zealand (NZ) provides a
98 valuable test-bed for the patterns, processes, and impacts of land use intensity because over the
99 past three decades pasture area has decreased but livestock densities and fertilizer inputs have
100 increased (MacLeod and Moller, 2006; StatsNZ, 2015). Like Finland and Switzerland, NZ has an
101 extensive long-term river water quality monitoring network, which has allowed many studies on
102 river water quality state and trends (Smith et al., 1996, 1997; Scarsbrook et al., 2003;
103 Scarsbrook, 2006; Ballantine and Davies-Colley, 2014) and effects of land use areal coverage
104 (Davies-Colley, 2013; Larned et al., 2004, 2016). However, this dataset has not been assessed as
105 regards changes in land use intensity that have occurred over the same period.

106 Here, we investigate long-term relationships among land use intensity, geomorphic
107 processes, and river water quality in NZ – which provides a particularly valuable case study
108 because: (1) it has had one of the highest rates of agricultural land intensification over recent
109 decades and thus serves as a potential indicator for countries that are also increasing agricultural
110 intensity; (2) it has a long, consistent, and comprehensive national water quality dataset; and (3)
111 it is physiographically-diverse. We examined monthly data for a suite of water quality variables
112 over a 26-year period for 77 diverse catchments. We then compared these states and trends of
113 river water quality to landscape data that characterized the catchments' geomorphology, soil
114 properties, and hydro-climatology; as well as temporal changes in land use areal coverage and
115 land use intensity, specifically livestock density and land disturbance, defined here as bare soil

116 resulting from vegetation loss. Altogether, these analyses reveal coincident spatiotemporal
117 patterns in land use intensity and water quality over a quarter of a century. Most of our analyses
118 were performed at the catchment scale which integrates the spatiotemporal changes that are
119 reflected in our water quality measurements, is the appropriate scale to analyze diffuse pollution,
120 and is the most appropriate spatial management unit (Howard-Williams et al., 2010).

121

122 2. Study area

123 New Zealand (*Aotearoa*, “Land of the long white cloud” in the language of indigenous
124 *Maori* people) is a small island nation (~268,000 km²) located between the South Pacific Ocean
125 to the east and the Tasman Sea to the west. Its two main islands (North Island and South Island)
126 are located between 34° and 47° S latitude. Being located on the active boundary between the
127 Australian and Pacific Plates, NZ’s geology and geomorphology are very diverse, including
128 active volcanoes, karst regions, a range of high fold mountains (the Southern Alps), large coastal
129 plains, and rolling hills across both hard- and soft-rocks. Being stretched latitudinally, with
130 nowhere more than about 150 km from the sea, between two major ocean waters combined with
131 its topographic variability, NZ also has a diverse climate with regional extremes, including sub-
132 tropical in the far north, temperate in the central North Island, extremely wet on the western side
133 of the Southern Alps (up to 10 m annually), and semi-arid in the rain shadow to the east of the
134 Southern Alps.

135 New Zealand is the last major habitable landmass to be settled by humans. Eastern
136 Polynesians first arrived around 1300 AD (Wilmshurst et al., 2008). Europeans first arrived in
137 the late-1700s, but large-scale settlement did not begin until the 1840s. Broad-scale agriculture
138 spread shortly after and has been intensifying since. While we address land use changes at the

139 national scale in this study, our water quality analyses focus on 77 diverse catchments across NZ
140 (Fig. 1), which cumulatively cover about half of NZ's land area.

141

142 3. Methods

143 3.1. Water quality data

144 Water quality data was obtained from NZ's National Rivers Water Quality Network
145 (NRWQN), which is operated and maintained by the National Institute of Water & Atmospheric
146 Research (NIWA). This network represents one of the world's most comprehensive river water
147 quality datasets: thirteen water quality and two biomonitoring variables have been measured
148 monthly (via in situ measurements and grab samples), with supporting flow estimation, from
149 1989-2014 at 77 sites whose catchments cumulatively drain approximately half of New
150 Zealand's land surface (Davies-Colley et al., 2011). Further, this dataset has been operationally
151 stable throughout its history, which allows us to calculate trends over this period. For this study,
152 we focused on eleven water quality variables and their coincident flow (Table 1). We did not
153 analyze ammoniacal nitrogen (NH_4) because early NH_4 samples were biased high by laboratory
154 contamination (Davies-Colley et al., 2011).

155 All water quality variables, except water temperature (T_w), were flow-normalized (for
156 each site separately) in JMP® Pro (v 11.2.1) with local polynomial regression (LOESS) using a
157 quadratic fit, a tri-cube weighting function, a smoothing window (alpha) of 0.67, and a four-pass
158 robustness to minimize the weights of outliers (Cleveland and Devlin, 1988); where, flow-
159 adjusted value = raw value – LOESS value + median value. With LOESS, there is no assumption
160 about the water quality variable's relationship with flow. For example, although visual clarity
161 usually decreases systematically with increasing flow (Smith et al., 1997), algae blooms at low

162 flows can sometimes reduce clarity. LOESS also allowed us to examine relative water quality
163 changes over long periods.

164

165 3.2. Physiographic data

166 Water quality metrics and trends were compared to a suite of landscape variables (Table
167 2). Catchment morphometrics (area, slope, ruggedness) were obtained from a 30-m digital
168 elevation model (DEM) that we rescaled (in order to align with other gridded spatial datasets)
169 from the 25-m DEM produced by Landcare Research. This 25-m DEM was interpolated from
170 20-m contours of the national TOPOBASE digital topographic dataset supplied by Land
171 Information New Zealand (LINZ; scale: 1:50,000). Catchment area (A) is the drainage area (in
172 km^2) above the NRWQN station, derived using Arc Hydro tools in ArcGIS 9.3.1 in combination
173 with the River Environment Classification (REC, v2.0), the national hydrography dataset derived
174 from a 30-m hydrologically correct DEM (Snelder et al., 2010). Mean catchment slope (S_c) was
175 derived from the same software package, using a 3x3 cell window. We defined ruggedness (R_r)
176 as the standard deviation of the 30-m slope grid for each catchment (*sensu* Grohmann et al.,
177 2011). Drainage density (D_d) was calculated from the ratio of the total length of REC streams to
178 catchment area (in km/km^2).

179 Soils data was obtained from the 1:50,000 Fundamental Soils Layers (FSL), which is
180 maintained by Landcare Research. Methods and data descriptions for this soils database are
181 described in Webb and Wilson (1995) and Newsome et al. (2008). Catchment-scale soil
182 variables (mean value across catchment) that we included in our analysis for being expected to
183 be related to water quality were: soil depth (Z_s), percent of catchment dominated by silty and
184 clayey surface soils ($SC\%$), soil pH (pH_s), cation exchange capacity (CEC), organic matter

185 percentage ($OM\%$), and phosphate retention (P_{ret}). Phosphate retention is a measure (in %) of the
186 amount of phosphate that is removed from solution by the soil via sorption (Saunders, 1965).
187 Thus, soils with high P_{ret} have low P-availability for plant growth.

188 Median annual precipitation (MAP), median annual temperature (MAT), and median
189 annual sunshine (MAS) averaged across each catchment was obtained from NIWA's National
190 Climate Database, which contained 5-km gridded daily weather data (Tait and Turner, 2005).
191 Our values for these three variables represent the median annual precipitation (total mm/y),
192 temperature (mean °C), and sunshine (hours/y) for the period 1981-2010. Relative water storage
193 (RWS) was calculated as the proportion of the annual catchment water yield (i.e. total volume of
194 water leaving the catchment in a year) stored in lakes and reservoirs. Reservoir/lake storage was
195 obtained from the Freshwater Ecosystems of New Zealand (FENZ) Database, described in
196 Snelder (2006). The last hydro-climatological variable we included in our analyses was the
197 median discharge (Q_{50}), which was calculated from the NRWQN 'flow stamping' at times of
198 water quality sampling from 1989-2014.

199

200 3.3. Land use areal coverage, intensity, and disturbance data

201 There are two national land use datasets for New Zealand. The Land-Use and Carbon
202 Analysis System (LUCAS) was developed by the NZ Ministry for the Environment (MfE, 2012)
203 for reporting and accounting of carbon fluxes and greenhouse gas emissions, as required by the
204 United Nations Framework on Climate Change and the Kyoto Protocol. Accordingly, LUCAS
205 uses 1990 as its reference year and maps land use in 12 classes for 2008 and 2012. The Land
206 Cover Database (LCDB) was developed by Landcare Research (LCR), with contributions from
207 MfE, Department of Conservation (DOC), Ministry for Primary Industries (MPI), and Regional

208 Councils (LCR, 2015). LCDB contains 35 land use classes for 1996, 2001, 2008, and 2012. Both
209 datasets use a minimum mapping area of 1 hectare, and use many of the same data and methods
210 to map land use. There are however, some key differences in their class designations and
211 classifications that are important to our analyses: (1) LUCAS includes Manuka/Kanuka as forest,
212 whereas LCDB designates Manuka/Kanuka as shrub; (2) LUCAS lumps all post-1989 forests
213 into one class, whereas LCDB differentiates between indigenous and plantation forests; (3)
214 LUCAS uses a conservative approach to mapping high-producing grasslands, whereas LCDB
215 uses phenological information to provide more accurate estimations of high-producing grassland.
216 Because of our focus on (water quality-impacting) plantation forests and high-producing
217 grasslands, we used the LCDB (v4.1) for our spatial and statistical analyses. We used LUCAS
218 only to quantify long-term changes from 1990 to 2012, before the LCDB was initiated in 1996.
219 Table 3 describes the land use classes we used in this research, which classes are included from
220 both datasets, and the national comparison between LUCAS and LCDB for 2012.

221 There are numerous metrics for land use intensity (Erb et al., 2013). At the catchment-
222 scale, we used livestock density as a metric for all grasslands; and we used land disturbance,
223 defined here as bare soil resulting from vegetation loss, as a metric for high-producing grasslands
224 and plantation forests. We also used national-scale annual fertilizer data (1989-2014) from
225 StatsNZ (2015) to compare long-term trends of river nutrient concentrations to nutrient inputs.
226 Livestock numbers for dairy cattle, beef cattle, sheep, and deer (at 1 ha resolution) for each
227 catchment were derived from maps provided by Ausseil et al. (2013), which is representative for
228 the year 2011. To assess total livestock impact on land disturbance, we multiplied each livestock
229 type by its AgriBase stock unit (SU) coefficient: sheep = 0.95 SU, deer = 1.9 SU, beef cattle =

230 5.3 SU, and dairy cattle = 6.65 SU (Woods et al., 2006). The total SU for each catchment was
231 then normalized by total catchment area, expressed as stock unit density (*SUD*) in SU/ha.

232 Changes in *SUD* from 1990 to 2012 (*SUD*₂₀₁₂₋₁₉₉₀) were assessed using district-level data
233 from StatsNZ (2015) on total numbers of sheep, deer, beef cattle, and dairy cattle. These
234 livestock numbers were then aggregated for each catchment and multiplied by their respective
235 SU coefficient. Stock units per hectare were then compared between 1990 and 2012 to assess
236 change in livestock impacts in each catchment. For Whakatane and Kawerau Districts, 1993 was
237 used because 1990 data was unavailable.

238 Land disturbance (i.e. bare soil resulting from vegetation loss) was quantified for all
239 high-producing grasslands (*D_{HG}*) and plantation forests (*D_{PF}*), as well as the whole catchment
240 (*D_C*) for the period 2000 - 2013. The methods for calculating and validating disturbance are
241 described in de Beurs et al. (2016). Briefly, MODIS BRDF corrected reflectance data
242 (MCD43A4) at 463 m spatial resolution and eight day temporal resolution was used to calculate
243 Tasseled Cap brightness, greenness and wetness based on the coefficients following Lobser and
244 Cohen (2007). These indices consist of linear combinations of all seven MODIS reflectance
245 bands to represent general image brightness which is comparable to albedo, image greenness
246 which is comparable to the better known vegetation indices such as NDVI and EVI, and image
247 wetness which is linked to the amount of water captured in the vegetation, most comparable to
248 normalized difference water indices. Missing pixels were ignored. We then calculated the mean
249 and standard deviation of each tasseled cap index for each combination of land cover class (LCR,
250 2015) and climatic region for each 8-day time period. We then used these measures to
251 standardize the calculated tasseled cap indices. To determine how disturbed each pixel was at
252 any point in time, we then calculated the forest and grassland disturbances. The forest

253 disturbance index is calculated as the standardized brightness minus the standardized greenness
254 and wetness. The idea is that disturbed forests appear brighter and less green and less wet than
255 undisturbed forests. The grassland index is the negative sum of all indices, indicating that
256 disturbed grasslands appear darker, less green and less wet than undisturbed grasslands. MODIS
257 disturbance data were visually validated against 7500 random pixels from Landsat imagery and
258 corresponding 15 high resolution Orbview-3 and Ikonos images. The overall accuracy of the
259 disturbance index based on Landsat data was 98%.

260

261 3.4. Statistical methods

262 We used nonparametric Spearman rank correlation coefficients (r_s) to look at
263 relationships between variables because many of the relationships were curvilinear. Statistical
264 significance was taken to be an alpha of 0.05. Bivariate comparisons between all variables
265 (Tables 1-3) were performed to explore for associations and identify correlated variables before
266 later multivariate analyses. Median values (from the 26-y monthly time-series) for water quality
267 variables at each site were used when compared to physiographic and land use variables of their
268 corresponding catchment. Stepwise regression was then used to rank-order the relative
269 contributions of multiple landscape variables associated with each major water quality variable.
270 Stepwise regression was used because it accounts for correlations among the independent
271 landscape variables. The order of variables in the stepwise regression model and the sign of their
272 coefficient (proportional [+] vs. inverse [-]) provides an objective measure of the contribution of
273 each landscape variable to river water quality. The level of entry into the model was set to $p =$
274 0.05. All the above statistical analyses were performed in JMP® Pro (v 11.2.1).

275 Temporal trends in water quality (1989 – 2014) and disturbance (2000 – 2013) data
276 were assessed with the seasonal Kendall test which was corrected for temporal autocorrelation
277 using the rkt R package; missing values were ignored. We also calculated the Seasonal Kendall
278 slope estimators (SKSE) using the same R package. Because some NRWQN sites had multiple
279 measurements in some months, a few records (no more than five) were removed from each site
280 in order to ensure 12 monthly values for each year for the SKSE test. There were also occasional
281 missing values for some variables throughout the time-series, particularly in the early years. Of
282 particular note, there were no *TN* values for 1994 as a result of contamination by leaking
283 ammonia refrigerant during storage of frozen subsamples. HV1 did not have data for 18 months
284 from 2012-2014.

285 In order to make trend comparisons among sites and derive an estimate of percent change
286 per year, we normalized SKSE values by dividing them by the raw data median to give the
287 relative SKSE (RSKSE) in percent change per year (Smith et al., 1996). Given that water
288 temperature (T_w) uses an arbitrary scale in °C, we only report SKSE values for this variable. We
289 also used the trend categories of Scarsbrook (2006): (1) no significant trend – the null hypothesis
290 for the Seasonal Kendall test was not rejected ($p > 0.05$); (2) significant increase/decrease – the
291 null hypothesis for the Seasonal Kendall test was rejected ($p < 0.05$); and (3) ‘meaningful’
292 increase/decrease – the trend was significant and the magnitude of the trend (RSKSE) was
293 greater than 1% per year. According to Ballantine and Davies-Colley (2014), a 1% change per
294 year translates to slightly more than 10% change per decade (due to compounding), a rate of
295 change that is easily detectable and observable.

296

297 4. Results

298 4.1. Physiographic characteristics

299 The 77 NRWQN catchments were physiographically diverse in terms of morphometric,
300 soil, and hydro-climatological variables (Table 4; Supplement Table 1). Most notable with
301 regards to its direct influence on runoff and water quality was median annual precipitation
302 (*MAP*), which ranged from 533 to 7,044 mm/y. When combined with the wide range of
303 catchment areas (*A*), median discharge (Q_{50}) varied over three orders of magnitude, from 0.4 to
304 515 m³/s, and annual water yield from 103 to 3,475 mm/y. In terms of soil, about a quarter of the
305 catchments had very sandy surface soils ($SC\% < 10$) and a quarter had fine-textured soils ($SC\%$
306 > 70). Phosphate retention (P_{ret}), an important variable for fertilizer management and
307 consequently water quality, was particularly high ($>57\%$; 10th percentile) for catchments HM2,
308 HM5, HM6, WA1, WA2, WA3, and WN5.

309 Several physiographic variables (Table 2) displayed strong latitudinal trends from North
310 to South and many were strongly correlated ($p < 0.001$; Supplement Fig. 1). In consideration of
311 these relationships and perceived importance for water quality (*sensu* Varanka and Luoto, 2012),
312 we used the following subset of minimally correlated physiographic variables for subsequent
313 multivariate analyses: catchment slope (S_c), silt-clay percentage ($SC\%$), phosphate retention
314 (P_{ret}), and median flow (Q_{50}).

315

316 4.2. Land use areal coverage and temporal changes

317 Land use in NZ, like physiography, varied widely; and our 77 catchments captured this
318 diversity (Fig. 1; Supplement Table 2). Thirteen catchments were dominated ($>50\%$) by non-
319 plantation forests (*NF*), with one (WN2) containing more than 94%. Thirteen other catchments
320 were dominated by shrub/grassland (*SG*) that was not intensively managed. The most dominant

321 land use was grasslands that were intensively managed (hereafter high-producing grasslands;
322 *HG*), covering the majority of the area for 31 catchments. Together, these three land uses made
323 up 84% of the catchments' areas. Plantation forest (*PF*) was the majority land use for three
324 catchments (RO3, RO5, RO2). Open water (*OW*) was the majority land use for one catchment
325 (RO1) and relatively high (>10%) for two others (RO6, DN10). Barren/other (*BO*), which was
326 largely bare rock, was relatively high (>10%) for 13 mountainous catchments. Urban (*UR*)
327 coverage rarely exceeded 1%, with only one catchment greater than 2% (WN1). Annual cropland
328 (*AC*) exceeded 1% in 11 catchments, but never exceeded 8%. Vegetated wetland (*VW*) and
329 perennial cropland (*PC*) were minimal in all catchments, each rarely exceeding 1%.

330 In general, *NF*, *SG* and *BO* areas dominated mountainous catchments with high S_c and
331 low Z_s ; while *HG* dominated most lowland catchments with low S_c , high Z_s , and high pH_s . Like
332 *HG*, *PF* mostly occurred on flat areas ($r_s = -0.48$ with S_c) with thick soils (0.35 with Z_s) that
333 were less acidic (0.31 with pH_s). Given the relative dominance of catchment land use,
334 relationships with physiographic variables, and potential effects on water quality in NZ rivers
335 (Davies-Colley, 2013; Howard-Williams et al., 2010), the land use variables used for subsequent
336 multivariate analyses were *NF*, *SG*, *HG*, *PF*, and *OW*.

337 Land use areal coverage did not change much from 1990 to 2012 across NZ (Fig. 2) or in
338 many catchments (Supplement Table 2). The greatest change was a 13.4% increase in *PF* in
339 GS1, which was almost entirely accounted for by a 13% decrease in *SG*. Thirteen other
340 catchments experienced small increases (3.0 - 6.6%) in *PF*, accounted for by decreases in *SG* or
341 *HG* or both. HM3 and HM4 had the greatest increases in *HG* at 3.4% and 2.0%, respectively.
342 High-producing grasslands (*HG*) for the other 75 catchments remained virtually unchanged (<

343 0.4%) or decreased. WH3 had the greatest decrease in *HG* at -4.8%. Land use areal coverage
344 change in other catchments was negligible.

345

346 4.3. Land use intensity and temporal changes

347 Changes in total stock unit density between 1990 and 2012 ($SUD_{2012-1990}$) were also
348 minor with only two catchments (AK1 and AK2: both -5.1 SU/ha owing to urban fringe
349 expansion) changing more than 1.6 SU/ha over this period (Supplement Table 3). Temporal
350 changes in $SUD_{2012-1990}$ for 56 of the 77 catchments were within the range of -1.0 to 1.0 SU/ha.
351 Although land use areal coverage and total livestock densities changed little 1990-2012,
352 livestock *types* changed considerably for many catchments (Supplement Table 3) and across NZ
353 (Fig. 2). The general pattern was dairy cattle replacing sheep. The number of dairy cattle from
354 1990 to 2012 increased in 72 catchments, with a mean increase of 0.6 SU/ha for all catchments;
355 while the number of sheep decreased in all 77 catchments (mean = -0.9 SU/ha). Deer and beef
356 cattle numbers changed little: 0.0 and -0.2 SU/ha, respectively.

357 When 2011 livestock densities were compared with physiographic variables, the
358 strongest relationships were found with combined SUD of dairy and beef cattle (hereafter
359 SUD_{cattle} ; Supplement Fig. 2). SUD_{cattle} decreased strongly with increasing slope, S_c ($r_s = -0.79$),
360 but increased with Z_s (0.43), pH_s (0.32), and P_{ret} (0.27). SUD_{cattle} also increased with MAT
361 (0.68) and MAS (0.42), but decreased with MAP (-0.34). Thus, highest cattle densities were
362 found in catchments such as WA3 (with the highest SUD_{cattle} at 15.7 SU/ha) that were relatively
363 flat, warm, sunny, and dry, with deep soils that had relatively high pH and high P-retention.
364 High-producing grasslands (*HG*) had similar, but less strong, correlations with these same
365 physiographic variables.

366 Catchment disturbance (D_C) varied widely over both space and time between 2000 and
367 2013 (Supplement Table 4). The maximum amount of D_C at one time was 35.7% for WN3 on
368 07-Apr-2003, almost entirely due to bare pastures. D_C exceeded 15% on six other occasions (264
369 days in total) in this catchment. In general, the North Island (Fig. 3) had a greater extent and
370 intensity of disturbance than the South Island (Fig. 4). The most intense disturbances occurred as
371 a result of plantation forest harvests, and these disturbances were on average visible for about 1.5
372 y up to about 4 y, with exceptions lasting more than 6 y. Indeed, D_C was strongly correlated to
373 PF coverage ($r_s = 0.51$). The catchment with the highest median D_C (10.5%) was RO3, which
374 had 69.8% of its catchment in PF and 17.7% in HG . Fourteen other catchments had D_C above
375 5%, and two-thirds of these were dominated by either PF or HG .

376 We also analyzed disturbance of plantation forests (D_{PF}) and high-producing grasslands
377 (D_{HG}) separately for each catchment. For catchments with at least 21.4-km² (100 MODIS pixels,
378 for the sake of statistical robustness) of plantation forest, the mean (\pm SD) D_{PF} (from 2000 to
379 2013) was $10.6 \pm 5.6\%$. The catchments with the highest D_{PF} were those with low mean annual
380 precipitation, MAP ($r_s = -0.42$). There were no significant relationships between D_{PF} and any of
381 the other physiographic variables. For catchments with at least 21.4-km² of high-producing
382 grasslands, the mean (\pm SD) D_{HG} was $6.0 \pm 6.4\%$. The catchments with the highest D_{HG} were
383 those with low mean annual sunshine (MAS ; $r_s = -0.25$), low mean annual temperature (MAT ; -
384 0.30), high catchment slope (S_c ; 0.25), and high ruggedness (R_r ; 0.31). The six catchments with
385 the highest D_{HG} (>15%) all had low phosphate retention (P_{ret} ; <32%). While it is assumed that
386 greater densities of livestock lead to greater pasture disturbance, we did not find a proportional
387 relationship between stock unit density (SUD) and D_{HG} among catchments. In fact, the highest
388 median D_{HG} was found for catchments with low SUD ($r_s = -0.45$). Over time however, we

389 observed a fairly strong trend ($r_s = 0.50$) of lower D_{HG} with decreasing SUD ($-SUD_{2012-1990}$). In
390 all there were seven catchments with significant or meaningful decreases in D_{HG} from 2000 to
391 2013 (assessed with Seasonal Kendall slope; SKSE), all of which had a negative $SUD_{2012-1990}$.

392

393 4.4. Water quality characteristics and trends

394 4.4.1. Catchment characteristics

395 Median monthly values of water quality variables for the 77 catchments ranged widely
396 (Table 5; Supplement Table 5). Some rivers had exceptional water quality all around, while
397 others had either current issues with multiple variables or worsening temporal trends (assessed
398 with SKSE from 1989 to 2014; Table 6). Because of the dependence of water quality on flow,
399 we first assessed temporal trends in Q . Only two catchments had significant increases in Q
400 (AX4, WH4), with the latter also being ‘meaningful.’ Three catchments had significant decreases
401 in Q (HM3, HM5, TU2) and five others also had ‘meaningful’ decreases in Q (CH2, GY4, HM4,
402 RO3, RO4).

403 Water temperatures (T_w) were not particularly high for any of the catchments; however,
404 21 rivers had significant increases in T_w , possibly the signature of climate change. The highest
405 rates of T_w increase ($0.04^\circ\text{C}/\text{y} < \text{SKSE} < 0.08^\circ\text{C}/\text{y}$) were for large alpine rivers in the central
406 South Island covered mostly by shrub/grasslands (TK3, TK4, TK6, AX3). Because of its strong
407 latitudinal trend (stronger than any land use effect), T_w was not analyzed further. Dissolved
408 oxygen (DO) was close to 100% for most catchments, but was particularly low (<90%) for two
409 catchments: RO2 which was affected by discharge from a large pulp mill at Kawerau, and AK2
410 which is on the Auckland fringe and thus affected by various peri-urban activities. DO was very
411 high (>110%) for one catchment (HV2) due to supersaturation from high periphyton in this

412 nutrient-enriched river. Temporal trends in *DO* from 1989 to 2014 were relatively minor
413 (RSKSE < 0.5%/y), except RO2 which had a significant increase (RSKSE = 0.7%/y) attributable
414 to progressive improvements in treatment of organic waste from its large pulp mill. Conductivity
415 (*COND*) was relatively low (<115 $\mu\text{S}/\text{cm}$) for all South Island catchments and varied
416 considerably for the North Island (54-528 $\mu\text{S}/\text{cm}$). Most catchments (52/77) experienced
417 significant or ‘meaningful’ increases in *COND* from 1989 to 2014. Water pH (pH_w) was neutral
418 to alkaline for all rivers, which have been described as calcium-sodium bicarbonate waters by
419 Close and Davies-Colley (1990), and only displayed minor changes (RSKSE < $\pm 0.1\%/y$) over
420 the 26-year study period.

421 Median visual water clarity (*CLAR*) was exceptionally high (>5 m) for seven catchments
422 and very low (<1 m) for 22 catchments. Since 1989, *CLAR* improved in almost half of the rivers,
423 and worsened in 4 rivers (Table 6; Supplement Table 5). *TURB* was strongly inversely
424 proportional to *CLAR* ($r_s = -0.97$) and generally followed opposite trends of *CLAR*. However,
425 fewer of its trends were significant and it had a disproportionately large number of ‘meaningful’
426 increases (17 catchments compared to only 2 ‘meaningful’ decreases in *CLAR*). *CDOM* was low
427 for most of the rivers, with only five catchments greater than 2.0 m^{-1} . Nineteen of the catchments
428 experienced significant or ‘meaningful’ decreases in *CDOM* since 1989, possibly due to the loss
429 of wetlands across NZ. Only one catchment had a ‘meaningful’ increase in *CDOM* (TK3).

430 Total nitrogen (*TN*) was relatively high (>455 mg/m^3) for almost a third of the
431 catchments, with the vast majority (17/23) of these being lowland catchments (<150 m in
432 elevation). Most of these catchments also had relatively high *NO_x*. Thirty-three catchments had
433 significant or ‘meaningful’ increases in *TN* from 1989 to 2014, while only five had significant or
434 ‘meaningful’ decreases in *TN* (Table 6). *NO_x* had a similar number of increasing temporal trends,

435 but also had ‘meaningful’ decreases for 12 catchments. Total phosphorus (*TP*) followed a similar
436 geographical pattern as *TN*. Eighteen of the 23 catchments with relatively high *TP* (>30 mg/m³)
437 were lowland catchments. Most of the catchments with relatively high *TP* (18/23) also had
438 relatively high *DRP* (>9.5 mg/m³). Seventeen catchments had ‘meaningful’ increases in *DRP*,
439 compared to only three with ‘meaningful’ decreases. There was more of a balance in temporal
440 trends of *TP*, with eight ‘meaningful’ increases and seven ‘meaningful’ decreases.

441 In addition to the expected correlations between *CLAR* and *TURB*, and among the
442 nitrogen and phosphorus constituents, several other significant relationships existed among the
443 water quality variables (Supplement Fig. 3). Taking into consideration this broad
444 multicollinearity, we focus our multivariate analyses on several key water quality variables,
445 particularly those that experienced the most changes from 1989 to 2014 (Table 6): *CLAR*, *TN*,
446 *NO_x*, *TP*, and *DRP*.

447

448 4.5. Water Quality relationships with physiography, land use, and disturbance

449 Visual water clarity (*CLAR*) generally decreased with *A* (-0.37; all following parentheses
450 in this section are *r_s* unless specified). Except for *TURB* (0.32), no other water quality variables
451 had significant relationships with catchment area. Several water quality variables correlated with
452 catchment slope (*S_c*), including: *TN* (-0.72), *TP* (-0.63), and *DRP* (-0.65), meaning N and P
453 concentrations were relatively high in lowland (low slope) catchments. *DRP* (0.65) and *TP* (0.61)
454 were directly proportional to mean annual temperature (*MAT*), but this association probably
455 arises because the highest phosphorus values occurred mainly in lowland catchments and some
456 of the northernmost catchments, temperature being strongly correlated with altitude and latitude.

457 *DRP* also had a (counterintuitive) significant relationship with soil phosphate retention, P_{ret}
458 (0.35). No other strong physiographic relationships emerged from our analyses.

459 The strongest relationships between water quality and land use areal coverage (Table 7)
460 included high-producing grasslands (*HG*), which had strong positive relationships with several
461 water quality variables except *CLAR* which decreased as *HG* increased. The lesser-managed
462 shrub/grasslands (*SG*) had generally opposite relationships with water quality, but note that *SG*
463 did not have significant relationships with *TURB* or *CLAR*. Non-plantation forest (*NF*) followed
464 the same trends as *SG*, but had fewer significant relationships with water quality. Plantation
465 forest (*PF*), on the other hand, followed the same trends as *HG*, with poorer water quality being
466 associated with greater coverage of *PF*; although correlations were not as strong as *HG*. *CDOM*,
467 *DRP*, and all N-constituents had significant negative correlations with open water (*OW*),
468 meaning that water quality improved with greater *OW* coverage, plausibly due to entrapment of
469 fine sediment and nutrients.

470 Water quality was significantly correlated with all stock unit density (SUD) metrics
471 (Table 7; Supplement Fig. 4), except deer (SUD_{de}) which only had relatively weak relationships
472 with *TN* and NO_x . The nutrients and CDOM had the strongest correlations with SUD_{cattle} , which
473 includes both dairy and beef cattle. *COND*, *CLAR*, and *TURB* had the strongest (slightly)
474 correlations with SUD_{be} . Overall, degraded water quality was strongly associated with high
475 livestock densities, even stronger than areal coverage of high-producing grasslands.

476 No significant correlations between water quality and total catchment disturbance (D_C)
477 were found; however, there were significant associations when disturbance was isolated by high-
478 producing grasslands (D_{HG}) and plantation forest (D_{PF} ; Table 7). Unexpectedly, *CLAR* and
479 *TURB* were not correlated to D_{HG} , and surprisingly, the rest of the water quality variables had a

480 significant *inverse* relationship with D_{HG} . Conversely, $CLAR$ was the only water quality variable
481 correlated to plantation forest disturbance, D_{PF} ($r_s = -0.27$). Some interesting results emerged
482 when temporal trends in water quality (via SKSE) were assessed for catchments with high
483 disturbance. Of the 15 catchments with D_c greater than 5%, six had ‘meaningful’ increases in
484 $TURB$ (RO3, HM4, RO6, WA6, HV6, HM2; all in North Island); while only one (HV5) had a
485 ‘meaningful’ decrease in $TURB$. Most of these 15 catchments also experienced significant
486 increases in TN (9 catchments; 7/9 also ‘meaningful’) and NO_x (10 catchments; 8/10 also
487 ‘meaningful’). Interestingly, TP and DRP significantly increased in only two of these highly
488 disturbed catchments.

489

490 4.6. Multivariate water quality relationships

491 In order to build on the above correlation analyses, the water quality variables of $CLAR$,
492 TN , NO_x , TP , and DRP were each assessed in a multivariate stepwise regression, using the
493 following ten physiographic and land use independent variables: S_c , $SC\%$, P_{ret} , Q_{50} , NF , SG ,
494 HG , PF , OW , and SUD_{cattle} (Table 8). The residual plots for all five water quality variables met
495 the assumptions of normality and linearity, but displayed heteroscedasticity with wide scatter for
496 high values. $CLAR$ was correlated to $-HG$, followed by $+OW$, $-Q_{50}$, and $-PF$, where signs
497 represent whether the relationship is positive (+) or inverse (-). Thus, water clarity was
498 predictably lower for larger rivers that drain larger areas of high-producing grasslands and/or
499 plantation forests, but improved with increased open water coverage (Fig. 5).

500 The combined stock unit density for beef and dairy cattle (SUD_{cattle}) was the primary
501 predictor for all four nutrient variables, with TN , TP , and DRP also being proportional to
502 plantation forest coverage (PF ; Table 8). Dissolved oxidized nitrogen (NO_x) was not

503 proportional to *PF*, or any other independent variable in the stepwise regression. Coverage of
504 high-producing grasslands (*HG*) and silt-clay surface soils (*SC%*) were also proportional factors
505 for TN. Whether intensity or areal coverage, land use was the primary and secondary predictor
506 for all five water quality variables (Fig. 5).

507

508 5. Discussion

509 5.1. River water quality states and trends

510 We characterized water quality states and trends for 77 river sites across New Zealand
511 (NZ) using a wide range of flows and water quality conditions for each site, including some
512 small floods. We acknowledge that our analyses did not fully capture large floods due to their
513 short durations, unlikelihood of occurring during the preset monthly sampling, and the fact that
514 we relied on grab samples. These episodic floods are particularly important for the water quality
515 of downstream waters such as lakes and estuaries (Stamm et al., 2014). The uncertainty
516 surrounding our lack of flood samples could have been mitigated by composite samples or
517 supplemental flood samples; however, our 26 years of monthly samples for each site ($n = 312$)
518 did allow us to confidently report median conditions and temporal trends in water quality
519 (Moosmann et al., 2014).

520 There was a wide range of water quality across NZ rivers (Table 5), with drastic
521 differences between upland and lowland rivers, distinguished by the 150 m elevation threshold.
522 For example, visual water clarity (*CLAR*), which is often used as a ‘master variable’ for overall
523 water quality (Davies-Colley et al., 2003; Julian et al., 2008), was high for upland rivers (mean =
524 3.2 m), with only two [alpine glacial flour-affected] rivers below the ANZECC (2000) guideline
525 of 0.6 m (CH3, AX3). Many of the upland rivers (7/33) had very high water clarity (> 5 m),

526 including one of the clearest non-lake-fed rivers in the world – Motueka River (NN2) with a
527 median *CLAR* of 9.8 m. The lowland rivers, in contrast, had a mean *CLAR* of 1.2 m, with 17
528 (39%) below the ANZECC guideline of 0.8 m. Note that these ANZECC (2000) guidelines,
529 which are statistical derivations (i.e. 20th-percentile of the first decade of the NRWQN record for
530 ‘reference’ sites), are merely ‘trigger values’ that when exceeded trigger a management response
531 to protect ecosystem health (Hart et al., 1999). Although these ‘trigger values’ are not effects-
532 based standards (which would be difficult to define for the wide variety of NZ ecosystems), they
533 do provide a useful reference for comparing water quality states and trends. Save for a few
534 borderline exceptions, the same sites that were below visual clarity guidelines also exceeded the
535 turbidity trigger values of 4.1 and 5.6 NTU for upland and lowland rivers, respectively.

536 Nine of the ten catchments with the highest TN (>740 mg/m³) were lowland catchments.
537 In all, 13 lowland catchments exceeded the ANZECC *TN* guideline of 614 mg/m³ and 8 upland
538 catchments exceeded the guideline of 295 mg/m³. Almost three quarters of these catchments
539 (15/21) also exceeded the *NO_x* guideline of 444 mg/m³ (lowland) and 167 mg/m³ (upland). There
540 were a similar number of sites exceeding ANZECC guidelines for *TP* (33/26 mg/m³ for
541 lowland/upland) and *DRP* (10/9 mg/m³ for lowland/upland), each with at least 20 and most of
542 these were corresponding. Our results on the state and trends of the 77 NRWQN catchments
543 generally accord with earlier NRWQN studies (e.g. Ballantine and Davies-Colley, 2014) and a
544 recent publication by Larned et al. (2016), which analyzed water quality states and trends for 461
545 NZ river sites for the period 2004-2013.

546 Based on ANZECC (2000) trigger values, we have organized the catchments into four
547 classes (Fig. 6): I. clean river with high visual water clarity (*CLAR*) and low dissolved inorganic
548 nutrients (*DIN*); II. sediment-impacted river with low *CLAR* and low *DIN*; III. nutrient-impacted

549 river with high *CLAR* and high DIN; and IV. sediment- and nutrient-impacted river with low
550 *CLAR* and high DIN. Note that the term ‘sediment-impacted’ is a connotation for total suspended
551 solids (TSS), which includes organic matter as well. In agriculture-dominated catchments, both
552 mineral sediment and particulate organic matter can greatly increase TSS (Julian et al., 2008).
553 We use *CLAR* as a preferred metric for suspended matter because TSS is not routinely measured
554 in the NRWQN (or other monitoring networks) while *CLAR* correlates strongly to TSS ($r = -$
555 0.92), and better than *TURB* ($r = 0.87$) (Ballantine et al., 2014). Further, *CDOM* in NZ rivers is
556 low with minimal impact on *CLAR*. We use NO_x as our preferred metric for DIN because it is
557 least affected by suspended sediment and soil properties (compared to *DRP*). However,
558 catchments that exceed ANZECC guidelines for *DRP* are indicated in Fig. 6 by grey-filled
559 markers.

560 When this classification is combined with the SKSE trend analyses (Table 6), we obtain a
561 clear picture of the current and potential state of NZ rivers (Fig. 6). Before individual rivers are
562 discussed, we first point out key differences between the upland and lowland catchments, which
563 will later be placed within the context of physiography and land use intensity. Most obvious, and
564 consistent with the findings of Larned et al. (2004), was that lowland rivers were much more
565 degraded, particularly by sediment. More than a third of the lowland catchments were either
566 Class II or IV (17/44); whereas, only two upland catchments were Class II. None of the upland
567 catchments were Class IV, and more than two-thirds were clean rivers (Class I). Both types had a
568 similar number of nutrient-impacted rivers (Class III). Particularly concerning is that almost half
569 of the lowland rivers (19/44) are currently experiencing ‘meaningful’ increases ($>1\%$ per year) in
570 NO_x , *DRP*, or both. The other striking trend is that many of the lowland rivers are becoming
571 clearer, with 18/44 experiencing ‘meaningful’ increases ($>1\%$ per year) in *CLAR* – which,

572 plausibly, has been attributed to increasing riparian fencing to exclude cattle from channels
573 (Davies-Colley, 2013; Ballantine and Davies-Colley, 2014; Larned et al., 2016).

574 While clearer rivers are seen as an improvement in water quality; when combined with
575 increasing nutrients, warmer water, and lower flows, the perfect recipe for toxic algae blooms is
576 created (Dodds and Welch, 2000; Hilton et al., 2006). Only recently has the widespread problem
577 of toxic algae blooms in NZ rivers been evidenced (Wood et al., 2015; McAllister et al., 2016),
578 and our results indicate that this problem could worsen given the increasing trends we found in
579 water temperatures, inorganic nutrients, and most influential in our opinion, water clarity.

580 Nutrient enrichment and global warming receive the most attention when it comes to degraded
581 water quality, but rivers have increasingly become light-limited (Hilton et al., 2006; Julian et al.,
582 2013) such that when clarity improves in warm, nutrient-rich rivers, algae can proliferate.

583 Particularly problematic for NZ is that its lowland catchments, which are warmer (mean median
584 T_w of 13.6 v 10.8 °C for upland rivers), have much greater *DRP* and *NO_x*, and have longer water
585 residence times, are the ones becoming appreciably clearer (Fig. 6). If droughts become more
586 frequent and intense in NZ, toxic algae blooms are also likely to become more frequent, more
587 widespread, and more problematic. However, this algae response is complex and depends on a
588 number of interacting factors such that the apparent potential for increasing algal nuisance might
589 not necessarily be realized in some rivers (Dodds and Welch, 2000; Hilton et al., 2006).

590

591 5.2. The role of physiography in dictating land use intensity across NZ

592 While physiography did not emerge as a significant independent variable in the
593 multivariate analyses (except *TN* with *SC%*), physiography is important because it largely
594 controls the location and intensity of agricultural land uses. The greatest coverages of high-

595 producing grasslands (HG) and the highest densities of cattle (SUD_{cattle}), the two primary
596 explanatory variables for all five major water quality variables (Table 8), were both found
597 predominantly in flat areas with deep soils located in warm, sunny, and relatively dry climates.
598 Livestock in NZ depend almost exclusively on pasture grasses and thus their productivity is
599 maximized when pasture productivity is maximized. The very large cattle are not well suited for
600 steep slopes, particularly dairy cattle which can weigh more than 500 kg. Deep soils are
601 important because they absorb and hold more water for plant uptake, and are not as susceptible
602 to waterlogging, especially in wetter climates. Year-round and intense grazing is best supported
603 by warm and sunny climates where pasture grasses are highly productive and recover quickly
604 following intense grazing such as strip/rotational grazing which is common in NZ dairy farms.

605 Another soil property we found to be positively correlated to SUD_{cattle} was phosphate
606 retention (P_{ret}). The highest dairy cow densities were found on Allophanic volcanic soils with
607 high P_{ret} , likely because these soils respond favorably to P-fertilizer and thus can be managed
608 more intensively. However, soils with high P_{ret} require more P-fertilizer, and thus generally have
609 higher export of DRP to rivers. Our finding of a significant positive correlation between these
610 two variables is consistent with this interpretation. Further, we found that high-producing
611 pastures with high P_{ret} had the lowest disturbance (D_{HG}), indicating that these intensively
612 managed pastures recover quickly following grazing. In a more comprehensive study of land
613 disturbance across the North Island of NZ, de Beurs et al. (2016) also found that Allophanic soils
614 had the least disturbance among all soil orders. Where high livestock densities occur in less than
615 ideal conditions, land disturbance is likely. Our catchment-scale analyses limit our interpretation
616 of specific situations, but based on our results, field observations and previous remote sensing
617 analyses, pasture disturbance in NZ will likely be highest during droughts on steep, south-facing

618 slopes with thin soils being heavily grazed by sheep. Under these conditions, grasses will be
619 grazed down to bare soil and recover very slowly.

620 Plantation forests (*PF*) in NZ also correlated with thick soils with relatively high P_{ret} on
621 flat areas, particularly the pumice soils of the central North Island. The porous nature of the
622 pumice soils allows them to efficiently hold and regulate nutrients, water, and air; while being
623 well-draining and resistant to compaction and flooding. Under these conditions, radiata pine (the
624 dominant *PF* species in NZ) grows rapidly (mean harvest cycle of 28 y) and can be harvested
625 year-round. Since 1990 however, many of the *PF* additions have occurred on steeper slopes in
626 response to carbon credit incentives, greater economic demand for wood products (PCE, 2013),
627 and the need for soil erosion control on steep pasture susceptible to land-sliding (Parkyn et al.,
628 2006).

629

630 5.3 Land use intensity and water quality in New Zealand rivers

631 5.3.1 High-producing pastures and livestock densities

632 High-producing grassland coverage (*HG*) was the primary explanatory variable for visual
633 clarity (*CLAR*; Table 8, Fig. 5). *CLAR* in NZ rivers is mostly influenced by mineral and organic
634 particulates (Davies-Colley et al., 2014). Livestock reduce visual clarity in multiple ways,
635 especially in NZ where high densities of multiple types of livestock tread year-round on
636 relatively steep slopes with highly erodible soils vegetated by shallow-root introduced grasses
637 which are susceptible to destabilization (McDowell et al., 2008). The year-round treading is
638 particularly important because most NZ regions during winter are very wet with short days,
639 which increases soil disturbance (pugging and compaction) and slows recovery times. Where

640 livestock have direct access to rivers, their trampling of riverbanks and instream disturbance is
641 often the main contributor to reduced *CLAR* (Trimble and Mendel, 1995; McDowell et al., 2008).

642 The lowland flatter areas in NZ have high *HG* coverage and high cattle stock densities
643 (*SUD_{cattle}*). These lowlands also have high drainage densities – often increased by artificial
644 drainage. The influence of *HG* on *CLAR* is thus exacerbated by this interaction of high *SUD_{cattle}*
645 and artificial drainage. Interestingly, *SUD_{cattle}* was not an explanatory variable for *CLAR* in the
646 stepwise regression, which is likely a result of two factors. First, *HG* and *SUD_{cattle}* are highly
647 correlated, and stepwise regression does not include secondary variables that are explaining the
648 same proportion of variance as the primary independent variable. Second, we found that *CLAR*
649 has actually *improved* in catchments where *SUD_{cattle}* is high and/or has increased (Fig. 6), which
650 we noted earlier could be a result of increased riparian fencing. In 2003, NZ implemented the
651 *Dairying and Clean Streams Accord*, which has led to the exclusion of dairy cattle from 87% (as
652 of 2012) of perennial rivers greater than 1 m in width (Bewsell et al., 2007; Howard-Williams et
653 al., 2010; Gunn and Rutherford, 2013). By excluding (dairy) cattle from channels and riparian
654 zones, the contribution of riverbank and bed erosion to degraded *CLAR* has likely been mitigated
655 and reduced over time (Trimble and Mendel, 1995; Hughes and Quinn, 2014). Indeed, *CLAR* has
656 been significantly and meaningfully improving in many of NZ's rivers (Table 6), even those with
657 increasing *SUD_{cattle}*, albeit from a fairly degraded condition. Of the 34 catchments with
658 significant increases in *CLAR*, all but 5 had increases in *SUD_{cattle}* from 1990 to 2012.

659 Another potential explanation for improved water clarity at numerous sites is the
660 considerable decrease in sheep density across the NZ landscape. NZ had 57.65 million sheep in
661 1990. By 2012, that number had been reduced by almost half, to 31.19 million (StatsNZ, 2015).
662 Although cattle are larger and have a greater treading impact per animal, the much greater

663 number of sheep means that stock unit density (SUD) may be broadly comparable as regards
664 environmental impact. Another difference is that sheep are generally placed on steeper, less
665 stable slopes in NZ, where headwater stream channels are located. Where there are breaks in
666 slope (even small ones), sheep create tracks of bare soil with their hooves and hillside scars with
667 their bodies (for scratching and shelter), both of which can enhance soil erosion (Evans, 1997).
668 Further, cattle (using their tongues) leave approximately half the grass height on the pasture after
669 grazing; whereas sheep (using their teeth) graze approximately 80% of grass height (down to
670 bare soil in dire conditions), leaving it exposed to erosion (Woodward, 1998). Considering all
671 these factors, sheep can have a greater impact on sediment runoff into rivers, and consequently
672 visual clarity, than suggested by their aversion to water *versus* cattle's attraction to water.
673 Although not isolated in our analyses, the particulate fractions of *TN* and *TP* have likely been
674 affected by similar processes as *CLAR* and may follow the same temporal trends (Ballantine and
675 Davies-Colley, 2014).

676 While *HG* was also strongly correlated to river nutrient concentrations (Table 7), the
677 primary explanatory variable for all four major nutrient metrics (Table 8, Fig. 5) was land use
678 intensity as measured by livestock density of beef and dairy cattle (SUD_{cattle}). The difference
679 between these two explanatory variables may seem trivial, however the distinction is important if
680 we want to understand future trends and effectiveness of water quality management strategies.
681 As we demonstrated, the area of land used for high-producing grasslands (*HG*) has not changed
682 much since 1990 (Fig. 2). In fact, it has decreased or stayed virtually the same in all but two of
683 the 77 catchments. Yet, nutrient concentrations have been increasing in many of the rivers (Table
684 6), which we attribute to (1) increasing numbers of cattle (mostly dairy) on both *HG* and *SG*, and
685 (2) legacy nutrients being slowly delivered to the rivers in groundwater. From 1990 to 2012, NZ

686 approximately doubled its number of dairy cattle, exceeding 6.4 million. (StatsNZ, 2015). This
687 enormous addition to a country that is only 268,000 km² in area, has been accompanied by more
688 than 1.426 million tonnes of P-based fertilizers and 335,000 tonnes of N-based fertilizers
689 annually (1990-2012 mean; StatsNZ, 2015). Of the nutrients consumed by lactating dairy cows,
690 approximately 79% of N and 66% of P are returned to the landscape in the form of urine and
691 feces (Monaghan et al., 2007). This results, potentially, in about 260,000 tonnes of N-based and
692 940,000 tonnes of P-based diffuse pollution. Some of these nutrients will be transported to rivers
693 during subsequent storms, but a majority will remain (building up) in the landscape to be slowly
694 added to rivers over decadal time-scales (Howard-Williams et al., 2010).

695

696 5.3.2. Plantation forests

697 All water quality variables were significantly correlated to plantation forest coverage
698 (*PF*; Table 7), with a negative relationship with *CLAR* but positive for all other variables. From
699 the stepwise regression, *PF* emerged as an explanatory variable for all major water quality
700 variables except *NO_x* (Table 8), suggesting that its dominant impact on river water quality was
701 from surface runoff. Plantation forestry activities can add a considerable amount of sediment and
702 nutrient pollution to rivers, especially during and immediately following harvesting (Fahey et al.,
703 2003; Croke and Hairsine, 2006; Davis, 2005). This harvesting period of maximum soil
704 disturbance usually lasts about two years (Fahey et al., 2003), but the land cover may remain
705 sparsely vegetated and susceptible to erosion for several years (but usually not more than 5 y; de
706 Beurs et al., 2016). The greatest *PF* impact on sediment runoff, and thus potentially *CLAR*, is
707 usually from road sidecast/runoff, shallow landslides, and channel scouring/gullyng (Fahey et
708 al., 2003; Motha et al., 2003; Fransen et al., 2001).

709 Rivers receive a pulse of nutrients during the forest harvest, but fertilizers are also
710 applied at time of re-planting and sometimes routinely to enhance growth (Davis, 2005). Radiata
711 pine in the pumice soils of the central North Island, the dominant area of *PF* in NZ, are
712 particularly responsive to both N- and P-fertilizers and thus likely receive ample supplements.
713 Like pasture fertilizers, some of these nutrients may be delivered to rivers during intense
714 precipitation, but there is also a legacy of nutrients left behind. Fertilizers have been applied to
715 plantation forests in NZ since the 1950s, with an intense period of application in the 1970s
716 (Davis, 2005). While fertilization rates (tonnes/ha/y) have decreased since 1980, the amount of
717 NO_x leaving catchments mostly covered in *PF* has significantly and ‘meaningfully’ increased
718 since 1989. None of these catchments had more than 17.7% *HG*, none had major increases in *HG*
719 ($< 0.3\%$), none had major increases in SUD_{cattle} (< 0.7 SU/ha), and none had a significant
720 increase in D_{PF} . What the catchments did have in common were all had gravelly/sandy pumice
721 soils (< 4.5 $SC\%$) and all were intensively managed as reported by Davis (2005) and as indicated
722 by high D_C ($> 6.8\%$). The extended periods of nonvegetated land due to weed control also
723 increases the amount of nutrients delivered to rivers over the long term (Davis, 2005).

724

725 5.3.3. Land disturbance and water quality

726 So far, we have discussed how land use affects water quality, with a focus on sediment
727 and nutrient runoff from high-producing grasslands (*HG*) and plantation forests (*PF*). When land
728 is disturbed (i.e. bare soil), sediment/nutrient mobilization can be enhanced. The most intense
729 and longest lasting disturbances occurred during plantation forest harvests. Following harvest,
730 we found that the land remained disturbed for 1-6 years, with a mean of 1.5 years. The overall
731 mean and median D_{PF} among all catchments was 10%, which means that plantation forestry

732 leaves large areas of disturbed land at any one time. When this bare land is exposed to intense
733 precipitation, large quantities of sediment and nutrients can be mobilized into the rivers. This
734 happened in the Motueka Catchment (NN1) in 2005 when a 50-y storm fell on some recently-
735 harvested plantation forests. For one of NN1's sub-catchments, the post-harvest disturbed land
736 caused a five-fold increase in sediment yield compared to pre-harvest events. Following this
737 event, sediment yields at NN1 were elevated by a factor of 2-3 over the next 3 years (Basher et
738 al., 2011). Similar sediment erosion events for plantation forests during the post-harvest
739 disturbance have been documented for other catchments across NZ (Hicks et al., 2000; Phillips et
740 al., 2005). Because these disturbances only last a few years, they typically do not show up as
741 temporal trends (via SKSE); however it is possible that they produce enough readily available
742 sediment to impact water quality for longer periods (Kamarinas et al., 2016).

743 The coincidence of rainstorms on disturbed pasture could have the same effect on
744 sediment/nutrient runoff if the pasture is connected to the stream network via steep slopes or
745 adjacent channels/canals (Dymond et al., 2010; Kamarinas et al., 2016). Pastures become
746 disturbed from overgrazing, strip grazing, pugging/soil compaction, tilling/reseeding,
747 cropping/harvesting, or landsliding on steep slopes. Given the high intensity of grazing
748 management in NZ, all of these are common. While D_{HG} was lower than D_{PF} on average, D_{HG}
749 had a higher maximum (Table 4). Spatiotemporal patterns in disturbance between these two land
750 uses were also different (de Beurs et al., 2016). D_{PF} covered large areas and lasted years at a
751 time; whereas D_{HG} had two patterns: (1) one related to dairy cattle strip grazing, which were
752 short-lived due to quick recovery times of grasses in fertilized soils; and (2) more widespread
753 and longer continuous disturbances occurring on steeper slopes grazed by sheep and beef cattle,
754 particularly following drought periods. Because our disturbance analyses had a spatial resolution

755 of 463 m, we likely missed some paddock-scale disturbances. Future work could use Landsat
756 imagery (30-m resolution) to assess disturbance (*sensu* de Beurs et al., 2016).

757 All six catchments with ‘meaningful’ increases in D_{HG} had large increases in dairy cattle
758 density 1990-2012 (mean of +1.0 SU/ha across the catchment). Not surprisingly, all six
759 catchments suffered impacts to water quality. Five of the six had ‘meaningful’ increases in DRP
760 and three had meaningful increases in NO_x and TN . One had a ‘meaningful’ increase in $TURB$
761 and three had significant reductions in DO . One of these catchments, in particular, may provide a
762 glimpse into NZ’s future if agricultural intensification continues. The Waingongoro River
763 catchment (WA3) is covered almost entirely by HG (91.2%), with practically all of this land
764 being used for intensive strip grazing. The SUD_{da} was 15.0 SU/ha in 1990 and increased to 15.4
765 SU/ha by 2012. The D_{HG} from 2000-2013 had a strong increasing trend of 9.8%/y RSKSE,
766 associated with the intensification of dairy operations (Wilcock et al., 2009). The result of all this
767 intensification was that WA3 had ‘meaningful’ increases in TP , DRP , and TN . The only reason
768 NO_x did not display a significant trend is because the catchment was already overloaded with a
769 median river concentration of 1,852 mg/m³. Noteworthy is that these significant trends of
770 increasing SUD_{da} , D_{HG} , and nutrients are occurring not only in lowland catchments on the North
771 Island (WA3, HV2), but also in upland catchments of the North Island (RO6), as well as both
772 lowland (TK1) and upland (CH3, TK2) catchments on the South Island.

773 While disturbance was not itself a strong predictor of water quality, it did help explain
774 outliers of land use-water quality relationships. For example, streams with high DRP (> 20
775 mg/m³; 10th percentile) had one of two dominant land uses, either plantation forest, PF (RO2,
776 RO3) or high-producing grassland, HG (HM5, WA3, WA9, HM4, HM2). The one exception was
777 RO4, which had relatively low coverage of PF (11.2%) and HG (2.9%). In fact, RO4 is

778 dominated by NF (79.1%). Upon closer examination, we found that the small areas of *PF* and
779 *HG* in RO4 were disturbed frequently. Further, most of the disturbed forestry occurred on steep
780 slopes and most of the disturbed pastures (practically all sheep and beef) occurred on hilly terrain
781 adjacent to stream channels. Our high temporal-resolution analyses of disturbance showed that
782 even though this catchment is mostly indigenous forest, intense disturbances on small
783 proportions of developed land can have a considerable impact on water quality. RO4 is also
784 experiencing significant increases in *TURB* and *TP*, as well as a significant decrease in *Q*.
785 Another outlier example was RO3, which was the only non-*HG*-dominated catchment with high
786 NO_x (634 mg/m³). RO3 was dominated by *PF* (69.8%), but it had the highest median disturbance
787 (10.5%) of all catchments. This catchment also exceeds ANZECC guidelines for *DRP* and has
788 experienced meaningful increases in *TURB*, *TN*, and NO_x .

789 We believe that land disturbance and consequently river eutrophication and reduced
790 visual clarity will continue to worsen in some NZ catchments based on the following. More
791 plantation forests were planted 1993-1997 (3,810 km²) than any other 5-y period in NZ history
792 (NZFFA, 2014). With a 28-y mean age of harvest, NZ will experience its greatest coverage and
793 intensity of forest disturbance around 2025, less than 10 years from now. When combined with
794 drought and intense storms, the potential for nutrient and sediment mobilization from these lands
795 into NZ's rivers is high, especially given that approximately 45% of these plantings occurred on
796 high-producing grasslands (NZFFA, 2014) where many of the legacy nutrients will be exported
797 to rivers during forest harvest (Davis, 2014). Many of these plantings also occurred on steep
798 slopes, which exacerbates sediment runoff. If carbon prices continue to stay low, there will be a
799 high likelihood that many of the harvested forests will be converted to pasture, adding even more
800 nutrients to NZ rivers (PCE, 2013). Given that the Central Government created a national policy

801 goal of nearly doubling the export to GDP ratio by the year 2025 (MBIE, 2015), NZ is likely to
802 see continued increases in livestock density, fertilizer usage, and supplemental feed to support
803 these extra livestock, all of which will add even more pressure and risks of eutrophication on
804 NZ's rivers.

805

806 **Conclusions**

807 This study had the overall goal of describing how changes in land use intensity impact
808 river water quality across broad scales and over long periods. To address this goal we used a
809 combination of 'brute force' statistical analyses (in terms of hundreds of analyses using a suite of
810 physiographic, land use, and water quality data for 77 catchments over 26 years) and careful
811 examination (using multi-resolution data to find patterns and relationships among these
812 variables). This goal was ambitious and we likely missed some relationships and details of water
813 quality changes. However, we found empirical evidence for several key relationships among land
814 use intensity, geomorphic processes, and water quality, which we now place into a broader
815 perspective.

816 The greatest negative impact on river water quality in New Zealand (NZ) in recent
817 decades has been high-producing pastures that require large amounts of fertilizer to support high
818 densities of livestock. While this claim has been previously published (Davies-Colley, 2013;
819 Howard-Williams et al., 2010; and references within), our results and supporting information
820 show that the relationship between high-producing pastures and water quality is complicated,
821 being dependent on livestock type/density, disturbance regime, and physiography, particularly
822 soil type. Dairy cattle receive much of the blame for degraded water quality because of their high
823 nutrient requirements (Howard-Williams et al., 2010), but beef cattle can also strongly degrade

824 water quality due to comparable required inputs and grazing on steeper land with a higher
825 potential for runoff (McDowell et al., 2008). Further, pasture designations/boundaries are
826 becoming increasingly blurred by modern cattle management, with greater movements of dairy
827 and beef cattle among pastures, greater use of high-producing pastures for beef, over-wintering
828 of dairy cattle on beef pastures, and cross-breeding (Morris, 2013). While riparian fencing has
829 plausibly improved the clarity of NZ rivers, the removal of millions of sheep from steep slopes
830 has also likely played a role that should be investigated further.

831 New Zealand is the global leading exporter of whole milk powder, butter, and sheep
832 products; and NZ's prominence in these industries is likely to continue over the next decade
833 (OECD/FAO, 2015). In this most recent environmental review by the Organisation for Economic
834 Co-operation and Development, NZ had the highest percent increase (1990-2005) in agricultural
835 production out of 29 OECD countries, the highest percent increase in N-fertilizer use, and the 2nd
836 highest increase in P-fertilizer use. This agricultural intensification over our study period is
837 reflected in overall nutrient enrichment of NZ rivers. If cattle continue to be added at the rates we
838 documented, additional fertilizers and supplemental feed will be needed. Even if best
839 management practices are adopted to reduce nutrient export to rivers, there is already a half-
840 century legacy of nutrients distributed across the NZ landscape that will continue to leak to the
841 rivers (Larned et al., 2016). Indeed, the full impact of agricultural intensification on river water
842 quality will not be fully appreciated for another several decades (Howard-Williams et al., 2010;
843 Vant and Smith, 2004). Having an extensive national network like the NRWQN to document and
844 study these water quality changes will be important.

845

846

847 Author contribution

848 J. Julian designed the study and performed most of the analyses. K. de Beurs developed the
849 disturbance dataset and performed all trend analyses, both with assistance from B. Owsley. R.
850 Davies-Colley provided water quality dataset and guidance on its use. A.-G. Ausseil developed
851 the stock unit density dataset and provided guidance on land use analyses. J. Julian prepared the
852 manuscript with contributions from all co-authors.

853

854 Acknowledgments

855 The inspiration for this research was J.P. Julian's Fulbright Senior Scholar Fellowship,
856 which was hosted by the National Institute of Water & Atmospheric Research (NIWA) in
857 Hamilton, NZ in 2012. This work was funded by NASA LCLUC grant NNX14AB77G and NSF
858 Geography grants #1359970 and #1359948 (Co-PIs Julian and de Beurs). Andrew Tait (Climate
859 Principal Scientist at NIWA) provided climate data. Agricultural production and other data
860 essential to this manuscript was collected by William Wright (Landcare Research; LCR). Many
861 other people in New Zealand provided expert advice, including Suzie Greenhalgh (LCR), Sandy
862 Elliot (NIWA), Andrew Hughes (NIWA), Deborah Ballantine (then of NIWA), Graham
863 McBride (NIWA), Murray Hicks (NIWA), David Hamilton (University of Waikato), Les Basher
864 (LCR), Ian Fuller (Massey University), Roger Young (Cawthron Institute), Rien Visser
865 (University of Canterbury), and David Lee-Jones (USDA FAS). Support for this project was also
866 provided by numerous Regional Councils/Districts, including Auckland, Canterbury, Horizons,
867 Tasman, and Waikato.

868

869

870 **References**

- 871 Ausseil, A. G. E., Dymond, J. R., Kirschbaum, M. U. F., Andrew, R. M., and Parfitt, R. L.:
872 Assessment of multiple ecosystem services in New Zealand at the catchment scale,
873 Environmental Modelling & Software, 43, 37-48, 10.1016/j.envsoft.2013.01.006, 2013.
- 874 Australian and New Zealand Environment and Conservation Council (ANZECC): Australian and
875 New Zealand Guidelines for Fresh and Marine Water Quality, Volume 1: The Guidelines,
876 National Water Quality Management Strategy, Paper No. 4, Onehunga, 314 pp., 2000.
- 877 Ballantine, D. J., and Davies-Colley, R. J.: Water quality trends in New Zealand rivers: 1989–
878 2009, Environmental Monitoring and Assessment, 186, 1939-1950, 10.1007/s10661-013-
879 3508-5, 2014.
- 880 Ballantine, D. J., Hughes, A. O., and Davies-Colley, R. J.: Mutual relationships of suspended
881 sediment, turbidity and visual clarity in New Zealand rivers, in: Proceedings of the
882 International Association of Hydrological Sciences, New Orleans, 265-271, 2014.
- 883 Baron, J. S., Poff, N. L., Angermeier, P. L., Dahm, C. N., Gleick, P. H., Hairston, N. G., Jackson,
884 R. B., Johnston, C. A., Richter, B. D., and Steinman, A. D.: Meeting ecological and
885 societal needs for freshwater, Ecological Applications, 12, 1247-1260, 2002.
- 886 Basher, L. R., Hicks, D. M., Clapp, B., and Hewitt, T.: Sediment yield response to large storm
887 events and forest harvesting, Motueka River, New Zealand, New Zealand Journal of
888 Marine and Freshwater Research, 45, 333-356, 2011.
- 889 Bewsell, D., Monaghan, R. M., and Kaine, G.: Adoption of stream fencing among dairy farmers
890 in four New Zealand catchments, Environmental Management, 40, 201-209, 2007.
- 891 Blüthgen, N., Dormann, C. F., Prati, D., Klaus, V. H., Kleinebecker, T., Hölzel, N., Alt, F.,
892 Boch, S., Gockel, S., Hemp, A., Müller, J., Nieschulze, J., Renner, S. C., Schöning, I.,

893 Schumacher, U., Socher, S. A., Wells, K., Birkhofer, K., Buscot, F., Oelmann, Y.,
894 Rothenwöhrer, C., Scherber, C., Tschardtke, T., Weiner, C. N., Fischer, M., Kalko, E. K.
895 V., Linsenmair, K. E., Schulze, E.-D., and Weisser, W. W.: A quantitative index of land-
896 use intensity in grasslands: Integrating mowing, grazing and fertilization, *Basic and*
897 *Applied Ecology*, 13, 207-220, <http://dx.doi.org/10.1016/j.baae.2012.04.001>, 2012.

898 Boesch, D. F.: Challenges and opportunities for science in reducing nutrient over-enrichment of
899 coastal ecosystems, *Estuaries*, 25, 886-900, [10.1007/bf02804914](https://doi.org/10.1007/bf02804914), 2002.

900 Brauman, K. A., Daily, G. C., Duarte, T. K., and Mooney, H. A.: The nature and value of
901 ecosystem services: An overview highlighting hydrologic services, in: *Annual Review of*
902 *Environment and Resources*, *Annual Review of Environment and Resources*, *Annual*
903 *Reviews*, Palo Alto, 67-98, 2007.

904 Brierley, G. P.: Landscape memory: the imprint of the past on contemporary landscape forms
905 and processes, *Area*, 42, 76-85, 2010.

906 Campbell, N., D'Arcy, B., Frost, A., Novotny, V., Sansom, A.: Diffuse pollution. An
907 introduction to the problems and solutions, IWA publishing, London, 322 pp., 2004.

908 Cleveland, W. S., and Devlin, S. J.: Locally weighted regression: An approach to regression
909 analysis by local fitting, *Journal of the American Statistical Association*, 83, 596-610,
910 1988.

911 Close, M. E., and Davies-Colley, R. J.: Baseflow water chemistry in New Zealand rivers 1.
912 Characterisation, *New Zealand Journal of Marine and Freshwater Research*, 24, 319-341,
913 [10.1080/00288330.1990.9516428](https://doi.org/10.1080/00288330.1990.9516428), 1990.

914 Croke, J. C., and Hairsine, P. B.: Sediment delivery in managed forests: a review, *Environmental*
915 *Reviews*, 14, 59-87, [10.1139/a05-016](https://doi.org/10.1139/a05-016), 2006.

916 Davies-Colley, R. J.: River water quality in New Zealand: An introduction and overview, in:
917 Ecosystem Services in New Zealand - Conditions and Trends, edited by: Dymond, J. R.,
918 Manaaki Whenua Press, Lincoln, 432-447, 2013.

919 Davies-Colley, R. J., Vant, W. N., and Smith, D. G.: Colour and Clarity of Natural Waters:
920 Science and Management of Optical Water Quality, Blackburn Press, Caldwell, 310 pp.,
921 2003.

922 Davies-Colley, R. J., Smith, D. G., Ward, R. C., Bryers, G. G., McBride, G. B., Quinn, J. M., and
923 Scarsbrook, M. R.: Twenty years of New Zealand's National Rivers Water Quality
924 Network: Benefits of careful design and consistent operation, Journal of the American
925 Water Resources Association, 47, 750-771, 2011.

926 Davies-Colley, R. J., Ballantine, D. J., Elliott, A. H., Swales, A., Hughes, A. O., and Gall, M. P.:
927 Light attenuation – a more effective basis for the management of fine suspended
928 sediment than mass concentration?, Water Science and Technology, 69, 1867-1874. doi:
929 10.2166/wst.2014.096, 2014.

930 Davis, M.: Nutrient losses from forestry in the Lake Taupo catchment, Environment Waikato
931 Technical Report 2005/37, Hamilton, 22, 2005.

932 Davis, M.: Nitrogen leaching losses from forests in New Zealand, New Zealand Journal of
933 Forestry Science, 44, 1-14, 10.1186/1179-5395-44-2, 2014.

934 de Beurs, K. M., Owsley, B. C., and Julian, J. P.: Disturbance analyses of forests and grasslands
935 with MODIS and Landsat in New Zealand, International Journal of Applied Earth
936 Observation and Geoinformation, 45, Part A, 42-54,
937 <http://dx.doi.org/10.1016/j.jag.2015.10.009>, 2016.

938 Dodds, W. K., and Welch, E. B.: Establishing nutrient criteria in streams, *Journal of the North*
939 *American Benthological Society*, 19, 186-196, 2000.

940 Dymond, J. R., Betts, H. D., and Schierlitz, C. S.: An erosion model for evaluating regional land-
941 use scenarios, *Environmental Modelling & Software*, 25, 289-298,
942 [10.1016/j.envsoft.2009.09.011](https://doi.org/10.1016/j.envsoft.2009.09.011), 2010.

943 Ekholm, P., Rankinen, K., Rita, H., Raike, A., Sjoblom, H., Raateland, A., Vesikko, L., Cano
944 Bernal, J. E., and Taskinen, A.: Phosphorus and nitrogen fluxes carried by 21 Finnish
945 agricultural rivers in 1985–2006, *Environmental Monitoring and Assessment*, 187, 1-17,
946 [10.1007/s10661-015-4417-6](https://doi.org/10.1007/s10661-015-4417-6), 2015.

947 Erb, K.-H., Haberl, H., Jepsen, M. R., Kuemmerle, T., Lindner, M., Muller, D., Verburg, P. H.,
948 and Reenberg, A.: A conceptual framework for analysing and measuring land-use
949 intensity, *Current Opinion in Environmental Sustainability*, 5, 464-470,
950 <http://dx.doi.org/10.1016/j.cosust.2013.07.010>, 2013.

951 Erb, K.-H., Fetzel, T., Haberl, H., Kastner, T., Kroisleitner, C., Lauk, C., Niedertscheider, M.,
952 and Plutzer, C.: Beyond Inputs and Outputs: Opening the Black-Box of Land-Use
953 Intensity, in: *Social Ecology: Society-Nature Relations across Time and Space*, edited by:
954 Haberl, H., Fischer-Kowalski, M., Krausmann, F., and Winiwarter, V., Springer
955 International Publishing, Cham, 93-124, 2016.

956 Evans, R.: Soil erosion in the UK initiated by grazing animals, *Applied Geography*, 17, 127-141,
957 [http://dx.doi.org/10.1016/S0143-6228\(97\)00002-7](http://dx.doi.org/10.1016/S0143-6228(97)00002-7), 1997.

958 Fahey, B. D., Marden, M., and Phillips, C. J.: Sediment yields from plantation forestry and
959 pastoral farming, coastal Hawke's Bay, North Island, New Zealand, *Journal of Hydrology*
960 (New Zealand), 42, 27-38, 2003.

961 Foley, J. A., DeFries, R., Asner, G. P., Barford, C., Bonan, G., Carpenter, S. R., Chapin, F. S.,
962 Coe, M. T., Daily, G. C., Gibbs, H. K., Helkowski, J. H., Holloway, T., Howard, E. A.,
963 Kucharik, C. J., Monfreda, C., Patz, J. A., Prentice, I. C., Ramankutty, N., and Snyder, P.
964 K.: Global consequences of land use, *Science*, 309, 570-574, 2005.

965 Forest Owners Association (FOA): New Zealand plantation forest industry: Facts and figures
966 2011/2012, Ministry for Primary Industries, Wellington, 46 pp., 2012.

967 Fransen, P. J. B., Phillips, C. J., and Fahey, B. D.: Forest road erosion in New Zealand:
968 overview, *Earth Surface Processes and Landforms*, 26, 165-174, 10.1002/1096-
969 9837(200102)26:2<165::AID-ESP170>3.0.CO;2-#, 2001.

970 Grohmann, C. H., Smith, M. J., and Riccomini, C.: Multiscale analysis of topographic surface
971 roughness in the Midland Valley, Scotland, *IEEE Transactions on Geoscience and*
972 *Remote Sensing*, 49, 1200-1213, 2011.

973 Gunn, A., and Rutherford, C.: The Dairying and Clean Streams Accord: Snapshot of Progress
974 2011/2012, Ministry for Primary Industries, Wellington, 12, 2013.

975 Hart, B. T., Maher, B., and Lawrence, I.: New generation water quality guidelines for ecosystem
976 protection, *Freshwater Biology*, 41, 347-359, 10.1046/j.1365-2427.1999.00435.x, 1999.

977 Hicks, D. M., Gomez, B., and Trustrum, N. A.: Erosion thresholds and suspended sediment
978 yields, Waipaoa River Basin, New Zealand, *Water Resources Research*, 36, 1129-1142,
979 10.1029/1999WR900340, 2000.

980 Hilton, J., O'Hare, M., Bowes, M. J., and Jones, J. I.: How green is my river? A new paradigm of
981 eutrophication in rivers, *Science of the Total Environment*, 365, 66-83, 2006.

982 Howard-Williams, C., Davies-Colley, R., Rutherford, K., and Wilcock, R.: Diffuse pollution and
983 freshwater degradation: New Zealand Perspectives, International Conference of the IWA
984 Diffuse Pollution Specialist Group, Beaupre, Quebec, 2010.

985 Hughes, A. O., and Quinn, J. M.: Before and After Integrated Catchment Management in a
986 Headwater Catchment: Changes in Water Quality, *Environmental Management*, 54,
987 1288-1305, 10.1007/s00267-014-0369-9, 2014.

988 Julian, J. P., Doyle, M. W., Powers, S. M., Stanley, E. H., and Riggsbee, J. A.: Optical water
989 quality in rivers, *Water Resources Research*, 44, W10411,
990 doi:10410.11029/12007WR006457, 2008.

991 Julian, J. P., Davies-Colley, R. J., Gallegos, C. L., and Tran, T. V.: Optical Water Quality of
992 Inland Waters: A Landscape Perspective, *Annals of the Association of American*
993 *Geographers*, 103, 309-318, 2013.

994 Kamarinas, I., Julian, J. P., Hughes, A. O., Owsley, B. C., and de Beurs, K. M.: Nonlinear
995 changes in land cover and sediment runoff in a New Zealand catchment dominated by
996 plantation forestry and livestock grazing, *Water*, 8, 436, doi:10.3390/w8100436, 2016.

997 Kronvang, B., Andersen, H. E., Børgesen, C., Dalgaard, T., Larsen, S. E., Bøgestrand, J., and
998 Blicher-Mathiasen, G.: Effects of policy measures implemented in Denmark on nitrogen
999 pollution of the aquatic environment, *Environmental Science & Policy*, 11, 144-152,
1000 <http://dx.doi.org/10.1016/j.envsci.2007.10.007>, 2008.

1001 Kuemmerle, T., Erb, K., Meyfroidt, P., Müller, D., Verburg, P. H., Estel, S., Haberl, H., Hostert,
1002 P., Jepsen, M. R., Kastner, T., Levers, C., Lindner, M., Plutzer, C., Verkerk, P. J., van der
1003 Zanden, E. H., and Reenberg, A.: Challenges and opportunities in mapping land use

1004 intensity globally, *Current Opinion in Environmental Sustainability*, 5, 484-493,
1005 <http://dx.doi.org/10.1016/j.cosust.2013.06.002>, 2013.

1006 Landcare Research (LCR): New Zealand Land Cover Database, available
1007 at: <http://www.lcdb.scinfo.org.nz/>, 2015.

1008 Larned, S. T., Scarsbrook, M. R., Snelder, T. H., Norton, N. J., and Biggs, B. J. F.: Water quality
1009 in low-elevation streams and rivers of New Zealand: recent state and trends in contrasting
1010 land-cover classes, *New Zealand Journal of Marine and Freshwater Research*, 38, 347-
1011 366, 2004.

1012 Larned, S., Snelder, T., Unwin, M., and McBride, G.: Water quality in New Zealand rivers:
1013 current state and trends, *New Zealand Journal of Marine and Freshwater Research*, DOI:
1014 10.1080/00288330.2016.1150309, 2016.

1015 Lobser, S. E., and Cohen, W.B.: MODIS tasseled cap: land cover characteristicsexpressed
1016 through transformed MODIS data, *Int. J. Remote Sens.*, 28,5079–5101, 2007.

1017 McAllister, T. G., Wood, S. A., and Hawes, I.: The rise of toxic benthic Phormidium
1018 proliferations: a review of their taxonomy, distribution, toxin content and factors
1019 regulating prevalence and increased severity, *Harmful Algae*, 55, 282-294, 2016.

1020 McDowell, R. W., Houlbrooke, D. J., Muirhead, R. W., Muller, K., Shepherd, M., and Cuttle, S.
1021 P.: *Grazed Pastures and Surface Water Quality*, Nova Science Publishers, New York, 238
1022 pp., 2008.

1023 Ministry of Business, Innovation & Employment (MBIE): *Business Growth Agenda: 2015*,
1024 available at: [http://www.mbie.govt.nz/info-services/business/business-growth-
1025 agenda/pdf-and-image-library/towards-2025/mb13078-1139-bga-report-01-export-
1026 markets-09sept-v17-fa-web.PDF](http://www.mbie.govt.nz/info-services/business/business-growth-agenda/pdf-and-image-library/towards-2025/mb13078-1139-bga-report-01-export-markets-09sept-v17-fa-web.PDF), 2015.

1027 Ministry for the Environment (MfE): Land-use and carbon analysis system (LUCAS): Satellite
1028 imagery interpretation guide for land-use classes (2nd edition), Ministry for the
1029 Environment, Wellington, 73 pp., 2012.

1030 Monaghan, R. M., Hedley, M. J., Di, H. J., McDowell, R. W., Cameron, K. C., and Ledgard, S.
1031 F.: Nutrient management in New Zealand pastures— recent developments and future
1032 issues, *N. Z. J. Agric. Res.*, 50, 181-201, 10.1080/00288230709510290, 2007.

1033 Moosmann, L., Müller, B., Gächter, R., Wüest, A., Butscher, E., and Herzog, P.: Trend-oriented
1034 sampling strategy and estimation of soluble reactive phosphorus loads in streams, *Water
1035 Resources Research*, 41, W01020, 10.1029/2004WR003539, 2005.

1036 Morris, S. T.: Sheep and beef cattle production systems, in: *Ecosystem Services in New Zealand
1037 - Conditions and Trends*, edited by: Dymond, J. R., Manaaki Whenua Press, Lincoln, 79-
1038 84, 2013.

1039 Motha, J. A., Wallbrink, P. J., Hairsine, P. B., and Grayson, R. B.: Determining the sources of
1040 suspended sediment in a forested catchment in southeastern Australia, *Water Resources
1041 Research*, 39, 1056, doi:10.1029/2001WR000794, 2003.

1042 Newsome, P. F. J., R.H., W., and Willoughby, E. J.: *Land Resource Information System Spatial
1043 Data Layers: Data Dictionary*, Palmerston North, 75, 2008.

1044 New Zealand Farm Forestry Association (NZFFA): *National Exotic Forest Description: as at 1
1045 April 2014*, Ministry for Primary Industries, Wellington, 77 pp., 2014.

1046 OECD/FAO: *OECD-FAO Agricultural Outlook 2015-2024*, OECD/Food and Agriculture
1047 Organization of the United Nations Paris, 148, 2015.

1048 Parkyn, S. M., Davies-Colley, R. J., Scarsbrook, M. R., Halliday, N. J., Nagels, J. W., Marden,
1049 M., and Rowan, D.: *Pine afforestation and stream health: a comparison of land-use in two*

1050 soft rock catchments, East Cape, New Zealand, New Zealand Natural Sciences, 31, 113-
1051 135, 2006.

1052 Parliamentary Commissioner for the Environment (PCE): Water quality in New Zealand: land
1053 use and nutrient pollution, Parliamentary Commissioner for the Environment,
1054 Wellington, 82 pp., available at: [http://www.pce.parliament.nz/media/1275/pce-water-](http://www.pce.parliament.nz/media/1275/pce-water-quality-land-use-web-amended.pdf)
1055 [quality-land-use-web-amended.pdf](http://www.pce.parliament.nz/media/1275/pce-water-quality-land-use-web-amended.pdf), 2013.

1056 Phillips C. J., Marden M., and Rowan D.: Sediment yield following plantation harvesting,
1057 Coromandel Peninsula, North Island, New Zealand, Journal of Hydrology (NZ), 44, 29-
1058 44, 2005.

1059 Ramankutty, N., Graumlich, L., Achard, F., Alves, D., Chhabra, A., DeFries, R. S., Foley, J. A.,
1060 Geist, H., Houghton, R. A., Goldewijk, K. K., Lambin, E. F., Millington, A., Rasmussen,
1061 K., Reid, R. S., and Turner, B. L.: Global Land-Cover Change: Recent Progress,
1062 Remaining Challenges, in: Land-Use and Land-Cover Change: Local Processes and
1063 Global Impacts, edited by: Lambin, E. F., and Geist, H., Springer Berlin Heidelberg,
1064 Berlin, Heidelberg, 9-39, 2006.

1065 Saunders, W. M. H.: Phosphate retention by New Zealand soils and its relationship to free
1066 sesquioxides, organic matter, and other soil properties, N. Z. J. Agric. Res., 8, 30-57,
1067 DOI: 10.1080/00288233.1965.10420021, 1965.

1068 Scarsbrook, M. R.: State and trends in the National River Water Quality Network (1989-2005),
1069 Ministry for the Environment, ME number 778, 48 pp., available
1070 at: <http://www.mfe.govt.nz/publications/ser/water-quality-network-nov06/index.html>,
1071 2006.

1072 Scarsbrook, M. R., McBride, C. G., McBride, G. B., and Bryers, G. G.: Effects of climate
1073 variability on rivers: Consequences for long term water quality analysis, Journal of the
1074 American Water Resources Association, 39, 1435-1447, 2003.

1075 Smith, D. G., McBride, G. B., Bryers, G. G., Wisse, J., and Mink, D. F. J.: Trends in New
1076 Zealand's National River Water Quality Network, New Zealand Journal of Marine and
1077 Freshwater Research, 30, 485-500, 10.1080/00288330.1996.9516737, 1996.

1078 Smith, D. G., Davies-Colley, R. J., Knoeff, J., and Slot, G. W. J.: Optical characteristics of New
1079 Zealand rivers in relation to flow, Journal of the American Water Resources Association,
1080 33, 301–312, 1997.

1081 Snelder, T. H.: Definition of a multivariate classification of New Zealand Lakes, National
1082 Institute of Water & Atmospheric Research Ltd, Christchurch, 32, 2006.

1083 Snelder, T., Biggs, B., and Weatherhead, M.: New Zealand River Environment Classification
1084 User Guide (REC, v2), Produced for the Ministry for the Environment by the National
1085 Institute of Water and Atmospheric Research (NIWA), Publication number ME 1026,
1086 2010.

1087 Stamm, C., Jarvie, H. P., and Scott, T.: What's More Important for Managing Phosphorus:
1088 Loads, Concentrations or Both?, Environmental science & technology, 48, 23-24,
1089 10.1021/es405148c, 2014.

1090 Statistics New Zealand (StatsNZ): 2012 Agricultural Census tables, available
1091 at: http://www.stats.govt.nz/browse_for_stats/industry_sectors/agriculture-horticulture-forestry/2012-agricultural-census-tables/livestock.aspx, 2015.

1093 Tait, A., and Turner, R.: Generating Multiyear Gridded Daily Rainfall over New Zealand,
1094 Journal of Applied Meteorology, 44, 1315-1323, 10.1175/JAM2279.1, 2005.

1095 Trimble, S. W., and Mendel, A. C.: The cow as a geomorphic agent—A critical review,
1096 *Geomorphology*, 13, 233–253, 1995.

1097 Vant, B., and Smith, P.: Nutrient concentrations and water ages in 11 streams flowing into
1098 Lake Taupo, Environment Waikato Technical Report 2002/18R, 26, 2004.

1099 Varanka, S., and Luoto, M.: Environmental determinants of water quality in boreal rivers based
1100 on partitioning methods, *River Research and Applications*, 28, 1034-1046,
1101 doi:10.1002/rra.1502, 2012.

1102 Vitousek, P. M., Mooney, H. A., Lubchenco, J., and Melillo, J. M.: Human domination of Earth's
1103 ecosystems, *Science*, 277, 494, 1997.

1104 Vorosmarty, C. J., McIntyre, P. B., Gessner, M. O., Dudgeon, D., Prusevich, A., Green, P.,
1105 Glidden, S., Bunn, S. E., Sullivan, C. A., Liermann, C. R., and Davies, P. M.: Global
1106 threats to human water security and river biodiversity, *Nature*, 467, 555-561, 2010.

1107 Webb, T. W., and Wilson, A. D.: *A Manual of Land Characteristics for Evaluation of Rural land*,
1108 Landcare Research New Zealand Ltd, Lincoln, 1995.

1109 Wilcock, R. J., Betteridge, K., Shearman, D., Fowles, C. R., Scarsbrook, M. R., Thorrold, B. S.,
1110 and Costall, D.: Riparian protection and on-farm best management practices for
1111 restoration of a lowland stream in an intensive dairy farming catchment: A case study,
1112 *New Zealand Journal of Marine and Freshwater Research*, 43, 803-818,
1113 doi:10.1080/00288330909510042, 2009.

1114 Wilmshurst, J. M., Anderson, A. J., Higham, T. F. G., Worthy, T. H.: Dating the late prehistoric
1115 dispersal of Polynesians to New Zealand using the commensal Pacific rat, *Proceedings of*
1116 *the National Academy of Sciences*, 105, 7676-7680, 2008.

1117 Wood, S., Hawes, I., McBride, G., Truman, P., and Dietrich, D.: Advice to inform the
1118 development of a benthic cyanobacteria attribute, Cawthron Institute, Report No. 2752,
1119 Nelson, 91, 2015.

1120 Woods, R., Bidwell, V., Clothier, B., Green, S., Elliott, S., Shankar, U., Harris, S., Hewitt, A.,
1121 Gibb, R., Parfitt, R., and Wheeler, D.: The CLUES Project: Predicting the effects of land-
1122 use on water quality - Stage II, National Institute of Water & Atmospheric Research,
1123 Christchurch, 2006.

1124 Woodward, S. J. R.: Bite mechanics of cattle and sheep grazing grass-dominant swards, Applied
1125 Animal Behaviour Science, 56, 203-222, 1998.

1126 Zobrist, J., and Reichert, P.: Bayesian estimation of export coefficients from diffuse and point
1127 sources in Swiss watersheds, Journal of Hydrology, 329, 207-223,
1128 <http://dx.doi.org/10.1016/j.jhydrol.2006.02.014>, 2006.

1129
1130
1131
1132
1133

1134 **Tables**

1135

1136 Table 1. Water quality variables measured by the National River Water Quality Network
 1137 (NRWQN) obtained from monthly grab samples from 1989 to 2014 for 77 catchments. Details
 1138 on analytical methods can be found in Davies-Colley et al. (2011).

Variable	Definition (units)
<i>Q</i>	Water discharge (m ³ /s)
<i>T_w</i>	Water temperature (°C)
<i>DO</i>	Dissolved oxygen (%)
<i>COND</i>	Water conductivity (µS/cm)
<i>pH_w</i>	Water pH (-log ₁₀ [H ⁺])
<i>CLAR</i>	Horizontal visual water clarity from black disc sighting range (m)
<i>TURB</i>	Water turbidity (NTU)
<i>CDOM</i>	Colored dissolved organic matter, measured as spectrophotometric absorbance of a membrane filtrate at 440 nm (m ⁻¹)
<i>TN</i>	Total nitrogen (mg/m ³)
<i>NO_x</i>	Oxidized nitrogen in nitrate and nitrite forms (mg/m ³)
<i>TP</i>	Total phosphorus (mg/m ³)
<i>DRP</i>	Dissolved reactive phosphorus (mg/m ³)

1139

1140

1141 Table 2. Landscape variables characterizing the 77 catchments of the National River Water
 1142 Quality Network (NRWQN). More details on sources for these data can be found in Methods
 1143 section.

Variable	Definition (units)	Source (resolution/scale)
Morphometric variables		
Area (A)	Total catchment area above monitoring site (km ²)	National Elevation Dataset (30 m)
Drainage density (D_d)	Total length of streams per catchment area (km/km ²)	River Environment Classification, v2 (1:24,000)
Catchment slope (S_c)	Mean slope across entire catchment (degrees)	National Elevation Dataset (30 m)
Ruggedness (R_r)	Standard deviation of catchment slope (degrees)	National Elevation Dataset (30 m)
Soil variables		
Silt-clay percentage ($SC\%$)	Percentage of catchment surface soils dominated by clayey or silty soils (%)	Fundamental Soil Layers (1:63,360)
Soil depth (Z_s)	Mean maximum potential rooting depth across catchment (m)	Fundamental Soil Layers (1:63,360)
Soil pH (pH_s)	Mean pH at 0.2-0.6 m depth across catchment ($-\log_{10}[H^+]$)	Fundamental Soil Layers (1:63,360)
Cation exchange capacity (CEC)	Weighted mean CEC at 0-0.6 m depth across catchment (cmoles [+)/kg)	Fundamental Soil Layers (1:63,360)
Organic matter percentage ($OM\%$)	Weighted mean of total carbon at 0-0.2 m depth across catchment (%)	Fundamental Soil Layers (1:63,360)
Phosphate retention (P_{ret})	Weighted mean of phosphate retention at 0-0.2 m depth across catchment (%)	Fundamental Soil Layers (1:63,360)
Hydro-climatological variables		
Median annual precipitation (MAP)	Median annual precipitation averaged across catchment (mm/y)	NIWA National Climate Database (5 km)
Median annual temperature (MAT)	Median annual temperature averaged across catchment (°C)	NIWA National Climate Database (5 km)
Median annual sunshine (MAS)	Median annual sunshine hours averaged across catchment (hours/y)	NIWA National Climate Database (5 km)
Median discharge (Q_{50})	Median discharge from NRWQN samples during 1989-2014 (m ³ /s)	NRWQN (catchment)

Relative water storage (<i>RWS</i>)	Proportion of annual Q_{50} stored in reservoirs/lakes (m^3/m^3)	Freshwater Environments New Zealand (1:50,000)
--	---	---

Land Use and Land Disturbance variables

Land use	Percent of catchment that is occupied by each land use (%); see Table 3 for land uses	Land Cover Database (LCDB, v 4.1), 2001 (1 ha)
High-producing pasture disturbance (<i>D_{HG}</i>)	Percent of high-producing grasslands within catchment that is disturbed (%), based on aggregate of 463-m pixels within catchment	de Beurs et al., 2016 (463 m; 8-day)
Plantation forestry disturbance (<i>D_{PF}</i>)	Percent of plantation forestry within catchment that is disturbed (%), based on aggregate of 463-m pixels within catchment	de Beurs et al., 2016 (463 m; 8-day)
Catchment disturbance (<i>D_C</i>)	Percent of catchment that is disturbed (%), based on aggregate of 463-m pixels within catchment	de Beurs et al., 2016 (463 m; 8-day)
Stock unit density (<i>SUD</i>)	Catchment-averaged stock unit density for dairy (<i>da</i>), beef (<i>be</i>), deer (<i>de</i>), and sheep (<i>sh</i>) in 2011 (SU/ha); subscripts are used to isolate SUD by livestock type	Ausseil et al., 2013 (1 ha)
Change in stock unit density (<i>SUD₂₀₁₂₋₁₉₉₀</i>)	Difference between SUD in 2012 and 1990 (SU/ha)	Statistics NZ (territorial authority)

1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161

1162 Table 3. Land use classification used in this study, aggregated from the LUCAS (v11) and
 1163 LCDB (v4.1) land use/cover datasets.

Class (abbreviation)	Description	LUCAS classes	LCDB classes	2012 national coverage (%) LUCAS / LCDB
Non-plantation forest (NF)	All non-plantation forests $\geq 5\text{m}$; does not include Manuka/Kanuka	71	68, 69	29.2 / 23.9
Plantation forest (PF)	All forests that are planted for the purpose of harvesting	72,73	64, 71	7.9 / 7.6
Shrub/Grassland (SG)	All shrubs $< 5\text{m}$ and grasses that are not intensively managed	74, 76	41-44, 50-58	33.0 / 25.4
High-producing grassland (HG)	High-quality pasture grasses that are intensively managed	75	40	21.6 / 33.0
Perennial cropland (PC)	Orchards and vineyards	77	33	0.4 / 0.4
Annual cropland (AC)	All annual crops and cultivated bare ground	78	30	1.4 / 1.4
Open water (OW)	Rivers, lakes/reservoirs, ponds, and estuaries	79	20-22	1.9 / 2.0
Vegetated wetland (VW)	Herbaceous or woody vegetation periodically flooded; includes mangroves	80	45-47, 70	0.5 / 0.7
Urban (UR)	Built-up areas, infrastructure, transportation networks, and urban parks/open spaces	81	1-5	0.8 / 0.9
Barren/Other (BO)	Bare rock, sand, gravel and other areas not dominated by vegetation; includes mining and permanent ice/snow	82	6-16	3.3 / 4.8

1164
 1165

1166 Table 4. Statistical description of landscape variables for the 77 NRWQN catchments. Refer to
 1167 Tables 2 and 3 for variable descriptions.

Variable	Units	Minimum	Median	Maximum	Mean \pm SD
Morphometric Variables					
Area (<i>A</i>)	km ²	26	1126	20539	2639 \pm 3714
Drainage density (<i>D_d</i>)	km/km ²	1.30	1.59	2.61	1.60 \pm 0.16
Catchment slope (<i>S_c</i>)	degrees	3.4	15.9	30.3	16.3 \pm 6.8
Ruggedness (<i>R_r</i>)	degrees	3.4	10.8	15.8	10.6 \pm 2.4
Soil Variables					
Silt-clay percentage (<i>SC%</i>)	%	0	47.3	98.7	44.0 \pm 31.6
Soil depth (<i>Z_s</i>)	m	0.55	0.96	1.50	1.02 \pm 0.22
Soil pH (<i>pH</i>)	$-\log_{10}[\text{H}^+]$	4.8	5.6	6.5	5.6 \pm 0.3
Cation exchange capacity (<i>CEC</i>)	cmoles [+]/kg	11.6	18.7	33.5	18.8 \pm 4.6
Organic matter percentage (<i>OM%</i>)	%	2.8	6.7	23.2	7.2 \pm 2.9
Phosphate retention (<i>P_{ret}</i>)	%	19.9	39.0	77.8	41.5 \pm 12.2
Hydro-climatological Variables					
Median annual precipitation (<i>MAP</i>)	mm/y	533	1652	7044	1778 \pm 873
Median annual temperature (<i>MAT</i>)	°C	5.0	9.9	15.1	9.9 \pm 2.4
Median annual sunshine (<i>MAS</i>)	hours/y	1325	1856	2116	1841 \pm 146
Median discharge (<i>Q₅₀</i>)	m ³ /s	0.4	26.0	515.0	69.6 \pm 112.6
Relative water storage (<i>RWS</i>)	m ³ /m ³	0	0	29.2	1.1 \pm 3.7
Land Use Variables					
Non-plantation forest (<i>NF</i>)	%	0.1	20.5	94.1	26.7 \pm 23.3
Plantation forest (<i>PF</i>)	%	0	3.3	69.8	8.2 \pm 12.3
Shrub/Grassland (<i>SG</i>)	%	0.4	21.7	82.3	26.6 \pm 20.2
High-producing grassland (<i>HG</i>)	%	0	21.6	91.2	30.9 \pm 26.2
Perennial cropland (<i>PC</i>)	%	0	0	1.3	0.1 \pm 0.2
Annual cropland (<i>AC</i>)	%	0	0.1	7.9	0.6 \pm 1.4
Open water (<i>OW</i>)	%	0	0.4	25.6	1.9 \pm 4.3

Vegetated wetland (<i>VW</i>)	%	0	0.1	2.2	0.3 ± 0.4
Urban (<i>UR</i>)	%	0	0.1	5.8	0.4 ± 0.7
Barren/Other (<i>BO</i>)	%	0	1.3	30.0	4.4 ± 6.5
Land Disturbance Variables					
Catchment disturbance (<i>D_C</i>)	%	0	3.4	10.5	3.6 ± 2.1
<i>HG</i> disturbance (<i>D_{HG}</i>)	%	0	4.4	34.9	6.0 ± 6.4
<i>PF</i> disturbance (<i>D_{PF}</i>)	%	0	9.9	27.8	10.4 ± 6.7
Stock unit density (<i>SUD</i>)	SU/ha	0	2.2	16.1	3.2 ± 3.1
Dairy <i>SUD</i> (<i>SUD_{da}</i>)	SU/ha	0	0.2	15.4	1.2 ± 2.4
Beef <i>SUD</i> (<i>SUD_{be}</i>)	SU/ha	0	0.5	3.5	0.7 ± 0.8
Sheep <i>SUD</i> (<i>SUD_{sh}</i>)	SU/ha	0	0.6	4.5	1.2 ± 1.3
Deer <i>SUD</i> (<i>SUD_{de}</i>)	SU/ha	0	0	0.2	0 ± 0

1168

1169

1170

1171 Table 5. Statistical description of medians of water quality variables for the 77 NRWQN
 1172 catchments. Note that the ratio of mean/median can be used as an index of data skewness.

Variable	Units	Minimum	Median	Maximum	Mean \pm SD
<i>T_w</i>	°C	7.2	12.2	16.9	12.4 \pm 2.4
<i>DO</i>	%	75.5	100.8	113.1	100.0 \pm 4.7
<i>COND</i>	μ S/cm	39	92	528	113 \pm 83
<i>pH_w</i>	$-\log_{10}[\text{H}^+]$	6.9	7.7	8.5	7.7 \pm 0.3
<i>CLAR</i>	m	0.1	1.5	9.8	2.1 \pm 1.8
<i>TURB</i>	NTU	0.3	2.1	82	4.2 \pm 9.4
<i>CDOM</i>	m^{-1}	0.1	0.7	4.6	0.9 \pm 0.8
<i>TN</i>	mg/m^3	40	259	2162	369 \pm 361
<i>NO_x</i>	mg/m^3	1	107	1852	230 \pm 302
<i>TP</i>	mg/m^3	3	15	115	24 \pm 24
<i>DRP</i>	mg/m^3	0.5	5.0	66.2	8.6 \pm 11.2

1173

Table 6. River water quality trends from 1989-2014. The table reports numbers of sites (out of 77) in different categories of water quality time trend. All variables were flow-adjusted except flow and water temperature. Significant trends were taken to be those with a p-value < 0.05 in the Seasonal Kendall test. Meaningful trends were taken to be those which also had a magnitude (RSKSE) greater than 1% per year.

Direction of trend	River Water Quality Variable (1989-2014)											
	<i>Q</i>	<i>T_w</i>	<i>DO</i>	<i>COND</i>	<i>pH_w</i>	<i>CLAR</i>	<i>TURB</i>	<i>CDOM</i>	<i>TP</i>	<i>DRP</i>	<i>TN</i>	<i>NO_x</i>
Meaningful Increase	1	0	0	4	0	29	17	1	8	17	27	24
Significant Increase	1	21	6	48	12	5	1	1	6	3	6	3
No Significant Trend	67	54	42	19	48	39	50	56	52	49	39	37
Significant Decrease	3	2	29	6	17	2	0	13	4	5	3	1
Meaningful Decrease	5	0	0	0	0	2	9	6	7	3	2	12

Table 7. Correlations of water quality (median values) vs. the major land uses, livestock densities, and median catchment disturbance of the 77 NRWQN catchments. All values represent Spearman correlation coefficients (r_s). Nonsignificant relationships ($p \geq 0.05$) are denoted by *NS*. T_w was not included because of its strong latitudinal trend. DO and pH_w were not included because they had no significant relationships with land use. SUD_{cattle} is the combination of dairy and beef cattle.

	<i>HG</i>	<i>SG</i>	<i>NF</i>	<i>PF</i>	<i>OW</i>	SUD_{da}	SUD_{be}	SUD_{cattle}	SUD_{sh}	SUD_{de}	D_C	D_{HG}	D_{PF}
<i>COND</i>	0.57	-0.53	<i>NS</i>	0.53	<i>NS</i>	0.44	0.63	0.60	0.35	<i>NS</i>	<i>NS</i>	-0.25	<i>NS</i>
<i>CLAR</i>	-0.45	<i>NS</i>	0.28	-0.31	<i>NS</i>	-0.41	-0.49	-0.49	-0.40	<i>NS</i>	<i>NS</i>	<i>NS</i>	-0.27
<i>TURB</i>	0.46	<i>NS</i>	-0.27	0.28	<i>NS</i>	0.38	0.50	0.48	0.40	<i>NS</i>	<i>NS</i>	<i>NS</i>	<i>NS</i>
<i>CDOM</i>	0.56	-0.55	<i>NS</i>	0.24	-0.29	0.48	0.53	0.57	0.24	<i>NS</i>	<i>NS</i>	-0.33	<i>NS</i>
<i>TN</i>	0.82	-0.56	-0.37	0.46	-0.25	0.79	0.75	0.85	0.60	0.26	<i>NS</i>	-0.40	<i>NS</i>
NO_x	0.70	-0.53	-0.25	0.44	-0.25	0.77	0.65	0.79	0.51	0.28	<i>NS</i>	-0.39	<i>NS</i>
<i>TP</i>	0.66	-0.54	-0.32	0.48	<i>NS</i>	0.58	0.66	0.72	0.42	<i>NS</i>	<i>NS</i>	-0.24	<i>NS</i>
<i>DRP</i>	0.59	-0.65	<i>NS</i>	0.50	-0.43	0.58	0.58	0.66	0.31	<i>NS</i>	<i>NS</i>	-0.32	<i>NS</i>

Table 8. Stepwise regressions of water quality variables (median values) on landscape descriptors (forward selection, $p < 0.05$). Signs of coefficients indicate whether the relationship is proportional (+) or inverse (-). Int is model intercept. Scatterplots that characterize the primary and secondary explanatory variables are displayed in Figure 5.

Water Quality Variable	Step	Landscape Variable	Model Estimate	Multivariate sequential r^2
<i>CLAR</i>	1	<i>HG</i>	-0.03	0.17
	2	<i>OW</i>	0.18	0.27
	3	<i>Q₅₀</i>	-0.01	0.35
	4	<i>PF</i>	-0.03	0.39
	Int		3.16	
<i>TN</i>	1	<i>SUD_{cattle}</i>	77.05	0.62
	2	<i>HG</i>	4.26	0.68
	3	<i>PF</i>	5.16	0.69
	4	<i>SC%</i>	1.80	0.72
	Int		-33.95	
<i>NO_x</i>	1	<i>SUD_{cattle}</i>	86.15	0.58
	Int		62.65	
<i>TP</i>	1	<i>SUD_{cattle}</i>	5.47	0.41
	2	<i>PF</i>	0.64	0.52
	Int		7.75	
<i>DRP</i>	1	<i>SUD_{cattle}</i>	2.23	0.31
	2	<i>PF</i>	0.38	0.48
	Int		1.14	

Figures

Figure 1. Land use and location of the 77 National River Water Quality Network (NRWQN) catchments. Catchment ID colors refer to dominant land use (>50%). Catchments with no dominant land use are black.

Figure 2. Changes in land use areal coverage, livestock, and fertilizer inputs across New Zealand 1989/1990 vs. 2011/2012. Nitrogen fertilizers include urea and ammonium sulphate. Phosphorus fertilizers include superphosphate and diammonium phosphate.

Figure 3. Disturbance frequency of North Island per 463-m pixel, based on interpretation of MODIS data 2000-2013.

Figure 4. Disturbance frequency of South Island per 463-m pixel, based on interpretation of MODIS data 2000-2013.

Figure 5. Multivariate relationships between major water quality variables (median value for each site) and land use variables. For each plot, the primary explanatory variable from the stepwise regression (Table 8) is the x-axis, with bubble color representing the secondary explanatory variable. Note that oxidized nitrogen (NO_x) did not have a secondary explanatory variable. Selected catchments discussed in the text are labeled.

Figure 6. River water quality classes for upland (A) and lowland (B) catchments in New Zealand: I. clean river with high visual water clarity (CLAR) and low dissolved inorganic nutrients (DIN); II. sediment-impacted river with low CLAR and low DIN; III. nutrient-impacted river with high CLAR and high DIN; and IV. sediment- and nutrient-impacted river with low CLAR and high DIN. Classes are organized by ANZECC (2000) trigger values for water clarity (x-axis) and NO_x (y-axis). Catchments that exceed ANZECC guidelines for DRP are indicated in by grey-filled markers. Arrows indicate direction of trend over the 26 years inclusive from 1989 if significant (dashed) or meaningful (solid). No arrow means the trend was not significant.




Figure 1. Land use and location of the 77 National River Water Quality Network (NRWQN) catchments. Catchment ID colors refer to dominant land use (>50%). Catchments with no dominant land use are black.




Figure 2. Changes in land use areal coverage, livestock, and fertilizer inputs across New Zealand 1989/1990 vs. 2011/2012. Nitrogen fertilizers include urea and ammonium sulphate. Phosphorus fertilizers include superphosphate and diammonium phosphate.




Figure 3. Disturbance frequency of North Island per 463-m pixel, based on MODIS data 2000-2013.




Figure 4. Disturbance frequency of South Island per 463-m pixel, based on MODIS data 2000-2013.









Figure 5. Multivariate relationships between major water quality variables (median value for each site) and land use variables. For each plot, the primary explanatory variable from the stepwise regression (Table 8) is the x-axis, with bubble color representing the secondary explanatory variable. Note that oxidized nitrogen (NO_x) did not have a secondary explanatory variable. Selected catchments discussed in the text are labeled.

A. Upland Catchments



B. Lowland Catchments




Figure 6. River water quality classes for upland (A) and lowland (B) catchments in New Zealand: I. clean river with high water clarity (CLAR) and low dissolved inorganic nutrients (DIN); II. sediment-impacted river with low CLAR and low DIN; III. nutrient-impacted river with high CLAR and high DIN; and IV. sediment- and nutrient-impacted river with low CLAR and high DIN. Classes are organized by ANZECC (2000) trigger values. DIN trigger values can be discriminated for NO_x (y-axis) and DRP (grey-filled markers). Arrows indicate whether the trend from 1989-2014 was significant (dashed) or meaningful (solid). No arrow means the trend was not significant.