

Anonymous Referee #1 Received and published: 16 November 2016

General Comments:

This manuscript investigated the water footprint of crop production for different crop structures in the HSP based on the statistics data of crop yield, crop acreage, fertilization and water withdrawal in 2012. The water footprint was decomposed into blue water footprint, green water footprint, and grey water footprint. Eight different crop structure planning scenarios were used for the assessment of water footprint for different crop structure. Although in my opinion the subject of research is interesting and may be helpful for the water resource management in the HSP, there are several important issues need to be addressed. So I recommend a major revision. Major points:

Response: Thanks for the reviewer's comments. After our careful modification for more than two months, we resubmitted the manuscript. The responses of the comments are as follows,

1. The language of the manuscript needs to be improved, since some sentences are too long and not well expressed. I would suggest the manuscript refined by a native speaker.

Response: We invited an Elsevier editorial company to modify this manuscript, and the proof file was attached.

2. In my opinion, the result in section 3 is rather brief, which is not robust enough for the publication in this high-quality journal. The study of water footprint for only one year (2012) is obviously lack of persuasion. I suggest extending the length of time series (such as 5 or 10 years) to compare the interannual variability of water footprint in the HSP.

Response: This advice is good, we have extended the time series from 2000 to 2012 and analyzed the temporal variability of WF in the HSP. Over the past 13 years (2000-2012), the total WF of crop production in the HSP was 604.8 km<sup>3</sup>, comprised of 288.5 km<sup>3</sup> WF<sub>blue</sub>, 141.3 km<sup>3</sup> WF<sub>green</sub> and 175.0 km<sup>3</sup> WF<sub>gray</sub>, and decreased by 22% (from 53.7 km<sup>3</sup> to 41.8 km<sup>3</sup>), 26% (from 26.5 km<sup>3</sup> to 19.7 km<sup>3</sup>), 14% (from 11.7 km<sup>3</sup> to 10.1 km<sup>3</sup>), and 23% (from 15.5 km<sup>3</sup> to 12.0 km<sup>3</sup>), respectively, from 2000 to 2012 (Fig. 3). The main reasons for the downtrend of the WF was due to the urbanization of farmland and the decrease of the winter wheat planting area. In addition, the total WF<sub>blue</sub> of these crops was approximately twice the amount of the total WF<sub>green</sub>, and the total WF<sub>gray</sub> was slightly more than the total WF<sub>green</sub>.

3. The scenarios setting of crop structure has a large impact on the results. Why choose eight scenarios rather than ten scenarios in this study? My question is whether or not these eight scenarios represent all possibilities of the crop structure. In addition, why cotton and peanut are not involved in the scenarios setting (Table 2)? Do they show little impact on water footprint in the HSP? Please clarify it.

Response: Good question. Taking into consideration the crop structure change from 2000 to 2012, the high ground-water usage for rice and winter wheat per unit and the local residents' pasta-based diet, eight different crop structure planning scenarios were formulated with the cotton, peanut and side-crops cultivating areas unchanged.

4. The conclusion (section 5) is too simple and less appealing to the readers. Please re-organize this part to highlight your innovation and new findings.

Response: The conclusion was modified and summarized the findings of this study. "This study analyzed the WF of crop production in the HSP and evaluated its temporal variation from 2000 to 2012. Over 13 years, the production of main crops consumed a total of approximately 604.8 km<sup>3</sup> of water, of which 288.5 km<sup>3</sup> of that was groundwater; additionally, the WF of the production of crops exhibited a downtrend yearly. Among the local main crops, winter wheat, summer maize and vegetables were the three leading crops in water consumption; their WF, WF<sub>blue</sub>, WF<sub>green</sub> and WF<sub>gray</sub> accounted for 76.2%, 73.7%, 74.2% and 81.6% of the total, respectively.

In this region, adjusting crop farming structures has been an important means to protect groundwater resources; therefore, we evaluated reasonable farming structures by analyzing scenarios of the main crops' WF in this plain and suggest that: scenario 6 with approximately 20% of the arable land in cultivation of winter wheat-summer maize in rotation, 40% of spring maize, 10% of vegetables, 10% of fruiters, 0% of rice and no change to other crops, will promote the sustainable development of agriculture in this region. This scenario, not only can protect approximately 14.5% of groundwater resources (compared to the baseline), but can also ensure the local supply of wheat, vegetables, and fruits."

Specific Comments:

Page 2, line 30: "has becoming. . ." should be "has become. . ."

Response: Ok.

Page 2, line 44: what is the meaning of "As s metric. . ."?

Response: Metric should be "method".

Page 3, line 60: please give the full name of "HSP", since it first appeared in the introduction of the paper.

Page 3, line 77: "are located in . . ." » "is located in . . ."

Page 4, line 80: it is better to use "from July to September"

Response: The above problems were modified.

Page 4, line 88: please check the number of weather stations in Figure 1. It seems to me that only 22-23 stations can be found. Please add the id number to the stations in Figure 1.

Response: Thanks for the reviewer's carefulness, the weather stations is 21 in figure 1.

Page 7, line 138: please move the sentence "ET<sub>c</sub> is crop actual evapotranspiration (mm)" to the front of the sentence "Pe is the effective . . ."

Page 10, line 204: please change to “indicated that vegetables and winter wheat. . .”

Response: We corrected line 138 and 204, thanks a lot.

Anonymous Referee #2 Received and published: 23 January 2017

This is an interesting manuscript, and the discussion of the water footprint of each kind of crops is beneficial to design the current crop structure to save agricultural water consumption. In my opinion, it can be accepted after moderate revision. The specific comments are below:

Response: Thanks for the reviewer’s comments. We resubmitted the manuscript after our careful modification. The responses of the comments are as follows,

1. The newly published papers as reference should be added, the newest papers are 2015 papers in the reference list.

Response: Ok, we have added some newest papers, which were published in 2015 and 2016.

2. The conclusions should be enriched according to the research aims given at the end of the discussion section. The research result of the first aim is missing, and should be added in the conclusion section.

Response: Good idea, we modified the discussion section further and summarized the findings of this study. “This study analyzed the WF of crop production in the HSP and evaluated its temporal variation from 2000 to 2012. Over 13 years, the production of main crops consumed a total of approximately 604.8 km<sup>3</sup> of water, of which 288.5 km<sup>3</sup> of that was groundwater; additionally, the WF of the production of crops exhibited a downtrend yearly. Among the local main crops, winter wheat, summer maize and vegetables were the three leading crops in water consumption; their WF, WF<sub>blue</sub>, WF<sub>green</sub> and WF<sub>gray</sub> accounted for 76.2%, 73.7%, 74.2% and 81.6% of the total, respectively.

In this region, adjusting crop farming structures has been an important means to protect groundwater resources; therefore, we evaluated reasonable farming structures by analyzing scenarios of the main crops’ WF in this plain and suggest that: scenario 6 with approximately 20% of the arable land in cultivation of winter wheat-summer maize in rotation, 40% of spring maize, 10% of vegetables, 10% of fruiters, 0% of rice and no change to other crops, will promote the sustainable development of agriculture in this region. This scenario, not only can protect approximately 14.5% of groundwater resources (compared to the baseline), but can also ensure the local supply of wheat, vegetables, and fruits.”

3. The authors gave eight scenarios, why? The authors should give the reason to give eight scenarios.

Response: Reasonable question. Taking into consideration the crop structure change from 2000 to 2012, the high ground-water usage for rice and winter wheat per unit and the local residents’ pasta-based diet, eight different crop structure planning scenarios were formulated with the cotton,

peanut and side-crops cultivating areas unchanged.

4. In the discussion section, that 4.3 the main shortcomings of this study is just uncertainties of the results, not shortcoming, so the title should C1 HESSD Interactive comment Printer-friendly version Discussion paper be corrected.

Response: Ok.

5. The authors discussed the water footprint for specific crop types. However, I cannot find the data source of water consumptions of each type of crop in “2.2 data source” section. It should be given.

Response: The water consumption of each crop was calculated by the WF equations, and WF can reflect the water consumption. And the data of the structure of crops was added in this section.

1 **Water footprint of crop production for different crop structures in the**  
2 **Hebei southern plain, North China**

3 **Yingmin Chu<sup>a,b,c</sup>, Yanjun Shen<sup>d</sup>, Zaijian Yuan<sup>e\*</sup>**

4 <sup>a</sup> School of Management, China University of Mining & Technology, Xuzhou 221116, Jiangsu, P.R.China;

5 <sup>b</sup> Guangdong Science and Technology Library, Guangzhou, 510070, P.R.China;

6 <sup>c</sup> College of Tourism, Hebei University of Economics & Business, Shijiazhuang 050018, Hebei, P.R.China;

7 <sup>d</sup> Center for Agricultural Resources Research, Institute of Genetics and Developmental Biology, Chinese Academy  
8 of Sciences, Shijiazhuang, Hebei 050021, P. R. China;

9 <sup>e</sup> Guangdong Key Laboratory of Agricultural Environmental Pollution Integrated Control, Guangdong  
10 Institute of Eco-environment Technology, Guangzhou, 510650, P. R. China

11  
12 \* Corresponding author. Tel: +86 13722793672 E-mail address: selfsurpass@163.com

13 **Abstract**

14 The North China Plain (NCP) ~~is has~~ serious ~~lack shortage~~ of fresh water resources, ~~while and~~  
15 crop production consumes ~~sd approximately about~~ 75% of the region's water. To estimate water  
16 consumption of different crops and crop structures in the NCP, the Hebei southern plain (HSP) was  
17 selected as a study area, ~~as because~~ it is a typical region of groundwater overdraft in the NCP. In this  
18 study, ~~the~~ water footprint (WF) ~~of crop production, was being used which was consisted~~ comprised  
19 of green, blue and grey ~~components~~ water footprints, and its annual variation was analyzed. The  
20 results ~~showed demonstrated the following:~~ (1) the WF ~~from the of the main crops~~ production of  
21 main crops was ~~about 451.80~~ km<sup>3</sup> in 2012 ~~and w.~~ Winter wheat, ~~summer maize vegetables~~ and  
22 ~~vegetables summer maize~~ were ~~in~~ the top ~~three leading among the main~~ water-consuming crops in  
23 the HSP, ~~while t.~~ The water footprint intensity (WFI) of cotton was the largest, and ~~for~~ vegetables, ~~it~~  
24 ~~was were~~ the smallest; (2) The total WF, WFblue, WFgreen and WFgray for 13 years (2000-2012)  
25 of crops production were 604.8 km<sup>3</sup>, 288.5 km<sup>3</sup>, 141.3 km<sup>3</sup> and 175.0 km<sup>3</sup>, respectively, with an  
26 annual downtrend from 2000 to 2012; (3) ~~W~~ winter wheat, ~~summer maize~~ and vegetables consumed  
27 the ~~main-most~~ groundwater, and their blue water footprint (WFblue) accounted for ~~6674.20~~ % of the

带格式的: 上标

带格式的: 上标

带格式的: 上标

带格式的: 上标

28 total WFblue in the HSP; (34) ~~the~~The crop structure scenarios analysis indicated that, with ~~about~~  
29 ~~approximately~~ 20% of arable land ~~cultivating~~~~cultivated with~~ winter wheat-summer maize in  
30 rotation, 40% spring maize, 10% vegetables and 10% fruiters, ~~can promote the~~ sustainable  
31 utilization of groundwater resources ~~can be promoted, at the same time can ensure~~and a sufficient  
32 supply of food, ~~including~~ vegetables and fruits, ~~can be ensured~~ in the HSP.

33 **Keywords:** ~~The~~ Hebei southern plain; water footprint; crop production; crop structure; scenario  
34 analysis

### 35 1 Introduction

36 ~~With the~~Due to excessive ~~consumption~~~~water usage~~~~of water by people~~, freshwater scarcity has  
37 becoming a threat to human society (Dong et al., 2013). ~~In the w~~Worldwide, the largest freshwater  
38 consumer is agriculture, ~~which consumed~~~~consuming~~ more than 70% of ~~the~~ world's freshwater  
39 (UNEP, 2007; Lucrezia et al., 2014). ~~Water resources have been heavily exploited by agriculture~~  
40 ~~worldwide (Konar et al., 2011) and t~~To ensure the increasing food demand, global water  
41 consumption ~~have~~~~has~~ almost doubled during the past 40 years (Gleick, 2003), and ~~water resources~~  
42 ~~have been heavily exploited for worldwide agriculture (Konar et al., 2011). In t~~Future, water use for  
43 food production will continue to ~~meet the~~be influenced by population growth and changes in dietary  
44 preferences (Rosegrant and Ringler, 2000). ~~This, which will lead to the consumptione of~~ more  
45 water resources. China is a freshwater ~~poor~~ country with ~~about~~~~approximately~~ 2100 m<sup>3</sup>/y ~~of~~ water  
46 resources per capita, accounting for only 28% of the world's per capita share. The spatial mismatch  
47 between water and arable land ~~strengthened~~~~reinforces~~ China's water challenge. ~~There is a~~About  
48 70% arable land in the north of the Yangze River ~~with~~~~contains~~ only ~~about~~~~approximately~~ 17% ~~of~~  
49 ~~the national total~~ water resources ~~of the national total in China. Due to the c~~Currently, as a water  
50 shortage ~~in the~~ area north of the Yangze River, the NCP is facing ~~the acutest its most severe~~ water  
51 scarcity issue, ~~The NCP presently contains~~accounting for only 1.3% of China's total available  
52 water with 225 m<sup>3</sup>/y per capita (White et al., 2015).

53 As ~~s a~~method~~ie~~ to assess ~~the~~ water use of ~~the~~ production systems, the water footprint (WF)

54 concept ~~has was been~~ proposed (Hoekstra, 2003), which includes ~~ed~~ direct and indirect water usage ~~ee~~  
55 of a consumer or producer (Hoekstra et al., 2009). In recent years, many researchers have used the  
56 WF to evaluate water use in agricultural production (Bocchiola et al., 2013; Chapagain and Hoekstra,  
57 2011; Chapagain and Orr, 2009; Gheewala et al., 2014; Jefferies et al., 2012; Lamastra et al., 2014;  
58 Mekonnen and Hoekstra, 2010, Shrestha et al., 2013; Wang et al., 2014; Xu et al., 2014; Zang et  
59 al., 2014; Wang et al., 2015; Suttayakul et al., 2016). The WF of crops reflects the water  
60 consumption of different crops, and ~~it can be~~ focused on local crop products. For ~~a certain~~ each crop,  
61 the blue WF (WFblue) refers to the volume of irrigation water ~~consumption~~ consumed, the green  
62 WF (WFgreen) is consistent with the effective rainfall for plants, and the grey WF (WFgrey)  
63 represents the volume of water required to dilute pollutants to the agreed maximum acceptable  
64 levels (Hoekstra and Chapagain, 2007). ~~For~~ Since the water consumption of each crop is different,  
65 the WF ~~of for~~ different crops ~~differ~~ varies greatly. Xu et al. (2014) analyzed the WF of six kinds of  
66 crops in Beijing from 1978 to 2012, and found maize accounts for 57% of the green WF and 46% of  
67 the grey WF ~~respectively~~, vegetables account for 45% of the blue WF, and wheat accounts for 26%  
68 of the total WF. Wang et al. (2015) found that winter wheat conserved ~~about~~ approximately  $1.9 \times$   
69  $10^9 \text{ m}^3 \text{ yr}^{-1}$  of WFblue ~~during from~~ 1998 ~~to~~ 2011 in the Hebei Plain.

70 The Hebei southern plain (HSP) was selected as the study area. It is located ~~at in~~ the northwest of  
71 the NCP ~~with about~~ and has approximately  $4.0 \times 10^4 \text{ km}^2$  of arable land (accounting for ~~about~~  
72 approximately 13% of the NCP and 3% of China's total). ~~In 2008, the HSP which~~ produced ~~about~~  
73 approximately  $2.7 \times 10^{10} \text{ kg}$  of grain ~~yield~~ (accounting for ~~about~~ approximately 5% of China's total)  
74 ~~with that had~~ a water consumption ~~about~~ approximately  $3.0 \times 10^{10} \text{ m}^3$  ~~in 2008~~ (Yuan and Shen,  
75 2013). The over-exploitation of groundwater in this region has had devastating consequences, ~~with~~  
76 the groundwater table ~~being has~~ decreased by more than 20 m within recent the past 30 years (Chen  
77 et al., 2003; Hu and Cheng, 2011). Because the WF of various crops is different and the crop  
78 structure of a region reflects the proportion of various crops growing ~~areas within that~~ region, the  
79 WF of the crop structure can illustrate the ~~whole entire~~ agricultural water consumption of ~~the that~~

80 region. ~~Study of the WF for crop structures can help to promote the sustainable utilization of water~~  
81 ~~resources for agriculture in the water shortage area.~~ The study of the WF for crop structures can help  
82 promote the sustainable utilization of water resources for agriculture, and can be particularly  
83 valuable for areas facing water shortage.

84 The main aims of this study were: (1) to quantify the WF of production of main crops ~~production~~  
85 in the HSP in 2012; ~~and~~ (2) to discuss ~~the a~~ reasonable crop structure based on the WF analysis  
86 for different crop structure scenarios. ~~Through~~ In this study, we propose a ~~most~~-suitable crop  
87 planting structure for this region, and ~~give~~ support ~~to~~ the development and implementation of  
88 policies on agricultural water management.

## 89 **2 Materials and methods**

### 90 2.1 Study area

91 The Hebei southern plain (114°20'E-119°25'E, 36°03'N-39°56'N), with an area of  
92 ~~about~~approximately 62,000 km<sup>2</sup>, are located in southern Beijing and Tianjin (Fig. 1). The climate in  
93 this region is temperate ~~continental~~ monsoon with a mean annual precipitation of 550 mm and a  
94 mean annual temperature of 11.5 °C. Precipitation has a non-uniform distribution throughout the  
95 year, and ~~about~~approximately 80% of the total precipitation occurs from July through September. In  
96 the HSP, most arable lands are irrigated by groundwater except ~~for~~ in the eastern part where there is  
97 saline shallow groundwater ~~restrains the irrigation~~. The ~~main~~ primary crops in the plain are wheat,  
98 maize (including summer maize and spring maize), cotton, and peanut; ~~the~~ main vegetable ~~species~~  
99 crops are Chinese cabbage, celery, cauliflower, onion, bean, rape, leek, coriander, fennel, and the  
100 main fruits are apple, pear, jujube and grape.





**Fig. 1 Location of the Hebei southern plain**

2.2 Data collection

The meteorological data ~~for~~ from 25-21 weather stations (Fig. 1) around the HSP, including daily maximum temperature, minimum temperature, average temperature, wind speed, relative humidity, precipitation, sunshine duration, vapor pressure, and atmospheric pressure, were obtained from the China Meteorological Data Sharing Service System (China Meteorological Administration, 2000-2012).

The statistics data for the plain in 2012, including crop yield, crop acreage and fertilization, were obtained from Hebei economic statistical yearbooks; and the data for water withdrawal were obtained from the water resources bulletins and relevant statistical yearbooks. The ~~l~~-and-use map ~~in~~ of the HSP for 2012 (Fig. 2) ~~of the plain were~~ was drawn based ~~on the off of~~ spot satellite images and ~~the a~~ topographic map (1:10000). The main land-use types include cropland, ~~construction land~~ urban, forestland, waters, orchard, wetland, grassland and shrub land (Table. 1).




Fig. 2 Land-use map of the Hebei southern plain

Table 1 Area of each land-use type and their ratios (%)

Land-use	Forestland	Shrub land	Grassland	Cropland	Orchard	Building land	Waters	Wetland	Total
Area( $10^5$ hm <sup>2</sup> )	3.66	0.31	0.72	42.80	2.61	7.02	2.91	1.83	61.85
%	5.91	0.49	1.17	69.20	4.21	11.35	4.70	2.95	100

The crop structure data were produced based on remote sensing data for this study area from 2000 to 2012 (Table. 2), which included MODIS NDVI (MOD13Q1), Terra/MODIS (MOD12Q1), and Landsat TM/ETM with spatial resolutions of 250 m, 1000 m and 30 m, respectively. Pan et al. (2015) and Wang et al. (2015) presented the details of this method. Compared with 2000, the crop planting area changed considerably; specifically, the planting area of winter maize-summer maize decreased by 34.76%, rice decreased by 31.61%, spring maize increased by 34.13%, vegetables increased by 26.05%, and fruiters increased by 33.04%, while cotton, peanut and others had a slight change, and the total cultivated area in HSP decreased by 12.58% in 2012 (Table. 2).

Table 2 Planting areas ( $10^5$  hm<sup>2</sup>) for the main crops and their percent change

Year	Winter wheat-summer maize	Spring maize	Vegetables	Fruiters	Cotton	Peanut	Rice	Others	Total
------	---------------------------	--------------	------------	----------	--------	--------	------	--------	-------

2000	29.54	3.89	6.72	1.96	5.59	2.42	0.44	1.38	51.94
2001	25.20	3.67	7.46	2.05	5.15	2.69	0.49	1.54	48.23
2002	22.79	3.91	6.85	2.26	7.27	2.47	0.45	1.41	47.41
2003	24.40	3.76	7.09	2.03	2.82	3.26	0.39	1.87	45.63
2004	24.11	4.89	7.27	1.86	3.45	2.86	0.34	1.64	46.42
2005	24.64	3.40	7.20	2.15	5.04	2.59	0.31	1.48	46.82
2006	24.69	4.41	6.96	1.76	4.24	2.51	0.30	1.43	46.31
2007	22.37	5.25	6.89	2.14	6.99	2.48	0.45	1.42	48.00
2008	24.31	4.18	7.43	2.36	4.62	2.68	0.32	1.53	47.43
2009	25.19	3.64	7.25	2.25	3.74	2.61	0.31	1.49	46.49
2010	23.24	4.85	7.20	2.12	3.99	2.59	0.31	1.48	45.79
2011	20.65	4.36	7.54	1.94	5.74	2.72	0.33	1.55	44.83
2012	19.27	5.22	8.47	2.61	5.61	2.50	0.30	1.43	45.41
Change	-34.76	34.13	26.05	33.04	0.39	2.64	-31.61	3.27	-12.58

129 2.3 Crop structure scenarios setting

130 The baseline for the crop structure (2012) in the HSP, consisted of 42.44% of winter  
131 wheat-summer maize rotation, 11.50% of spring maize, 18.65% of vegetables, 5.75% of fruiters,  
132 12.35% of cotton, 5.51% of peanut, 0.66% of rice, and 3.15% of others (side crops i.e., millet,  
133 sorghum, sweet potato and others). Taking into consideration the crop structure change from 2000  
134 to 2012, the high ground-water usage for rice and winter wheat per unit and the local residents'  
135 pasta-based diet. Eight different crop structure planning scenarios were formulated according to  
136 the main crops of the baseline with the cotton, peanut and side-side crops cultivating areas  
137 unchanged (Table 23). These scenarios involved reducing winter wheat-summer maize and rice  
138 cultivating area to 40% and 0% separately respectively; and increasing spring maize cultivating area  
139 to 13.94% (scenario 1); reducing winter wheat-summer maize to 30% and increasing spring maize  
140 to 23.94% (scenario 2); reducing winter wheat-summer maize to 20% and increasing spring maize  
141 to 33.94% (scenario 3); reducing winter wheat-summer maize to 10% and increasing spring maize  
142 to 43.94% (scenario 4); reducing winter wheat-summer maize to 0 and increasing spring maize to  
143 53.94% (scenarios 5); reducing winter wheat-summer maize to 20% and increasing spring maize to  
144 38.99%, and adjusting vegetables and fruiters to 10% (scenario 6); reducing winter wheat-summer  
145 maize to 20%, and increasing spring maize to 28.99%, vegetables to 20% and fruiters to 10%  
146 (scenario 7); reducing winter wheat-summer maize to 20% and increasing spring maize to 28.99%,  
147 decreasing vegetables to 10% and increasing fruiters to 20% (scenario 8).

**Table 2-3 Crop structure planning scenarios for the Hebei southern plain**

Crop structure		Winter wheat- summer maize	Spring maize	Vegetables	Fruiters	Cotton	peanut	Rice	Others	Total
Baseline	Area(10 <sup>5</sup> hm <sup>2</sup> )	19.27	5.22	8.47	2.61	5.61	2.50	0.30	1.43	45.41
	%	42.44	11.50	18.65	5.75	12.35	5.51	0.66	3.15	100
Scenario 1	Area(10 <sup>5</sup> hm <sup>2</sup> )	18.16	6.33	8.47	2.61	5.61	2.50	0	1.43	45.41
	%	40.00	13.94	18.65	5.75	12.35	5.51	0	3.15	100
Scenario 2	Area(10 <sup>5</sup> hm <sup>2</sup> )	13.62	10.87	8.47	2.61	5.61	2.50	0	1.43	45.41
	%	30.00	23.94	18.65	5.75	12.35	5.51	0	3.15	100
Scenario 3	Area(10 <sup>5</sup> hm <sup>2</sup> )	9.08	15.41	8.47	2.61	5.61	2.50	0	1.43	45.41
	%	20.00	33.94	18.65	5.75	12.35	5.51	0	3.15	100
Scenario 4	Area(10 <sup>5</sup> hm <sup>2</sup> )	4.54	19.95	8.47	2.61	5.61	2.50	0	1.43	45.41
	%	10.00	43.94	18.65	5.75	12.35	5.51	0	3.15	100
Scenario 5	Area(10 <sup>5</sup> hm <sup>2</sup> )	0	24.49	8.47	2.61	5.61	2.50	0	1.43	45.41
	%	0	53.94	18.65	5.75	12.35	5.51	0	3.15	100
Scenario 6	Area(10 <sup>5</sup> hm <sup>2</sup> )	9.08	17.71	4.54	4.54	5.61	2.50	0	1.43	45.41
	%	20.00	38.99	10.00	10.00	12.35	5.51	0	3.15	100
Scenario 7	Area(10 <sup>5</sup> hm <sup>2</sup> )	9.08	13.16	9.08	4.54	5.61	2.50	0	1.43	45.41
	%	20.00	28.99	20.00	10.00	12.35	5.51	0	3.15	100
Scenario 8	Area(10 <sup>5</sup> hm <sup>2</sup> )	9.08	13.16	4.54	9.08	5.61	2.50	0	1.43	45.41
	%	20.00	28.99	10.00	20.00	12.35	5.51	0	3.15	100

149 2.4 *WF evaluation*

150 The WF of a crop production is the sum of the green, blue and grey components-water footprints  
151 (Chapagain et al., 2006). The WF of 7-seven main primary type kinds of crops planted in the HSP is  
152 calculated separately:

$$153 \quad WF = \sum_{a=1}^n WF_a \quad (1)$$

$$154 \quad WF = WF_{blue} + WF_{green} + WF_{grey} \quad (2)$$

155 where  $WF$  is the total water footprint ( $m^3 yr^{-1}$ );  $WF_a$  is the water footprint of each type of crop in the  
156 Hebei plain;  $WF_{blue}$  is the blue water footprint ( $m^3 yr^{-1}$ );  $WF_{green}$  is the green water footprint ( $m^3$   
157  $yr^{-1}$ ), and  $WF_{grey}$  is the grey water footprint ( $m^3 yr^{-1}$ ).

158 The WF intensity ( $WFI$ ) of a crop production is evaluated by dividing  $WF$  with crop yield:

$$159 \quad WFI_a = WF_a / Y_a \quad (3)$$

160 where  $WFI_a$  is the WF intensity of a certain crop ( $m^3 ton^{-1}$ ) and  $Y_a$  is the yield of that kind of crop  
161 (ton).

162 ~~Green-The green~~ water footprint was represented by crop evaporation or effective rainfall:

$$163 \quad WF_{blue} = 10 \times ET_{blue} \times A \quad (4)$$

164  $WF_{green} = 10 \times ET_{green} \times A$  (5)

165  $ET_{blue} = \max\{0, ET_c - P_e\}$  (6)

166  $ET_{green} = \min\{P_e, ET_c\}$  (7)

167 where  $ET_{blue}$  is the blue water evapotranspiration during the growth period of crops (mm);  $ET_{green}$  is  
 168 green water evapotranspiration (mm);  $A$  is the acreage of the ~~calculating-calculated~~ crop (hm<sup>2</sup>);  $ET_c$   
 169 is the actual crop evapotranspiration (mm);  $P_e$  is the effective precipitation (mm), which can be  
 170 calculated using ~~the~~ Soil Conservation Service Method developed by the U.S. Department of  
 171 Agriculture (USDA);  ~~$ET_c$  is crop actual evapotranspiration (mm).~~

172 
$$P_e = \begin{cases} P \times (125 - 0.6P) / 125 & P \leq 250 / 3 \\ 125 / 3 + 0.1P & P > 250 / 3 \end{cases}$$
 (8)

173 where  $P$  is the precipitation (mm).

174  $ET_c$  can be calculated based on the reference evapotranspiration ( $ET_0$ ) which is estimated  
 175 according to the FAO56-PM model (Allen et al., 1998);

176  $ET_c = K_c \times ET_0$  (9)

177 
$$ET_0 = \frac{0.408\Delta(R_n - G) + \gamma \frac{900}{T_{em} + 273} u_2 (e_s - e_a)}{\Delta + \gamma(1 + 0.34u_2)}$$
 (10)

178 where  $K_c$  is the crop coefficient, and the  $K_c$  of the crops was determined according to their growing  
 179 stage (Duan, 2004);  $R_n$  is the net radiation at the vegetation surface (MJ m<sup>-2</sup> d<sup>-1</sup>);  $G$  is the soil heat  
 180 flux density (MJ m<sup>-2</sup> d<sup>-1</sup>);  $T_{em}$  is the daily average temperature (°C);  $u_2$  is the wind speed at a 2 m  
 181 height (m s<sup>-1</sup>);  $e_s$  is the vapor pressure of the air at saturation (kPa);  $e_a$  is the actual vapor pressure  
 182 (kPa);  $\Delta$  is the slope of the vapor pressure curve (kPa °C<sup>-1</sup>); and  $\gamma$  is the psychrometric constant  
 183 (kPa °C<sup>-1</sup>). A complete set of equations is proposed by Allen et al. (1998) to compute the variables  
 184 in Eq. (10) according to available weather data and the time step computation, which constitute the  
 185 FAO-PM method.  $G$  can be ignored for daily time step computations.

186 ~~For~~ Due to a ~~lack~~ of ~~accessed-accessible~~ data, the grey WF of crops only ~~assimilate-assesses~~  
 187 nitrogen contamination without considering the effect of pesticides and other fertilizers, ~~which and~~

was calculated ~~as by~~ the following equation (Hoekstra et al., 2009):

$$WF_{grey} = (\delta \times U_N \times 10^6) / \rho_0 \quad (11)$$

where  $U_N$  is the applied amount of N fertilizer (ton).  $\delta$  represents the leaching rate to freshwater with values 5-15% (Zhang and Zhang, 1998) and we use ambient water quality standard for nitrogen (10 mg L<sup>-1</sup>) as the permissible concentration ( $\rho_0$ ). Due to a lack of ~~accessed-accessible~~ data, we ignored pesticides and other fertilizer ~~heres~~.

### 3 Results

#### 3.1 WF and WFI of crop production in 2012

~~In 2012, the total~~The WF of crops production in 2012 was analyzed, and the results were taken as the baseline for the crop structure analysis. The total WF of the production of crops in the HSP was ~~about~~approximately ~~5141.08~~ km<sup>3</sup>, of which ~~27.64%~~ was WFgreen (~~1410.1~~ km<sup>3</sup>), ~~48.47%~~ was WFblue (~~2419.7~~ km<sup>3</sup>) and ~~24.09%~~ was WFgrey (~~12.20~~ km<sup>3</sup>), ~~respectively~~ (Table 34). We found large differences ~~of among~~ the WF, WFgreen, WFblue and WFgrey ~~within for~~ the main crops: ~~among these crops (wheat, maize, cotton, peanut, rice, vegetables, fruiters), w.~~ Winter wheat, summer maize and vegetables and summer maize were the three leading crops in water consumption, taking ~~28.89%~~ (~~1412.70~~ km<sup>3</sup>), ~~23.84%~~ (~~1310.1~~ km<sup>3</sup>) and ~~19.38%~~ (~~107.76~~ km<sup>3</sup>) of the total WF, respectively. ~~The WF of spring maize, cotton, peanut, rice, fruiters and others was were~~ ~~32.90~~ km<sup>3</sup> (~~5.47%~~), ~~43.08~~ km<sup>3</sup> (~~7.29%~~), 1.6 km<sup>3</sup> (~~2.94%~~), 0.3 km<sup>3</sup> (~~0.61%~~), ~~2.76~~ km<sup>3</sup> (~~4.96%~~) and ~~10.90~~ km<sup>3</sup> (~~1.82%~~), respectively. The WFgreen of these crops was ~~2.01~~ km<sup>3</sup> (~~accounted-accounting~~ ~~for 14.221%~~ of the total WFgreen), ~~4.4-3.6~~ km<sup>3</sup> (~~31.5%~~), ~~10.94~~ km<sup>3</sup> (~~10.09%~~), ~~1.70.9~~ km<sup>3</sup> (~~12.09%~~), ~~0.7.5~~ km<sup>3</sup> (~~4.95%~~), 0.1 km<sup>3</sup> (~~0.71%~~), ~~31.0.3~~ km<sup>3</sup> (~~21.013%~~), ~~0.6.5~~ km<sup>3</sup> (~~4.95%~~), and 0.2 km<sup>3</sup> (~~1.42%~~), respectively, ~~in of~~ which ~~vegetables summer maize~~ was the largest, ~~then was summer maize followed by winter wheat~~. The WFblue of these crops was ~~75.8.1~~ km<sup>3</sup> (~~accounted-accounting~~ ~~for 31.826%~~ of the total WFblue), ~~3.3.7~~ km<sup>3</sup> (~~13.49%~~), ~~01.9.3~~ km<sup>3</sup> (~~5.47%~~), ~~12.6.1~~ km<sup>3</sup> (~~7.211%~~), ~~0.7.8~~ km<sup>3</sup> (~~2.94%~~), 0.2 km<sup>3</sup> (~~0.61%~~), ~~84.5.9~~ km<sup>3</sup> (~~23.85%~~), 1.1 km<sup>3</sup> (~~4.96%~~), ~~0.6.5~~ km<sup>3</sup> (~~1.83%~~), respectively, ~~in of~~ which ~~vegetables winter wheat~~ was the largest, ~~then was winter wheat followed~~

214 by vegetables. The WFgrey of these crops was 4.8 km<sup>3</sup> (accounting for ~~39.440%~~ of the total  
 215 WFgrey), 2.9 km<sup>3</sup> (~~23.74%~~), 0.7 km<sup>3</sup> (~~5.4%~~), 0.7 km<sup>3</sup> (~~5.7%~~), ~~0.3-2~~ km<sup>3</sup> (~~2.9%~~), 0.1 km<sup>3</sup> (~~0.51%~~),  
 216 ~~1.7-5~~ km<sup>3</sup> (~~13.93%~~), 1.0 km<sup>3</sup> (~~8.0%~~), 0.2 km<sup>3</sup> (~~1.52%~~), respectively, ~~in of~~ which winter wheat was  
 217 the largest, ~~then followed by was~~ summer maize.


218 The situation of the WFI was totally different from the situation of the WF (Table 34). Among  
 219 these crops, the WFI of cotton was the largest and for vegetables, it was the smallest, and the  
 220 former was about approximately eight-six times as much as that of greater than the latter; the WFI of  
 221 winter wheat summer maize was basically equal to which that of peanuts spring maize.

222 **Table 3-4 The WF (km<sup>3</sup>) and the WFI (m<sup>3</sup> ton<sup>-1</sup>) of each crop**

Crop types	WFgreen	WFblue	WFgrey	WF	WFI
Winter wheat	<del>2.01</del>	<del>75.81</del>	4.8	<del>1412.70</del>	<del>4086887.90</del>
Summer maize	<del>3.644</del>	<del>3.733</del>	<del>2.929</del>	<del>10.1406</del>	<del>701.77360</del>
Spring maize	<del>1.314</del>	<del>1.309</del>	<del>0.707</del>	<del>2.930</del>	<del>691.17091</del>
Cotton	<del>0.917</del>	<del>2.116</del>	<del>0.707</del>	<del>3.840</del>	<del>1493.015733</del>
Peanut	<del>0.507</del>	<del>0.807</del>	<del>0.203</del>	<del>1.616</del>	<del>1043.210828</del>
Rice	<del>0.101</del>	<del>0.202</del>	<del>0.101</del>	<del>0.403</del>	<del>903.19139</del>
Vegetables	<del>1.330</del>	<del>4.985</del>	<del>1.517</del>	<del>7.7131</del>	<del>183.62077</del>
Fruiters	<del>0.506</del>	<del>1.111</del>	<del>1.010</del>	<del>2.627</del>	<del>246.42575</del>
Others	<del>0.202</del>	<del>0.506</del>	<del>0.202</del>	<del>0.910</del>	<del>1030.811470</del>
Total	<del>10.1141</del>	<del>19.7247</del>	<del>12.0122</del>	<del>41.8510</del>	

223 3.2 Annual WF of crop production

224 Over the past 13 years (2000-2012), the total WF of crop production in the HSP was 604.8 km<sup>3</sup>,  
 225 comprised of 288.5 km<sup>3</sup> WFblue, 141.3 km<sup>3</sup> WFgreen and 175.0 km<sup>3</sup> WFgray, and decreased by  
 226 22% (from 53.7 km<sup>3</sup> to 41.8 km<sup>3</sup>), 26% (from 26.5 km<sup>3</sup> to 19.7 km<sup>3</sup>), 14% (from 11.7 km<sup>3</sup> to 10.1  
 227 km<sup>3</sup>), and 23% (from 15.5 km<sup>3</sup> to 12.0 km<sup>3</sup>), respectively, from 2000 to 2012 (Fig. 3). The main  
 228 reasons for the downtrend of the WF was due to the urbanization of farmland and the decrease of  
 229 the winter wheat planting area. In addition, the total WFblue of these crops was approximately  
 230 twice the amount of the total WFgreen, and the total WFgray was slightly more than the total  
 231 WFgreen.



**Fig. 3. WF of crop production in the HSP from 2000 to 2012**

Table 5 shows the WF of each crop over 13 years. Winter wheat, summer maize and vegetables are the three leading crops for water usage, taking 33%, 28% and 16% of the total WF, respectively. Notably, summer maize accounted for 42% of WFgreen, 22% of WFblue and 27% of WFgrey; winter wheat accounted for 21% of WFgreen, 31% of WFblue and 44% of WFgrey; and vegetables accounted for 11% of WFgreen, 22% of WFblue and 11% of WFgrey (Table 5).

**Table 5 WF (km³) of each crop in the HSP from 2000 to 2012**

Crop types	WFgreen	WFblue	WFgrey	WF
Winter wheat	30.0	89.1	77.6	196.7
Summer maize	58.8	62.6	46.6	168.0
Spring maize	10.5	13.8	6.9	31.2
Cotton	11.9	24.4	8.0	44.3
Peanut	6.2	12.8	3.4	22.4
Rice	0.8	3.5	0.9	5.2
Vegetables	15.4	62.3	18.7	96.4
Fruiters	4.8	12.5	10.3	27.7
Others	2.8	7.5	2.6	12.9
Total	141.4	288.5	175.1	604.8

### 3.2.3 Scenario analysis of WF for different crop structure

Results from the ~~seven-eight~~ scenarios (Table 4~~6~~) illustrated ~~that~~that the following: (1) the WF (~~including-comprised of~~ WFgreen, WFblue and WFgrey) of all the scenarios was smaller than the baseline, and those of scenario 5 were the smallest in the eight scenarios; (2) the WF of scenario 3 and scenario ~~8-6 was-were~~ essentially equal, ~~and which of as were~~ scenario 7 was slightly larger than



245 ~~them and scenario 6 was slightly larger than~~ scenario 4~~8~~; (3) the WF (~~including WFGreen, WFblue~~  
 246 ~~and WFGrey) was getting smaller and smaller~~reduced from scenario 1 to scenario 5 ~~with as~~ the  
 247 planting area of winter wheat and summer maize rotation decreased to zero and spring maize  
 248 increased to 53.94%~~-%~~; (4) the WFGreen of ~~the scenario 2,3,6,7 and scenario 8 were all the~~  
 249 ~~scenarios was~~ nearly equal, and the value was approximately ~~12.9~~9 km<sup>3</sup>.

250 **Table 4-6 WF (km<sup>3</sup>) of different crop structure scenarios in the Hebei southern plain**

Crop structure	WFGreen	WFblue	WFGrey	WF
Baseline	<del>10.144.4</del>	<del>19.724.7</del>	<del>12.042.2</del>	<del>41.854.0</del>
Scenario 1	<del>9.912.7</del>	<del>19.224.2</del>	<del>11.711.9</del>	<del>40.848.7</del>
Scenario 2	<del>9.412.4</del>	<del>18.322.2</del>	<del>10.410.6</del>	<del>38.145.3</del>
Scenario 3	<del>8.912.1</del>	<del>17.420.4</del>	<del>9.29.4</del>	<del>35.441.9</del>
Scenario 4	<del>8.411.8</del>	<del>16.418.5</del>	<del>7.98.1</del>	<del>32.738.5</del>
Scenario 5	<del>7.911.5</del>	<del>15.616.7</del>	<del>6.76.9</del>	<del>30.335.1</del>
Scenario 6	<del>9.112.4</del>	<del>16.817.7</del>	<del>9.69.6</del>	<del>35.539.6</del>
Scenario 7	<del>9.112.1</del>	<del>18.620.1</del>	<del>9.910.3</del>	<del>37.642.6</del>
Scenario 8	<del>9.212.2</del>	<del>17.618.8</del>	<del>10.710.7</del>	<del>37.541.7</del>

带格式表格

## 251 4 Discussions

### 252 4.1 Crop water consumption

253 In the HSP, ~~irrigation water has been the primary source of water for agricultural needs~~~~the~~  
 254 ~~agricultural water consumption mainly came from irrigation~~ (Yuan and Shen, 2013), ~~which was~~  
 255 ~~confirmed this study and this study also proved this point~~. According to the above analysis, ~~the~~ water  
 256 consumption of ~~the~~ crops (~~except maize, cotton and peanut~~) mainly came from irrigation, and their  
 257 WFblue accounted for ~~about~~approximately 50% of the WF (Table ~~34 and Table 5~~). Although  
 258 irrigation can directly increase ~~yield of the crop yields~~, it also usually ~~increased~~increases the crop  
 259 WF (da Silva et al., 2013). In ~~areas of~~ water shortage ~~area~~, improving water use efficiency to reduce  
 260 groundwater exploitation is imperative, ~~and~~ ~~deficit~~Deficit irrigation ~~was~~has been widely used to  
 261 save groundwater resources in the NCP (Ma et al., 2013), ~~which by~~ ~~took~~taking better account of  
 262 crop yield and water consumption.

263 During the 13 years, ~~the~~ WFblue of ~~vegetables winter wheat~~ was the largest ~~in of~~ these crops,  
 264 ~~and then was winter wheat~~ followed by summer maize, and then vegetables; which ~~indicated~~  
 265 ~~indicates that winter wheat, summer maize and~~ vegetables ~~and winter wheat~~ consumed a large  
 266 amount of groundwater. The WFblue of the crops, apart from summer maize and spring maize, was

267 ~~more than double their WFgreen; furthermore, the WFblue of rice and vegetables was more than~~  
268 ~~quadruple their WFgreen.~~The WFgreen of ~~both summer maize and spring maize were~~  
269 ~~approximately more than its WFblue, and the WFgreen of cotton and peanut was approximately~~  
270 ~~equal to their WFblue. This was,~~ because the rapid growth stage of ~~these four cropmaizes~~ was ~~from~~  
271 ~~June to August, this period was~~ basically synchronized with ~~the~~ rainy season (July to September) in  
272 this region, and the precipitation ~~can basically was able to~~ meet the needs of crop growth in this  
273 period. ~~So-Therefore,~~ in arid and ~~semi-semi~~-arid areas, cultivating rain fed crops is an effective  
274 approach to save groundwater. While for ~~wheat, rice, vegetables, fruiters and other crops,~~ the  
275 ~~precipitation cannot meet their needs~~WFblue was significantly more than their WFgreen. ~~The main~~  
276 ~~reason was the precipitation can not meet the needs of these crops, and the; therefore,~~ water  
277 ~~consumption of~~for these crops ~~needs to come mainly~~mainly came from irrigation.

#### 278 4.2 WF responses to crop structure

279 Crop structure affects the water consumption directly. ~~From-~~The above analysis ~~shows that,~~ with  
280 the decrease of winter wheat-summer maize rotation planting area and the increase of spring maize  
281 (scenario 1 to scenario 5), the WF (~~including-comprised of~~ WFgreen, WFblue and WFGrey)  
282 decreased (Table 4(6)-). ~~S~~pecifically, when the area of winter wheat-summer maize decreased 10%  
283 and spring maize increased 10% (~~relative to the total farmland area~~), the average WF, WFgreen,  
284 WFblue and WFGrey decreased 7.92%, 25.54%, 85.91% and 12.78%, respectively. However,  
285 ~~people consumed flour as the major~~since wheat is a staple food in the HSP and ~~wheat is~~a ration  
286 crop ~~here, and this region needs to guarantee~~we should plant a certain area of winter wheat to  
287 ~~guarantee the~~food self-sufficiency ~~in this region,~~ areas should still be planted with winter wheat,  
288 ~~despite in spite of it consumed a lot of water resourees~~its large consumption of water resources. ~~In~~  
289 ~~per unit area, the water consumption of v~~vegetables ~~had a low-level WFI; however, the water~~  
290 ~~consumption of vegetables per unit area,~~ was much more than ~~with~~ other crops (scenario 6 to  
291 scenario 7) ~~in spite of its WFI was low level. Despite this, but~~ the HSP should protect the basic  
292 supply of vegetables and fruits for Beijing, Tianjin and ~~the~~ Hebei province, ~~p~~. Planting ~~and keeping~~

293 a certain areas ~~of-with~~ vegetables and fruiters is necessary.

294 ~~Changes to c~~Crop structure ~~changing~~ directly affects irrigation ~~water~~ consumption (or WFblue)  
295 and indirectly affects the emissions of environmental pollutants ~~which that were closely linked~~  
296 ~~with~~can be measured by WFGrey. In the study area, ~~crop~~ water consumption ~~for crops is primarily~~  
297 ~~attributable to mainly came from~~ groundwater irrigation. ~~i~~ It is ~~imperative to an urgency to~~  
298 ~~find~~identify out a reasonable crop structure by considering the sustainable use of groundwater and  
299 ~~the lifestyle of~~ local people's ~~daily life~~. According to the above scenario analysis, we found the crop  
300 structure of scenario 6 ~~was to be~~ reasonable. Because this structure can guarantee ~~the~~ regional  
301 self-sufficient ~~of~~cy food, ~~including~~ vegetables, fruits, cotton, ~~and peanut etes at the same time, and~~  
302 the groundwater consumption of this structure was acceptable. In addition, policies on agricultural  
303 crop structure optimization should be encouraged, with the aim of relieving the pressure on  
304 groundwater for crop production and ensuring food security in this region. In recent years, winter  
305 wheat and summer maize were being replaced by spring crops in many places of the HSP, this was  
306 ~~been~~ called "the spring corn planting belt phenomenon" (Feng et al., 2007; Huang et al., 2012;  
307 Wang et al., 2014). ~~Undoubtedly~~Clearly, this phenomenon can help ~~to in~~ the restoration of  
308 groundwater resources in this region.

309 4.3 ~~The m~~Main ~~shortcomings uncertainties~~ of this study

310 Firstly, the estimation of WF (~~including comprised of~~ WFGreen, WFblue and WFGrey) was  
311 affected by crop distribution, ~~in regards to the spatial differences of for the~~ underlying surface  
312 conditions, climatic conditions and irrigation conditions ~~have spatial difference, but t~~. The crop  
313 ~~distribution of the baseline mainly came from land use map and statistical data and the crop~~  
314 structure scenarios ~~only considered the crop planting areas and~~ did not take into account the crop  
315 distribution, ~~this study only considered the crop planting area and ignored its distribution~~. Secondly,  
316 the scenarios ~~setting~~ had a certain ~~degree of~~ randomness ~~without considering~~since there was  
317 ~~no consideration to planting area~~ changes ~~of in~~ planting areas of cotton, peanut and others (Table 2);  
318 ~~in fact, due to~~. For example, ~~with cotton lacking a high market value and having the difficulty~~

319 ~~difficulties in its management of cotton management~~ (e.g., ~~requiring it needs~~ artificial picking–)  
320 ~~without high price~~, its growing area was likely shrinking, and its distribution was changing. Thirdly,  
321 due to the ~~development of~~ urbanization in this region, the area of arable land has been shrinking, ~~at~~  
322 ~~the same time; likewise~~, some arable land was abandoned because many rural young people went to  
323 work in cities ~~in many rural communities, but~~. Our scenario analysis, however, did not take into  
324 account these ~~phenomenons–phenomena, as we for–lacked~~ the corresponding data. Fourthly,  
325 climatic variability has major effects on crop WF (Sun et al., 2010; Bocchiola et al., 2013; Yang et  
326 al., 2013), and many researchers have found that this region has undergone an upward trend of  
327 temperatures and a declining trend of precipitation since the 1960s (Hu et al., 2002; Yuan et al.,  
328 2009; Sun et al., 2010). If precipitation continues to decline ~~and while~~ temperature increases ~~in the~~  
329 ~~future over time~~, these climatic developments will certainly affect the WF for crop production  
330 ~~certainly and~~. These ~~questions–effects~~ are worth an in-depth ~~analyzing~~ analysis, which ~~can~~ could  
331 provide valuable information for water resource management.

## 332 5 Conclusions

333 This study analyzed the WF of crop production in the HSP and evaluated its temporal variation  
334 from 2000 to 2012. Over 13 years, the production of main crops consumed a total of approximately  
335 604.8 km<sup>3</sup> of water, of which 288.5 km<sup>3</sup> of that was groundwater; additionally, the WF of the  
336 production of crops exhibited a downtrend yearly. Among the local main crops, winter wheat,  
337 summer maize and vegetables were the three leading crops in water consumption; their WF,  
338 WF<sub>blue</sub>, WF<sub>green</sub> and WF<sub>gray</sub> accounted for 76.2%, 73.7%, 74.2% and 81.6% of the total,  
339 respectively.

340 In this region, adjusting crop farming structures was has been an important means to protect  
341 groundwater resources ~~in the HSP. This study; therefore, we~~ evaluated ~~the~~ reasonable farming  
342 structures by analyzing scenarios ~~analysis~~ of the main crops' WF in this plain and ~~suggested~~ that:  
343 scenario 6–with about approximately 20% of ~~the~~ arable land in cultivating–cultivation of winter  
344 wheat-summer maize in rotation, 40% of cultivating–spring maize, 10% cultivating of vegetables,

345 10% ~~cultivating~~ of fruiters, ~~without~~ 0% of rice and ~~no change to~~ other crops, ~~unchanging~~ (i.e.  
346 ~~scenario 6~~) were available to ~~will~~ promote the sustainable development of agriculture in this region,  
347 ~~which~~. This scenario not only can protect ~~approximately 14.5% of groundwater resources~~ (the  
348 ~~groundwater resources~~ compared to the baseline), but ~~also~~ can ~~also~~ ensure the local supply of  
349 ~~food~~ wheat, vegetables, and fruits.

350 **Acknowledgements** ~~The~~ This paper was supported by the Natural Science Foundation of China  
351 (40901130, ~~41471027~~), the high-level leading talent introduction program of GDAS (2060599), ~~the~~  
352 ~~Innovation-driven Development Capacity Construction Program (2017GDASCX-0861)~~, ~~the~~  
353 ~~Scientific Platform and Innovation Capability Construction Program of GDAS~~  
354 ~~(2016GDASPT-0304)~~, ~~the Research Team of the Guangdong Natural Science Foundation~~  
355 ~~(S2012030006144)~~, the Science and Technology Project of Hebei Province (16454203D) and the  
356 Social Science Foundation of Hebei Province (HB15GL083). We are also grateful to the reviewers  
357 and editors.

### 358 **References**

- 359 Bocchiola, D., Nana, E., Soncini, A., 2013. Impact of climate change scenarios on crop yield and  
360 water footprint of maize in the Po valley of Italy. *Agric. Water Manag.* 116, 50-61.
- 361 ~~Castellanos, M.T., Cartagena, M.C., Requejo, M.I., et al., 2016. Agronomic concepts in water~~  
362 ~~footprint assessment: A case of study in a fertirrigated melon crop under semiarid conditions.~~  
363 ~~Agric. Water Manag., 170, 81-90.~~
- 364 Chapagain, A.K., Hoekstra, A.Y., 2011. The blue, green and grey water footprint of rice from  
365 production and consumption perspectives. *Ecol. Econ.* 70, 749-758.
- 366 Chapagain, A.K., Orr, S., 2009. An improved water footprint methodology linking global  
367 consumption to local water resources: a case of Spanish tomatoes. *J. Environ. Manag.* 90,  
368 1219-1228.
- 369 ~~Daniela, L., Jacopo, B., Marco, F., 2016. Water Footprint of crop productions: A review. *Sci. Total*~~  
370 ~~Environ., 548-549:236-251.~~

371 da Silva, V.d.P.R., da Silva, B.B., Albuquerque, W.G., Borges, C.J.R., de Sousa, I.F. Neto, J.D.,  
372 2013. Crop coefficient, water requirements, yield and water use efficiency of sugarcane growth  
373 in Brazil. *Agric. Water Manag.* 128(10), 102-109.

374 Dong, H., Geng, Y., Sarkis, J., Fujita, T., Okadera, T., Xue, B., 2013. Regional water footprint  
375 evaluation in China: A case of Liaoning. *Science of the Total Environment*, 442, 215-224.

376 [Duan, A.W., 2004. Irrigation water quota of main crops in North China. Beijing: Agricultural](#)  
377 [Science and Technology Press of China.](#)

378 Feng, Z.M., Liu, D.W., Zhang, Y.H., 2007. Water requirements and irrigation scheduling of spring  
379 maize using GIS and CropWat model in Beijing-Tianjin-Hebeiregion. *Chin. Geogr. Sci.* 17,  
380 56-63.

381 Gheewala, S.H., Silalertruksa, T., Nilsalab, P., Mungkung, R., Perret, S.R., Chaiyawannakarn, N.,  
382 2014. Water footprint and impact of water consumption for food, feed, fuel crops production in  
383 Thailand. *Water*, 6, 1698-1718.

384 Gleick, P.H., 2003. Global freshwater resources: soft-path solutions for the 21st century. *Science*  
385 302, 1524-1528

386 Hoekstra, A.Y. (Ed.), 2003. *Virtual Water Trade: Proceedings of the International Expert Meeting*  
387 *on Virtual Water Trade*. Delft, The Netherlands, 12-13 December 2002, Value of Water Research  
388 Report Series No. 12. UNESCO-IHE, Delft, The Netherlands, Available at:  
389 [www.waterfootprint.org/Reports/Report12.pdf](http://www.waterfootprint.org/Reports/Report12.pdf)

390 Hoekstra, A.Y., Chapagain, A.K., 2007. Water footprints of nations: water use by people as a  
391 function of their consumption pattern. *Water Resour. Manag.* 21, 35-48.

392 Hoekstra, A.Y., Chapagain, A.K., Aldaya, M.M., Mekonnen, M.M., 2009. *Water Footprint Manual:*  
393 *State of the Art 2009*.

394 Huang, J., Bradley, G.R., Xu, C.C., Zhang, H.L., Chen, F., 2012. Cropping pattern modifications  
395 change water resource demands in the Beijing metropolitan area. *J. Integr. Agric.* 11, 1914-1923.

396 Hu, C.S., Zhang, X.Y., Cheng, Y.S., Pei, D., 2002. An analysis on dynamics of water table and

397 overdraft in the piedmont of Mt Taihang. *Syst. Sci. Compr. Stud. Agric.* 18,89–91 (in Chinese).

398 Jefferies, D., Muñoz, I., Hodges, J., King, V.J., Aldaya, M., Ercin, A.E., Canals, L.M.,  
399 Hoekstra, A.Y., 2012. Water footprint and life cycle assessment as approaches to assess potential  
400 impacts of products on water consumption. Key learning points from pilot studies on tea and  
401 margarine. *J. Clean. Prod.* 33, 155-166.

402 Konar, M., Dalin, C., Suweis, S., Hanasaki, N., Rinaldo, A., Rodriguez-Iturbe, I., 2011. Water for  
403 food: the global virtual water trade network. *Water Resour. Res.*  
404 47, <http://dx.doi.org/10.1029/2010WR010307>.

405 Lamastra, L., Suciù, N.A., Novelli, E., Trevisan, M., 2014. A new approach to assessing the water  
406 footprint of wine: an Italian case study. *Sci. Total Environ.* 490, 748-756.

407 Lucrezia, L., Nicoleta, A.S., Elisa, N., Marco, T., 2014. A new approach to assessing the water  
408 footprint of wine: An Italian case study. *Science of the Total Environment*, 490, 748-756.

409 Ma, Y., Feng, S., Song, X., 2013. A root zone model for estimating soil water balance and crop  
410 yield responses to deficit irrigation in the North China Plain. *Agric. Water Manag.* 127(28),  
411 13-24.

412 Mekonnen, M.M., Hoekstra, A.Y., 2010. The green, blue and grey water footprint of crops and  
413 derived crop products. Value of water research report series No. 47. UNESCO-IHE, Delft, The  
414 Netherlands.

415 [Pan, X. P., Li, G.X., Liu, F. G., Wu, X.F., Kondoh, A., Shen, Y.J., 2015. Using remote sensing to](#)  
416 [determine spatio-temporal variations in winter wheat growing area in the North China Plain.](#)  
417 [Chinese Journal of Eco-Agriculture, 23\(4\): 497-505 \(in Chinese\).](#)

418 Postel, S., 2000. Entering an era of water scarcity: the challenges ahead. *Ecological Applications*,  
419 10 (4), 941-948.

420 Ridoutt, B., Eady, S., Sellahewa, J., Simons, L., Bektash, R., 2009. Water footprinting at the  
421 product brand level: case study and future challenges. *Journal of Cleaner Production* 17 (13),  
422 1228-1235.

- 423 Rosegrant, M.W., Ringler, C., 2000. Impact on food security and rural development of transferring  
424 water out of agriculture. *Water Policy* 1, 567-586.
- 425 ~~Suttayakul, P., H-Kittikun, A., Suksaroj, C., Mungkalasiri, J., Wisansuwannakorn, R., Musikavong,~~  
426 ~~C., 2016. Water footprints of products of palm plantations and palm oil mills in Thailand.~~  
427 ~~*Science of the Total Environment*, 542, 521-529.~~
- 428 Shrestha, S., Pandey, V.P., Chanamai, C., Ghosh, D.K., 2013. Green, blue and grey water footprints  
429 of primary crops production in Nepal. *Water Resour. Manag.* 5223-5243.
- 430 Sun, H., Shen, Y., Yu, Q., Flerchinger, G.N., Zhang, Y., Liu, C., Zhang, X., 2010. Effect  
431 of precipitation change on water balance and WUE of the winter wheat-summermaize rotation in  
432 the North China Plain. *Agric. Water Manage.* 97,1139-1145.
- 433 Suttayakul, P., H-Kittikun, A., Suksaroj, C., Mungkalasiri, J., Wisansuwannakorn, R., Musikavong,  
434 C., 2016. Water footprints of products of oil palm plantations and palm oil mills in Thailand.  
435 *Science of the Total Environment*, 542, 521-529.
- 436 UNEP, 2007. *Global Environment Outlook e Geo 4: Environment for Development*. United  
437 Nations Environment Programme, Valletta, Malta.
- 438 ~~Wang, H.Y., Pan, X.P., Luo, J.M., Luo, Z.P., Chang, C.P., Shen, Y.J., 2015. Using remote sensing to~~  
439 ~~analyze spatiotemporal variations in crop planting in the North China Plain. *Chinese Journal of*~~  
440 ~~*Eco-Agriculture*, 23(9):1199-1209 (in Chinese).~~
- 441 Wang, X., Li, X.B., Fischer, G., 2015. Impact of the changing area sown to winter wheat on crop  
442 water footprint in the North China Plain. *Ecological Indicators*, 57:100-109.
- 443 Wang, Y.B., Wu, P.T., Engel, B.A., Sun, S.K., 2014. Application of water footprint combined with  
444 unified virtual crop pattern to evaluate crop water productivity in grain production in China. *Sci.*  
445 *Total Environ.*, 497-498, 1-9.
- 446 Wang, X., Li, X., Xin, L., 2014. Impact of the shrinking winter wheat sown area on agricultural  
447 water consumption in the Hebei Plain. *J. Geogr. Sci.* 24, 313-330.
- 448 White, D.J., Feng, K.S., Sun, L.X., Hubacek, K., 2015. A hydro-economic MRIO analysis of the



- 449 Haihe River Basin's water footprint and water stress. *Ecological Modelling*, 318, 157-167.
- 450 Xu, Y.J., Huang, K., Yu, Y.J., Wang, X.M., 2014. Changes in water footprint of crop production in  
451 Beijing from 1978 to 2012: a logarithmic mean Divisia index decomposition analysis. *Journal*  
452 *of Cleaner Production*, 87:180-187.
- 453 Yang, X., Gao, W., Shi, Q., Chen, F., Chu, Q., 2013. Impact of climate change on the water  
454 requirement of summer maize in the Huang–Huai–Hai farming region. *Agric. Water Manag.* 124,  
455 20–27.
- 456 Yuan, Z.J., Shen, Y.J. Estimation of Agricultural Water Consumption from Meteorological and  
457 Yield Data: A Case Study of Hebei, North China. *PLOS ONE*, 2013, 8(3), e58685.
- 458 Yuan, Z.J., Shen, Y.J., Chu, Y.M., et al, 2009. Variations and distribution of temperature and  
459 precipitation of Haihe River Basin in recent 40 years. *Research of Soil and Water Conservation*,  
460 16(3):24-26 (in Chinese).
- 461 Zang, T., Xie, X., Huang, Z., 2014. Life cycle water footprints of nonfood biomass fuels in China.  
462 *Environ. Sci. Technol.* 48, 4137-4144.



**ELSEVIER**

**Language Editing Services**

*Registered Office:*  
Elsevier Ltd  
The Boulevard, Langford Lane,  
Kidlington, OX5 1GB, UK.  
Registration No. 331566771

### **To whom it may concern**

The paper "Water footprint of crop production for different crop structures in the Hebei southern plain, North China" by Yingmin Chu was edited by Elsevier Language Editing Services.

Kind regards,

Biji Mathilakath  
**Elsevier Webshop Support**