

Interactive comment on “The probability distribution of daily streamflow in the conterminous United States” by Annalise G. Blum et al.

Anonymous Referee #1

Received and published: 18 October 2016

The paper fits theoretical distributions to a large dataset of empirical streamflow observations covering the conterminous US. The study finds that median annual flow duration curves (FDC), which portray flow distribution in a typical year, can be reasonably fitted to three-parameter distributions. In contrast, period of record FDC that incorporate extreme streamflow variations over numerous years cannot be appropriately fitted, even to more complex theoretical distributions. The authors explore the implications of that finding on predictions in ungauged catchments using linear regressions in case studies. Predicting streamflow signatures (particularly FDC) in ungauged basins is both extremely useful and challenging and the findings of this study are interesting and important, particularly the insight that mAFDC might be both easier to predict and

C1

more practically relevant than PoRFDC. However, there are two points that I would like to see further discussed before publication, as well as the few minor comments listed below.

First, the study is an impressive effort to fit FDCs to a very large dataset of unregulated catchments – this is definitely a key contribution of the paper. However, by covering the whole conterminous US, the dataset covers a wide variety of climates, catchment characteristics and flow regimes, and it would have been interesting to explore how the fit to specific distributions varies regionally. The shape of FDCs depicts the local flow regime, which are themselves related to climate and catchment characteristics (see e.g., Botter 2013). It would be nice to see whether there is a link between flow regimes, climate/catchment characteristics and the best fitted theoretical distribution. It would also be nice to discuss how the best-fit distributions relate to the distributions that might be expected from process-based models (Botter 2007, Botter 2009, Muller 2014, Muneeppeerakul 2010, etc), given the dominant flow processes in particular catchments.

Second, while I appreciate the effort to extend an already complex and large scale analysis to prediction in ungauged basins, I would like to see more details on how the regression models were obtained (i.e. how the regression covariates were selected for Eqn 7-9), and a discussion on whether these regression models have a physical interpretation. Specifically, I am concerned about using linear regressions to estimate distribution parameters, which arguably have a more ambiguous physical interpretation as moments. Mean flow (first moment) for instance can be argued to be a linear combination of observable characteristics like mean rainfall, as per the water balance equation. The issue in regressing GPA3 parameters is that they are not linear combinations of the moments of the distribution, so using linear regressions to estimate the parameters does not allow moments to be linearly related. In other words, in this specific case, linearly regressed parameters are not compatible with a linear water balance relation on mean flow. To address this issue, please either apply linear regressions on

C2

the moments of the distributions instead of the parameters (and discuss the physical interpretation of the linear models when appropriate), or make the case that Eqn 7-9 are not incompatible with water balance principles.

[To illustrate my point on linear regressions, let's assume the simplest linear model possible, where predictions are simply taken as the mean of the observed sample (this can happen in the specific case of the water balance model above if all catchments have an identical mean rainfall). Let's say that we have a sample of three catchments with the following GPA3 parameters and mean flow (computed from the parameters):

Basin | location param | scale param | shape param | Mean

1 | 0 | 100 | 0.3 | 143

2 | 0 | 1 | 0.05 | 1

3 | 0 | 20 | 0.1 | 22

The predicted mean flow in a fourth catchment obtained from the observed mean flows (i.e. by taking the mean of the mean) is 47, whereas the mean flow computed from predicted GPA3 parameters (i.e. computed from the mean value of each parameter) is 55.]

Minor comments:

p5 l 17: please define GOF.

p12 l14-19: I have seen this issue most often addressed by taking the logarithm of flow quantiles before computing the NSE. Is there any reason why you preferred the selected approach?

p14 l13-15: I agree that modelling errors on FDCs are best assessed graphically and appreciate the effort of showing fits for particular basins with low, median and high NSE. Error duration curves (e.g., Muller 2016) are a great way of visualizing performance fits over large samples (as opposed to individual basins), and it would be informative in my

C3

opinion to display the relevant EFDCs for the whole dataset.

p.16 l3-4: I realize that concerns of overfitting regression models are to some extent addressed in the LOO cross validation analysis, but please display covariates statistics (e.g., quartiles and range or boxplot) to show that there is enough variability in the samples to credibly argue that the LOO performance is externally valid.

p.17 l2-4: Have you tested for serial correlation? Serial correlation affects the estimation of OLS standard errors and are particularly likely to occur between flow-connected gauges (i.e. gauges located on the same river).

p.18 l.11-20: The paper makes the great case that MAFDCs are easier to fit and have more practical relevance than PoRFDC. However, I would appreciate a more complete discussion of the tradeoff involved: mAFDC loses information on inter-annual variability, hence their better fit to "simpler" distributions. This is an important caveat that has strong implications for practical applications and should be made clearer in the discussion/conclusion in my opinion.

Table C1: I have trouble understanding how the BFI_AVE is a regression covariate for prediction in ungauged basins. My understanding is that flow observations are necessary to compute the BFI in the first place.

References:

Botter, G., Basso, S., Rodriguez-Iturbe, I., and Rinaldo, A.: Resilience of river flow regimes, *Proceedings of the National Academy of Sciences*, 110, doi:10.1073/pnas.1311920 110, 2013.

Botter, G., Porporato, A., Rodriguez-Iturbe, I., and Rinaldo, A.: Basin-scale soil moisture dynamics and the probabilistic characterization of carrier hydrologic flows: Slow, leaching prone components of the hydrologic response, *Water Resources Res.*, 43, doi:10.1029/2006WR005 043, 2007.

Botter, G., Porporato, A., Rodriguez-Iturbe, I., and Rinaldo, A.: Nonlinear storage-

C4

discharge relations and catchment streamflow regimes, *Water resources research*, 45, doi: 10.1029/2008WR007 658, 2009.

Müller, M. F., Dralle, D. N., and Thompson, S. E.: Analytical model for flow duration curves in seasonally dry climates, *Water Resources Research*, 50, doi:10.1002/2014WR015 301, 2014. Muneeppeerakul, R., Azaele, S., Botter, G., Rinaldo, A., and Rodriguez-Iturbe, I.: Daily streamflow analysis based on a two-scaled gamma pulse model, *Water Resources Research*, 46, doi: 10.1029/2010WR009 286, 2010. Müller, M. F. and Thompson SE.: Comparing statistical and process-based flow duration curve models in ungauged basins and changing rain regimes, *Hydrology and Earth System Sciences* 20.2 669-683, 2016

Interactive comment on *Hydrol. Earth Syst. Sci. Discuss.*, doi:10.5194/hess-2016-460, 2016.