

1 *Supplementary material for*

2 **On Coupled Unsaturated-Saturated Flow Processes Induced by Vertical,**
3 **Horizontal and Slant Wells in Unconfined Aquifers**

4 Xiuyu Liang^a, Hongbin Zhan^b, You-Kuan Zhang^c, Jin Liu^a

5 ^aSchool of Earth Sciences and Engineering, Nanjing University,
6 Nanjing, Jiangsu 210093, P.R. China (xyliang@nju.edu.cn)

7 ^bDepartment of Geology & Geophysics, Texas A&M University, College Station, TX 77843-
8 3115, USA. (zhan@geos.tamu.edu)

9 ^cSchool of Environment Sciences and Engineering,
10 South University of Sciences and Technology of China,
11 Shenzhen, Guangdong 518055, P.R. China

12

13

14 **S1. Dimensionless transform for Eqs. (1)-(4)**

15 For the purpose of mathematical convenience, we define the following dimensionless variables,

$$\begin{aligned}
 16 \quad s_D &= \frac{4\pi K d}{Q} s, t_D = \frac{K}{S_s d^2} t, x_D = \alpha_x \frac{x}{d}, y_D = \alpha_y \frac{y}{d}, z_D = \alpha_z \frac{z}{d}, x_{0D} = \alpha_x \frac{x_0}{d}, y_{0D} = \alpha_y \frac{y_0}{d}, z_{0D} = \\
 17 \quad &\alpha_z \frac{z_0}{d}, \alpha_x = (K/K_x)^{1/2}, \alpha_y = (K/K_y)^{1/2}, \alpha_z = (K/K_z)^{1/2}, K = (K_x K_y K_z)^{1/3}, u_D = \\
 18 \quad &\frac{4\pi K d}{Q} u, \kappa_D = \frac{d}{\alpha_z} \kappa, \beta = \frac{\kappa_D}{\sigma}, \sigma = \frac{S_s d}{S_y \alpha_z}, b_D = \alpha_z \frac{b}{d}, \quad (S1)
 \end{aligned}$$

19 where the subscript D denotes the dimensionless terms. Substituting above dimensionless variables
 20 into Eqs. (1)-(4), one obtains the following dimensionless forms of the governing equations for the
 21 saturated zone,

$$22 \quad \frac{\partial^2 s_D}{\partial x_D^2} + \frac{\partial^2 s_D}{\partial y_D^2} + \frac{\partial^2 s_D}{\partial z_D^2} + 4\pi \delta(x_D - x_{0D}) \delta(y_D - y_{0D}) \delta(z_D - z_{0D}) = \frac{\partial s_D}{\partial t_D}, \quad 0 \leq z_D < \alpha_z, (S2a)$$

$$23 \quad s_D(x_D, y_D, z_D, 0) = 0, \quad (S2b)$$

$$24 \quad \frac{\partial s_D}{\partial z_D}(x_D, y_D, z_D, t_D)|_{z_D=0} = 0, \quad (S2c)$$

$$25 \quad \lim_{x_D \rightarrow \pm\infty} s_D(x_D, y_D, z_D, t_D) = \lim_{y_D \rightarrow \pm\infty} s_D(x_D, y_D, z_D, t_D) = 0, \quad (S2d)$$

26 and for the unsaturated zone,

$$27 \quad \frac{\partial^2 u_D}{\partial x_D^2} + \frac{\partial^2 u_D}{\partial y_D^2} + \frac{\partial^2 u_D}{\partial z_D^2} - \kappa_D \frac{\partial u_D}{\partial z_D} = \beta \frac{\partial u_D}{\partial t_D}, \quad \alpha_z \leq z_D < \alpha_z + b_D, \quad (S3a)$$

$$28 \quad u_D(x_D, y_D, z_D, 0) = 0, \quad (S3b)$$

$$29 \quad \frac{\partial u_D}{\partial z_D}(x_D, y_D, z_D, t_D)|_{z_D=\alpha_z+b_D} = 0, \quad (S3c)$$

$$30 \quad \lim_{x \rightarrow \pm\infty} u_D(x_D, y_D, z_D, t_D) = \lim_{y \rightarrow \pm\infty} u_D(x_D, y_D, z_D, t_D) = 0, \quad (S3d)$$

31 and at the interface,

$$32 \quad s_D - u_D = 0, \quad z_D = \alpha_z, \quad (S4a)$$

$$33 \quad \frac{\partial s_D}{\partial z_D} - \frac{\partial u_D}{\partial z_D} = 0, \quad z_D = \alpha_z. \quad (S4b)$$

34

35 **S2: Laplace domain solution of the unsaturated zone**

36 The Laplace transformation of Eqs. (S3) and (S4) are written as

37
$$\frac{\partial^2 \bar{u}_D}{\partial x_D^2} + \frac{\partial^2 \bar{u}_D}{\partial y_D^2} + \frac{\partial^2 \bar{u}_D}{\partial z_D^2} - \kappa_D \frac{\partial \bar{u}_D}{\partial z_D} = \beta p \bar{u}_D, \quad \alpha_z \leq z_D < \alpha_z + b_D, \quad (\text{S5a})$$

38
$$\frac{\partial \bar{u}_D}{\partial z_D} (x_D, y_D, z_D, p) \Big|_{z_D = \alpha_z + b_D} = 0, \quad (\text{S5b})$$

39
$$\lim_{x_D \rightarrow \pm\infty} \bar{u}_D(x_D, y_D, z_D, p) = \lim_{y_D \rightarrow \pm\infty} \bar{u}_D(x_D, y_D, z_D, p) = 0, \quad (\text{S5c})$$

40 and

41
$$\bar{s}_D - \bar{u}_D = 0, \quad z_D = \alpha_z, \quad (\text{S6a})$$

42
$$\frac{\partial \bar{s}_D}{\partial z_D} - \frac{\partial \bar{u}_D}{\partial z_D} = 0, \quad z_D = \alpha_z, \quad (\text{S6b})$$

43 where p is the Laplace transform parameter and the overbar indicates a variable in the Laplace
 44 domain. The Eq. (S5) in a cylindrical coordinate system can be written as following

45
$$\frac{1}{r_D} \frac{\partial \bar{u}_D}{\partial r} + \frac{\partial^2 \bar{u}_D}{\partial r^2} + \frac{\partial^2 \bar{u}_D}{\partial z_D^2} - \kappa_D \frac{\partial \bar{u}_D}{\partial z_D} = \beta p \bar{u}_D, \quad \alpha_z \leq z_D < \alpha_z + b_D, \quad (\text{S7a})$$

46
$$\frac{\partial \bar{u}_D}{\partial z_D} (r_D, z_D, p) = 0, \quad z_D = \alpha_z + b_D, \quad (\text{S7b})$$

47
$$\bar{u}_D(\infty, z_D, p) = 0. \quad (\text{S7c})$$

48 Based on the methods of the separation variables and Eq. (5), the solution of Eq. (S7a) with
 49 boundary condition Eq. (S7c) will be (Rezaei et al., 2016):

50
$$\bar{u}_D(r_D, z_D, p) = \sum_{n=0}^{\infty} \frac{8 \cos(\omega_n z_{0D})}{p \Psi(\omega_n)} K_0(\Omega_n |r_D - r_{0D}|) \mathcal{H}_n(z_D, p). \quad (\text{S8})$$

51 Substituting Eq. (S8) into Eq. (S7) yields:

52
$$\frac{\partial^2 \mathcal{H}_n}{\partial z_D^2} - \kappa_D \frac{\partial \mathcal{H}_n}{\partial z_D} - (\beta p - \Omega_n^2) \mathcal{H}_n = 0, \quad \alpha_z \leq z_D < \alpha_z + b_D, \quad (\text{S9a})$$

53
$$\frac{\partial \mathcal{H}_n}{\partial z_D} (z_D, p) = 0, \quad z_D = \alpha_z + b_D. \quad (\text{S9b})$$

54 The general solution of Eq. (S9a) is

$$55 \quad \mathcal{H}_n = \begin{cases} e^{Mz_D} [C_1 e^{Nz_D} + C_2 e^{-Nz_D}], & \text{if } \Delta > 0, \\ e^{Mz_D} [C_1 \sin(N_1 z_D) + C_2 \cos(N_1 z_D)], & \text{if } \Delta < 0, \\ e^{Mz_D} [C_1 z_D + C_2], & \text{if } \Delta = 0, \end{cases} \quad (\text{S10})$$

56 where $M = \kappa_D/2$; $N = \sqrt{\Delta}$; $N_1 = \sqrt{-\Delta}$; $\Delta = \kappa_D^2/4 + \beta p - \Omega_n^2$. C_1 and C_2 are determined on
57 the basis of Eqs. (S6a) and (S9b), then substituting them into Eq. (S10) yields

$$58 \quad \mathcal{H}_n = \begin{cases} \cos(\omega_n \alpha_z) \frac{(M+N) \exp[2N(\alpha_z+b_D)+(M-N)z_D] - (M-N) \exp[(M+N)z_D]}{(M+N) \exp[2N(\alpha_z+b_D)+(M-N)\alpha_z] - (M-N) \exp[(M+N)\alpha_z]}, & \text{if } \Delta > 0, \\ \cos(\omega_n \alpha_z) \exp(Mz_D - M\alpha_z) \frac{[N_1 \tan(N_1(\alpha_z+b_D)) - M] \sin(N_1 z_D) + [M \tan(N_1(\alpha_z+b_D)) + N_1] \cos(N_1 z_D)}{[N_1 \tan(N_1(\alpha_z+b_D)) - M] \sin(N_1 \alpha_z) + [M \tan(N_1(\alpha_z+b_D)) + N_1] \cos(N_1 \alpha_z)}, & \text{if } \Delta < 0, \\ \cos(\omega_n \alpha_z) \exp(Mz_D - M\alpha_z) \frac{1+M(\alpha_z+b_D)-Mz_D}{1+M(\alpha_z+b_D)-M\alpha_z}, & \text{if } \Delta = 0. \end{cases} \quad (\text{S11})$$

59

60 S3: Evaluation for eigenvalues ω_n

61 Substituting Eqs. (5) and (7) into Eq. (S6b) yields

$$62 \quad \begin{cases} -\omega_n \tan(\omega_n \alpha_z) = \frac{1 - \exp(-2Nb_D)}{1/(M-N) - \exp(-2Nb_D)/(M+N)}, & \text{if } \Delta > 0 \\ -\omega_n \tan(\omega_n \alpha_z) = M + \frac{[N_1 \tan(N_1(\alpha_z+b_D)) - M]N_1 - [M \tan(N_1(\alpha_z+b_D)) + N_1]N_1 \tan(N_1 \alpha_z)}{[N_1 \tan(N_1(\alpha_z+b_D)) - M] \tan(N_1 \alpha_z) + [M \tan(N_1(\alpha_z+b_D)) + N_1]}, & \text{if } \Delta < 0 \\ \omega_n = (\kappa_D^2/4 + (\beta - 1)p)^{1/2}. & \text{if } \Delta = 0 \end{cases} \quad (\text{S12})$$

63 The eigenvalues of the finite cosine Fourier transform ω_n is the positive root of Eq. (S12).

64 For $\Delta > 0$ the solution domain of ω_n is separated into an infinite series of sub-domain intervals

65 with a period of π , i.e., $\left(\frac{(2i-1)\pi}{2\alpha_z}, \frac{(2i+1)\pi}{2\alpha_z}\right)$, $i = 0, 1, 2, \dots$. Each solution of ω_n in an individual

66 sub-domain is obtained using the Newton-Raphson method. For $\Delta < 0$ the solution domain of

67 ω_n is separated into an infinite series of irregular sub-domains due to the complex formula on the

68 right side of the equation. These irregular sub-domains can be identified by seeking the singular

69 points of function $F(\omega) = M + \frac{[N_1 \tan(N_1(\alpha_z+b_D)) - M]N_1 - [M \tan(N_1(\alpha_z+b_D)) + N_1]N_1 \tan(N_1 \alpha_z)}{[N_1 \tan(N_1(\alpha_z+b_D)) - M] \tan(N_1 \alpha_z) + [M \tan(N_1(\alpha_z+b_D)) + N_1]} + \omega \tan(\omega \alpha_z)$, where ω is a variable

70 of function $Y = F(\omega)$ and the roots of $F(\omega) = 0$ are eigenvalues ω_n . Specifically, the

71 eigenvalues ω_n are intersection points of curve $Y = F(\omega)$ with $Y = 0$. These intersection

72 points are separated into an infinite series of irregular sub-domains in which every sub-domain is
73 composed of two adjacent singular points ($\omega = \omega_s$) of function $Y = F(\omega)$. The ω_n in an
74 individual sub-domain is obtained using the Newton-Raphson method. It is shown in Eqs. (5) and
75 (7) that the saturated zone solution involving the components of unsaturated zone is only
76 represented in the ω_n terms. Changes of the unsaturated zone parameters lead to different ω_n
77 values, which affect groundwater flow in the saturated zone.

78

79 **References**

80 Rezaei, A., Zare, M., and Zhan, H. B.: Aquitard Horizontal Dispersion on Reactive Solute
81 Transport in an Aquifer-Aquitard System, *Transport Porous Med*, 113, 695-716,
82 10.1007/s11242-016-0719-6, 2016.

83