

Inter-comparison of daily precipitation products for large-scale hydro-climatic applications over Canada

Jefferson S. Wong^{1*}, Saman Razavi¹, Barrie R. Bonsal², Howard S. Wheeler¹, & Zilefac E. Asong¹

¹ *Global Institute for Water Security and School of Environment and Sustainability, University of Saskatchewan, 11 Innovation Blvd, Saskatoon, SK, Canada S7N 3H5*

² *Environment and Climate Change Canada, 11 Innovation Blvd, Saskatoon, SK, Canada S7N 3H5*

* Corresponding author:

Email: jefferson.wong@usask.ca

Phone: +1 306 966 7816

Abstract

A number of global and regional gridded climate products based on multiple data sources are available that can potentially provide reliable estimates of precipitation for climate and hydrological studies. However, research into the consistency of these products for various regions has been limited and in many cases non-existent. This study inter-compares several gridded precipitation products over 15 terrestrial ecozones in Canada for different seasons. The spatial and temporal variability of the errors (relative to station observations) was quantified over the period of 1979 to 2012 at a 0.5° and daily spatiotemporal resolution. These datasets were assessed in their ability to represent the daily variability of precipitation amounts by four performance measures: percentage of bias, root-mean-square-error, correlation coefficient, and standard deviation ratio. Results showed that most of the datasets were relatively skillful in central Canada. However, they tended to overestimate precipitation amounts in the west and underestimate in the north and east, with the underestimation being particularly dominant in northern Canada (above 60° N). The global product by WATCH Forcing Data ERA-Interim (WFDEI) augmented by Global Precipitation Climatology Centre (GPCC) data (WFDEI [GPCC]) performed best with respect to different metrics. The Canadian Precipitation Analysis (CaPA) product performed comparably with WFDEI [GPCC], however it only provides data starting in 2002. All the datasets performed best in summer, followed by autumn, spring, and winter in order of decreasing quality. Findings from this study can provide guidance to potential users regarding the performance of different precipitation products for a range of geographical regions and time periods.

Keywords: precipitation; evaluation and comparison; datasets; ecozones; hydro-climatology; Canada

1. Introduction

The availability of accurate data, especially precipitation, is essential for understanding the climate system and hydrological processes since it is a vital element of the water and energy cycles and a key forcing variable for driving hydrological models. Reliable precipitation measurements provide valuable information for meteorologists, climatologists, hydrologists, and other decision makers in many applications, including climate and/or land-use change studies (e.g. Cuo et al., 2011;Huisman et al., 2009;Dore, 2005), agricultural and environmental research (e.g. Zhang et al., 2012;Hively et al., 2006), natural hazards (e.g. Taubenbock et al., 2011;Kay et al., 2009;Blenkinsop and Fowler, 2007), and hydrological and water resources planning (e.g. Middelkoop et al., 2001;Hong et al., 2010). With respect to land-surface hydrology, the increasing sophistication of distributed hydrological modeling has urged the requirement of better and more reliable gridded precipitation estimates at a minimum, daily temporal resolution. Before incorporating precipitation measurements, quantifying their uncertainty becomes an essential prerequisite for hydrological applications and is increasingly critical for potential users who are left without guidance and/or confidence in the myriad of products for their specific hydrological problems over different geographical regions. This study attempts to address this issue by comparing and examining the error characteristics of different types of gridded precipitation products and assessing how these products perform geographically and temporally over Canada.

Precipitation measurements and their limitations

With technological and scientific advancements over the past three decades, tremendous progress has been made in the various methods of precipitation measurement, each one with its own strengths and limitations. Rain gauges provide the direct physical readings with relatively accurate measurements at specific points. However, such measurements are subject to various errors arising from wind effects (Nešpor et al., 2000;Ciach, 2003), evaporation (Strangeways, 2004;Mekis and Hogg, 1999), undercatch (Yang et al., 1998;Adam and Lettenmaier, 2003;Mekis and Hogg, 1999), and instrumental problems. Moreover, rain-gauge measurements are often spatially interpolated, which may not capture the true spatial variability of precipitation fields due to sparse gauge networks. Ground-based radar measurements can estimate precipitation over a relatively large area (radius of 200 to 300 km) but are prone to inaccuracies as a result of beam spreading, curvature of the earth, and terrain blocking (Dinku et al., 2002;Young et al., 1999), and errors in the rain rate-

reflectivity relationship, range effects, and clutter (Jameson and Kostinski, 2002; Villarini and Krajewski, 2010). Development of satellite-based precipitation estimates such as the Global Precipitation Measurement (GPM) mission (Hou et al., 2014) has provided excellent spatial coverage but also contain inaccuracies resulting primarily from temporal sampling errors, instrumental errors, and algorithm errors (Nijssen and Lettenmaier, 2004; Gebremichael et al., 2005). Recognizing the limitations in the various precipitation observation methods, a number of attempts to combine information from multiple sources have been undertaken (Xie and Arkin, 1996; Maggioni et al., 2014; Shen et al., 2010). Reanalysis data provide an alternative source of precipitation estimates by assimilating all available data (rain-gauge stations, aircraft, satellite, etc.) into a background forecast physical model. However, accuracies in reanalysis precipitation are dependent on the specific analysis-forecast systems and the choice of physical parameterizations (Betts et al., 2006). Numerical climate models including Atmosphere-Ocean General Circulation Models (AOGCMs) and Regional Climate Models (RCMs) offer another potential source of precipitation estimates, as well as future precipitation simulations. Precipitation estimates from climate models, however, remain relatively coarse in resolution and often produce systematic bias due to imperfect model-conceptualization, discretization and spatial averaging within grid cells (Teutschbein and Seibert, 2010; Xu et al., 2005).

Scope and Objectives

Numerous previous evaluation efforts among the precipitation products have been limited into three groups of inter-comparison of (1) satellite-derived products (e.g. Adler et al., 2001; Xie and Arkin, 1995; Turk et al., 2008); (2) reanalysis data (e.g. Janowiak et al., 1998; Bosilovich et al., 2008; Betts et al., 2006; Bukovsky and Karoly, 2007); and (3) climate model simulations (e.g. Covey et al., 2003; Christensen et al., 2007; Mearns et al., 2006; 2012). Despite the aforementioned efforts, few studies have conducted a detailed inter-comparison among different types of precipitation products. Gottschalck et al. (2005) compared seasonal total precipitation of several satellite-derived, rain-gauge-based, and model-simulated datasets over contiguous United States and showed the spatial root mean square error of seasonal total precipitation and mean correlation of daily precipitation between each product and the impacts of these errors on land surface modelling. Additionally, Ebert et al. (2007) examined 12 satellite-derived precipitation products and four numerical weather prediction models over the United States, Australia, and northwestern

Europe and found that satellite-derived estimates performed best in summer and model-induced ones were best in winter. However, a number of questions regarding the reliability of the precipitation products remained in doubt, including: to what extent do the users have the knowledge about the error information associated with all these different types of precipitation products; how do the error distribution of precipitation products vary by location and season; and which product(s) should the users have more confidence for their regions of interest. Answering these questions is therefore a crucial first step in quantifying the spatial and temporal variability of the precipitation products so as to better understand their reliability as forcing inputs in hydrological modelling and other related studies.

Given the emergence of various products derived from different methods and sources (Tapiador et al., 2012), accuracy comparison studies of precipitation products have been reported over several regions; examples include the globe (e.g. Gebregiorgis and Hossain, 2015; Adler et al., 2001; Tian and Peters-Lidard, 2010), Europe (e.g. Frei et al., 2006; Chen et al., 2006; Kidd et al., 2012), Africa (e.g. Dinku et al., 2008; Asadullah et al., 2008), North America (e.g. Tian et al., 2009; West et al., 2007), South America (e.g. Vila et al., 2009), and China (e.g. Shen et al., 2010; Wetterhall et al., 2006). However, less attention has been paid to high-latitude regions such as Canada where a considerable proportion of precipitation is in the form of snow (Behrangi et al., 2016). In many regions of Canada, precipitation-gauge stations are sparsely distributed and the information required for hydrological modelling may not be available at the site of interest. This is especially true in northern areas (north of 60° N) and over mountainous regions where precipitation-gauge stations are usually 500 to 700 km apart or at low elevations (Wang and Lin, 2015). Meanwhile, the decline and closure of manual observing precipitation-gauge stations further reduced the spatial coverage and availability of long-term precipitation measurements (Metcalf et al., 1997; Mekis and Hogg, 1999; Rapaic et al., 2015). Of additional concern, the observations for solid precipitation (snow, snow pellets, ice pellets, and ice crystals) and precipitation phase (liquid or solid) changes make accurate measurement of precipitation more difficult and challenging, and the measurement errors have been found to range from 20 to 50 % for automated systems (Rasmussen et al., 2012). The Meteorological Service of Canada has implemented a network of 31 radars (radar coverage at full range of 256 km) along southern Canada (see Fortin et al. (2015b) Fig. 1 for spatial distribution). This Canadian radar network has been employed as an additional source of observations in generating a gridded product for Canada

(see Sect. 3.2.2 for details). Yet, the shortcomings of using the radar data are twofold: (1) many areas of the country (north of 60° N) are not covered by this network; and (2) the implementation of the network began in 1997 and thus did not have sufficient lengths of data for any long-term hydro-climatic studies. The availability, coverage, and quality of precipitation-gauge measurements are thus obstacles to effective hydrological modelling and water management in Canada. However, the availability of several global and regional gridded precipitation products which provide complete coverage of the whole country at applicable time and spatial scales may provide a viable alternative for regional- to national-scale hydrological applications in Canada.

Given the aforementioned, this study aims to (1) inter-compare various daily gridded precipitation products against the best available precipitation-gauge observations; and (2) characterize the error distributions of different types of precipitation products over time and different geographical regions in Canada. Such inter-comparison will in turn help assess the performance of the precipitation products over specific climatic/hydrological regions.

The rest of this paper is organized as follows: a brief description of the study area and precipitation data is provided in Sect. 2 and 3. The methodology for evaluating precipitation products against the precipitation-gauge station observations is described in Sect. 4. Results and discussion are provided in Sect. 5 and 6, respectively, with a summary and conclusion following in Sect. 7.

2. Study Area

Canada, which covers a land area of 9.9 million km², extends from 42° N to 83° N latitude and spans between 141° W to 52° W longitude. With substantial variations over its landmass, the country can be divided into many regions according to aspects such as climate, topography, vegetation, soil, geology, and land use. The National Ecological Framework for Canada classified ecologically distinct areas with four hierarchical levels of generalization (15 ecozones, 53 ecoprovinces, 194 ecoregions, and 1021 ecodistricts from broadest to the smallest) (Ecological Stratification Working Group, 1996; Marshall et al., 1999). Similarly, the Standard Drainage Area Classification (SDAC) was developed to delineate hydrographic areas to cover all the land and interior freshwater lakes of the country with three levels of classification (11 major drainage areas, 164 sub-drainage areas, and 974 sub-sub-drainage areas) (Brooks et al., 2002; Pearse et al., 1985). The precipitation comparisons in this study incorporated both the ecological and hydrological delineations. This involved classifying the Canadian landmass into 15 ecozones for the main study

(Fig. 1) and 14 major drainage areas (the Arctic Major Drainage Area was further divided into Arctic and Mackenzie, whereas the St. Lawrence Major Drainage Area was further split into St. Lawrence, Great Lakes, and Newfoundland). Results are based on the ecozone classification, while those based on drainage areas are reported in the supplementary material.

3. Precipitation Data

3.1. Precipitation-gauge station observations

In Canada, climate data collection is coordinated by the Federal government, which is made available by the National Climate Data Archive of Environment and Climate Change Canada (NCDA). These data provide the basis for all available quality controlled climate observations. There are a total of 1499 precipitation-gauge stations (as of 2012) across Canada. However, given the frequent addition and subtraction of climate stations, these numbers have greatly varied through time with peak reporting in the 1970s followed by a general decline to the present (see Hutchinson et al. (2009) Figs. 1 and 2 for details). Furthermore, the existing precipitation observations are often subject to various errors, with gauge undercatch being of significant concern (Mekis and Hogg, 1999). To account for various measurement issues, Mekis and Hogg (1999) first produced the Adjusted and Homogenized Canadian Climate Data (AHCCD) including adjusted daily rainfall and snowfall values and Mekis and Vincent (2011) then updated the data for a subset of 464 stations over Canada. The data extend back to 1895 for a few long-term stations and run through 2014. As a result of adjustments, total rainfall amounts were on the order of 5 to 10 % higher in southern Canada and more than 20 % in the Canadian Arctic when compared to the original observations. Adjustments to snowfall were even larger and varied throughout the country. These adjusted values are widely considered as better estimates of actual precipitation and therefore have been used in numerous analyses (e.g. Nalley et al., 2012; Shook and Pomeroy, 2012; Wan et al., 2013; Asong et al., 2015). Given the lack of an adjusted daily gridded precipitation product for Canada, the AHCCD station precipitation is considered to be the best available data for Canada and thus is used as the benchmark for all gridded precipitation product comparisons.

3.2. Gridded precipitation products

Seven precipitation datasets were chosen for assessment based on the following criteria: (1) a complete coverage of Canada; (2) minimum of daily temporal and 0.5° (~50 km) spatial

resolutions; (3) sufficient length of data (>30 years) for long-term study including recent years up to 2012; and (4) representing a range of sources/methodologies (e.g. station based, remote sensing, model, blended products). Table 1 summarizes these datasets, including their full names and original spatial and temporal resolutions for the versions used.

3.2.1. Station-based product – ANUSPLIN

Hutchinson et al. (2009) used the Australian National University Spline (ANUSPLIN) model to develop a dataset of daily precipitation, and daily minimum and maximum air temperature over Canada at a spatial resolution of 300 arc-seconds (0.0833° or ~ 10 km) for the period of 1961 to 2003. All available NCDA stations (that ranged from 2000 to 3000 for any given year during this period) were used as an input to the gridding procedure. To retain maximum spatial coverage, the smaller number of stations in AHCCD were not incorporated (i.e. only unadjusted archive values were used). Interpolation procedures included incorporation of tri-variate thin-plate smoothing splines using spatially continuous functions of latitude, longitude, and elevation. Hopkinson et al. (2011) subsequently extended this original dataset to the period 1950 to 2011. The Canadian ANUSPLIN has now further been updated to 2013 and has recently been used as the basis of ‘observed’ data for evaluating different climate datasets (e.g. Eum et al., 2012) and for assessing the effects of different climate products in hydro-climatological applications (e.g. Eum et al., 2014; Bonsal et al., 2013; Shrestha et al., 2012a).

3.2.2. Station-based multiple-source product – CaPA

In November 2003, the Canadian Precipitation Analysis (CaPA) was developed to produce a dataset of 6-hourly precipitation accumulation over North America in real-time at a spatial resolution of 15 km (from 2002 onwards) (Mahfouf et al., 2007). Data were generated using an optimum interpolation technique (Daley, 1993), which required a specification of error statistics between observations and a background field (e.g. Bhargava and Danard, 1994; Garand and Grassotti, 1995). For Canada, the short-term precipitation forecasts from the Canadian Meteorological Centre (CMC)’s regional Global Environmental Multiscale (GEM) model (Cote et al., 1998a; 1998b) were used as the background field with the rain-gauge measurements from NCDA as the observations to generate an analysis error at every grid point. CaPA became operational at the CMC in April 2011, with updates in the statistical interpolation method (Lespinas et al., 2015) and increase of spatial resolution to 10 km. The assimilation of Quantitative

Precipitation Estimates from the Canadian Weather Radar Network is also used as an additional source of observations (Fortin et al., 2015b). With its continuous improvement and different configurations, CaPA has been employed in Canada for various environmental prediction applications (e.g. Eum et al., 2014;Fortin et al., 2015a;Pietroniro et al., 2007;Carrera et al., 2015). However, the study period of these applications only start in 2002.

3.2.3. **Reanalysis-based multiple-source products – Princeton, WFDEI, and NARR**

Princeton

The Terrestrial Hydrology Research Group at the Princeton University initially developed a dataset of 3-hourly near-surface meteorology with global coverage at 1.0° spatial resolution (~120 km) from 1948 to 2000 for driving land surface models and other terrestrial systems (Sheffield et al., 2006). This dataset (called hereafter “Princeton”) was constructed based on the National Centers for Environmental Prediction-National Center for Atmospheric Research (NCEP-NCAR) reanalysis (2.0° and 6-hourly) (Kalnay et al., 1996;Kistler et al., 2001), combined with a suite of global observation-based data including the Climatic Research Unit (CRU) monthly climate variables (New et al., 1999, 2000), the Global Precipitation Climatology Project (GPCP) daily precipitation (Huffman et al., 2001), the Tropical Rainfall Measuring Mission (TRMM) 3-hourly precipitation (Huffman et al., 2002), and the NASA Langley Research Center monthly surface radiation budget (Gupta et al., 1999). With the inclusion of additional temperature and precipitation data (e.g. Willmott et al., 2001), Princeton has been updated and is currently available with two versions. This study used the 1901-2012 experimental version at 0.5° at daily time steps. Studies employing Princeton to examine different hydrological aspects have been carried out over different parts of Canada (Wang et al., 2013;Kang et al., 2014;Wang et al., 2014).

WFDEI

To simulate the terrestrial water cycle using different land surface models and general hydrological models, the European Union Water and Global Change (WATCH) Forcing Data (WFD) were created to provide datasets of sub-daily (3- and 6-hourly) and daily meteorological data with global coverage at 0.5° spatial resolution (~50 km) from 1901 to 2001 (Weedon et al., 2011). Similar to Princeton, the WFD were derived from the 40-year European Centre for Medium-Range Weather Forecasts (ECMWF) Re-Analysis (ERA-40) (1.0° and 3-hourly) (Uppala et al., 2005) and

combined with the CRU monthly variables and the Global Precipitation Climatology Centre (GPCC) monthly data (Rudolf and Schneider, 2005; Schneider et al., 2008; Fuchs, 2009). The WATCH Forcing Data methodology applied to ERA-Interim (WFDEI) dataset has further been developed covering the period of 1979 to 2012 (Weedon et al., 2014). The WFDEI used the same methodology as the WFD, but was based on the ERA-Interim (Dee et al., 2011) with higher spatial resolution (0.7°). As for the WFD, the WFDEI had two sets of rainfall and snowfall data generated by using either CRU or GPCC precipitation totals. Both sets of data were used in this study (hereafter known as WFDEI [CRU] and WFDEI [GPCC], respectively). To date, specific studies using the WFDEI related to Canada have been limited to the investigation of permafrost changes in the Arctic regions (e.g. Chadburn et al., 2015; Park et al., 2015; Park et al., 2016).

NARR

With the aim of evaluating spatial and temporal water availability in the atmosphere, the North American Regional Reanalysis (NARR) was developed to provide datasets of 3-hourly meteorological data for the North America domain at a spatial resolution of 32 km ($\sim 0.3^\circ$) covering the period of 1979 to 2003 as the retrospective system and is being continued in near real-time (currently up to 2015) as the Regional Climate Data Assimilation System (R-CDAS) (Mesinger et al., 2006). The components in generating NARR included the NCEP-DOE reanalysis (Kanamitsu et al., 2002), the NCEP regional Eta Model (Mesinger et al., 1988; Black, 1988) and the Noah land-surface model (Mitchell et al., 2004; Ek et al., 2003), and the use of numerous additional data sources (see Mesinger et al., 2006 Table 2). For hydrological modelling in Canada, Choi et al. (2009) found that NARR provided reliable climate inputs for northern Manitoba while Woo and Thorne (2006) concluded that NARR had a cold bias resulting in later snowmelt peaks in subarctic Canada. In addition, Eum et al. (2012) identified a structural break point in the NARR dataset beginning in January 2004 over the Athabasca River basin due to the assimilation of station observations over Canada being discontinued in 2003.

3.2.4. GCM statistically downscaled products – PCIC

The Pacific Climate Impacts Consortium (PCIC), which is a regional climate service centre at the University of Victoria, British Columbia, Canada, has offered datasets of statistically downscaled daily precipitation and daily minimum and maximum air temperature under three different Representative Concentration Pathways (RCPs) scenarios (RCP 2.6, 4.5, and 8.5) (Meinshausen

et al., 2011) over Canada at a spatial resolution of 300 arc-seconds (0.833° or ~ 10 km) for the historical and projected period of 1950 to 2100 (Pacific Climate Impacts Consortium; University of Victoria, Jan 2014). These downscaled datasets were a composite of 12 GCM projections from the Coupled Model Inter-comparison Project Phase 5 (CMIP5) (Taylor et al., 2012) and the ANUSPLIN dataset. The historical 1950 to 2005 period of the ANUSPLIN was used for bias-correction and downscaling of the GCMs. Two different methods were used to downscale to a finer resolution (Werner and Cannon, 2016). These included Bias Correction Spatial Disaggregation (BCSD) (Wood et al., 2004) following Maurer and Hidalgo (2008) and Bias Correction Constructed Analogues (BCCA) with Quantile mapping reordering (BCCAQ), which was a post-processed version of BCCA (Maurer et al., 2010). The ensemble of the PCIC dataset has currently been used in studying the hydrological impacts of climate change on river basins mainly in British Columbia (e.g. Shrestha et al., 2011; Shrestha et al., 2012b; Schnorbus et al., 2014) and Alberta (e.g. Kienzle et al., 2012; Forbes et al., 2011) in Canada. In this study, only four GCMs with two respective statistical downscaling methods were chosen for comparison (see Table 2 for details). The choice of the four GCMs was to match those available in the NA-CORDEX dataset (see next section for details).

3.2.5. GCM-driven RCM dynamically downscaled products – NA-CORDEX

Sponsored by the World Climate Research Programme (WCRP), the COordinated Regional climate Downscaling EXperiment (CORDEX) over North America domain (NA-CORDEX) provides dynamically downscaled datasets of 3-hourly or daily meteorological data over most of North America (below 80° N) at spatial resolutions of 0.22° and 0.44° (~ 25 and ~ 50 km) under RCP 4.5 and 8.5 for the historical (1950 – 2005) and future (2006 – 2100) period (Giorgi et al., 2009). Drawing from the strengths of the North American Regional Climate Change Assessment Program (NARCCAP) (Mearns et al., 2012), a matrix of six GCMs from the CMIP5 driving six different RCMs was selected to compare and characterize the uncertainties of RCMs and thus provided climate scenarios for further impact and adaption studies. Current studies using NA-CORDEX datasets were mainly focused on evaluating the model performance of different GCM-driven RCM simulations over North America (e.g. Lucas-Picher et al., 2013; Martynov et al., 2013; Separovic et al., 2013). In this study, two GCMs and three RCMs were chosen for comparison due to the availability of the NA-CORDEX dataset (see Table 3 for details).

4. Methodology

4.1. Pre-processing

Due to the different spatial and temporal resolutions of the various precipitation products, the first step was to re-grid each onto a common $0.5^\circ \times 0.5^\circ$ resolution to match the lowest-resolution dataset. It was acknowledged that re-gridding products onto a common spatial resolution might introduce more errors or uncertainties and the number of interpolation steps should be minimized. However, the main focus of this study was to inter-compare various gridded precipitation products using precipitation-gauge station data as a reference/benchmark but not to assess the individual accuracy of each product against the reference dataset. Therefore, upscaling to a common resolution provided a direct and more consistent inter-comparison. Such methodology was consistent with similar studies in the literature (e.g. Janowiak et al., 1998; Rauscher et al., 2010; Kimoto et al., 2005). All data were accumulated to daily time scale for comparison. Two common time spans were selected since CaPA covered a shorter timeframe compared to the rest of the products: (1) long-term comparison from January 1979 to December 2012 with the exclusion of CaPA (from January 1979 to December 2005 for PCIC and NA-CORDEX as the historical period of the datasets ends in 2005); and (2) short-term comparison from January 2002 to December 2012 when CaPA data are available. Daily values were summed over the four standard seasons (spring: March to May – MAM, summer: June to August – JJA, autumn: September to November – SON, and winter: December to February – DJF) to inter-compare the precipitation products at a seasonal scale.

To identify the most consistent gridded dataset corresponding to different seasons and regions, comparisons of each dataset with direct precipitation-gauge station data from the aforementioned AHCCD were carried out. For the period of 1979 to 2012, only 169 of the original 464 stations across Canada were available. This drastic drop was due to 271 stations ending before or after early 2000s and 23 not having a complete year of 2012. Subsequently, any of the 169 stations where the percentage of missing values exceeded 10 % during the study period were also eliminated. This resulted in 145 and 137 stations across Canada for long-term and short-term comparison respectively (see Fig. 1 for locations). Note that most of the stations are located in southern Canada with only 15 stations above 60° N.

Gridded-based precipitation estimates at the coordinates of the precipitation-gauge stations were then extracted by employing an inverse-distance-square weighting method (Cressman, 1959), which has been used to interpolate climate data for simple and efficient applications (Eum et al., 2014; Shen et al., 2001). This method assumes that an interpolated point is solely influenced by the nearby gridded points based on the inverse of the distance between the interpolated point and the gridded points. The interpolations were carried out on an individual ecodistrict basis and were based on both the number of precipitation-gauge stations and number of $0.5^\circ \times 0.5^\circ$ grid cells within the ecodistrict in question. For instance, when a single precipitation-gauge station was located within an ecodistrict, the value of the interpolated point was calculated by using all of the gridded points within that ecodistrict. When two or more precipitation-gauge stations were within the same ecodistrict, their interpolated values were calculated by using the same numbers of gridded points but with different weightings based on inverse distance. In the case when an ecodistrict contained one grid cell, no weighting was used and the interpolated value was equal to the nearest grid point.

4.2. Comparison of probability distributions using Kolmogorov-Smirnov test

A two-sample, non-parametric Kolmogorov-Smirnov (K-S) test was used to compare the cumulative distribution functions (CDFs) of gridded precipitation products with the AHCCD. The null hypothesis (H_0) was that the two datasets came from same population. For each season, monthly total precipitation data were used to avoid commonly known issues of numerous zero values in the daily precipitation data that might affect significance. The K-S test was repeated independently for all precipitation-gauge stations at 5 % significance level ($\alpha = 0.05$). A measure of reliability (in percent) was calculated based on counting the number of stations that do not reject the null hypothesis (any p -values greater than 0.05) over the total number of stations (145 and 137 stations in long-term and short-term comparison respectively), as shown in Eq. (1).

$$\% \text{ of reliability} = \frac{\text{number of stations that support } H_0}{\text{total number of precipitation gauge stations}} \cdot 100 \quad (1)$$

4.3. Comparison of gridded precipitation data using performance measures

Since the generation of the climate model-based precipitation products (PCIC dataset and NA-CORDEX dataset) only preserved the statistical properties without considering the day-by-day sequencing of precipitation events in the observational record, these two datasets were excluded

from the following comparison, which only focused on the station-based and reanalysis-based gridded products. In particular, these products were assessed in their ability to represent the daily variability of precipitation amounts in different ecozones by four performance measures: percentage of bias (P_{Bias}) (P_{Bias}), root-mean-square-error ($RMSE$) (E_{rms}), correlation coefficient (r), and standard deviation ratio (σ_G/σ_R), as shown by Eqs. (2) to (5), respectively.

$$P_{Bias;s} = \frac{\sum_i^N (G_i - R_i)}{\sum_i^N (R_i)} \cdot 100 \quad (2)$$

$$E_{rms;s} = \sqrt{\frac{\sum_i^N (G_i - R_i)^2}{N}} \quad (3)$$

$$r_s = \frac{\sum_i^N (G_i - \bar{G})(R_i - \bar{R})}{\sqrt{\sum_i^N (G_i - \bar{G})^2} \sqrt{\sum_i^N (R_i - \bar{R})^2}} \quad (4)$$

$$(\sigma_G/\sigma_R)_s = \frac{\sqrt{\frac{\sum_i^N (G_i - \bar{G})^2}{N}}}{\sqrt{\frac{\sum_i^N (R_i - \bar{R})^2}{N}}} \quad (5)$$

where s is the season, G and R are the spatial average of the daily gridded precipitation product and the reference observation dataset (precipitation-gauge stations) respectively, \bar{G} and \bar{R} are the daily mean of gridded precipitation product and point station data over the time spans (1979-2012 and 2002-2012), respectively, i is the i -th day of the season, and N is the total numbers of day in the season. These four performance measures examined different aspects of the gridded precipitation products, with P_{Bias} for accuracy of product estimation, $RMSE$ for magnitude of the errors, r for strength and direction of the linear relationship between gridded products and precipitation-gauge station data, and σ_G/σ_R for amplitude of the variations.

5. Results

5.1. Reliability of precipitation products

The percentage of reliability of each precipitation dataset during every season for the periods of 1979 to 2012 and 2002 to 2012 across Canada is shown in Fig. 2. The higher the percentage, the more reliable the precipitation dataset in question. In general, for long-term comparison (Fig. 2 left panel), WFDEI [GPCC] provided the highest percentage of reliability for the individual

seasons (from spring to winter: 72.5 %, 81.4 %, 70.3 %, and 50.3 %) while NARR had the lowest percentage (24.8 %, 45.5 %, 27.6 %, and 11.7 %). Therefore in spring, WFDEI [GPCC] is not significantly different for 72.5 % of the 145 precipitation-gauge stations while for NARR it is only 24.8 %. ANUSPLIN is second in spring and summer (56.6 % and 73.1 %) and WFDEI [CRU] in autumn and winter (63.4 % and 45.5 %).

Regarding the PCIC ensembles, the different GCMs provided a range of reliabilities for the individual seasons. MPI-ESM-LR performed the best in summer (70.2 %) and CanESM2 in autumn (45.5 %). GFDL-ESM2G generally gave more reliable estimates in spring and winter (57.4 % and 41.7 %). Overall, the performance of MPI-ESM-LR (52.0 %) was the best among the GCMs, followed by GFDL-ESM2G (50.1 %), CanESM2 (47.8 %), and HadGEM2 (36.2 %). In terms of statistical downscaling methods, the BCCAQ was on average slightly better than BCSD (49.5 % versus 44.0 %) with the former having a greater similarity in spring and summer as opposed to autumn and winter. These small differences therefore suggest that both methods are similar. With respect to the NA-CORDEX ensembles, the CRCM5 RCM gave the most reliable estimates in summer and autumn regardless of the GCM used. CanRCM4 had the best reliability in spring (49.4 %) whereas RegCM4 had the poorest reliability in spring and summer (24.4 % and 34.0 %). Overall, the reliability of MPI-ESM-LR (44.7 %) was better than that of CanESM2 (42.5 %) regardless of the RCMs used whereas the reliability of CRCM5 (43.6 %) was the best among the RCMs, followed by CanRCM4 (41.2 %), and RegCM4 (32.5 %). It should also be noted that in all cases, the gridded station-based and reanalysis-based products outperformed the climate model-simulated products.

With regard to the short-term comparison (Fig. 2 middle panel), ANUSPLIN showed better performance in summer with 94.1 % of reliability among the 137 precipitation-gauge stations while CaPA indicated better skill in winter with 68.6 % of reliability. Again, WFDEI [GPCC] in general provided the most consistent and reliable estimates with over 65 % of reliability in all four seasons. It is interesting to note that for the most part there is a higher percentage of reliability in short-term period compared to long-term period. Reasons for this are not clear but can be partly attributed to the fact that the power of K-S test (i.e. the probability of rejecting the null hypothesis when the alternative is true) decreases with the number of samples.

Figures 3, 4 and 5 display the seasonal distributions of p -values using the K-S test for long-term and short-term comparison, respectively. Due to the uneven distribution of precipitation-gauge stations across Canada, the number of stations in each ecozone are different (Table 4), with no stations in Region 1 (Arctic Cordillera), and Regions 2 to 5, 10, 12, and 15 have less than 10 stations. As a result, regions having more than or equal to 10 stations (6 to 9 and 13, 14) were only shown in box-whisker plots for illustration. Different colours in the figures corresponded to the various precipitation products. The higher the p -values (> 0.05) in each ecozone (represented by a thick black line in box-whisker plots towards 1 in y-axis in Figs. 3, 4 and 5), the more confidence we attribute to each gridded precipitation datasets in that ecozone.

From 1979 to 2012 (Fig. 3), the consistency of each type of precipitation products is explored by assessing the median of the p -values. Overall, all the precipitation products showed very low reliability and consistency in winter among these ecozones and in every season in Regions 13 and 14 (Pacific Maritime and Montane Cordillera) as the medians were close to zero, despite a couple of locations having higher chance of same CDFs as in the precipitation-gauge station data. The WFDEI [GPCC] dataset provided the highest consistency in the remaining three seasons except for Region 7 (Atlantic Maritime) where ANUSPLIN showed higher medians (0.51 and 0.46) than WFDEI [GPCC] (0.42 and 0.42) in spring and autumn respectively. Noticeably NARR provided the lowest median among the reanalysis-based datasets in all four seasons in Regions 6 to 8 but gave fairly consistent estimates in Regions 9 and 10, especially in summer in Region 9 (Boreal Plain) where it came second after WFDEI [GPCC]. The medians of Princeton were similar with those of ANUSPLIN on average in these regions except for summer in which ANUSPLIN offered higher medians than Princeton. WFDEI [CRU] generally showed consistent estimates among these ecozones with medians well above 0.05 except for Region 7 (Atlantic Maritime) in spring and autumn. From 1979 to 2005 (Fig. 5), the PCIC ensembles and the NA-CORDEX ensembles showed different degrees of consistency among their GCM members with generally higher p -values using BCCAQ method than BCSD method in spring and summer regardless of GCMs in the PCIC datasets. CanESM2 was generally having higher consistency and reliable estimates than MPI-ESM-LR in spring and summer but opposite case in autumn in the NA-CORDEX ensembles. In addition, almost all the precipitation products had lower chance of having same CDFs as the precipitation-gauge stations in ecozones above 60° N (Regions 2 to 5, 11, and 12) (figure not shown).

For the shorter time period of 2002 to 2012 (Fig. 4), CaPA showed the highest consistency in winter in Regions 6, 8, 9, and 13 whereas ANUSPLIN was the highest in summer in Regions 8, 13, and 14, echoing the results found in Fig. 2. However, the reliability and consistency of CaPA in summer was not particularly high, especially in Regions 8 and 13 where the medians were approaching zero. In addition, in ecozones above 60° N, similar performances were seen among the precipitation products in the period of 2002 to 2012 as compared with the long-term performance.

5.2. Daily variability of precipitation (station- and reanalysis-based products)

The accuracy (*PBias*), magnitude of the errors (*RMSE*), strength and direction of the relationship between gridded products and precipitation-gauge station data (*r*), and amplitude of the variations (σ_G/σ_R) are shown in Figs. 6 and 7 for the period of 1979 to 2012. In general, the gridded precipitation products that agree well with the precipitation-gauge station data should have relatively high correlation and low RMSE, low bias and similar standard deviation (light grey or dark grey squares in Figs. 6 and 7).

In terms of accuracy (Fig. 6 left panel), all precipitation products tended to generally overestimate total precipitation in Regions 12 to 14, while Region 14 (Montane Cordillera) had the overall highest positive *PBias* for the individual seasons (from spring to winter: >20.9 %, >6.24 %, >14.4 %, and >26.8 %). On the other hand, all products mostly underestimated the precipitation amounts in Regions 3 to 6, 9, and 10. This was especially worse in Region 3 (Southern Arctic) where the underestimation of precipitation amounts for the individual seasons were >-22.6 %, >-2.2 %, >-10.2 %, and >-28.1 %, respectively. In particular, ANUSPLIN was associated with a generally negative *PBias* for all the ecozones in four seasons, except for Regions 12 (Boreal Cordillera) and 14 (Montane Cordillera). The accuracy of ANUSPLIN was the worst in winter, with underestimation of precipitation amounts ranging from -7.8 % in Region 13 (Pacific Maritime) to -38.7 % in Region 3 (Southern Arctic). WFDEI [CRU] and WFDEI [GPCC] had similar performances across different regions. They performed particularly well in summer in Regions 2 to 9 where the accuracy was within -4.6 % to 4.2 %. With the exception of Regions 13 and 14, Princeton and NARR generally provided the overall largest and second largest underestimation of precipitation amounts across different ecozones. NARR performed the worst in Regions 7

(Atlantic Maritime) and 8 (Mixedwood Plain) where the precipitation amounts for the individual seasons were underestimated by $>-42.0\%$, $>-33.1\%$, $>-38.8\%$, and $>-59.7\%$.

When examining the magnitude of errors (Fig. 7 left panel), all products showed very high magnitude of errors in Regions 6 to 8, and 13, while Region 13 (Pacific Maritime) had the greatest *RMSE* for the individual seasons (from spring to winter: >5.35 mm/day, >3.74 mm/day, >7.82 mm/day, and >8.24 mm/day). Specifically, ANUSPLIN showed generally better correspondence with precipitation-gauge station data, providing the overall lowest *RMSE* across ecozones in four seasons (2.50 mm/day, 3.24 mm/day, 2.79 mm/day, and 2.45 mm/day) with the only exception in spring in Region 15 (Hudson Plain). Moreover, referring to Fig. 7 (right panel), ANUSPLIN had the overall highest *r* across ecozones in four seasons (0.75, 0.78, 0.80, and 0.74). On the contrary, Princeton had the worst performance in both magnitude of errors and correlation with observations irrespective of ecozone or season, with the grand *RMSE* and *r* of 5.65 mm/day and 0.17 respectively. The performances of WFDEI [CRU], WFDEI [GPCC], and NARR were in between ANUSPLIN and Princeton and they shared similar *RMSE* and *r* across different regions and seasons. The resulting values of the *RMSE* metric in Regions 7 (Atlantic Maritime) and 13 (Pacific Maritime) tended to be larger than that of other ecozones. However, the other metrics such as *PBias* and *r* showed better performance in these regions. This suggests that higher *RMSE* values can be mainly attributed to the fact that precipitation amounts are higher in the maritime regions.

Regarding the amplitude of variations (Fig. 6 right panel), all datasets generally had variations that were much smaller than precipitation-gauge station data in Regions 3, 4, and 11 in four seasons. In particular, ANUSPLIN and NARR were consistently having too little variability across different ecozones, especially in winter in which σ_G/σ_R ranged from 0.41 in Region 15 (Hudson Plain) to 0.76 in Region 13 (Pacific Maritime). WFDEI [CRU] and WFDEI [GPCC] had the most similar standard deviations as that of precipitation-gauge station data in Regions 5 to 8 in autumn and winter, while Princeton estimated σ_G/σ_R the best in Regions 4 to 10 in summer. However, Princeton had much larger variability in Regions 12 to 14 in spring and Regions 6 to 8 in autumn.

Concerning the short-term comparison (Table 5), CaPA performed the best in spring and autumn in terms of accuracy, with the lowest positive *PBias* of 0.7 % and the lowest negative *PBias* of -1.3 % respectively. The performance of CaPA generally resembled that of ANUSPLIN regarding the magnitude of errors and correlation with observations, which were the second lowest *RMSE*

for the individual seasons (from spring to winter: 2.70 mm/day, 3.74 mm/day, 3.35 mm/day, and 3.05 mm/day) and the second highest r (0.72, 0.73, 0.75, and 0.70), respectively. Despite its better performances in $RMSE$ and r , CaPA was generally not able to capture satisfactorily the amplitude of variations, with consistently lower values in four seasons (0.83, 0.82, 0.85, and 0.72). However, CaPA showed more skill compared to ANUSPLIN (0.72, 0.76, 0.74, and 0.64) and NARR (0.75, 0.75, 0.72, and 0.63). In addition, the five gridded products in the long-term comparison performed similarly in the period of 2002 to 2012, with ANUSPLIN having the lowest annual $RMSE$ and highest annual r of 3.00 mm/day and 0.79 and Princeton being the worst again with the highest annual $RMSE$ and lowest annual r of 6.33 mm/day and 0.17 respectively.

6. Discussion

The preceding has provided insight into the relative performance of various gridded precipitation products over Canada relative to gauge measurements over different seasons and ecozones. Results showed that there is no particular product that is superior for all performance measures although some datasets are consistently better. Based on the performances, one could broadly characterize the station- and reanalysis-based precipitation products into four groups: (1) ANUSPLIN and CaPA with negative $PBias$, low $RMSE$, high r , and small σ_G/σ_R ; (2) WFDEI [CRU] and WFDEI [GPCC], with relatively small $PBias$, high $RMSE$, fair r , and similar standard deviation; (3) Princeton, with negative $PBias$, high $RMSE$, low r , and a mixture of large and small σ_G/σ_R ; and (4) NARR, with negative $PBias$, high $RMSE$, fair r , and small σ_G/σ_R . Among the reanalysis-based gridded products, Princeton performed the worst in all seasons and regions in terms of minimizing error magnitudes (Figs. 8 and 9). Princeton was especially poor in winter (Fig. 8) and showed significant underestimation in regions above 60° N (Fig. 9). This could be due to the use of the NCEP-NCAR reanalysis as the basis to generate the dataset, which have been shown to be less accurate than NCEP-DOE reanalysis (used in NARR) and ERA-40 reanalysis (used in WFD) (Sheffield et al., 2006). The better performance of NARR in capturing the timings and amounts of precipitation compared to Princeton was probably because NCEP-DOE reanalysis was a major improvement upon the earlier NCEP-NCAR reanalysis in both resolution and accuracy. However, the overall reliability of NARR was among the poorest mainly because of non-assimilation of gauge precipitation observations over Canada from 2004 onwards, as reported by Mesinger et al. (2006). ANUSPLIN and CaPA performed well in capturing the timings and minimizing the error

magnitudes of the precipitation, despite their general underestimation across Canada ($PBias$ ranging from -7.7 % (Region 13) to -40.7 % (Region 3) and -2.0 % (Region 15) to -17.1 % (Region 8) in the period of 2002 to 2012) (Fig. 9) and too little variability (grand σ_G/σ_R of 0.72 and 0.80 of the same period). This was not surprising given that the generation of the products was based on the unadjusted precipitation-gauge stations where the total rainfall amounts were increased after adjustment (Mekis and Vincent, 2011). WFDEI [CRU] and WFDEI [GPCC], on the other hand, performed well in estimating the accuracy and amplitude of variations, but not the timings and error magnitudes of the precipitation. This could probably due to the positive bias offsetting the negative bias resulting in small mean bias, but was picked up by $RMSE$ that gives more weights to the larger errors. The larger errors could result from a mismatch of occurrence of precipitation in the time series, as reflected by the fair correlation coefficients (grand r of 0.52 and 0.50 for WFDEI [CRU], 0.54 and 0.53 for WFDEI [GPCC], for time periods of 1979 to 2012 and 2002 to 2012 respectively).

By matching the statistical properties of the adjusted gauge measurements at monthly time scale, one could establish the confidence in using the climate model-simulated products for long-term hydro-climatic studies. Comparing the overall reliability of the PCIC and NA-CORDEX datasets, it was found that for the individual seasons the PCIC ensembles (spring, summer, and winter: 54.0 %, 64.7 %, and 35.7 %) outperformed the NA-CORDEX ensembles (39.1 %, 45.0 %, and 31.3 %) except in autumn when the NA-CORDEX ensembles (45.5 %) provided slightly higher reliability than the PCIC ensembles (45.2 %). The better reliability of the PCIC datasets could be due to the use of ANUSPLIN to train the GCMs and thus, the statistical properties of the downscaled outputs are guided by those of the ANUSPLIN. Similarly, for ecozones where more than 10 precipitation-gauge stations could be found (Regions 6 to 9, 13 and 14), the PCIC ensembles (reliability ranging from 35.7 % to 64.4 %) also outperformed the NA-CORDEX ensembles (from 17.2 % to 61.6 %). This would suggest that the PCIC ensembles may be the preferred choice for long-term climate change impact assessment over Canada, although further research is required.

The evaluations of this comparison were impacted by the spatial distribution of adjusted precipitation-gauge stations (Mekis and Vincent, 2011), which were assumed to be the best representation of reality owing to efforts in improving the raw archive of the precipitation-gauge stations. However, the major limitation of this dataset was the number of precipitation-gauge

stations that could be used for comparison. As aforementioned, due to temporal coverage not encompassing the entire study period and not having a complete year of 2012, over half of the precipitation-gauge stations were discarded from the analysis. Although the locations of the remaining stations covered much of Canada, there are only one or a few stations located in some of the ecozones (e.g. Region 3 to 5, 11, and 15). Even in Region 10 (Prairie) there are only nine precipitation-gauge stations for analysis. While the reliability of different types of gridded products could be tested in these ecozones, the consistency of the performance of each gridded product could not be established due to small sample sizes.

In addition, results from the above analysis should be interpreted with care because the precipitation-gauge station data are point measurements whereas the gridded precipitation products are areal averages, of which the accuracy and precision of the estimates can be very different given the non-linear responses of precipitation (Ebert et al., 2007). When comparing point measurements and areal-average estimates, fundamental challenges occur because of the sampling errors arising from different sampling schemes and errors related to gauge instrumentation (Bowman, 2005). It is therefore difficult to have perfect spatial matching between point measurements (gauge stations) and areal-averaged estimates (gridded products) (Sapiano and Arkin, 2009; Hong et al., 2007). However, in the absence of a sufficiently dense precipitation gauge network in Canada, the options for assessing different gridded products are limited. The only gridded product that is basically representing areal averages of precipitation (via interpolation) based on ground observations is ANUSPLIN. As aforementioned (see Sect. 3.2.1), this product has its own limitations and may not be qualified to be considered as the “ground truth”. Therefore, ANUSPLIN is also included in the pool of gridded products to be evaluated. Notwithstanding the issues, using the selected gauge measurements would remain the best way for the evaluation of the multiple gridded products because the set of gauges used had been adjusted (e.g. for undercatch) and are the most accurate source of information on precipitation in Canada (although small with limited spatial coverage). Also, given that all the gridded products are compared against this common set of station observations, it is assumed that the bias that the difference between point and areal data introduces into the analysis is consistent for all the products. Therefore, given the current data situation, the preceding methods could be used for comparing the performance of different daily gridded precipitation products.

7. Conclusion

A number of gridded climate products incorporating multiple sources of data have recently been developed with the aim of providing better and more reliable measurements for climate and hydrological studies. There is a pressing need for characterizing the quality and error characteristics of various precipitation products and assessing how they perform at different spatial and temporal scales. This is particularly important in light of the fact that these products are the main driver of hydrological models in many regions, including Canadian watersheds where precipitation-gauge network is typically limited and sparse. This study was conducted to inter-compare several gridded precipitation products of their probability distributions and quantify the spatial and temporal variability of the errors relative to station observations in Canada, so as to provide some insights for potential users in selecting the products for their particular interests and applications. Based on the above analysis, the following conclusions can be drawn:

- In general, all the products performed best in summer, followed by autumn, spring, and winter in order of decreasing quality. The lower reliability in winter is likely the result of difficulty in accurately capturing solid precipitation.
- Overall, WFDEI [GPCC] and CaPA performed best with respect to different performance measures. WFDEI [GPCC], however, may be a better choice for long-term analyses as it covers a longer historical period. ANUSPLIN and WFDEI [CRU] also performed comparably, with considerably lower quality than WFDEI [GPCC] and CaPA. Princeton and NARR demonstrated the lowest quality in terms of different performance measures.
- Station-based and reanalysis-based products tended to underestimate total precipitation across Canada except in southwestern regions (Pacific Maritime and Montane Cordillera) where the tendency was towards overestimation. This may be due to the fact that the majority of precipitation-gauge stations are located at lower altitudes which might not accurately reflect areal precipitation due to topographic effect.
- In southern Canada, WFDEI [GPCC] and CaPA demonstrated their best performance in the western cold interior (Boreal Plain, Prairie, Montane Cordillera) in terms of timing and magnitude of daily precipitation.

- In northern Canada (above 60° N), the different products tended to moderately (ranging from -0.6 % to -40.3 %) and in cases significantly (up to -60.3 % in Taiga Cordillera) underestimate total precipitation, while reproducing the timing of daily precipitation rather well. It should be noted that this assessment was based on only a limited number of precipitation-gauges in the north.
- Comparing the climate model-simulated products, PCIC ensembles generally performed better than NA-CORDEX ensembles in terms of reliability and consistency in four seasons across Canada.
- In terms of statistical downscaling methods, the BCCAQ method was slightly more reliable than the BCSD method across Canada on the annual basis.
- Regarding GCMs, MPI-ESM-LR provides the highest reliability, followed by GFDL-ESM2G, CanESM2, and HadGEM2. With respect to RCMs, CRCM5 performed the best regardless of the GCM used, followed by CanRCM4, and RegCM4.

The findings from this analysis provide additional information for potential users to draw inferences about the relative performance of different gridded products. Although no clear-cut product was shown to be superior, researchers/users can use this information for selecting or excluding various datasets depending on their purpose of study. It is realized that this investigation only focused on the daily time scale at a relatively coarse 0.5° x 0.5° resolution suitable for large-scale hydro-climatic studies. Further research is thus required towards performance assessment of various products with respect to precipitation extremes, which often have the greatest hydro-climatic impacts. As new products become available, similar comparisons should be conducted to assess their reliability.

Acknowledgements

The financial support from the Canada Excellence Research Chair in Water Security is gratefully acknowledged. Thanks are due to Melissa Bukovsky and Katja Winger from the NA-CORDEX modelling group for providing access to RegCM4 and CRCM5 data used in this study. The authors are also grateful to the various organizations that made the datasets freely available to the scientific community.

References

- Adam, J. C., and Lettenmaier, D. P.: Adjustment of global gridded precipitation for systematic bias, *J Geophys Res-Atmos*, 108, Artn 4257 10.1029/2002jd002499, 2003.
- Adler, R. F., Kidd, C., Petty, G., Morissey, M., and Goodman, H. M.: Intercomparison of global precipitation products: The third Precipitation Intercomparison Project (PIP-3), *B Am Meteorol Soc*, 82, 1377-1396, Doi 10.1175/1520-0477(2001)082<1377:Iogppt>2.3.Co;2, 2001.
- Asadullah, A., McIntyre, N., and Kigobe, M.: Evaluation of five satellite products for estimation of rainfall over Uganda, *Hydrolog Sci J*, 53, 1137-1150, DOI 10.1623/hysj.53.6.1137, 2008.
- Asong, Z. E., Khaliq, M. N., and Wheeler, H. S.: Regionalization of precipitation characteristics in the Canadian Prairie Provinces using large-scale atmospheric covariates and geophysical attributes, *Stoch Env Res Risk A*, 29, 875-892, 10.1007/s00477-014-0918-z, 2015.
- Behrangi, A., Christensen, M., Lebsock, M. R., Stephens, G., Huffman, G. J., Bolvin, D., Adler, R. F., Gardner, A., Lambriksen, B., and Fetzer, E.: Status of High latitude precipitation estimates from observations and reanalyses, *Journal of Geophysical Research: Atmospheres*, 2016.
- Betts, A. K., Zhao, M., Dirmeyer, P. A., and Beljaars, A. C. M.: Comparison of ERA40 and NCEP/DOE near-surface data sets with other ISLSCP-II data sets, *J Geophys Res-Atmos*, 111, Artn D22s04 10.1029/2006jd007174, 2006.
- Bhargava, M., and Danard, M.: Application of Optimum Interpolation to the Analysis of Precipitation in Complex Terrain, *J Appl Meteorol*, 33, 508-518, Doi 10.1175/1520-0450(1994)033<0508:Aooitt>2.0.Co;2, 1994.
- Black, T. L.: The step-mountain, eta coordinate regional model: A documentation, National Meteorological Center, Development Division, 1988.
- Blenkinsop, S., and Fowler, H. J.: Changes in European drought characteristics projected by the PRUDENCE regional climate models, *Int J Climatol*, 27, 1595-1610, 10.1002/joc.1538, 2007.
- Bonsal, B. R., Aider, R., Gachon, P., and Lapp, S.: An assessment of Canadian prairie drought: past, present, and future, *Clim Dynam*, 41, 501-516, 10.1007/s00382-012-1422-0, 2013.
- Bosilovich, M. G., Chen, J. Y., Robertson, F. R., and Adler, R. F.: Evaluation of global precipitation in reanalyses, *J Appl Meteorol Clim*, 47, 2279-2299, 10.1175/2008jamc1921.1, 2008.
- Bowman, K. P.: Comparison of TRMM precipitation retrievals with rain gauge data from ocean buoys, *J Climate*, 18, 178-190, Doi 10.1175/Jcli3259.1, 2005.
- Brooks, R., Harvey, K., Kirk, D., Soulard, F., Paul, P., and Murray, A.: Building a Canadian Digital Drainage Area Framework, 55 th Annual CWRA Conference, Winnipeg, Manitoba, Canada, 2002,
- Bukovsky, M. S., and Karoly, D. J.: A brief evaluation of precipitation from the North American Regional Reanalysis, *J Hydrometeorol*, 8, 837-846, 10.1175/Jhm595.1, 2007.
- Carrera, M. L., Belair, S., and Bilodeau, B.: The Canadian Land Data Assimilation System (CaLDAS): Description and Synthetic Evaluation Study, *J Hydrometeorol*, 16, 1293-1314, 10.1175/Jhm-D-14-0089.1, 2015.
- Chadburn, S. E., Burke, E. J., Essery, R. L. H., Boike, J., Langer, M., Heikenfeld, M., Cox, P. M., and Friedlingstein, P.: Impact of model developments on present and future simulations of permafrost in a global land-surface model, *Cryosphere*, 9, 1505-1521, 10.5194/tc-9-1505-2015, 2015.

Chen, D. L., Achberger, C., Raisanen, J., and Hellstrom, C.: Using statistical downscaling to quantify the GCM-related uncertainty in regional climate change scenarios: A case study of Swedish precipitation, *Adv Atmos Sci*, 23, 54-60, DOI 10.1007/s00376-006-0006-5, 2006.

Choi, W., Kim, S. J., Rasmussen, P. F., and Moore, A. R.: Use of the North American Regional Reanalysis for Hydrological Modelling in Manitoba, *Can Water Resour J*, 34, 17-36, 2009.

Christensen, J. H., Carter, T. R., Rummukainen, M., and Amanatidis, G.: Evaluating the performance and utility of regional climate models: the PRUDENCE project, *Climatic Change*, 81, 1-6, 10.1007/s10584-006-9211-6, 2007.

Ciach, G. J.: Local random errors in tipping-bucket rain gauge measurements, *J Atmos Ocean Tech*, 20, 752-759, Doi 10.1175/1520-0426(2003)20<752:Lreith>2.0.Co;2, 2003.

Cote, J., Desmarais, J. G., Gravel, S., Methot, A., Patoine, A., Roch, M., and Staniforth, A.: The operational CMC-MRB Global Environmental Multiscale (GEM) model. Part II: Results, *Monthly Weather Review*, 126, 1397-1418, Doi 10.1175/1520-0493(1998)126<1397:Tocmge>2.0.Co;2, 1998a.

Cote, J., Gravel, S., Methot, A., Patoine, A., Roch, M., and Staniforth, A.: The operational CMC-MRB Global Environmental Multiscale (GEM) model. Part I: Design considerations and formulation, *Monthly Weather Review*, 126, 1373-1395, Doi 10.1175/1520-0493(1998)126<1373:Tocmge>2.0.Co;2, 1998b.

Covey, C., AchutaRao, K. M., Cubasch, U., Jones, P., Lambert, S. J., Mann, M. E., Phillips, T. J., and Taylor, K. E.: An overview of results from the Coupled Model Intercomparison Project, *Global Planet Change*, 37, 103-133, 10.1016/S0921-8181(02)00193-5, 2003.

Cressman, G. P.: An operational objective analysis system, *Monthly Weather Review*, 87, 367-374, 1959.

Cuo, L., Beyene, T. K., Voisin, N., Su, F. G., Lettenmaier, D. P., Alberti, M., and Richey, J. E.: Effects of mid-twenty-first century climate and land cover change on the hydrology of the Puget Sound basin, Washington, *Hydrol Process*, 25, 1729-1753, 10.1002/hyp.7932, 2011.

Daley, R.: *Atmospheric data analysis*, 2, Cambridge university press, 1993.

Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Holm, E. V., Isaksen, L., Kallberg, P., Kohler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J. J., Park, B. K., Peubey, C., de Rosnay, P., Tavolato, C., Thepaut, J. N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Q J Roy Meteor Soc*, 137, 553-597, 10.1002/qj.828, 2011.

Dinku, T., Anagnostou, E. N., and Borga, M.: Improving radar-based estimation of rainfall over complex terrain, *J Appl Meteorol*, 41, 1163-1178, Doi 10.1175/1520-0450(2002)041<1163:Irbeor>2.0.Co;2, 2002.

Dinku, T., Connor, S. J., Ceccato, P., and Ropelewski, C. F.: Comparison of global gridded precipitation products over a mountainous region of Africa, *Int J Climatol*, 28, 1627-1638, 10.1002/joc.1669, 2008.

Dore, M. H. I.: Climate change and changes in global precipitation patterns: What do we know?, *Environ Int*, 31, 1167-1181, 10.1016/j.envint.2005.03.004, 2005.

Ebert, E. E., Janowiak, J. E., and Kidd, C.: Comparison of near-real-time precipitation estimates from satellite observations and numerical models, *B Am Meteorol Soc*, 88, 47-+, 10.1175/Bams-88-1-47, 2007.

729 Ecological Stratification Working Group: A national ecological framework for Canada, Centre for Land and
730 Biological Resources Research, 1996.

731 Ek, M. B., Mitchell, K. E., Lin, Y., Rogers, E., Grunmann, P., Koren, V., Gayno, G., and Tarpley, J. D.:
732 Implementation of Noah land surface model advances in the National Centers for Environmental Prediction
733 operational mesoscale Eta model, *J Geophys Res-Atmos*, 108, Artn 8851 10.1029/2002jd003296, 2003.

734 Eum, H. I., Gachon, P., Laprise, R., and Ouarda, T.: Evaluation of regional climate model simulations versus gridded
735 observed and regional reanalysis products using a combined weighting scheme, *Clim Dynam*, 38, 1433-1457,
736 10.1007/s00382-011-1149-3, 2012.

737 Eum, H. I., Dibike, Y., Prowse, T., and Bonsal, B.: Inter-comparison of high-resolution gridded climate data sets and
738 their implication on hydrological model simulation over the Athabasca Watershed, Canada, *Hydrol Process*, 28, 4250-
739 4271, 10.1002/hyp.10236, 2014.

740 Forbes, K. A., Kienzie, S. W., Coburn, C. A., Byrne, J. M., and Rasmussen, J.: Simulating the hydrological response
741 to predicted climate change on a watershed in southern Alberta, Canada, *Climatic Change*, 105, 555-576,
742 10.1007/s10584-010-9890-x, 2011.

743 Fortin, V., Jean, M., Brown, R., and Payette, S.: Predicting Snow Depth in a Forest-Tundra Landscape using a
744 Conceptual Model Allowing for Snow Redistribution and Constrained by Observations from a Digital Camera, *Atmos*
745 *Ocean*, 53, 200-211, 10.1080/07055900.2015.1022708, 2015a.

746 Fortin, V., Roy, G., Donaldson, N., and Mahidjiba, A.: Assimilation of radar quantitative precipitation estimations in
747 the Canadian Precipitation Analysis (CaPA), *J Hydrol*, 531, 296-307, 2015b.

748 Frei, C., Scholl, R., Fukutome, S., Schmidli, J., and Vidale, P. L.: Future change of precipitation extremes in Europe:
749 Intercomparison of scenarios from regional climate models, *J Geophys Res-Atmos*, 111, Artn D06105
750 10.1029/2005jd005965, 2006.

751 Fuchs, T.: GPCC annual report for year 2008: Development of the GPCC data base and analysis products, DWD Rep,
752 2009.

753 Garand, L., and Grassotti, C.: Toward an Objective Analysis of Rainfall Rate Combining Observations and Short-
754 Term Forecast Model Estimates, *J Appl Meteorol*, 34, 1962-1977, Doi 10.1175/1520-
755 0450(1995)034<1962:Taoaor>2.0.Co;2, 1995.

756 Gebregiorgis, A. S., and Hossain, F.: How well can we estimate error variance of satellite precipitation data around
757 the world?, *Atmos Res*, 154, 39-59, 10.1016/j.atmosres.2014.11.005, 2015.

758 Gebremichael, M., Krajewski, W. F., Morrissey, M. L., Huffman, G. J., and Adler, R. F.: A detailed evaluation of
759 GPCP 1 degrees daily rainfall estimates over the Mississippi river basin, *J Appl Meteorol*, 44, 665-681, Doi
760 10.1175/Jam2233.1, 2005.

761 Giorgi, F., Jones, C., and Asrar, G. R.: Addressing climate information needs at the regional level: the CORDEX
762 framework, *World Meteorological Organization (WMO) Bulletin*, 58, 175, 2009.

763 Gottschalck, J., Meng, J., Rodell, M., and Houser, P.: Analysis of multiple precipitation products and preliminary
764 assessment of their impact on global land data assimilation system land surface states, *J Hydrometeorol*, 6, 573-598,
765 Doi 10.1175/Jhm437.1, 2005.

766 Gupta, S. K., Ritchey, N. A., Wilber, A. C., Whitlock, C. H., Gibson, G. G., and Stackhouse, P. W.: A climatology of
 767 surface radiation budget derived from satellite data, *J Climate*, 12, 2691-2710, Doi 10.1175/1520-
 768 0442(1999)012<2691:Acosrb>2.0.Co;2, 1999.
 769 Hively, W. D., Gerard-Marchant, P., and Steenhuis, T. S.: Distributed hydrological modeling of total dissolved
 770 phosphorus transport in an agricultural landscape, part II: dissolved phosphorus transport, *Hydrol Earth Syst Sc*, 10,
 771 263-276, 2006.
 772 Hong, Y., Gochis, D., Cheng, J. T., Hsu, K. L., and Sorooshian, S.: Evaluation of PERSIANN-CCS rainfall
 773 measurement using the NAME Event Rain Gauge Network, *J Hydrometeorol*, 8, 469-482, 10.1175/Jhm574.1, 2007.
 774 Hong, Y., Adler, R. F., Huffman, G. J., and Pierce, H.: Applications of TRMM-Based Multi-Satellite Precipitation
 775 Estimation for Global Runoff Prediction: Prototyping a Global Flood Modeling System, *Satellite Rainfall Applications*
 776 *for Surface Hydrology*, 245-265, 10.1007/978-90-481-2915-7_15, 2010.
 777 Hopkinson, R. F., McKenney, D. W., Milewska, E. J., Hutchinson, M. F., Papadopol, P., and Vincent, L. A.: Impact
 778 of Aligning Climatological Day on Gridding Daily Maximum-Minimum Temperature and Precipitation over Canada,
 779 *J Appl Meteorol Clim*, 50, 1654-1665, 10.1175/2011jamec2684.1, 2011.
 780 Hou, A. Y., Kakar, R. K., Neeck, S., Azarbarzin, A. A., Kummerow, C. D., Kojima, M., Oki, R., Nakamura, K., and
 781 Iguchi, T.: The Global Precipitation Measurement Mission, *B Am Meteorol Soc*, 95, 701-+, 10.1175/Bams-D-13-
 782 00164.1, 2014.
 783 Huffman, G. J., Adler, R. F., Morrissey, M. M., Bolvin, D. T., Curtis, S., Joyce, R., McGavock, B., and Susskind, J.:
 784 Global precipitation at one-degree daily resolution from multisatellite observations, *J Hydrometeorol*, 2, 36-50, Doi
 785 10.1175/1525-7541(2001)002<0036:Gpaodd>2.0.Co;2, 2001.
 786 Huffman, G. J., Adler, R. F., Stocker, E., Bolvin, D. T., and Nelkin, E. J.: Analysis of TRMM 3-hourly multi-satellite
 787 precipitation estimates computed in both real and post-real time, 2002.
 788 Huisman, J. A., Breuer, L., Bormann, H., Bronstert, A., Croke, B. F. W., Frede, H. G., Graff, T., Hubrechts, L.,
 789 Jakeman, A. J., Kite, G., Lanini, J., Leavesley, G., Lettenmaier, D. P., Lindstrom, G., Seibert, J., Sivapalan, M., Viney,
 790 N. R., and Willems, P.: Assessing the impact of land use change on hydrology by ensemble modeling (LUCHEM) III:
 791 Scenario analysis, *Adv Water Resour*, 32, 159-170, 10.1016/j.advwatres.2008.06.009, 2009.
 792 Hutchinson, M. F., Mckenney, D. W., Lawrence, K., Pedlar, J. H., Hopkinson, R. F., Milewska, E., and Papadopol,
 793 P.: Development and Testing of Canada-Wide Interpolated Spatial Models of Daily Minimum-Maximum Temperature
 794 and Precipitation for 1961-2003, *J Appl Meteorol Clim*, 48, 725-741, 10.1175/2008jamc1979.1, 2009.
 795 Jameson, A. R., and Kostinski, A. B.: Spurious power-law relations among rainfall and radar parameters, *Q J Roy*
 796 *Meteor Soc*, 128, 2045-2058, Doi 10.1256/003590002320603520, 2002.
 797 Janowiak, J. E., Gruber, A., Kondragunta, C. R., Livezey, R. E., and Huffman, G. J.: A comparison of the NCEP-
 798 NCAR reanalysis precipitation and the GPCP rain gauge-satellite combined dataset with observational error
 799 considerations, *J Climate*, 11, 2960-2979, Doi 10.1175/1520-0442(1998)011<2960:Acotnn>2.0.Co;2, 1998.
 800 Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen,
 801 J., Zhu, Y., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Leetmaa,

802 A., Reynolds, R., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-year reanalysis project, *B Am Meteorol Soc*, 77,
 803 437-471, Doi 10.1175/1520-0477(1996)077<0437:Tnyrp>2.0.Co;2, 1996.
 804 Kanamitsu, M., Ebisuzaki, W., Woollen, J., Yang, S. K., Hnilo, J. J., Fiorino, M., and Potter, G. L.: NCEP-DOE
 805 AMIP-II reanalysis (R-2), *B Am Meteorol Soc*, 83, 1631-1643, 10.1175/Bams-83-11-1631, 2002.
 806 Kang, D. H., Shi, X. G., Gao, H. L., and Dery, S. J.: On the Changing Contribution of Snow to the Hydrology of the
 807 Fraser River Basin, Canada, *J Hydrometeorol*, 15, 1344-1365, 10.1175/Jhm-D-13-0120.1, 2014.
 808 Kay, A. L., Davies, H. N., Bell, V. A., and Jones, R. G.: Comparison of uncertainty sources for climate change impacts:
 809 flood frequency in England, *Climatic Change*, 92, 41-63, 10.1007/s10584-008-9471-4, 2009.
 810 Kidd, C., Bauer, P., Turk, J., Huffman, G. J., Joyce, R., Hsu, K. L., and Braithwaite, D.: Intercomparison of High-
 811 Resolution Precipitation Products over Northwest Europe, *J Hydrometeorol*, 13, 67-83, 10.1175/Jhm-D-11-042.1,
 812 2012.
 813 Kienzie, S. W., Nemeth, M. W., Byrne, J. M., and MacDonald, R. J.: Simulating the hydrological impacts of climate
 814 change in the upper North Saskatchewan River basin, Alberta, Canada, *J Hydrol*, 412, 76-89,
 815 10.1016/j.jhydrol.2011.01.058, 2012.
 816 Kimoto, M., Yasutomi, N., Yokoyama, C., and Emori, S.: Projected Changes in Precipitation Characteristics around
 817 Japan under the Global Warming, *Sola*, 1, 85-88, 10.2151/sola.2005.023, 2005.
 818 Kistler, R., Kalnay, E., Collins, W., Saha, S., White, G., Woollen, J., Chelliah, M., Ebisuzaki, W., Kanamitsu, M.,
 819 Kousky, V., van den Dool, H., Jenne, R., and Fiorino, M.: The NCEP-NCAR 50-year reanalysis: Monthly means CD-
 820 ROM and documentation, *B Am Meteorol Soc*, 82, 247-267, Doi 10.1175/1520-
 821 0477(2001)082<0247:Tnnym>2.3.Co;2, 2001.
 822 Lespinas, F., Fortin, V., Roy, G., Rasmussen, P., and Stadnyk, T.: Performance Evaluation of the Canadian
 823 Precipitation Analysis (CaPA), *J Hydrometeorol*, 16, 2045-2064, 10.1175/Jhm-D-14-0191.1, 2015.
 824 Lucas-Picher, P., Somot, S., Deque, M., Decharme, B., and Alias, A.: Evaluation of the regional climate model
 825 ALADIN to simulate the climate over North America in the CORDEX framework, *Clim Dynam*, 41, 1117-1137,
 826 10.1007/s00382-012-1613-8, 2013.
 827 Maggioni, V., Sapiano, M. R. P., Adler, R. F., Tian, Y. D., and Huffman, G. J.: An Error Model for Uncertainty
 828 Quantification in High-Time-Resolution Precipitation Products, *J Hydrometeorol*, 15, 1274-1292, 10.1175/Jhm-D-
 829 13-0112.1, 2014.
 830 Mahfouf, J. F., Brasnett, B., and Gagnon, S.: A Canadian precipitation analysis (CaPA) project: Description and
 831 preliminary results, *Atmos Ocean*, 45, 1-17, 2007.
 832 Marshall, I., Schut, P., and Ballard, M.: A national ecological framework for Canada: Attribute data. Ottawa, Ontario:
 833 Environmental Quality Branch, Ecosystems Science Directorate, Environment Canada and Research Branch,
 834 Agriculture and Agri-Food Canada, 1999.
 835 Martynov, A., Laprise, R., Sushama, L., Winger, K., Separovic, L., and Dugas, B.: Reanalysis-driven climate
 836 simulation over CORDEX North America domain using the Canadian Regional Climate Model, version 5: model
 837 performance evaluation, *Clim Dynam*, 41, 2973-3005, 10.1007/s00382-013-1778-9, 2013.

838 Maurer, E. P., and Hidalgo, H. G.: Utility of daily vs. monthly large-scale climate data: an intercomparison of two
839 statistical downscaling methods, *Hydrol Earth Syst Sc*, 12, 551-563, 2008.

840 Maurer, E. P., Hidalgo, H. G., Das, T., Dettinger, M. D., and Cayan, D. R.: The utility of daily large-scale climate data
841 in the assessment of climate change impacts on daily streamflow in California, *Hydrol Earth Syst Sc*, 14, 1125-1138,
842 10.5194/hess-14-1125-2010, 2010.

843 Mearns, L., Arritt, R., Biner, S., Bukovsky, M. S., McGinnis, S., Sain, S., Caya, D., Correia, J., Flory, D., Gutowski,
844 W., Takle, E. S., Jones, R., Leung, R., Moufouma-Okia, W., McDaniel, L., Nunes, A. M. B., Qian, Y., Roads, J.,
845 Sloan, L., and Snyder, M.: Overview of the North American regional climate change assessment program, NOAA
846 RISA-NCAR Meeting, 2006,

847 Mearns, L. O., Arritt, R., Biner, S., Bukovsky, M. S., McGinnis, S., Sain, S., Caya, D., Correia, J., Flory, D., Gutowski,
848 W., Takle, E. S., Jones, R., Leung, R., Moufouma-Okia, W., McDaniel, L., Nunes, A. M. B., Qian, Y., Roads, J.,
849 Sloan, L., and Snyder, M.: THE NORTH AMERICAN REGIONAL CLIMATE CHANGE ASSESSMENT
850 PROGRAM Overview of Phase I Results, *B Am Meteorol Soc*, 93, 1337-1362, 2012.

851 Meinshausen, M., Smith, S. J., Calvin, K., Daniel, J. S., Kainuma, M. L. T., Lamarque, J. F., Matsumoto, K., Montzka,
852 S. A., Raper, S. C. B., Riahi, K., Thomson, A., Velders, G. J. M., and van Vuuren, D. P. P.: The RCP greenhouse gas
853 concentrations and their extensions from 1765 to 2300, *Climatic Change*, 109, 213-241, 10.1007/s10584-011-0156-z,
854 2011.

855 Mekis, E., and Hogg, W. D.: Rehabilitation and analysis of Canadian daily precipitation time series, *Atmos Ocean*,
856 37, 53-85, 1999.

857 Mekis, E., and Vincent, L. A.: An Overview of the Second Generation Adjusted Daily Precipitation Dataset for Trend
858 Analysis in Canada, *Atmos Ocean*, 49, 163-177, Pii 938569134 10.1080/07055900.2011.583910, 2011.

859 Mesinger, F., Janjic, Z. I., Nickovic, S., Gavrillov, D., and Deaven, D. G.: The Step-Mountain Coordinate - Model
860 Description and Performance for Cases of Alpine Lee Cyclogenesis and for a Case of an Appalachian Redevelopment,
861 *Monthly Weather Review*, 116, 1493-1518, Doi 10.1175/1520-0493(1988)116<1493:Tsmcmd>2.0.Co;2, 1988.

862 Mesinger, F., DiMego, G., Kalnay, E., Mitchell, K., Shafran, P. C., Ebisuzaki, W., Jovic, D., Woollen, J., Rogers, E.,
863 Berbery, E. H., Ek, M. B., Fan, Y., Grumbine, R., Higgins, W., Li, H., Lin, Y., Manikin, G., Parrish, D., and Shi, W.:
864 North American regional reanalysis, *B Am Meteorol Soc*, 87, 343-360, 10.1175/Bams-87-3-343, 2006.

865 Metcalfe, J. R., Routledge, B., and Devine, K.: Rainfall measurement in Canada: Changing observational methods
866 and archive adjustment procedures, *J Climate*, 10, 92-101, Doi 10.1175/1520-0442(1997)010<0092:Rmicco>2.0.Co;2,
867 1997.

868 Middelkoop, H., Daamen, K., Gellens, D., Grabs, W., Kwadijk, J. C. J., Lang, H., Parmet, B. W. A. H., Schadler, B.,
869 Schulla, J., and Wilke, K.: Impact of climate change on hydrological regimes and water resources management in the
870 rhine basin, *Climatic Change*, 49, 105-128, Doi 10.1023/A:1010784727448, 2001.

871 Mitchell, K. E., Lohmann, D., Houser, P. R., Wood, E. F., Schaake, J. C., Robock, A., Cosgrove, B. A., Sheffield, J.,
872 Duan, Q. Y., Luo, L. F., Higgins, R. W., Pinker, R. T., Tarpley, J. D., Lettenmaier, D. P., Marshall, C. H., Entin, J. K.,
873 Pan, M., Shi, W., Koren, V., Meng, J., Ramsay, B. H., and Bailey, A. A.: The multi-institution North American Land

874 Data Assimilation System (NLDAS): Utilizing multiple GCIP products and partners in a continental distributed
875 hydrological modeling system, *J Geophys Res-Atmos*, 109, Artn D07s90 10.1029/2003jd003823, 2004.

876 Nalley, D., Adamowski, J., and Khalil, B.: Using discrete wavelet transforms to analyze trends in streamflow and
877 precipitation in Quebec and Ontario (1954-2008), *J Hydrol*, 475, 204-228, 10.1016/j.jhydrol.2012.09.049, 2012.

878 Nešpor, V., Krajewski, W. F., and Kruger, A.: Wind-induced error of raindrop size distribution measurement using a
879 two-dimensional video disdrometer, *J Atmos Ocean Tech*, 17, 1483-1492, 2000.

880 New, M., Hulme, M., and Jones, P.: Representing twentieth-century space-time climate variability. Part I:
881 Development of a 1961-90 mean monthly terrestrial climatology, *J Climate*, 12, 829-856, Doi 10.1175/1520-
882 0442(1999)012<0829:Rtcstc>2.0.Co;2, 1999.

883 New, M., Hulme, M., and Jones, P.: Representing twentieth-century space-time climate variability. Part II:
884 Development of 1901-96 monthly grids of terrestrial surface climate, *J Climate*, 13, 2217-2238, Doi 10.1175/1520-
885 0442(2000)013<2217:Rtcstc>2.0.Co;2, 2000.

886 Nijssen, B., and Lettenmaier, D. P.: Effect of precipitation sampling error on simulated hydrological fluxes and states:
887 Anticipating the Global Precipitation Measurement satellites, *J Geophys Res-Atmos*, 109, Artn D02103
888 10.1029/2003jd003497, 2004.

889 Pacific Climate Impacts Consortium; University of Victoria: Statistically Downscaled Climate Scenarios, in, 20th
890 April 2016 ed., Downloaded from <https://www.pacificclimate.org/data/statistically-downscaled-climate-scenarios> on
891 20th April 2016, Jan 2014.

892 Park, H., Fedorov, A. N., Zheleznyak, M. N., Konstantinov, P. Y., and Walsh, J. E.: Effect of snow cover on pan-
893 Arctic permafrost thermal regimes, *Clim Dynam*, 44, 2873-2895, 10.1007/s00382-014-2356-5, 2015.

894 Park, H., Yoshikawa, Y., Oshima, K., Kim, Y., Thanh, N. D., Kimball, J. S., and Yang, D. Q.: Quantification of
895 Warming Climate-Induced Changes in Terrestrial Arctic River Ice Thickness and Phenology, *J Climate*, 29, 1733-
896 1754, 10.1175/Jcli-D-15-0569.1, 2016.

897 Pearse, P. H., Bertrand, F., and MacLaren, J. W.: Currents of change; Final Report: inquiry on Federal water policy,
898 Inquiry on Federal Water Policy, 1985.

899 Pietroniro, A., Fortin, V., Kouwen, N., Neal, C., Turcotte, R., Davison, B., Versegny, D., Soulis, E. D., Caldwell, R.,
900 Evora, N., and Pellerin, P.: Development of the MESH modelling system for hydrological ensemble forecasting of the
901 Laurentian Great Lakes at the regional scale, *Hydrol Earth Syst Sc*, 11, 1279-1294, 2007.

902 Rapaic, M., Brown, R., Markovic, M., and Chaumont, D.: An Evaluation of Temperature and Precipitation Surface-
903 Based and Reanalysis Datasets for the Canadian Arctic, 1950-2010, *Atmos Ocean*, 53, 283-303,
904 10.1080/07055900.2015.1045825, 2015.

905 Rasmussen, R., Baker, B., Kochendorfer, J., Meyers, T., Landolt, S., Fischer, A. P., Black, J., Theriault, J. M., Kucera,
906 P., Gochis, D., Smith, C., Nitu, R., Hall, M., Ikeda, K., and Gutmann, E.: HOW WELL ARE WE MEASURING
907 SNOW? The NOAA/FAA/NCAR Winter Precipitation Test Bed, *B Am Meteorol Soc*, 93, 811-829, 10.1175/Bams-
908 D-11-00052.1, 2012.

909 Rauscher, S. A., Coppola, E., Piani, C., and Giorgi, F.: Resolution effects on regional climate model simulations of
910 seasonal precipitation over Europe, *Clim Dynam*, 35, 685-711, 10.1007/s00382-009-0607-7, 2010.

911 Rudolf, B., and Schneider, U.: Calculation of gridded precipitation data for the global land-surface using in-situ gauge
 912 observations, Proc. Second Workshop of the Int. Precipitation Working Group, 2005, 231-247,
 913 Sapiano, M. R. P., and Arkin, P. A.: An Intercomparison and Validation of High-Resolution Satellite Precipitation
 914 Estimates with 3-Hourly Gauge Data, J Hydrometeorol, 10, 149-166, 10.1175/2008jhm1052.1, 2009.
 915 Schneider, U., Fuchs, T., Meyer-Christoffer, A., and Rudolf, B.: Global precipitation analysis products of the GPCC,
 916 Global Precipitation Climatology Centre (GPCC), DWD, Internet Publikation, 112, 2008.
 917 Schnorbus, M., Werner, A., and Bennett, K.: Impacts of climate change in three hydrologic regimes in British
 918 Columbia, Canada, Hydrol Process, 28, 1170-1189, 10.1002/hyp.9661, 2014.
 919 Separovic, L., Alexandru, A., Laprise, R., Martynov, A., Sushama, L., Winger, K., Tete, K., and Valin, M.: Present
 920 climate and climate change over North America as simulated by the fifth-generation Canadian regional climate model,
 921 Clim Dynam, 41, 3167-3201, 10.1007/s00382-013-1737-5, 2013.
 922 Sheffield, J., Goteti, G., and Wood, E. F.: Development of a 50-year high-resolution global dataset of meteorological
 923 forcings for land surface modeling, J Climate, 19, 3088-3111, Doi 10.1175/Jcli3790.1, 2006.
 924 Shen, S. S. P., Dzikowski, P., Li, G. L., and Griffith, D.: Interpolation of 1961-97 daily temperature and precipitation
 925 data onto Alberta polygons of ecodistrict and soil landscapes of Canada, J Appl Meteorol, 40, 2162-2177, Doi
 926 10.1175/1520-0450(2001)040<2162:Iodtap>2.0.Co;2, 2001.
 927 Shen, Y., Xiong, A. Y., Wang, Y., and Xie, P. P.: Performance of high-resolution satellite precipitation products over
 928 China, J Geophys Res-Atmos, 115, Artn D02114 10.1029/2009jd012097, 2010.
 929 Shook, K., and Pomeroy, J.: Changes in the hydrological character of rainfall on the Canadian prairies, Hydrol Process,
 930 26, 1752-1766, 10.1002/hyp.9383, 2012.
 931 Shrestha, R., Berland, A., Schnorbus, M., and Werner, A.: Climate change impacts on hydro-climatic regimes in the
 932 Peace and Columbia watersheds, British Columbia, Canada, Pacific climate impacts consortium, University of
 933 Victoria, 37, 2011.
 934 Shrestha, R. R., Dibike, Y. B., and Prowse, T. D.: Modelling of climate-induced hydrologic changes in the Lake
 935 Winnipeg watershed, J Great Lakes Res, 38, 83-94, 10.1016/j.jglr.2011.02.004, 2012a.
 936 Shrestha, R. R., Schnorbus, M. A., Werner, A. T., and Berland, A. J.: Modelling spatial and temporal variability of
 937 hydrologic impacts of climate change in the Fraser River basin, British Columbia, Canada, Hydrol Process, 26, 1841-
 938 1861, 10.1002/hyp.9283, 2012b.
 939 Strangeways, I.: Improving precipitation measurement, Int J Climatol, 24, 1443-1460, 10.1002/joc.1075, 2004.
 940 Tapiador, F. J., Turk, F. J., Petersen, W., Hou, A. Y., Garcia-Ortega, E., Machado, L. A. T., Angelis, C. F., Salio, P.,
 941 Kidd, C., Huffman, G. J., and de Castro, M.: Global precipitation measurement: Methods, datasets and applications,
 942 Atmos Res, 104, 70-97, 10.1016/j.atmosres.2011.10.021, 2012.
 943 Taubenbock, H., Wurm, M., Netzband, M., Zwenzner, H., Roth, A., Rahman, A., and Dech, S.: Flood risks in
 944 urbanized areas - multi-sensoral approaches using remotely sensed data for risk assessment, Nat Hazard Earth Sys,
 945 11, 431-444, 10.5194/nhess-11-431-2011, 2011.
 946 Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An Overview of Cmp5 and the Experiment Design, B Am Meteorol
 947 Soc, 93, 485-498, 10.1175/Bams-D-11-00094.1, 2012.

948 Teutschbein, C., and Seibert, J.: Regional climate models for hydrological impact studies at the catchment scale: a
 949 review of recent modeling strategies, *Geography Compass*, 4, 834-860, 2010.
 950 Tian, Y. D., Peters-Lidard, C. D., Eylander, J. B., Joyce, R. J., Huffman, G. J., Adler, R. F., Hsu, K. L., Turk, F. J.,
 951 Garcia, M., and Zeng, J.: Component analysis of errors in satellite-based precipitation estimates, *J Geophys Res-*
 952 *Atmos*, 114, Artn D24101 10.1029/2009jd011949, 2009.
 953 Tian, Y. D., and Peters-Lidard, C. D.: A global map of uncertainties in satellite-based precipitation measurements,
 954 *Geophys Res Lett*, 37, Artn L24407 10.1029/2010gl046008, 2010.
 955 Turk, F. J., Arkin, P., Ebert, E. E., and Sapiaro, M. R. P.: Evaluating High-Resolution Precipitation Products, *B Am*
 956 *Meteorol Soc*, 89, 1911-1916, 10.1175/2008bams2652.1, 2008.
 957 Uppala, S. M., Kållberg, P., Simmons, A., Andrae, U., Bechtold, V. d., Fiorino, M., Gibson, J., Haseler, J., Hernandez,
 958 A., and Kelly, G.: The ERA-40 re-analysis, *Q J Roy Meteor Soc*, 131, 2961-3012, 2005.
 959 Vila, D. A., de Goncalves, L. G. G., Toll, D. L., and Rozante, J. R.: Statistical Evaluation of Combined Daily Gauge
 960 Observations and Rainfall Satellite Estimates over Continental South America, *J Hydrometeorol*, 10, 533-543,
 961 10.1175/2008jhm1048.1, 2009.
 962 Villarini, G., and Krajewski, W. F.: Review of the Different Sources of Uncertainty in Single Polarization Radar-
 963 Based Estimates of Rainfall, *Surv Geophys*, 31, 107-129, 10.1007/s10712-009-9079-x, 2010.
 964 Wan, H., Zhang, X. B., Zwiers, F. W., and Shiogama, H.: Effect of data coverage on the estimation of mean and
 965 variability of precipitation at global and regional scales, *J Geophys Res-Atmos*, 118, 534-546, 10.1002/jgrd.50118,
 966 2013.
 967 Wang, S., Yang, Y., Luo, Y., and Rivera, A.: Spatial and seasonal variations in evapotranspiration over Canada's
 968 landmass, *Hydrol Earth Syst Sc*, 17, 3561-3575, 10.5194/hess-17-3561-2013, 2013.
 969 Wang, S. S., Huang, J. L., Li, J. H., Rivera, A., McKenney, D. W., and Sheffield, J.: Assessment of water budget for
 970 sixteen large drainage basins in Canada, *J Hydrol*, 512, 1-15, 10.1016/j.jhydrol.2014.02.058, 2014.
 971 Wang, X. L. L., and Lin, A.: An algorithm for integrating satellite precipitation estimates with in situ precipitation
 972 data on a pentad time scale, *J Geophys Res-Atmos*, 120, 3728-3744, 10.1002/2014jd022788, 2015.
 973 Weedon, G. P., Gomes, S., Viterbo, P., Shuttleworth, W. J., Blyth, E., Osterle, H., Adam, J. C., Bellouin, N., Boucher,
 974 O., and Best, M.: Creation of the WATCH Forcing Data and Its Use to Assess Global and Regional Reference Crop
 975 Evaporation over Land during the Twentieth Century, *J Hydrometeorol*, 12, 823-848, 10.1175/2011jhm1369.1, 2011.
 976 Weedon, G. P., Balsamo, G., Bellouin, N., Gomes, S., Best, M. J., and Viterbo, P.: The WFDEI meteorological forcing
 977 data set: WATCH Forcing Data methodology applied to ERA-Interim reanalysis data, *Water Resour Res*, 50, 7505-
 978 7514, 10.1002/2014wr015638, 2014.
 979 Werner, A. T., and Cannon, A. J.: Hydrologic extremes - an intercomparison of multiple gridded statistical
 980 downscaling methods, *Hydrol Earth Syst Sc*, 20, 1483-1508, 10.5194/hess-20-1483-2016, 2016.
 981 West, G. L., Steenburgh, W. J., and Cheng, W. Y. Y.: Spurious grid-scale precipitation in the North American regional
 982 reanalysis, *Monthly Weather Review*, 135, 2168-2184, 10.1175/Mwr3375.1, 2007.
 983 Wetterhall, F., Bardossy, A., Chen, D. L., Halldin, S., and Xu, C. Y.: Daily precipitation-downscaling techniques in
 984 three Chinese regions, *Water Resour Res*, 42, Artn W11423 10.1029/2005wr004573, 2006.

Willmott, C. J., Matsuura, K., and Legates, D.: Terrestrial air temperature and precipitation: monthly and annual time series (1950–1999), Center for climate research version, 1, 2001.

Woo, M. K., and Thorne, R.: Snowmelt contribution to discharge from a large mountainous catchment in subarctic Canada, *Hydrol Process*, 20, 2129-2139, 10.1002/hyp.6205, 2006.

Wood, A. W., Leung, L. R., Sridhar, V., and Lettenmaier, D. P.: Hydrologic implications of dynamical and statistical approaches to downscaling climate model outputs, *Climatic Change*, 62, 189-216, DOI 10.1023/B:CLIM.0000013685.99609.9e, 2004.

Xie, P. P., and Arkin, P. A.: An Intercomparison of Gauge Observations and Satellite Estimates of Monthly Precipitation, *J Appl Meteorol*, 34, 1143-1160, Doi 10.1175/1520-0450(1995)034<1143:Aiogoa>2.0.Co;2, 1995.

Xie, P. P., and Arkin, P. A.: Global monthly precipitation: An intercomparison of several datasets based on gauge observations, satellite estimates and model predictions, *Eighth Conference on Satellite Meteorology and Oceanography*, 225-229, 1996.

Xu, C. Y., Widen, E., and Halldin, S.: Modelling hydrological consequences of climate change - Progress and challenges, *Adv Atmos Sci*, 22, 789-797, Doi 10.1007/Bf02918679, 2005.

Yang, D. Q., Goodison, B. E., Metcalfe, J. R., Golubev, V. S., Bates, R., Pangburn, T., and Hanson, C. L.: Accuracy of NWS 8" standard nonrecording precipitation gauge: Results and application of WMO intercomparison, *J Atmos Ocean Tech*, 15, 54-68, Doi 10.1175/1520-0426(1998)015<0054:Aonsnp>2.0.Co;2, 1998.

Young, C. B., Nelson, B. R., Bradley, A. A., Smith, J. A., Peters-Lidard, C. D., Kruger, A., and Baeck, M. L.: An evaluation of NEXRAD precipitation estimates in complex terrain, *J Geophys Res-Atmos*, 104, 19691-19703, Doi 10.1029/1999jd900123, 1999.

Zhang, Q., Sun, P., Singh, V. P., and Chen, X. H.: Spatial-temporal precipitation changes (1956-2000) and their implications for agriculture in China, *Global Planet Change*, 82-83, 86-95, 10.1016/j.gloplacha.2011.12.001, 2012.

List of Tables

Table 1 Precipitation products used in this study.

Dataset	Full Name	Type	Spatial Resolution	Temporal Resolution	Duration	Coverage	Reference
ANUSPLIN	Australian National University Spline	Station-based Interpolated	300 arc-second (~0.0833°/ ~10 km)	24 hr	1950 – 2013	Canada	Hutchinson et al. (2009)
CaPA	Canadian Precipitation Analysis	Station-based Model-derived	10 km (~0.0833°)	6 hr	2002 – 2014	North America	Mahfouf et al. (2007)
Princeton	Global dataset at the Princeton University	Reanalysis-based multiple source	0.5° (~50 km)	3 hr	1901 – 2012	Global	Sheffield et al. (2006)
WFDEI [CRU]	Water and Global Change Forcing Data methodology applied to ERA-Interim [Climate Research Unit]	Reanalysis-based multiple source	0.5° (~50 km)	3 hr	1979 – 2012	Global	Weedon et al. (2014)
WFDEI [GPCC]	Water and Global Change Forcing Data methodology applied to ERA-Interim [Global Precipitation Climatology Centre]	Reanalysis-based multiple source	0.5° (~50 km)	3 hr	1979 – 2012	Global	Weedon et al. (2014)
NARR	North American Regional Reanalysis	Reanalysis-based multiple source	32 km (0.3°)	3 hr	1979 – 2015	North America	Mesinger et al. (2006)
PCIC	Pacific Climate Impacts Consortium	Station-driven GCM	300 arc-second (~0.0833°/ ~10 km)	24 hr	Historical: 1950 – 2005 Projected: 2006 – 2100	Canada	Pacific Climate Impacts Consortium; University of Victoria (Jan 2014)
NA-CORDEX	North America COordinated Regional climate Downscaling EXperiment	GCM-driven RCM	0.22° (25 km)	3 hr	Historical: 1950 – 2005 Projected: 2006 – 2100	North America	Giorgi et al. (2009)

Table 2 GCMs chosen in the Pacific Climate Impacts Consortium (PCIC) dataset.

PCIC	Full Name	Country	Statistical Downscaling Method
GFDL-ESM2G_BCCAQ	Geophysical Fluid Dynamics	USA	Bias Correction Constructed Analogues with Quantile mapping reordering
GFDL-ESM2G_BCSD	Laboratory Earth System Model 2G		Bias Correction Spatial Disaggregation
HadGEM2-ES_BCCAQ	Hadley Global Environmental Model	UK	Bias Correction Constructed Analogues with Quantile mapping reordering
HadGEM2-ES_BCSD	2 – Earth System		Bias Correction Spatial Disaggregation
CanESM2_BCCAQ	Second generation Canadian Earth	Canada	Bias Correction Constructed Analogues with Quantile mapping reordering
CanESM2_BCSD	System Model		Bias Correction Spatial Disaggregation
MPI-ESM-LR_BCCAQ	Max-Planck-Institute Earth System	Germany	Bias Correction Constructed Analogues with Quantile mapping reordering
MPI-ESM-LR_BCSD	Model running on low resolution		Bias Correction Spatial Disaggregation

Table 3 GCMs-RCMs chosen in the North America COordinated Regional climate Downscaling EXperiment (NA-CORDEX) dataset.

NA-CORDEX	Full Name	
	Global Circulation Model (GCM)	Regional Climate Model (RCM)
CanESM2 – CanRCM4	Second generation Canadian Earth System Model	Fourth generation Canadian Regional Climate Model
CanESM2 – CRCM5_UQAM		Fifth generation Canadian Regional Climate Model
MPI-ESM-LR – CRCM5_UQAM	Max-Planck-Institute Earth System Model running on low resolution	Fourth generation Regional Climate Model
MPI-ESM-LR – RegCM4		

Table 4 Number of precipitation-gauge stations within each Ecozone.

Region (Ecozone)		Number of Precipitation-gauge Stations	
		1979 – 2012	2002 – 2012
1	Arctic Cordillera	0	0
2	Northern Arctic	4	4
3	Southern Arctic	1	1
4	Taiga Plain	2	2
5	Taiga Shield	4	5
6	Boreal Shield	31	29
7	Atlantic Maritime	10	9
8	Mixedwood Plain	18	16
9	Boreal Plain	14	14
10	Prairie	9	7
11	Taiga Cordillera	1	0
12	Boreal Cordillera	6	6
13	Pacific Maritime	15	15
14	Montane Cordillera	28	26
15	Hudson Plain	2	3
Total		145	137

Table 5 Performance measures (accuracy (PBias), magnitude of the errors (RMSE), strength and direction of relationship between gridded products and precipitation-gauge stations (r), and amplitude of the variations (σ_G/σ_R)) of each type of gridded precipitation products when evaluating against the precipitation-gauge station data over Canada in four seasons for the time period of 2002 to 2012.

Performance Measure	Season	Precipitation Product					
		ANUSPLIN	Princeton	WFDEI [CRU]	WFDEI [GPCC]	NARR	CaPA
PBias (%)	Spring	-14.2	-12.9	3.1	1.0	5.7	0.7
	Summer	-9.3	-4.7	2.6	0.8	-1.3	-4.4
	Autumn	-16.1	-16.0	-3.1	-2.7	-9.3	-1.3
	Winter	-19.9	-22.4	-3.3	-1.2	-11.9	-8.6
	Annual	-14.7	-13.6	-1.3	-1.4	-5.7	-4.2
RMSE (mm/day)	Spring	2.39	5.30	3.68	3.64	3.42	2.70
	Summer	3.41	7.18	5.33	5.12	5.17	3.74
	Autumn	3.00	6.76	4.82	4.70	4.46	3.35
	Winter	2.70	5.24	3.95	3.98	3.61	3.05
	Annual	3.00	6.33	4.61	4.51	4.35	3.34
r (--)	Spring	0.78	0.16	0.53	0.55	0.55	0.72
	Summer	0.78	0.13	0.45	0.49	0.46	0.73
	Autumn	0.80	0.18	0.53	0.56	0.55	0.75
	Winter	0.76	0.17	0.51	0.53	0.54	0.70
	Annual	0.79	0.17	0.50	0.54	0.51	0.74
σ_G/σ_R (--)	Spring	0.72	1.04	0.91	0.95	0.75	0.83
	Summer	0.76	0.97	0.80	0.84	0.75	0.82
	Autumn	0.74	1.02	0.91	0.95	0.72	0.85
	Winter	0.64	0.97	0.96	1.06	0.63	0.72
	Annual	0.74	0.99	0.86	0.92	0.72	0.82

List of Figures




Figure 1. 15 terrestrial ecozones of Canada with numerical codes indicating Region from 1 Arctic Cordillera to 15 Hudson Plain. Big (a total of 145) and small (a total of 137) white dots are the extracted precipitation-gauge stations from the Canadian adjusted and homogenized precipitation datasets of Mekis and Vincent (2011) for the period of 1979 to 2012 and 2002 to 2012 respectively. Black dots are major cities in Canada.




Figure 2. The percentage of reliability, calculated by the Eq. (1), of each precipitation dataset in four seasons for the period of 1979 to 2012 (left panel), 2002 to 2012 (middle panel), and 1979 to 2005 (right panel) across Canada. The higher the percentage, the more reliable the precipitation dataset. Different colours represent different precipitation products, with magenta representing the whole PCIC datasets and cyan representing the whole NA-CORDEX datasets. The full names of the precipitation products are provided in Tables 1, 2, and 3.




Figure 3. Distributions of p -value of the K-S test in four seasons for the period of 1979 to 2012 (long-term comparison without CaPA). Note that the numbers of precipitation-gauge stations in each ecozone are different (see Table 4). The p -values of Regions 6 to 9, and 13 to 14 (R6-R9, and R13-R14), which have more than or equal to 10 stations, were only shown for illustration in box-whisker plots with bottom, band (black thick line) and top of the box indicating the 25th, 50th (median), and 75th percentiles, respectively.




Figure 4. Distributions of p -value of the K-S test in four seasons for the period of 2002 to 2012 (short-term comparison with the inclusion of CaPA). Note that the numbers of precipitation-gauge stations in each ecozone are different (see Table 4). The p -values of Regions 6, 8 to 9, and 13 to 14 (R6, R8-R9, and R13-R14), which have more than or equal to 10 stations, were only shown for illustration in box-whisker plots with bottom, band (black thick line) and top of the box indicating the 25th, 50th (median), and 75th percentiles, respectively.




Figure 5. Distributions of p -value of the K-S test in four seasons for the period of 1979 to 2005 (long-term comparison of PCIC and NA-CORDEX). Note that the numbers of precipitation-gauge stations in each ecozone are different (see Table 4). The p -values of Regions 6 to 9, and 13 to 14 (R6-R9, and R13-R14), which have more than or equal to 10 stations, were only shown for illustration in box-whisker plots with bottom, band (black thick line) and top of the box indicating the 25th, 50th (median), and 75th percentiles, respectively.




Figure 6. Portrait diagram showing the accuracy ($PBias$) (left), and amplitude of the variations (σ_G/σ_R) (right) of each type of gridded precipitation products when evaluating against the precipitation-gauge station data in each ecozone (Region 1 to 15) in four seasons for the time period of 1979 to 2012. Each column indicates one gridded precipitation product and each row represents one ecozone with numerical code corresponding to region shown in Fig. 1. White indicates that no data are available due to no precipitation-gauge stations existing in that region.




Figure 7. Portrait diagram showing magnitude of the errors (RMSE) (left), and strength and direction of relationship between gridded products and precipitation-gauge stations (r) (right) of each type of gridded precipitation products when evaluating against the precipitation-gauge station data in each ecozone (Region 1 to 15) in four seasons for the time period of 1979 to 2012. Each column indicates one gridded precipitation product and each row represents one ecozone with numerical code corresponding to region shown in Fig. 1. White indicates that no data are available due to no precipitation-gauge stations existing in that region.




Figure 8. Scatter plots showing absolute PBIAS (x-axis) versus RMSE (y-axis) of each precipitation dataset in four seasons and the entire year for the period of 1979 to 2012 (left panel) and 2002 to 2012 (right panel). Each hollow circle represents one ecozone and the solid stars indicate the overall average across ecozones.




Figure 9. Bar graphs showing the annual accuracy (PBias) (first row) and magnitude of the errors (RMSE) (second row) of each precipitation dataset for the period of 1979 to 2012 (left panel) and 2002 to 2012 (right panel) in different ecozones. The white bar shows the scale of the bars with number beside it indicating the value of the bar.