

Supplement: Supporting Information

Table S1

Sequence of treatments in Watershed 6, a north-facing 8.9 hectare catchment.

| Year | Treatment | References |
|---------------|--|---|
| 1909-1923 | The Coweeta Basin, including WS6, was commercially cut with ca. 30% reduction in basal area. Cutting progressed from low to high elevation; WS6 would have been the first area cut. | Elliott and Swank, 2008 |
| 1934 | A weir was installed at the base of the watershed, and nine vegetation plots were measured across the east-side of the watershed. | Hewlett, 1961; Douglass and Swank, 1972 |
| 1928-1940 | Chestnut blight (<i>Endothia parasitica</i> Murr.) killed the remaining <i>Castanea dentata</i> trees. | |
| 1942 | 12% of the catchment (1.1 ha) along the stream was cut to determine the effect of cutting streamside vegetation on water yield. | Dunford and Fletcher, 1947 |
| 1958 , summer | Vegetation was measured in 37 plots prior to grass conversion. | |
| 1958, fall | All merchantable timber was cut and removed. All remaining vegetation was cut, piled and burned. A seedbed was prepared by burning and soil scarification. | Hibbert ,1969 |
| 1959 | Grass conversion by application of 1100 kg ha ⁻¹ limestone, 366 kg ha ⁻¹ NPK fertilizer, and 3.7 kg ha ⁻¹ fescue (<i>Festuca octiflora</i> Walter) grass seed across the entire watershed. | Hibbert, 1969 |
| 1960-1965 | Woody sprouts were foliar sprayed with 2, 4, 5-T herbicide, annually for five years. | |
| 1960 | Application of 37 kg ha ⁻¹ NH ₄ -NO ₃ . | |
| 1965 | Application of 110 kg ha ⁻¹ NPK fertilizer and 27 kg ha ⁻¹ 60% potash. | |
| 1966- early | Three herbicides were used to eliminate the grass cover. | Hibbert, 1969 |
| 1967 | Atrazine, paraquat, and 2,4-D were applied on the entire watershed except for a 3 m buffer strip on each side of the perennial stream. | |
| 1967 | No further treatments, vegetation was allowed to revert to successional vegetation. For first several growing seasons, the entire area was dominated by a dense cover of herbaceous species. | Burt and Swank, 1992 |
| 1968 | Vegetation was measured in 34 plots, the first growing season following treatment abandonment. | Burt and Swank, 1992 |
| 1982 | 15-year-old forest | Elliott et al., 1998 |
| 1995 | 28-year-old forest | Elliott et al., 1998 |
| 2012 | 45-year-old forest | This study. |

Table S2

Common tree species across the eastern United States classified by hydrotype (mesophytic, semi-mesophytic, xerophytic)^a, xylem anatomy (diffuse-porous, ring-porous, semi-ring-porous, or tracheid)^b, and stomatal control (isohydric or anisohydric)^c.

| Species | Hydrotype ^a | Xylem anatomy ^b | Stomatal control ^c |
|---|------------------------|----------------------------|-------------------------------|
| <i>Acer rubrum</i> | Mesophytic | Diffuse | Isohydric |
| <i>Acer saccharum</i> | Mesophytic | Diffuse | Isohydric |
| <i>Acer pensylvanicum</i> | Mesophytic | Diffuse | Isohydric |
| <i>Betula alleghaniensis</i> | Mesophytic | Diffuse | Isohydric |
| <i>Betula lenta</i> | Mesophytic | Diffuse | Isohydric |
| <i>Betula papyrifera</i> | Mesophytic | Diffuse | Isohydric |
| <i>Carya</i> spp. | Semi-mesophytic | Semi-ring | Intermediate |
| <i>Castanea dentata</i> | Semi-mesophytic | Semi-ring | Anisohydric |
| <i>Fagus grandifolia</i> | Mesophytic | Diffuse | Isohydric |
| <i>Fraxinus</i> spp. (<i>americana</i> , <i>pennsylvanica</i>) | Mesophytic | Ring | Anisohydric |
| <i>Liriodendron tulipifera</i> | Mesophytic | Diffuse | Isohydric |
| <i>Magnolia</i> spp. (<i>acuminata</i> , <i>fraseri</i>) | Mesophytic | Diffuse | Isohydric |
| <i>Nyssa sylvatica</i> | Semi-mesophytic | Diffuse | Intermediate |
| <i>Oxydendron arboreum</i> | Semi-mesophytic | Ring | Intermediate |
| <i>Pinus</i> spp. (<i>rigida</i> , <i>taeda</i> , <i>echinata</i> , <i>pungens</i> , <i>virginiana</i>) | Xerophytic | Tracheid | Isohydric |
| <i>Pinus palustris</i> | Xerophytic | Tracheid | Isohydric |
| <i>Pinus strobus</i> | Semi-mesophytic | Tracheid | Isohydric |
| <i>Prunus serotina</i> | Semi-mesophytic | Ring | Anisohydric |
| <i>Quercus alba</i> | Xerophytic | Ring | Anisohydric |
| <i>Quercus coccinea</i> | Xerophytic | Ring | Anisohydric |
| <i>Quercus montana</i> | Xerophytic | Ring | Anisohydric |
| <i>Quercus rubra</i> | Semi-mesophytic | Ring | Anisohydric |
| <i>Quercus velutina</i> | Semi-mesophytic | Ring | Anisohydric |
| <i>Robinia pseudoacacia</i> | Semi- mesophytic | Ring | Isohydric |
| <i>Tilia americana</i> | Mesophytic | Diffuse | Isohydric |
| <i>Tsuga canadensis</i> | Mesophytic | Tracheid | Anisohydric |

Classifications were extracted from the literature; ^ahydrotype (Flatley et al., 2015), ^bxylem anatomy (Ewers et al., 2007) and ^cstomatal control (Roman et al., 2015); ^cstomatal control of *Castanea dentata* based on *C. sativa* (Gomes-Laranjo et al., 2012).

Table S3

Mean density, basal area, total aboveground biomass, and leaf area index (LAI) for the treated (WS6) and reference (WS14, WS18) watersheds overtime. Standard errors are in parentheses.

| | | WS6 | | | | |
|---|--|---------------------|---------------------|---------------------|----------------------|---------------------|
| | | 1934 N = 5 | 1958 N = 37 | 1982 N = 34 | 1995 N = 34 | 2012 N = 34 |
| Density (stems ha ⁻¹) | | 4302 (213) | 5942 (291) | 1712 (121) | 1338 (142) | 1174 (90) |
| Basal area (m ² ha ⁻¹) | | 28.39 (2.61) | 15.96 (1.38) | 11.21 (0.82) | 21.00 (1.45) | 27.36 (1.81) |
| Biomass (Mg ha ⁻¹) | | 209.855 (20.713) | 99.508 (12.110) | 45.564 (3.760) | 126.550 (9.617) | 201.470 (14.665) |
| LAI (m ² m ⁻²) | | 6.447 (0.603) | 3.273 (0.293) | 1.823 (0.128) | 3.616 (0.241) | 4.848 (0.330) |
| | | WS14 | | | | |
| | | 1934 N = 30 | 1969 N = 30 | 1988-1992 N = 30 | 2009-2010 N = 30 | |
| Density (stems ha ⁻¹) | | 2877 (158) | 6178 (472) | 3872 (576) | 3672 (322) | |
| Basal area (m ² ha ⁻¹) | | 27.21 (1.55) | 26.18 (1.11) | 29.52 (1.525) | 36.93 (1.85) | |
| Biomass (Mg ha ⁻¹) | | 224.260 (20.374) | 178.486 (12.920) | 206.608 (14.840) | 244.290 (17.428) | |
| LAI (m ² m ⁻²) | | 5.808 (0.340) | 5.115 (0.241) | 5.728 (0.298) | 6.384 (0.348) | |
| | | WS18 | | | | |
| | | 1934 N = 8 | 1953 N=8 | 1969 N = 8 | 1992 N = 7 | 2009 N = 8 |
| Density (stems ha ⁻¹) | | 3029 (326) | 8025 (1611) | 7220 (1727) | 5196 (1337) | 4636 (1289) |
| Basal area (m ² ha ⁻¹) | | 26.04 (2.00) | 25.75 (1.10) | 28.13 (0.92) | 35.76 (3.28) | 34.37 (3.06) |
| Biomass (Mg ha ⁻¹) | | 190.844 (28.758) | 148.424 (12.837) | 183.742 (9.641) | 218.356 (29.9885) | 208.079 (13.518) |
| LAI (m ² m ⁻²) | | 5.432 (0.360) | 5.001 (0.208) | 5.376 (0.195) | 6.151 (0.551) | 5.951 (0.394) |

In 1934, only percent cover of evergreen shrubs (*Kalmia latifolia* and *Rhododendron maximum*) was estimated for each plot. For all other years, diameter of evergreen shrubs was measured. The higher density in years 1953 and 1958 versus 1934 is partly due to evergreen shrub diameters not being measured or counted in the permanent plots in 1934.

Table S4

Watershed 14: mean density, basal area, aboveground biomass, and leaf area index (LAI) of species over time [1934, 1969, 1988-1992, and 2009-2010]. Standard errors are in parentheses.

| Species in 1934 | Density | Basal area | Aboveground biomass (kg ha ⁻¹) | | LAI |
|--------------------------------|---------------------------|------------------------------------|--|-----------------|-----------------------------------|
| | (stems ha ⁻¹) | (m ² ha ⁻¹) | Leaf | Total | (m ² m ⁻²) |
| <i>Castanea dentata</i> | 764 (119) | 12.50 (1.44) | 1,509 (175) | 120,549 (17775) | 2.506 (0.291) |
| <i>Quercus montana</i> | 269 (43) | 3.79 (0.79) | 576 (129) | 43,036 (11489) | 0.818 (0.184) |
| <i>Carya spp.</i> | 228 (30) | 1.62 (0.30) | 196 (44) | 11,991 (3710) | 0.369 (0.084) |
| <i>Quercus coccinea</i> | 164 (37) | 1.74 (0.36) | 488 (97) | 11,225 (2596) | 0.605 (0.122) |
| <i>Acer rubrum</i> | 234 (24) | 1.22 (0.17) | 222 (28) | 6,664 (1171) | 0.381 (0.048) |
| <i>Nyssa sylvatica</i> | 108 (14) | 0.96 (0.22) | 68 (11) | 5,791 (1814) | 0.166 (0.028) |
| <i>Quercus velutina</i> | 56 (12) | 0.79 (0.19) | 101 (27) | 5,112 (1501) | 0.126 (0.034) |
| <i>Quercus alba</i> | 29 (15) | 0.45 (0.22) | 59 (25) | 3,347 (2162) | 0.087 (0.037) |
| <i>Robinia pseudoacacia</i> | 71 (9) | 0.58 (0.09) | 68 (11) | 2,888 (604) | 0.090 (0.015) |
| <i>Quercus rubra</i> | 40 (9) | 0.46 (0.13) | 58 (19) | 2,850 (1146) | 0.084 (0.026) |
| <i>Tsuga canadensis</i> | 32 (15) | 0.48 (0.26) | 100 (48) | 2,512 (1384) | 0.077 (0.038) |
| <i>Betula lenta</i> | 128 (56) | 0.41 (0.15) | 21 (9) | 1,865 (820) | 0.055 (0.024) |
| <i>Liriodendron tulipifera</i> | 181 (70) | 0.48 (0.16) | 41 (13) | 1,429 (444) | 0.075 (0.024) |
| <i>Oxydendrum arboreum</i> | 50 (8) | 0.32 (0.07) | 29 (5) | 1,139 (270) | 0.073 (0.014) |
| <i>Cornus florida</i> | 121 (24) | 0.32 (0.07) | 35 (8) | 761 (186) | 0.076 (0.015) |
| <i>Sassafras albidum</i> | 112 (16) | 0.27 (0.04) | 32 (5) | 723 (165) | 0.051 (0.009) |
| <i>Hamamelis virginiana</i> | 113 (35) | 0.23 (0.07) | 27 (8) | 517 (159) | 0.056 (0.017) |
| <i>Castanea pumila</i> | 63 (16) | 0.14 (0.04) | 17 (4) | 353 (101) | 0.027 (0.007) |
| <i>Tilia americana</i> | 2 (1) | 0.04 (0.04) | 5 (4) | 306 (300) | 0.009 (0.008) |
| <i>Pinus rigida</i> | 2 (2) | 0.08 (0.08) | 4 (4) | 300 (300) | 0.009 (0.009) |
| <i>Amelanchier arborea</i> | 31 (10) | 0.08 (0.03) | 9 (3) | 213 (81) | 0.015 (0.005) |
| <i>Acer pensylvanicum</i> | 28 (11) | 0.08 (0.03) | 14 (6) | 204 (88) | 0.029 (0.010) |
| <i>Magnolia acuminata</i> | 2 (1) | 0.03 (0.03) | 4 (3) | 180 (174) | 0.011 (0.010) |
| <i>Magnolia fraseri</i> | 6 (4) | 0.02 (0.02) | 3 (4) | 104 (84) | 0.023 (0.014) |
| <i>Fagus grandifolia</i> | 3 (2) | 0.01 (0.01) | 2 (1) | 62 (56) | 0.003 (0.002) |
| <i>Symplocos tinctoria</i> | 12 (7) | 0.02 (0.01) | 3 (2) | 51 (31) | 0.004 (0.003) |
| <i>Aesculus flava</i> | 9 (5) | 0.02 (0.01) | 2 (1) | 50 (24) | 0.004 (0.002) |
| <i>Fraxinus americana</i> | 5 (3) | 0.01 (0.01) | 1 (0.7) | 24 (12) | 0.005 (0.003) |

| | | | | | |
|-----------------------------|------------|---------------|-------------|-----------------|-----------------|
| <i>Quercus falcata</i> | 0.8 (0.6) | 0.002 (0.001) | 2 (1) | 5 (3) | 0.002 (0.001) |
| <i>Diospyros virginiana</i> | 0.8 (0.8) | 0.002 (0.002) | 0.2 (0.2) | 4 (4) | 0.0003 (0.0003) |
| <i>Salix nigra</i> | 5 (4) | 0.01 (0.01) | 0.03 (0.02) | 0.03 (0.02) | 0.002 (0.001) |
| <i>Linder benzoin</i> | 2 (2) | 0.005 (0.004) | 0.01 (0.01) | 0.01 (0.01) | <0.0001 |
| All <i>Quercus</i> | 560 (65) | 7.24 (0.93) | 1,284 (168) | 65,576 (11941) | 1.736 (0.231) |
| Total | 2877 (158) | 27.21 (1.55) | 3,699 (213) | 224,260 (20374) | 5.808 (0.340) |

Species in 1969

| | | | | | |
|--------------------------------|------------|-------------|-----------|----------------|---------------|
| <i>Quercus montana</i> | 215 (34) | 5.59 (1.12) | 836 (175) | 56,570 (13468) | 1.188 (0.248) |
| <i>Quercus coccinea</i> | 74 (15) | 2.60 (0.57) | 418 (85) | 19,432 (4697) | 0.535 (0.108) |
| <i>Acer rubrum</i> | 216 (22) | 2.33 (0.32) | 388 (49) | 15,618 (2539) | 0.678 (0.086) |
| <i>Carya spp.</i> | 148 (28) | 1.56 (0.37) | 201 (58) | 13,069 (4446) | 0.330 (0.095) |
| <i>Quercus rubra</i> | 49 (10) | 1.66 (0.34) | 230 (51) | 12,420 (3080) | 0.295 (0.066) |
| <i>Liriodendron tulipifera</i> | 98 (26) | 1.39 (0.40) | 116 (34) | 7,270 (2190) | 0.214 (0.062) |
| <i>Betula lenta</i> | 93 (33) | 0.99 (0.34) | 69 (24) | 6,332 (2235) | 0.188 (0.066) |
| <i>Quercus velutina</i> | 14 (3) | 0.76 (0.25) | 112 (41) | 6,284 (2465) | 0.143 (0.052) |
| <i>Tsuga canadensis</i> | 48 (22) | 1.11 (0.48) | 226 (97) | 5,688 (2537) | 0.180 (0.078) |
| <i>Kalmia latifolia</i> | 2361 (508) | 1.40 (0.29) | 239 (51) | 5,268 (1124) | 0.184 (0.038) |
| <i>Quercus alba</i> | 20 (7) | 0.63 (0.28) | 78 (32) | 5,020 (2667) | 0.115 (0.046) |
| <i>Rhododendron maximum</i> | 2092 (411) | 1.51 (0.29) | 220 (43) | 5,539 (1102) | 0.170 (0.033) |
| <i>Robinia pseudoacacia</i> | 34 (6) | 0.78 (0.16) | 93 (20) | 4,640 (1074) | 0.126 (0.027) |
| <i>Nyssa sylvatica</i> | 45 (7) | 0.65 (0.17) | 44 (9) | 3,780 (1353) | 0.111 (0.024) |
| <i>Oxydendrum arboreum</i> | 48 (8) | 0.81 (0.15) | 60 (11) | 3,619 (668) | 0.158 (0.028) |
| <i>Cornus florida</i> | 152 (37) | 0.57 (0.14) | 65 (16) | 1,718 (470) | 0.131 (0.033) |
| <i>Magnolia acuminata</i> | 5 (2) | 0.16 (0.07) | 19 (8) | 1,142 (517) | 0.064 (0.027) |
| <i>Castanea dentata</i> | 104 (24) | 0.29 (0.08) | 33 (9) | 1,069 (552) | 0.055 (0.015) |
| <i>Amelanchier arborea</i> | 66 (17) | 0.23 (0.06) | 26 (7) | 747 (228) | 0.044 (0.012) |
| <i>Fagus grandifolia</i> | 7 (4) | 0.12 (0.09) | 14 (11) | 656 (530) | 0.024 (0.018) |
| <i>Tilia americana</i> | 1 (1) | 0.08 (0.08) | 10 (9) | 619 (606) | 0.016 (0.015) |
| <i>Pinus rigida</i> | 2 (2) | 0.14 (0.14) | 10 (10) | 597 (597) | 0.019 (0.019) |
| <i>Acer pensylvanicum</i> | 27 (9) | 0.11 (0.04) | 22 (8) | 485 (188) | 0.039 (0.013) |
| <i>Sassafras albidum</i> | 23 (4) | 0.13 (0.04) | 15 (4) | 472 (119) | 0.024 (0.007) |
| <i>Hamamelis virginiana</i> | 47 (21) | 0.11 (0.05) | 12 (6) | 256 (119) | 0.025 (0.012) |
| <i>Aesculus flava</i> | 4 (2) | 0.04 (0.02) | 5 (3) | 215 (125) | 0.009 (0.005) |
| <i>Magnolia fraseri</i> | 8 (3) | 0.04 (0.02) | 5 (2) | 155 (64) | 0.017 (0.006) |

| | | | | | |
|------------------------------|------------|--------------|-------------|-----------------|-----------------|
| <i>Castanea pumila</i> | 16 (7) | 0.04 (0.02) | 5 (2) | 112 (60) | 0.008 (0.004) |
| <i>Fraxinus americana</i> | 2 (1) | 0.02 (0.02) | 3 (2) | 102 (73) | 0.005 (0.003) |
| <i>Diospyros virginiana</i> | 8 (7) | 0.02 (0.02) | 3 (3) | 84 (80) | 0.005 (0.004) |
| <i>Symplocos tinctoria</i> | 5 (2) | 0.01 (0.004) | 1 (0.4) | 22 (8) | 0.002 (0.001) |
| <i>R. calendulaceum</i> | 79 (21) | 0.16 (0.04) | 0.4 (0.1) | 0.4 (0.1) | 0.001 (0.0002) |
| <i>Pyrularia pubera</i> | 28 (11) | 0.06 (0.02) | 0.15 (0.06) | 0.15 (0.06) | 0.0002 (0.0001) |
| <i>Calycanthus floridana</i> | 5 (3) | 0.01 (0.01) | 0.02 (0.02) | 0.03 (0.02) | < 0.0001 |
| <i>Ilex opaca</i> | 2 (2) | 0.01 (0.01) | 0.02 (0.02) | 0.02 (0.02) | < 0.0001 |
| All <i>Quercus</i> | 372 (43) | 11.25 (1.26) | 1,674 (197) | 99,728 (13973) | 2.275 (0.273) |
| Total | 6178 (472) | 26.18 (1.11) | 3,587 (168) | 178,486 (12920) | 5.114 (0.241) |

Species in 1988-1992

| | | | | | |
|--------------------------------|------------|-------------|-----------|----------------|---------------|
| <i>Quercus montana</i> | 129 (20) | 5.40 (0.98) | 808 (156) | 53,796 (12688) | 1.147 (0.221) |
| <i>Acer rubrum</i> | 147 (19) | 3.70 (0.54) | 574 (79) | 28,991 (4924) | 1.004 (0.138) |
| <i>Quercus rubra</i> | 25 (5) | 2.14 (0.52) | 321 (83) | 18,536 (5138) | 0.411 (0.106) |
| <i>Quercus coccinea</i> | 46 (11) | 2.39 (0.61) | 342 (80) | 18,224 (4963) | 0.438 (0.103) |
| <i>Tsuga canadensis</i> | 45 (16) | 2.53 (1.10) | 423 (175) | 13,731 (6000) | 0.357 (0.150) |
| <i>Liriodendron tulipifera</i> | 62 (15) | 2.11 (0.56) | 175 (47) | 13,006 (3573) | 0.324 (0.086) |
| <i>Betula lenta</i> | 70 (19) | 1.60 (0.51) | 120 (39) | 11,006 (3595) | 0.325 (0.105) |
| <i>Carya spp.</i> | 62 (16) | 0.89 (0.24) | 115 (34) | 7,273 (2424) | 0.188 (0.055) |
| <i>Quercus velutina</i> | 11 (4) | 0.61 (0.31) | 91 (49) | 5,289 (2920) | 0.117 (0.062) |
| <i>Oxydendrum arboreum</i> | 43 (7) | 1.09 (0.19) | 76 (13) | 5,202 (924) | 0.197 (0.034) |
| <i>Robinia pseudoacacia</i> | 21 (3) | 0.68 (0.16) | 81 (19) | 4,333 (1164) | 0.111 (0.026) |
| <i>Nyssa sylvatica</i> | 44 (8) | 0.78 (0.20) | 54 (12) | 4,127 (1282) | 0.137 (0.029) |
| <i>Rhododendron maximum</i> | 1300 (350) | 1.51 (0.31) | 152 (37) | 3,443 (832) | 0.117 (0.029) |
| <i>Kalmia latifolia</i> | 1476 (434) | 0.89 (0.25) | 150 (43) | 3,313 (965) | 0.116 (0.034) |
| <i>Cornus florida</i> | 113 (34) | 0.76 (0.23) | 88 (27) | 3,300 (1291) | 0.177 (0.055) |
| <i>Magnolia acuminata</i> | 5 (2) | 0.27 (0.16) | 32 (21) | 2,170 (1390) | 0.107 (0.066) |
| <i>Tilia americana</i> | 4 (3) | 0.29 (0.18) | 35 (22) | 2,088 (1298) | 0.058 (0.036) |
| <i>Amelanchier arborea</i> | 32 (8) | 0.29 (0.08) | 35 (10) | 1,402 (472) | 0.058 (0.016) |
| <i>Quercus alba</i> | 9 (5) | 0.19 (0.12) | 26 (16) | 1,133 (760) | 0.039 (0.024) |
| <i>Acer saccharum</i> | 10 (6) | 0.17 (0.10) | 38 (17) | 1,050 (665) | 0.049 (0.030) |
| <i>Sassafras albidum</i> | 15 (4) | 0.20 (0.07) | 23 (9) | 964 (422) | 0.039 (0.015) |

| | | | | | |
|-----------------------------|------------|---------------|-------------|-----------------|-----------------|
| <i>Pinus rigida</i> | 2 (2) | 0.16 (0.15) | 16 (14) | 830 (808) | 0.027 (0.027) |
| <i>Castanea dentata</i> | 37 (12) | 0.15 (0.07) | 17 (9) | 824 (638) | 0.029 (0.014) |
| <i>Hamamelis virginiana</i> | 37 (21) | 0.13 (0.06) | 15 (7) | 404 (184) | 0.030 (0.014) |
| <i>Aesculus flava</i> | 6 (2) | 0.08 (0.03) | 9 (4) | 378 (180) | 0.015 (0.007) |
| <i>Magnolia fraseri</i> | 9 (3) | 0.08 (0.03) | 19 (10) | 350 (136) | 0.032 (0.011) |
| <i>Acer pensylvanicum</i> | 12 (6) | 0.07 (0.03) | 14 (6) | 342 (169) | 0.024 (0.010) |
| <i>Diospyros virginiana</i> | 2 (2) | 0.04 (0.03) | 5 (4) | 295 (267) | 0.009 (0.007) |
| <i>Fraxinus americana</i> | 2 (2) | 0.01 (0.01) | 1 (0.8) | 30 (25) | 0.002 (0.001) |
| <i>Castanea pumila</i> | 4 (2) | 0.01 (0.005) | 0.94 (0.53) | 18 (10) | 0.002 (0.001) |
| <i>Fagus grandifolia</i> | 0.4 (0.4) | 0.001 (0.001) | 0.10 (0.10) | 2 (2) | 0.0002 (0.0002) |
| <i>R. calendulaceum</i> | 22 (14) | 0.05 (0.03) | 0.12 (0.08) | 0.13 (0.08) | 0.0002 (0.0001) |
| <i>Pyrularia pubera</i> | 20 (11) | 0.04 (0.02) | 0.11 (0.06) | 0.12 (0.07) | 0.0002 (0.0001) |
| All <i>Quercus</i> | 220 (28) | 10.72 (1.34) | 1,589 (204) | 96,981 (15036) | 2.153 (0.281) |
| Total | 3872 (576) | 29.52 (1.52) | 3,861 (195) | 206,608 (14840) | 5.728 (0.298) |

Species in 2009-2010

| | | | | | |
|--------------------------------|------------|-------------|------------|----------------|---------------|
| <i>Quercus montana</i> | 111 (19) | 6.96 (1.17) | 1060 (185) | 72,528 (14898) | 1.505 (0.263) |
| <i>Acer rubrum</i> | 144 (18) | 4.61 (0.60) | 709 (89) | 36,058 (5304) | 1.240 (0.156) |
| <i>Quercus coccinea</i> | 42 (8) | 3.12 (0.62) | 387 (78) | 24,268 (5203) | 0.496 (0.100) |
| <i>Quercus rubra</i> | 22 (5) | 2.08 (0.50) | 311 (79) | 17,890 (4931) | 0.399 (0.102) |
| <i>Liriodendron tulipifera</i> | 49 (12) | 2.48 (0.65) | 204 (53) | 17,740 (4697) | 0.379 (0.098) |
| <i>Betula lenta</i> | 43 (11) | 1.71 (0.51) | 155 (47) | 14,730 (4513) | 0.422 (0.128) |
| <i>Carya spp.</i> | 55 (14) | 1.19 (0.33) | 156 (47) | 9,996 (3243) | 0.255 (0.077) |
| <i>Kalmia latifolia</i> | 1728 (342) | 4.35 (0.83) | 319 (62) | 7,915 (1528) | 0.250 (0.048) |
| <i>Oxydendrum arboreum</i> | 38 (6) | 1.42 (0.23) | 93 (15) | 7,422 (1227) | 0.240 (0.038) |
| <i>Rhododendron maximum</i> | 1159 (206) | 4.47 (0.82) | 257 (46) | 6,477 (1165) | 0.201 (0.035) |
| <i>Nyssa sylvatica</i> | 39 (8) | 0.90 (0.21) | 60 (13) | 4,828 (1233) | 0.152 (0.034) |
| <i>Quercus alba</i> | 8 (4) | 0.50 (0.28) | 58 (29) | 4,352 (2819) | 0.086 (0.043) |
| <i>Magnolia acuminata</i> | 4 (2) | 0.36 (0.20) | 42 (24) | 3,185 (1803) | 0.143 (0.081) |
| <i>Amelanchier arborea</i> | 40 (10) | 0.54 (0.14) | 64 (17) | 2,876 (887) | 0.106 (0.029) |
| <i>Quercus velutina</i> | 6 (3) | 0.38 (0.21) | 48 (26) | 2,493 (1407) | 0.064 (0.036) |
| <i>Magnolia fraseri</i> | 19 (4) | 0.35 (0.11) | 41 (13) | 2,344 (905) | 0.139 (0.045) |
| <i>Robinia pseudoacacia</i> | 11 (2) | 0.36 (0.10) | 42 (12) | 2,210 (736) | 0.058 (0.017) |
| <i>Tilia americana</i> | 1 (1) | 0.16 (0.12) | 18 (14) | 1,333 (1079) | 0.031 (0.024) |
| <i>Sassafras albidum</i> | 10 (2) | 0.15 (0.05) | 18 (6) | 748 (280) | 0.029 (0.009) |

| | | | | | |
|-----------------------------|------------|---------------|-------------|-----------------|-----------------|
| <i>Cornus florida</i> | 28 (6) | 0.18 (0.05) | 21 (6) | 637 (198) | 0.042 (0.011) |
| <i>Tsuga canadensis</i> | 1 (1) | 0.10 (0.09) | 17 (16) | 569 (555) | 0.015 (0.014) |
| <i>Fagus grandifolia</i> | 3 (2) | 0.08 (0.06) | 9 (7) | 511 (431) | 0.015 (0.012) |
| <i>Fraxinus americana</i> | 5 (3) | 0.08 (0.05) | 10 (6) | 502 (308) | 0.016 (0.009) |
| <i>Acer pensylvanicum</i> | 31 (20) | 0.11 (0.05) | 22 (10) | 508 (199) | 0.039 (0.018) |
| <i>Aesculus flava</i> | 6 (3) | 0.09 (0.05) | 11 (6) | 457 (238) | 0.018 (0.009) |
| <i>Acer saccharum</i> | 4 (4) | 0.05 (0.05) | 8 (8) | 258 (258) | 0.014 (0.014) |
| <i>Hamamelis virginiana</i> | 25 (12) | 0.08 (0.04) | 10 (4) | 241 (112) | 0.019 (0.009) |
| <i>Diospyros virginiana</i> | 1 (1) | 0.02 (0.02) | 2 (2) | 70 (70) | 0.003 (0.003) |
| <i>Pinus rigida</i> | 1 (1) | 0.02 (0.02) | 1 (1) | 69 (48) | 0.002 (0.001) |
| <i>Castanea dentata</i> | 8 (2) | 0.01 (0.003) | 1 (0.4) | 20 (7) | 0.002 (0.001) |
| <i>Symplocos tinctoria</i> | 3 (2) | 0.01 (0.003) | 1 (0.4) | 16 (10) | 0.001 (0.001) |
| <i>Castanea pumila</i> | 2 (2) | 0.004 (0.004) | 0.5 (0.5) | 11 (11) | 0.001 (0.001) |
| <i>Halesia carolina</i> | 0.4 (0.4) | 0.002 (0.002) | 0.2 (0.2) | 5 (5) | 0.0003 (0.0003) |
| All <i>Quercus</i> | 190 (25) | 13.04 (1.48) | 1,868 (222) | 122,732 (17166) | 2.550 (0.307) |
| Total | 3672 (322) | 36.93 (1.85) | 4,170 (227) | 244,290 (17428) | 6.384 (0.348) |

In 1934, only percent cover of evergreen shrubs (*Kalmia latifolia* and *Rhododendron maximum*) was estimated for each plot. For all other years, diameter of evergreen shrubs was measured.

Table S5

Watershed 18: mean density, basal area, aboveground biomass, and leaf area index (LAI) of species over time [1934, 1953, 1969, 1992, and 2009]. Standard errors are in parentheses.

| Species in 1934 | Density (stems ha ⁻¹) | Basal area (m ² ha ⁻¹) | Aboveground biomass (kg ha ⁻¹) | LAI (m ² m ⁻²) |
|--------------------------------|--------------------------------------|--|--|--|
| | | | Leaf | Total |
| <i>Castanea dentata</i> | 581 (149) | 7.99 (1.51) | 963 (186) | 76,843 (23586) |
| <i>Quercus montana</i> | 273 (48) | 3.54 (1.19) | 525 (193) | 34,656 (15047) |
| <i>Quercus rubra</i> | 45 (17) | 2.45 (1.13) | 359 (169) | 20,131 (9619) |
| <i>Quercus coccinea</i> | 226 (36) | 2.71 (0.70) | 716 (152) | 16,980 (4885) |
| <i>Nyssa sylvatica</i> | 167 (30) | 1.76 (0.76) | 112 (30) | 12,220 (7618) |
| <i>Pinus rigida</i> | 66 (47) | 1.37 (1.02) | 94 (74) | 5,785 (4459) |
| <i>Carya</i> spp. | 344 (119) | 1.27 (0.39) | 120 (40) | 5,551 (2191) |
| <i>Quercus velutina</i> | 197 (742) | 1.30 (0.53) | 132 (55) | 5,275 (2211) |
| <i>Robinia pseudoacacia</i> | 95 (16) | 0.76 (0.27) | 91 (32) | 5,031 (2281) |
| <i>Acer rubrum</i> | 270 (64) | 0.65 (0.16) | 140 (33) | 2,237 (744) |
| <i>Cornus florida</i> | 305 (145) | 0.83 (0.39) | 93 (44) | 2,135 (1001) |
| <i>Oxydendrum arboreum</i> | 109 (24) | 0.57 (0.13) | 56 (12) | 1,789 (450) |
| <i>Castanea pumila</i> | 211 (104) | 0.43 (0.21) | 50 (24) | 928 (465) |
| <i>Quercus alba</i> | 9 (6) | 0.16 (0.15) | 24 (22) | 738 (723) |
| <i>Sassafras albidum</i> | 62 (15) | 0.13 (0.04) | 14 (4) | 275 (67) |
| <i>Hamamelis virginiana</i> | 56 (43) | 0.11 (0.10) | 13 (12) | 248 (232) |
| <i>Liriodendron tulipifera</i> | 28 (24) | 0.06 (0.04) | 5 (4) | 120 (105) |
| <i>Amelanchier arborea</i> | 2 (2) | 0.003 (0.003) | 0.04 (0.04) | 7 (7) |
| <i>Diospyros tinctoria</i> | 2 (2) | 0.003 (0.003) | 0.04 (0.04) | 7 (7) |
| <i>Linder benzoin</i> | 6 (6) | 0.01 (0.01) | 0.03 (0.03) | < 1 |
| All <i>Quercus</i> | 752 (105) | 10.16 (1.22) | 1,757 (199) | 77,782 (15104) |
| Total | 3030 (326) | 26.04 (2.00) | 3,503 (221) | 190,845 (28758) |
| Species in 1953 | | | | |
| <i>Quercus montana</i> | 293 (92) | 3.73 (1.33) | 546 (214) | 35,634 (17030) |
| <i>Quercus rubra</i> | 31 (10) | 2.62 (1.26) | 395 (197) | 22,667 (11671) |
| <i>Quercus coccinea</i> | 153 (36) | 3.34 (0.79) | 664 (136) | 22,541 (6158) |

| | | | | | |
|--------------------------------|-------------|---------------|-------------|-----------------|---------------|
| <i>Kalmia latifolia</i> | 5173 (1833) | 2.87 (0.91) | 522 (180) | 11,464 (3935) | 0.402 (0.139) |
| <i>Carya</i> spp. | 362 (124) | 1.90 (0.65) | 202 (76) | 10,643 (4624) | 0.331 (0.125) |
| <i>Pinus rigida</i> | 66 (42) | 2.16 (1.38) | 161 (108) | 9,690 (6329) | 0.319 (0.210) |
| <i>Quercus velutina</i> | 145 (742) | 1.45 (0.53) | 157 (69) | 6,653 (3079) | 0.202 (0.089) |
| <i>Nyssa sylvatica</i> | 172 (24) | 1.15 (0.30) | 94 (19) | 5,247 (1738) | 0.239 (0.048) |
| <i>Acer rubrum</i> | 301 (71) | 1.05 (0.31) | 208 (54) | 4,482 (1940) | 0.365 (0.096) |
| <i>Robinia pseudoacacia</i> | 45 (6) | 0.70 (0.21) | 84 (25) | 4,425 (1765) | 0.114 (0.034) |
| <i>Cornus florida</i> | 326 (148) | 1.20 (0.56) | 136 (63) | 3,661 (1702) | 0.274 (0.128) |
| <i>Oxydendrum arboreum</i> | 106 (19) | 0.95 (0.16) | 82 (14) | 3,483 (617) | 0.216 (0.038) |
| <i>Liriodendron tulipifera</i> | 81 (60) | 0.56 (0.41) | 47 (35) | 2,240 (1550) | 0.087 (0.064) |
| <i>Quercus marilandica</i> | 12 (12) | 0.31 (0.31) | 37 (37) | 1,695 (1695) | 0.047 (0.047) |
| <i>Quercus alba</i> | 16 (11) | 0.25 (0.24) | 37 (35) | 1,354 (1333) | 0.055 (0.051) |
| <i>Castanea pumila</i> | 136 (52) | 0.28 (0.11) | 33 (13) | 635 (261) | 0.054 (0.022) |
| <i>Symplocos tinctoria</i> | 101 (34) | 0.21 (0.07) | 25 (8) | 483 (165) | 0.041 (0.014) |
| <i>Castanea dentata</i> | 102 (14) | 0.21 (0.03) | 24 (3) | 448 (61) | 0.039 (0.005) |
| <i>Sassafras albidum</i> | 72 (32) | 0.16 (0.06) | 18 (7) | 353 (140) | 0.030 (0.012) |
| <i>Hamamelis virginiana</i> | 73 (72) | 0.15 (0.14) | 17 (16) | 324 (316) | 0.035 (0.033) |
| <i>Rhododendron maximum</i> | 73 (27) | 0.10 (0.06) | 9 (4) | 207 (92) | 0.007 (0.001) |
| <i>Juglans cinerea</i> | 12 (7) | 0.02 (0.01) | 2 (1) | 53 (30) | 0.003 (0.001) |
| <i>Tsuga canadensis</i> | 2 (2) | 0.003 (0.003) | 2 (2) | 12 (12) | 0.001 (0.001) |
| <i>Acer saccharum</i> | 2 (2) | 0.003 (0.003) | 1 (1) | 9 (9) | 0.001 (0.001) |
| <i>Amelanchier arborea</i> | 2 (2) | 0.003 (0.003) | 0.4 (0.4) | 7 (7) | 0.001 (0.001) |
| <i>Diospyros virginiana</i> | 2 (2) | 0.003 (0.003) | 0.4 (0.4) | 7 (7) | 0.001 (0.001) |
| <i>Rhodo. calendulaceum</i> | 162 (105) | 0.33 (0.21) | 0.9 (0.5) | 0.9 (0.6) | 0.001 (0.001) |
| All <i>Quercus</i> | 651 (119) | 11.71 (0.84) | 1,838 (165) | 90,547 (11999) | 2.437 (0.233) |
| Total | 8025 (1611) | 25.75 (1.10) | 3,507 (207) | 148,424 (12837) | 5.001 (0.208) |

Species in 1969

| | | | | | |
|-------------------------|-------------|-------------|-----------|----------------|---------------|
| <i>Quercus montana</i> | 236 (66) | 4.09 (1.36) | 605 (219) | 40,050 (17662) | 0.860 (0.310) |
| <i>Quercus rubra</i> | 38 (9) | 3.71 (1.87) | 584 (307) | 34,908 (19010) | 0.747 (0.393) |
| <i>Quercus coccinea</i> | 100 (23) | 4.48 (1.21) | 664 (161) | 33,432 (10073) | 0.850 (0.206) |
| <i>Kalmia latifolia</i> | 5216 (1829) | 2.73 (0.93) | 518 (180) | 11,352 (3957) | 0.399 (0.139) |
| <i>Pinus rigida</i> | 50 (32) | 2.22 (1.48) | 185 (125) | 10,641 (7143) | 0.356 (0.239) |
| <i>Quercus velutina</i> | 62 (26) | 1.79 (0.94) | 221 (121) | 10,417 (5990) | 0.282 (0.155) |
| <i>Carya</i> spp. | 153 (62) | 1.47 (0.69) | 171 (91) | 9,729 (5870) | 0.281 (0.149) |

| | | | | | |
|--------------------------------|-------------|---------------|-------------|-----------------|-----------------|
| <i>Nyssa sylvatica</i> | 72 (12) | 1.07 (0.26) | 72 (14) | 5839 (1679) | 0.182 (0.036) |
| <i>Acer rubrum</i> | 265 (63) | 1.21 (0.31) | 229 (54) | 5837 (1938) | 0.402 (0.095) |
| <i>Oxydendrum arboreum</i> | 131 (31) | 1.44 (0.27) | 118 (22) | 5645 (1044) | 0.308 (0.058) |
| <i>Liriodendron tulipifera</i> | 52 (36) | 0.99 (0.72) | 83 (60) | 5186 (3520) | 0.153 (0.111) |
| <i>Robinia pseudoacacia</i> | 34 (12) | 0.62 (0.22) | 74 (25) | 3660 (1515) | 0.101 (0.035) |
| <i>Cornus florida</i> | 248 (102) | 1.03 (0.48) | 116 (54) | 3272 (1568) | 0.235 (0.109) |
| <i>Quercus alba</i> | 12 (9) | 0.30 (0.29) | 42 (40) | 1677 (1662) | 0.062 (0.059) |
| <i>Rhododendron maximum</i> | 194 (84) | 0.20 (0.10) | 22 (10) | 499 (224) | 0.017 (0.007) |
| <i>Castanea dentata</i> | 72 (36) | 0.14 (0.07) | 17 (8) | 317 (158) | 0.028 (0.014) |
| <i>Acer pensylvanicum</i> | 14 (12) | 0.07 (0.06) | 14 (13) | 310 (299) | 0.024 (0.022) |
| <i>Sassafras albidum</i> | 52 (23) | 0.12 (0.05) | 14 (6) | 299 (140) | 0.023 (0.010) |
| <i>Symplocos tinctoria</i> | 50 (23) | 0.11 (0.05) | 12 (6) | 257 (126) | 0.021 (0.010) |
| <i>Hamamelis virginiana</i> | 56 (51) | 0.11 (0.10) | 13 (12) | 248 (225) | 0.026 (0.024) |
| <i>Castanea pumila</i> | 30 (15) | 0.06 (0.03) | 7 (3) | 131 (66) | 0.012 (0.006) |
| <i>Tsuga canadensis</i> | 2 (2) | 0.003 (0.003) | 2 (2) | 12 (12) | 0.001 (0.001) |
| <i>Amelanchier arborea</i> | 2 (2) | 0.003 (0.003) | 0.4 (0.4) | 7 (7) | 0.001 (0.001) |
| <i>Magnolia acuminata</i> | 2 (2) | 0.003 (0.003) | 0.4 (0.4) | 7 (7) | 0.001 (0.001) |
| <i>Magnolia fraseri</i> | 2 (2) | 0.003 (0.003) | 0.4 (0.4) | 7 (7) | 0.001 (0.001) |
| <i>R. calendulaceum</i> | 71 (55) | 0.16 (0.04) | 0.4 (0.3) | 0.4 (0.3) | 0.001 (0.001) |
| <i>Pyrularia pubera</i> | 3 (3) | 0.05 (0.02) | 0.14 (0.06) | 0.15 (0.06) | 0.0002 (0.0001) |
| All <i>Quercus</i> | 448 (83) | 14.37 (1.10) | 2,116 (155) | 120,485 (11264) | 2.802 (0.208) |
| Total | 7220 (1727) | 28.13 (0.92) | 3,786 (169) | 183,742 (9641) | 5.376 (0.195) |

Species in 1992

| | | | | | |
|--------------------------------|-------------|-------------|-----------|----------------|---------------|
| <i>Quercus montana</i> | 96 (33) | 4.54 (1.50) | 703 (245) | 51,478 (21338) | 0.998 (0.349) |
| <i>Quercus coccinea</i> | 52 (13) | 3.73 (1.33) | 460 (124) | 30,313 (12505) | 0.589 (0.159) |
| <i>Quercus rubra</i> | 16 (6) | 2.12 (1.17) | 323 (185) | 18,582 (10984) | 0.413 (0.237) |
| <i>Acer rubrum</i> | 304 (52) | 2.98 (0.64) | 516 (102) | 17,895 (4906) | 0.902 (0.178) |
| <i>Kalmia latifolia</i> | 3859 (1309) | 6.72 (2.06) | 620 (202) | 14,823 (4798) | 0.477 (0.155) |
| <i>Carya</i> spp. | 71 (26) | 1.75 (0.91) | 228 (127) | 14,277 (8671) | 0.374 (0.208) |
| <i>Pinus rigida</i> | 48 (30) | 2.94 (1.73) | 240 (140) | 14,025 (8224) | 0.468 (0.274) |
| <i>Liriodendron tulipifera</i> | 36 (28) | 2.04 (1.50) | 169 (124) | 13,383 (9420) | 0.312 (0.231) |
| <i>Quercus velutina</i> | 30 (14) | 1.79 (1.19) | 238 (166) | 12,074 (8800) | 0.305 (0.213) |
| <i>Oxydendrum arboreum</i> | 93 (24) | 2.08 (0.45) | 153 (34) | 9,124 (1911) | 0.400 (0.088) |
| <i>Nyssa sylvatica</i> | 64 (17) | 1.14 (0.26) | 78 (16) | 5,962 (1634) | 0.199 (0.040) |

| | | | | | |
|-----------------------------|-------------|---------------|-------------|-----------------|---------------|
| <i>Robinia pseudoacacia</i> | 30 (10) | 0.74 (0.32) | 88 (38) | 4,602 (2271) | 0.120 (0.052) |
| <i>Pinus strobus</i> | 4 (4) | 0.62 (0.62) | 65 (65) | 3,474 (3474) | 0.119 (0.119) |
| <i>Quercus alba</i> | 9 (22) | 0.40 (0.39) | 54 (71) | 2,392 (2383) | 0.079 (0.077) |
| <i>Cornus florida</i> | 96 (33) | 0.65 (0.32) | 74 (26) | 2,213 (1156) | 0.149 (0.074) |
| <i>Castanea dentata</i> | 89 (12) | 0.34 (0.14) | 39 (9) | 1,221 (732) | 0.066 (0.029) |
| <i>Sassafras albidum</i> | 43 (4) | 0.15 (0.09) | 17 (9) | 451 (258) | 0.029 (0.018) |
| <i>Hamamelis virginiana</i> | 91 (20) | 0.20 (0.16) | 23 (7) | 447 (358) | 0.046 (0.037) |
| <i>Rhododendron maximum</i> | 38 (24) | 0.38 (0.26) | 14 (9) | 359 (241) | 0.011 (0.007) |
| <i>Amelanchier arborea</i> | 14 (8) | 0.08 (0.05) | 9 (10) | 233 (164) | 0.015 (0.010) |
| <i>Acer pensylvanicum</i> | 5 (4) | 0.03 (0.02) | 6 (4) | 133 (86) | 0.011 (0.007) |
| <i>Tsuga canadensis</i> | 5 (4) | 0.03 (0.02) | 11 (9) | 123 (108) | 0.008 (0.006) |
| <i>Castanea pumila</i> | 20 (20) | 0.04 (0.04) | 5 (5) | 86 (86) | 0.008 (0.008) |
| <i>Magnolia fraseri</i> | 9 (3) | 0.02 (0.01) | 3 (1) | 54 (29) | 0.009 (0.004) |
| <i>Fagus grandifolia</i> | 4 (2) | 0.01 (0.01) | 16 (10) | 16 (10) | 0.001 (0.001) |
| <i>Magnolia acuminata</i> | 5 (2) | 0.004 (0.004) | 0.4 (0.4) | 8 (8) | 0.001 (0.001) |
| <i>Pyrularia pubera</i> | 4 (4) | 0.01 (0.01) | 0.1 (0.06) | 0.02 (0.02) | <0.001 |
| <i>R. calendulaceum</i> | 4 (4) | 0.01 (0.01) | 0.02 (0.02) | 0.02 (0.02) | <0.001 |
| All <i>Quercus</i> | 204 (42) | 12.59 (3.14) | 1,778 (394) | 114,840 (29105) | 2.385 (0.521) |
| Total | 5196 (1337) | 35.76 (3.28) | 4,163 (422) | 218,356 (29985) | 6.151 (0.551) |

Species in 2009

| | | | | | |
|--------------------------------|-------------|-------------|-----------|----------------|---------------|
| <i>Quercus montana</i> | 86 (19) | 3.21 (0.84) | 475 (130) | 30,050 (10165) | 0.676 (0.185) |
| <i>Quercus rubra</i> | 22 (10) | 3.02 (1.61) | 468 (248) | 27,514 (14470) | 0.599 (0.317) |
| <i>Acer rubrum</i> | 257 (39) | 3.84 (0.71) | 632 (107) | 25,849 (6205) | 1.106 (0.187) |
| <i>Quercus coccinea</i> | 45 (15) | 3.17 (1.10) | 405 (133) | 25,051 (9099) | 0.518 (0.171) |
| <i>Pinus rigida</i> | 41 (25) | 3.69 (2.27) | 337 (210) | 18,836 (11662) | 0.636 (0.395) |
| <i>Liriodendron tulipifera</i> | 39 (27) | 2.53 (1.92) | 209 (158) | 18,812 (13746) | 0.387 (0.293) |
| <i>Carya</i> spp. | 69 (21) | 2.08 (0.83) | 276 (120) | 17,712 (8396) | 0.454 (0.196) |
| <i>Kalmia latifolia</i> | 3378 (1231) | 5.88 (1.97) | 542 (191) | 12,977 (4546) | 0.418 (0.147) |
| <i>Oxydendrum arboreum</i> | 94 (20) | 2.09 (0.46) | 152 (32) | 9,444 (2092) | 0.396 (0.085) |
| <i>Nyssa sylvatica</i> | 69 (20) | 1.31 (0.24) | 86 (13) | 7,210 (1698) | 0.220 (0.033) |
| <i>Robinia pseudoacacia</i> | 20 (5) | 0.56 (0.27) | 67 (33) | 3,691 (1980) | 0.091 (0.045) |
| <i>Quercus alba</i> | 6 (5) | 0.53 (0.53) | 67 (66) | 3,689 (3678) | 0.099 (0.097) |

| | | | | | |
|-----------------------------|-------------|---------------|-------------|-----------------|---------------|
| <i>Quercus velutina</i> | 6 (3) | 0.31 (0.16) | 38 (21) | 1,841 (1059) | 0.049 (0.027) |
| <i>Cornus florida</i> | 28 (6) | 0.42 (0.18) | 48 (21) | 1,457 (688) | 0.097 (0.042) |
| <i>Rhododendron maximum</i> | 233 (131) | 0.96 (0.59) | 52 (31) | 1,303 (767) | 0.040 (0.024) |
| <i>Amelanchier arborea</i> | 19 (7) | 0.54 (0.11) | 31 (13) | 1,264 (586) | 0.051 (0.022) |
| <i>Sassafras albidum</i> | 45 (41) | 0.15 (0.14) | 18 (18) | 455 (450) | 0.030 (0.029) |
| <i>Acer pensylvanicum</i> | 12 (5) | 0.06 (0.05) | 11 (7) | 270 (171) | 0.020 (0.012) |
| <i>Magnolia fraseri</i> | 19 (6) | 0.06 (0.11) | 7 (3) | 177 (84) | 0.024 (0.011) |
| <i>Symplocos tinctoria</i> | 14 (6) | 0.05 (0.03) | 5 (3) | 142 (99) | 0.009 (0.005) |
| <i>Hamamelis virginiana</i> | 40 (34) | 0.05 (0.04) | 6 (5) | 110 (93) | 0.013 (0.011) |
| <i>Castanea dentata</i> | 20 (11) | 0.03 (0.02) | 4 (2) | 70 (40) | 0.006 (0.003) |
| <i>Betula lenta</i> | 3 (2) | 0.02 (0.01) | 1 (1) | 60 (53) | 0.002 (0.002) |
| <i>Magnolia acuminata</i> | 5 (3) | 0.02 (0.01) | 2 (2) | 54 (45) | 0.007 (0.005) |
| <i>Fagus grandifolia</i> | 5 (2) | 0.01 (0.005) | 1 (1) | 21 (14) | 0.002 (0.001) |
| <i>Fraxinus americana</i> | 2 (2) | 0.003 (0.003) | 0.4 (0.4) | 8 (8) | 0.001 (0.001) |
| <i>Pinus strobus</i> | 3 (2) | 0.003 (0.002) | 0.02 (0.01) | 4 (3) | <0.001 |
| <i>Castanea pumila</i> | 2 (2) | 0.002 (0.002) | 0.2 (0.2) | 3 (3) | <0.001 |
| <i>R. calendulaceum</i> | 17 (8) | 0.02 (0.01) | 0.06 (0.03) | 1 (0.04) | <0.001 |
| All <i>Quercus</i> | 167 (29) | 10.25 (1.74) | 1,455 (235) | 88,145 (15363) | 1.941 (0.304) |
| Total | 4635 (1289) | 34.37 (3.06) | 3,944 (312) | 208,079 (13518) | 5.951 (0.394) |

In 1934, only percent cover of evergreen shrubs (*Kalmia latifolia* and *Rhododendron maximum*) was estimated for each plot. For all other years, diameter of evergreen shrubs was measured.

Table S6

Watershed 6: mean density, basal area, aboveground biomass, and leaf area index (LAI) of species over time: before (1934, 1958), during (1961-1965), and after grass-to-forest-conversion (1968, 1982, 1995, 2012). Standard errors are in parentheses. Proportional contributions of dominant species to the watershed totals are in brackets.

| Species in 1934 | Density (stems ha ⁻¹) | Basal area (m ² ha ⁻¹) | Aboveground biomass (kg ha ⁻¹) | | LAI (m ² m ⁻²) |
|-----------------------------|--------------------------------------|--|--|-----------------|--|
| | | | Leaf | Total | |
| <i>Castanea dentata</i> | 600 (162) | 11.99 (3.57) | 1,443 (430) | 102,386 (31647) | 2.395 (0.714) |
| <i>Quercus montana</i> | 320 (118) | 3.77 (1.20) | 568 (209) | 40,294 (18605) | 0.806 (0.296) |
| <i>Quercus coccinea</i> | 322 (76) | 3.35 (1.12) | 908 (214) | 23,031 (8900) | 1.162 (0.274) |
| <i>Carya</i> spp. | 268 (53) | 1.95 (0.61) | 238 (85) | 14,615 (5688) | 0.391 (0.140) |
| <i>Pinus rigida</i> | 25 (17) | 1.98 (1.23) | 179 (112) | 10,040 (6241) | 0.339 (0.211) |
| <i>Acer rubrum</i> | 498 (184) | 1.33 (0.36) | 279 (79) | 4,914 (1378) | 0.488 (0.138) |
| <i>Nyssa sylvatica</i> | 230 (80) | 1.04 (0.48) | 92 (27) | 4,659 (3411) | 0.234 (0.070) |
| <i>Oxydendrum arboreum</i> | 180 (28) | 0.77 (0.11) | 81 (12) | 2,167 (296) | 0.212 (0.031) |
| <i>Quercus velutina</i> | 50 (9) | 0.35 (0.11) | 36 (14) | 1,466 (633) | 0.047 (0.018) |
| <i>Robinia pseudoacacia</i> | 40 (20) | 0.28 (0.14) | 32 (17) | 1,047 (593) | 0.044 (0.023) |
| <i>Cornus florida</i> | 135 (56) | 0.35 (0.18) | 39 (20) | 804 (470) | 0.078 (0.041) |
| <i>Quercus alba</i> | 75 (25) | 0.25 (0.12) | 54 (24) | 698 (420) | 0.079 (0.035) |
| <i>Amelanchier arborea</i> | 52 (14) | 0.11 (0.03) | 12 (3) | 231 (61) | 0.020 (0.005) |
| <i>Quercus rubra</i> | 20 (9) | 0.06 (0.03) | 4 (2) | 136 (70) | 0.006 (0.003) |
| <i>Quercus falcata</i> | 8 (5) | 0.03 (0.02) | 19 (11) | 97 (68) | 0.024 (0.014) |
| <i>Castanea pumila</i> | 15 (9) | 0.03 (0.02) | 4 (2) | 66 (40) | 0.006 (0.004) |
| <i>Sassafras albidum</i> | 10 (5) | 0.02 (0.01) | 2 (1) | 44 (21) | 0.004 (0.002) |
| <i>Diospyros virginiana</i> | 5 (5) | 0.03 (0.01) | 1 (1) | 22 (22) | 0.002 (0.002) |
| All <i>Quercus</i> | 787 (104) | 7.78 (2.12) | 1,569 (404) | 65,626 (24887) | 2.122 (0.559) |
| Total | 4302 (213) | 28.39 (2.61) | 4,134 (377) | 209,855 (20713) | 6.447 (0.603) |
| Species in 1958 | | | | | |
| <i>Quercus montana</i> | 165 (26) | 3.14 (0.65) | 470 (103) | 32,321 (8492) | 0.667 (0.146) |
| <i>Quercus coccinea</i> | 88 (16) | 2.21 (0.47) | 418 (71) | 15,557 (1079) | 0.536 (0.091) |
| <i>Acer rubrum</i> | 284 (42) | 1.38 (0.27) | 244 (41) | 8,593 (2526) | 0.428 (0.072) |

| | | | | | |
|--------------------------------|------------|---------------|-------------|----------------|-----------------|
| <i>Carya</i> spp. | 186 (26) | 1.27 (0.26) | 145 (34) | 8,182 (2194) | 0.238 (0.056) |
| <i>Pinus rigida</i> | 14 (6) | 1.13 (0.52) | 98 (46) | 5,621 (2617) | 0.189 (0.088) |
| <i>Quercus velutina</i> | 41 (9) | 0.89 (0.27) | 112 (37) | 5,394 (1856) | 0.143 (0.047) |
| <i>Rhododendron maximum</i> | 1893 (270) | 1.00 (0.15) | 188 (27) | 4,128 (592) | 0.145 (0.021) |
| <i>Quercus alba</i> | 80 (24) | 0.80 (0.28) | 126 (39) | 4,112 (1757) | 0.185 (0.057) |
| <i>Kalmia latifolia</i> | 1827 (338) | 0.94 (0.17) | 181 (33) | 3,692 (732) | 0.139 (0.026) |
| <i>Nyssa sylvatica</i> | 98 (18) | 0.68 (0.17) | 51 (10) | 3,592 (1292) | 0.131 (0.025) |
| <i>Oxydendrum arboreum</i> | 105 (14) | 0.63 (0.11) | 57 (9) | 2,203 (440) | 0.148 (0.024) |
| <i>Quercus rubra</i> | 20 (7) | 0.28 (0.14) | 36 (18) | 1,802 (912) | 0.046 (0.023) |
| <i>Cornus florida</i> | 240 (40) | 0.50 (0.09) | 56 (11) | 1,317 (305) | 0.113 (0.021) |
| <i>Robinia pseudoacacia</i> | 32 (10) | 0.17 (0.04) | 19 (5) | 697 (210) | 0.026 (0.007) |
| <i>Liriodendron tulipifera</i> | 138 (44) | 0.18 (0.05) | 16 (4) | 454 (136) | 0.029 (0.008) |
| <i>Betula lenta</i> | 215 (69) | 0.20 (0.05) | 6 (2) | 435 (110) | 0.015 (0.004) |
| <i>Hamamelis virginiana</i> | 153 (65) | 0.13 (0.05) | 15 (5) | 236 (89) | 0.030 (0.011) |
| <i>Fagus grandifolia</i> | 18 (6) | 0.05 (0.03) | 6 (4) | 222 (165) | 0.010 (0.006) |
| <i>Sassafras albidum</i> | 55 (27) | 0.06 (0.02) | 7 (3) | 206 (131) | 0.012 (0.005) |
| <i>Castanea dentata</i> | 31 (9) | 0.06 (0.02) | 6 (2) | 119 (39) | 0.009 (0.003) |
| <i>Amelanchier arborea</i> | 22 (5) | 0.04 (0.01) | 5 (1) | 111 (32) | 0.011 (0.003) |
| <i>Tsuga canadensis</i> | 7 (4) | 0.02 (0.01) | 11 (6) | 102 (59) | 0.007 (0.004) |
| <i>Symplocos tinctoria</i> | 10 (4) | 0.02 (0.01) | 2 (1) | 36 (17) | 0.003 (0.001) |
| <i>Diospyros virginiana</i> | 2 (1) | <0.01 (0.01) | 1 (0.6) | 27 (18) | 0.002 (0.001) |
| <i>Magnolia acuminata</i> | 7 (2) | 0.01 (0.004) | 1 (0.5) | 24 (9) | 0.004 (0.002) |
| <i>Fraxinus americana</i> | 12 (7) | 0.01 (0.01) | 1 (1) | 18 (11) | 0.002 (0.001) |
| <i>Salix nigra</i> | 10 (10) | 0.01 (0.01) | 0.59 (0.59) | 7 (7) | 0.001 (0.001) |
| <i>Castanea pumila</i> | 1 (0.8) | <0.01 (0.01) | 0.33 (0.18) | 6 (3) | 0.001 (0.001) |
| <i>Juglans cinerea</i> | 2 (1) | <0.01 (0.01) | 0.16 (0.14) | 5 (4) | 0.0003 (0.0002) |
| <i>Acer pensylvanicum</i> | 1 (1) | 0.01 (0.01) | 0.20 (0.20) | 3 (3) | 0.001 (0.001) |
| <i>R. calendulaceum</i> | 97 (24) | 0.05 (0.01) | 0.14 (0.04) | 0.16 (0.04) | 0.0002 (0.0001) |
| <i>Cleathra acuminata</i> | 62 (33) | 0.03 (0.02) | 0.09 (0.05) | 0.10 (0.05) | 0.0001 (0.0001) |
| <i>Pyrularia pubera</i> | 14 (7) | 0.03 (0.01) | 0.07 (0.04) | 0.08 (0.04) | 0.0001 (0.0001) |
| <i>Calycanthus florid</i> a | 6 (4) | <0.01 (0.01) | 0.01 (0.01) | 0.01 (0.01) | 0.0001 (0.0001) |
| <i>Vaccinium stamineum</i> | 6 (4) | <0.01 (0.01) | 0.01 (0.01) | 0.01 (0.01) | 0.0001 (0.0001) |
| <i>Cornus alternifolia</i> | 5 (2) | <0.01 (0.003) | 0.65 (0.32) | 0.01 (0.01) | 0.001 (0.001) |
| All <i>Quercus</i> | 393 (34) | 7.33 (1.05) | 1,162 (160) | 59,187 (11048) | 1.577 (0.220) |
| Total | 5942 (292) | 15.96 (1.38) | 2,283 (206) | 99,508 (12110) | 3.273 (0.294) |

| | | | | |
|-------------------------------|---|-------|--------------|---------------|
| 1961 – Grass | SLA = 253; LWR = 0.51 from Poorter and Remkes (1990) for <i>Festuca ovina</i> | 3,723 | 7,300 (276) | 9.419 |
| 1962 – Grass | | 3,100 | 6,080 (201) | 7.843 |
| 1963 – Grass | | 2,892 | 5,670 (932) | 7.317 |
| 1965 – Grass | | 3,310 | 6,490 (1067) | 8.374 |
| 1968 - Grasses & forbs | | | 3,918 (390) | 6.462 (0.644) |

Species in 1982

| | | | | | |
|--------------------------------|----------|---------------|-----------|---------------|-----------------|
| <i>Robinia pseudoacacia</i> | 868 (93) | 7.13 (0.72) | 840 (84) | 29,501 (3466) | 1.142 (0.115) |
| <i>Liriodendron tulipifera</i> | 166 (33) | 2.16 (0.40) | 181 (33) | 9,745 (1853) | 0.336 (0.062) |
| <i>Betula lenta</i> | 99 (48) | 0.53 (0.25) | 27 (12) | 2,284 (1032) | 0.072 (0.032) |
| <i>Acer rubrum</i> | 90 (16) | 0.22 (0.05) | 46 (10) | 732 (178) | 0.080 (0.017) |
| <i>Juglans cinerea</i> | 1 (1) | 0.09 (0.09) | 12 (12) | 707 (707) | 0.019 (0.019) |
| <i>Sassafras albidum</i> | 49 (21) | 0.14 (0.06) | 16 (7) | 372 (171) | 0.027 (0.012) |
| <i>Carya</i> spp. | 40 (14) | 0.13 (0.04) | 10 (4) | 367 (128) | 0.017 (0.006) |
| <i>Pinus rigida</i> | 10 (7) | 0.11 (0.08) | 3 (2) | 300 (212) | 0.009 (0.006) |
| <i>Prunus serotina</i> | 13 (5) | 0.08 (0.04) | 9 (5) | 273 (148) | 0.015 (0.007) |
| <i>Diospyros virginiana</i> | 56 (21) | 0.10 (0.04) | 12 (5) | 254 (107) | 0.015 (0.006) |
| <i>Amelanchier arborea</i> | 6 (5) | 0.06 (0.04) | 6 (5) | 217 (154) | 0.011 (0.008) |
| <i>Oxydendrum arboreum</i> | 43 (11) | 0.09 (0.03) | 11 (3) | 200 (66) | 0.030 (0.009) |
| <i>Quercus montana</i> | 10 (5) | 0.06 (0.03) | 6 (4) | 164 (106) | 0.009 (0.006) |
| <i>Cornus florida</i> | 32 (10) | 0.05 (0.02) | 6 (2) | 98 (38) | 0.011 (0.004) |
| <i>Quercus alba</i> | 6 (6) | 0.03 (0.03) | 6 (6) | 85 (85) | 0.009 (0.009) |
| <i>Symplocos tinctoria</i> | 18 (8) | 0.03 (0.01) | 3 (2) | 62 (34) | 0.005 (0.002) |
| <i>Rhododendron maximum</i> | 22 (16) | 0.01 (0.01) | 2 (2) | 52 (38) | 0.002 (0.001) |
| <i>Quercus rubra</i> | 10 (5) | 0.02 (0.01) | 1 (1) | 37 (24) | 0.002 (0.001) |
| <i>Pinus strobus</i> | 3 (2) | 0.02 (0.01) | 0.4 (0.3) | 35 (34) | 0.001 (0.001) |
| <i>Quercus velutina</i> | 7 (3) | 0.01 (0.01) | 1 (1) | 29 (15) | 0.001 (0.001) |
| <i>Salix nigra</i> | 12 (12) | 0.01 (0.01) | 1 (1) | 21 (21) | 0.002 (0.002) |
| <i>Platanus occidentalis</i> | 3 (2) | 0.01 (0.004) | 0.6 (0.5) | 12 (10) | 0.001 (0.001) |
| <i>Fraxinus americana</i> | 3 (3) | 0.003 (0.003) | 0.3 (0.3) | 5 (5) | 0.001 (0.001) |
| <i>Hamamelis virginiana</i> | 4 (4) | 0.003 (0.003) | 0.4 (0.4) | 5 (5) | 0.001 (0.001) |
| <i>Kalmia latifolia</i> | 1 (1) | 0.001 (0.001) | 0.1 (0.1) | 3 (3) | 0.0001 (0.0001) |
| <i>Tilia americana</i> | 1 (1) | 0.002 (0.002) | 0.2 (0.2) | 2 (2) | 0.0003 (0.0003) |
| <i>Vitis</i> spp. | 106 (22) | 0.08 (0.02) | 0.2 (0.1) | 0.3 (0.1) | 0.0004 (0.0001) |

| | | | | | |
|----------------------------|------------|---------------|-------------|---------------|-----------------|
| <i>Rhus copallina</i> | 22 (22) | 0.02 (0.02) | 0.1 (0.1) | 0.1 (0.1) | 0.0001 (0.0001) |
| <i>Sambucus canadensis</i> | 4 (4) | 0.004 (0.004) | 0.01 (0.01) | 0.01 (0.01) | <0.0001 |
| <i>Malus angustifolia</i> | 1 (1) | 0.005 (0.005) | 0.01 (0.01) | 0.01 (0.01) | <0.0001 |
| All <i>Quercus</i> | 34 (9) | 0.12 (0.05) | 15 (7) | 314 (137) | 0.021 (0.011) |
| Total | 1712 (121) | 11.21 (0.83) | 1204 (88) | 45,563 (3761) | 1.823 (0.128) |

Species in 1995

| | | | | | |
|--------------------------------|----------|---------------|-------------|---------------|-----------------|
| <i>Liriodendron tulipifera</i> | 166 (30) | 7.86 (1.07) | 651 (89) | 52,132 (7157) | 1.205 (0.164) |
| <i>Robinia pseudoacacia</i> | 204 (24) | 7.33 (0.88) | 875 (105) | 45,123 (5872) | 1.190 (0.142) |
| <i>Betula lenta</i> | 81 (39) | 1.26 (0.55) | 86 (36) | 7,820 (3267) | 0.234 (0.098) |
| <i>Acer rubrum</i> | 194 (27) | 1.29 (0.25) | 235 (43) | 6,498 (1415) | 0.412 (0.075) |
| <i>Juglans cinerea</i> | 1 (1) | 0.29 (0.29) | 47 (47) | 3,565 (3565) | 0.077 (0.077) |
| <i>Prunus serotina</i> | 13 (5) | 0.44 (0.22) | 53 (27) | 2,832 (1574) | 0.088 (0.044) |
| <i>Carya spp.</i> | 31 (9) | 0.31 (0.10) | 31 (10) | 1,354 (473) | 0.050 (0.017) |
| <i>Sassafras albidum</i> | 26 (13) | 0.28 (0.16) | 33 (18) | 1,179 (683) | 0.056 (0.031) |
| <i>Quercus montana</i> | 6 (4) | 0.18 (0.13) | 25 (18) | 1,136 (847) | 0.035 (0.025) |
| <i>Amelanchier arborea</i> | 6 (4) | 0.15 (0.10) | 17 (12) | 796 (554) | 0.029 (0.020) |
| <i>Cornus florida</i> | 44 (14) | 0.24 (0.08) | 28 (9) | 740 (246) | 0.056 (0.018) |
| <i>Oxydendrum arboreum</i> | 42 (12) | 0.24 (0.08) | 24 (8) | 731 (266) | 0.063 (0.021) |
| <i>Diospyros virginiana</i> | 12 (7) | 0.12 (0.06) | 14 (8) | 519 (316) | 0.023 (0.012) |
| <i>Rhododendron maximum</i> | 147 (60) | 0.17 (0.06) | 20 (8) | 468 (184) | 0.015 (0.006) |
| <i>Quercus alba</i> | 7 (6) | 0.10 (0.09) | 18 (16) | 370 (354) | 0.026 (0.024) |
| <i>Pinus strobus</i> | 4 (2) | 0.11 (0.07) | 5 (3) | 358 (224) | 0.011 (0.007) |
| <i>Kalmia latifolia</i> | 94 (60) | 0.10 (0.07) | 12 (8) | 268 (174) | 0.009 (0.006) |
| <i>Fraxinus americana</i> | 7 (4) | 0.06 (0.04) | 8 (5) | 255 (182) | 0.013 (0.009) |
| <i>Symplocos tinctoria</i> | 18 (6) | 0.04 (0.02) | 5 (3) | 147 (109) | 0.008 (0.005) |
| <i>Platanus occidentalis</i> | 3 (2) | 0.02 (0.02) | 3 (2) | 95 (85) | 0.005 (0.004) |
| <i>Quercus rubra</i> | 4 (2) | 0.02 (0.01) | 2 (1) | 63 (47) | 0.002 (0.001) |
| <i>Pinus rigida</i> | 1 (1) | 0.01 (0.01) | 0.4 (0.4) | 39 (39) | 0.001 (0.001) |
| <i>Quercus rubra</i> | 3 (2) | 0.01 (0.01) | 0.9 (0.8) | 29 (26) | 0.001 (0.001) |
| <i>Hamamelis virginiana</i> | 7 (6) | 0.01 (0.01) | 1 (1) | 22 (21) | 0.002 (0.002) |
| <i>Halesia carolina</i> | 1 (1) | 0.003 (0.003) | 0.3 (0.3) | 6 (6) | 0.001 (0.001) |
| <i>Acer pensylvanicum</i> | 1 (1) | 0.001 (0.001) | 0.2 (0.2) | 2 (2) | 0.0004 (0.0004) |
| <i>Vitis spp.</i> | 144 (26) | 0.17 (0.03) | 0.4 (0.07) | 0.5 (0.1) | 0.001 (0.0001) |
| <i>Clethra acuminata</i> | 12 (8) | 0.01 (0.01) | 0.04 (0.02) | 0.4 (0.3) | 0.0001 (0.0001) |

| | | | | | |
|----------------------------|------------|--------------|-------------|----------------|-----------------|
| <i>Rhus copallina</i> | 40 (24) | 0.12 (0.07) | 0.3 (0.2) | 0.3 (0.1) | 0.0005 (0.0003) |
| <i>Malus angustifolia</i> | 1 (1) | 0.02 (0.02) | 0.04 (0.04) | 0.04 (0.04) | 0.0001 (0.0001) |
| <i>Pyrularia pubera</i> | 10 (7) | 0.01 (0.01) | 0.02 (0.01) | 0.02 (0.02) | 0.0001 (0.0001) |
| <i>Sambucus canadensis</i> | 1 (1) | <0.001 | <0.01 | <0.01 | <0.0001 |
| All <i>Quercus</i> | 21 (7) | 0.32 (0.15) | 46 (23) | 1,598 (900) | 0.065 (0.034) |
| Total | 1338 (142) | 21.00 (1.45) | 2196 (142) | 126,998 (9618) | 3.616 (0.241) |

Species in 2012

| | | | | | |
|--------------------------------|----------|---------------|-------------|-----------------|-----------------|
| <i>Liriodendron tulipifera</i> | 144 (19) | 14.06 (1.53) | 1157 (126) | 115,093 (12904) | 1.840 (0.200) |
| <i>Robinia pseudoacacia</i> | 82 (13) | 4.64 (0.71) | 555 (85) | 32,210 (5287) | 0.689 (0.106) |
| <i>Acer rubrum</i> | 260 (30) | 2.93 (0.49) | 496 (73) | 18,314 (3584) | 0.714 (0.113) |
| <i>Betula lenta</i> | 60 (25) | 1.71 (0.64) | 134 (50) | 12,329 (4562) | 0.166 (0.061) |
| <i>Juglans cinerea</i> | 1 (1) | 0.49 (0.49) | 85 (85) | 7,221 (7221) | 0.140 (0.140) |
| <i>Prunus serotina</i> | 7 (4) | 0.64 (0.40) | 77 (48) | 5,225 (3465) | 0.100 (0.063) |
| <i>Quercus montana</i> | 2 (2) | 0.19 (0.19) | 29 (29) | 1,838 (1838) | 0.036 (0.036) |
| <i>Oxydendrum arboreum</i> | 41 (13) | 0.48 (0.16) | 41 (14) | 1,772 (616) | 0.081 (0.027) |
| <i>Sassafras albidum</i> | 16 (7) | 0.25 (0.18) | 30 (21) | 1,248 (990) | 0.038 (0.028) |
| <i>Carya</i> spp. | 19 (8) | 0.24 (0.11) | 25 (12) | 1,194 (602) | 0.036 (0.017) |
| <i>Fraxinus Americana</i> | 9 (4) | 0.21 (0.11) | 25 (14) | 1,057 (607) | 0.032 (0.018) |
| <i>Diospyros virginiana</i> | 7 (6) | 0.12 (0.09) | 14 (10) | 650 (535) | 0.018 (0.013) |
| <i>Rhododendron maximum</i> | 150 (51) | 0.33 (0.11) | 27 (9) | 650 (218) | 0.008 (0.003) |
| <i>Pinus strobus</i> | 3 (2) | 0.15 (0.11) | 9 (6) | 599 (430) | 0.019 (0.014) |
| <i>Cornus florida</i> | 12 (5) | 0.12 (0.05) | 13 (6) | 429 (204) | 0.027 (0.012) |
| <i>Cornus alternifolia</i> | 3 (3) | 0.06 (0.06) | 13 (6) | 429 (204) | 0.027 (0.012) |
| <i>Quercus alba</i> | 3 (2) | 0.08 (0.07) | 8 (8) | 387 (387) | 0.016 (0.016) |
| <i>Hamamelis virginiana</i> | 25 (17) | 0.08 (0.05) | 9 (6) | 203 (141) | 0.012 (0.008) |
| <i>Kalmia latifolia</i> | 54 (25) | 0.06 (0.03) | 7 (3) | 175 (80) | 0.002 (0.001) |
| <i>Platanus occidentalis</i> | 1 (1) | 0.03 (0.03) | 7 (3) | 175 (80) | 0.002 (0.001) |
| <i>Quercus rubra</i> | 6 (3) | 0.05 (0.03) | 5 (3) | 169 (101) | 0.006 (0.003) |
| <i>Acer pensylvanicum</i> | 1 (1) | 0.003 (0.003) | 4 (4) | 141 (141) | 0.006 (0.006) |
| <i>Symplocos tinctoria</i> | 7 (4) | 0.02 (0.01) | 3 (2) | 73 (46) | 0.005 (0.003) |
| <i>Malus angustifolia</i> | 1 (1) | 0.02 (0.02) | 2 (2) | 62 (62) | 0.003 (0.003) |
| <i>Ligustrum sinense</i> | 7 (7) | 0.02 (0.02) | 2 (2) | 43 (43) | 0.003 (0.003) |
| <i>Castanea dentata</i> | 4 (4) | 0.005 (0.005) | 0.58 (0.58) | 9 (9) | 0.001 (0.001) |
| <i>Clethra acuminata</i> | 24 (24) | 0.04 (0.04) | 0.11 (0.11) | 9 (9) | 0.0002 (0.0002) |

| | | | | | |
|------------------------------------|-----------|---------------|-------------|-----------------|-----------------|
| <i>Juniperus virginiana</i> | 3 (2) | 0.005 (0.004) | 0.05 (0.04) | 7 (6) | 0.0002 (0.0001) |
| <i>Vitis</i> spp. | 201 (29) | 0.31 (0.12) | 0.82 (0.12) | 0.9 (0.1) | 0.001 (0.0002) |
| <i>Aristolochia macrophylla</i> | 3 (3) | 0.002 (0.002) | 0.01 (0.01) | 0.01 (0.01) | <0.0001 |
| <i>R. calendulaceum</i> | 6 (5) | 0.006 (0.005) | 0.02 (0.01) | 0.02 (0.01) | <0.0001 |
| <i>Parthenocissus quinquefolia</i> | 4 (2) | 0.006 (0.004) | 0.02 (0.01) | 0.02 (0.01) | <0.0001 |
| <i>Pyrularia pubera</i> | 3 (3) | 0.003 (0.003) | 0.01 (0.01) | 0.01 (0.01) | <0.0001 |
| All <i>Quercus</i> | 10 (4) | 0.31 (0.20) | 45 (30) | 2,367 (1854) | 0.064 (0.043) |
| Total | 1174 (90) | 27.36 (1.81) | 2770 (185) | 201,470 (14665) | 4.848 (0.330) |

In 1934, only percent cover of evergreen shrubs (*Kalmia latifolia* and *Rhododendron maximum*) was estimated for each plot. For all other years, diameter of evergreen shrubs was measured.

Figure S1

Mean (\pm SE bars) aboveground biomass for the dominant species in the **(a)** treated WS6, **(b)** reference WS14, and **(c)** reference WS18 over time. Dominant species were *Castanea dentata*, *Quercus (montana, coccinea, rubra, velutina, alba)*, *Carya spp.*, *Acer rubrum*, *Liriodendron tulipifera*, *Robinia pseudoacacia*, and *Betula lenta*.




Figure S2

Change in evapotranspiration (D_{ET} , calculated as $P - \hat{Q}_T$; where, $D_{ET} = [P - Q_T] - ([P - \hat{Q}_T]; M1, M2 = 0)$) for the treated WS6 over time (bars). Solid lines are the standard errors of the mean prediction. We used the paired-watershed approach with WS18 as the reference. The year of harvest, conversion to Kentucky-31 fescue grass (*Festuca octiflora*) cover, fertilize, herbicide, and abandonment to allow forest regeneration are denoted by dashed lines.

