

Author's Response

Hess-2016-555

Dear,

Thanks for the editors and reviewers for their time and comments. The manuscript has already been revised to respond to each of the comments.

Sincerely yours

Yonggang Yang and Bojie Fu

Hess-2016-555-RC1

The subject of this manuscript is interesting and worthy of publication. I believe that the topic will be of broad interest to readers. In my opinion, the key contribution of the manuscript is that it aims to investigate soil water migration processes using isotopes, integrated with sampling in the field, experimental observation and laboratory analysis, and to integrate data from several different compartments of the hydrological system, i.e. precipitation, soil water, plant xylem and root etc.

However, I think some minor modifications are required before the manuscript's publication.

2.2 Sample collection and field experiment "Samples of precipitation, soil, xylem, stem, leaf and root were collected in the study area." "Stem samples (about 0.5 to 2 cm diameter and 5 cm length) were obtained." While no linkage of analysis about stem and leaf was found with the previous and the subsequent discussions. This paragraph should be moved and edited.

Response: Done. Page5, lines 117-119. We have revised it. The statement of "stem and leaf" has been removed. The analysis about stem and leaf will be the research point in the future.

4.1.1 Evidences for piston-type flow there are two continuous rainfall events on 20 July 2014 and 24 September 2014, which infiltrating uniformly into deep depths, as shown by the reducing D and ^{18}O values with time after the precipitation. Here how many days did the two continuous rainfall events last respectively? What about the precipitation values? Please give specific explanation.

Response: Done. Page12, lines 267-270. We have revised it. The rainfall event on 20 July 2014, with the precipitation of 25mm, lasted for three days. The rainfall event on 24 September 2014, with the precipitation of 16mm, lasted for two days.

4.2.1 Water origin of water sources are uptake by plants “The results show that isotope composition of xylem water of various plants is different”. In the paper, what part can explain this point? There is no such comparative analysis about different plants, so please give further explanation or delete. In addition, please give explanation about the reason of choosing 130cm as the maximum sampling depth. What about groundwater? Why no discussion in this study?

Response: Done. Page13, lines 307-308. We have revised it. In this paper, our discussion focuses on water origin of *caraganakorshinshiikom* xylem. Therefore, we delete the statement of “The results show that isotope composition of xylem water of various plants is different.”

Characterized by dry climate, less precipitation, more evaporation and thicker soil layers, and groundwater buried deeper, groundwater in the study area is difficult to use due to the depth of water table in the northwestern Loess Plateau. The thickness of the soil layer in vadose zone is typically Tens of meters, even a few hundred meters. Many researchers have shown that groundwater in the study area has no effect on plants. Therefore in this study the influence of groundwater does not exist. Thus, soil water is almost the only water resource in the study area. *Caraganakorshinshiikom* in the study area are artificial planting, old small, root system distribution within the 100 cm. There is little more than 100 cm depth. Therefore, we choose 130cm as the maximum sampling depth.

Hess-2016-555-RC2

The purpose of this paper is to probe into the migration process soil water, and identify each potential water source, which is well justified. There are also good data sets. To my knowledge, this would be the comprehensive study of soil water migration in the unsaturated zone in the western China. It does

look like that the data are very good and useful, especially the isotopic data. Publication of this study could advance our understanding of alpine hydrology and also be of interest to hydrologists. There are some problems with the paper, and it will be revised to get it for publication in the journal. They are as follows:

- (1) Add an inset map (e.g., a small China map) showing where the basin is relative to Beijing and Dingxi.

Response: Done. We have revised it. We have added an inset map (China map), which shows where the basin is relative to Beijing and Dingxi. Fig.1 Location of the sampling sites in the study area

- (2) On page 13 it is stated that isotope composition of xylem water of various plants are different. Soil water at the 40-60cm depth is mainly used by caraganakorshinshikom. Groundwater? Precipitation? Why? I am not saying that the recharge is not possible but the stronger hydrogeological background is needed for it to be plausible. The Area description, especially the Hydrological Background, is also required.

Response: Done. Page4, lines 91-105. We have revised it.

Characterized by dry climate, less precipitation, more evaporation and thicker soil layers, and groundwater buried deeper, groundwater in the study area is difficult to use due to the depth of water table in the northwestern Loess Plateau. The thickness of the soil layer in vadose zone is typically Tens of meters, even a few hundred meters.

The climate is semi-arid, with an average annual temperature of 6.3°C, annual accumulated temperature of 2239°C, extreme maximum temperature 34.3°C, extreme minimum temperature of -27.1 °C in the study area. The annual mean precipitation is 420mm. The annual mean evaporation is 1510mm. The aridity index is 1.15. Precipitation is low and unevenly distributed in temporal and spatial. The summer rainfall accounted for over 60% of the annual precipitation. This area is a part of the typical semi-arid in the hilly and gully region of Loess Plateau, with the altitude ranging from 1900m to 2250m.

The watershed area is 8.91km², which belongs to the hilly and gully region of Loess Plateau. Gully density is 3.14km/km², and the ditch depth ranges from 30 to 50m. Soil type is yellow loessal soil and saline soil, and the average thickness ranges from 40 to 60m. The soil density of soil layer

ranges from 1.1 to 1.4 g/cm³, average soil porosity is 55%. Soil structure has a vertical joint, and the nature of soil is loose and its wet collapsibility is serious. The grassland and shrubland ecosystems are the most extensively dominant ecosystems in the Anjiugou river basin. As it is a representative area of Loess Plateau area, the Anjiugou River basin is a suitable area for soil water study in semi-arid region.

The study area has broken terrain and serious soil erosion, with the terrain being loess long beam and terraces, and gully valley landscape. Geological structure is the uplift zone between the eastern part of Qilian fold system and the west Qinling fold system, at an altitude of 1700 m ~ 2580 m, with the gully density being 3 ~ 5 km/km², ditch slope being 5 ~ 10%, and the mountain slope being generally 20 to 50. The sunny slope is steep, while the shade slope is relatively flat.

The soil parent material is quaternary eolian loess, and the zonal soil mainly is yellow spongy soils, sierozem, which belongs to the typical semiarid loess hilly-gully region. It has soft soil, homogeneous structure, thicker soil layer, good water performance, and the widest distribution. The average thickness is 40 ~ 60 m. Clay soil is between 33.12% ~ 42.17%, organic matter content is between 0.37%~1.34%, soil bulk density is 1.17 g/cm³, wilting moisture content is 7.3%, and the saturated moisture content is 21.9% at the 0 ~ 20 cm. The soil bulk density is 1.09~1.36g cm⁻³, and the porosity is 50%~55% at the 2 m soil layer. The soil has vertical joint and strong collapsibility, so the soil erosion is easily happened, and the soil erosion modulus is 5000 t/km² · a.

(3) Vacuum distillation was considered as a reliable and accepted method for soil water and plant water abstraction in this paper. Please give a description on detailed procedure, such as time and temperature selection.

Response: Done. Page6, lines 131-135. We have revised it.

Water was extracted from soil, leaf, branch, xylem and root by cryogenic vacuum distillation method. The moisture in the soil or plants under the condition of vacuum (vacuum below60MT), was heated by heat set to 105 °C after evaporation. Water vapor of evaporation in -50 °C (liquid nitrogen) was collected with frozen water collecting pipe (top-down frozen, in order to increase the collecting rate), and the precision extract of δD and δ¹⁸ are ±3‰ and ±0.3‰, respectively.

(4) On page 12, “Through analyzing the D and ^{18}O in xylem water of caraganakorshinshiikom, and the D and ^{18}O in potential source such as soil water and precipitation in Anjiagou Gully basin”. Here what is the meaning of “in Anjiagou Gully basin”.

Response: Done. Page13, lines 303-304. We have revised it. The correct expression should be “in Anjiagou River basin”

(5) How about terrain and vegetation?

Response: Done. Page4, lines 91-111. We have revised it.

The vegetation type belongs to arid grassland vegetation type, with less distribution of natural arbor. The grassland and shrubland ecosystems were the most extensive dominant ecosystems. Woodland area is less; with most being open forest land. The area of the crown density being greater than 0.2 are only *Caragana intermedia* and *Pinus tabulaeformis*, and species are single. The distribution of natural herbs mainly are *Ben's s. grandis*, thyme, cold pole, camel peng, with vegetation coverage 10 ~ 20%. Natural coverage is commonly 25% ~ 35% in the sunny slope, 50% ~ 60% in the shade slope. Natural vegetations are mainly compositae Asteraceae, Leguminosae, Gramineae, etc. The vegetation is sparse, and species are relatively poor because of long-term influence by human activities. The artificial forest vegetations mainly are *Caragana intermedia*, *Hippophae rhamnoides*, *Pinus tabulaeformis*, *Platycladus orientalis*, *Stipa bungeana*, etc.

The study area has broken terrain and serious soil erosion, with the terrain being loess long beam and terraces, and gully valley landscape. Geological structure is the uplift zone between the eastern part of Qilian fold system and the west Qinling fold system, at an altitude of 1700 m ~ 2580 m, with the gully density being 3 ~ 5 km/km², ditch slope being 5 ~ 10%, and the mountain slope being generally 20 to 50. The sunny slope is steep, while the shade slope is relatively flat. Clay soil is between 33.12% ~ 42.17%, organic matter content is between 0.37% ~ 1.34%, soil bulk density is 1.17 g/cm³, wilting moisture content is 7.3%, and the saturated moisture content is 21.9% at the 0 ~ 20 cm. The soil bulk density is 1.09~1.36g cm⁻³, and the porosity is 50% ~ 55% at the 2 m soil layer. The soil has vertical joint and strong collapsibility, so the soil erosion is easily happened, and the soil erosion modulus is 5000 t/km² · a.

Hess-2016-555-RC3

This paper examined how water infiltration, evaporation, and transpiration affected the isotope distribution in soil profile. They used water isotopes assisted by the water content measurements. They also used water sourcing approach to identify how much water come from different soil depths and precipitation. Stable isotope is very effective for this research. A lot of data is acquired, and the new knowledge is obtained. I suggest some modifications before publication.

The following are my suggestions.

1, To add description of soil and vegetation in the study area;

Response: Done. Page4, lines 91-111. We have revised it.

2, To add a description of soil water sampling time;

Response: Done. Page5, lines 114-128. We have revised it.

Soil samples in the unsaturated zone were collected from May 2013 to October 2015. Soil was sampled at 10 cm intervals for the first 40 cm, 20 cm intervals from 40 to 100 cm, 30 cm intervals from 100 to 130 cm. Maximum depths of sampling ranged up to 130 cm (Plant root is rarely found below 100 cm in the study area). At each sampling site, soil moisture (volumetric soil water content) was obtained with time domain reflectometry (TDR) in the field manually at 0-10, 10-20, 20-30, 30-40, 40-60, 60-80, 80-100and 100-130 cm. Soil moisture content was determined by oven drying method simultaneously.

3, Keywords should be hydrogen and oxygen stable isotope;

Response: Done. Page2, lines 28.We have revised it.

4, How many precipitation samples were there to define LMWL? Please add sample number.

Response: Done. Page7, lines 162-164. 34 precipitation samples were collected from May 2013 to October 2015.

5, To increase the new relevant references.

Response: Done. Page15, lines 344-435. We have added new relevant references

Hess-2016-555-RC4

Soil moisture for vegetation growth is very important in Loess Plateau. Soil water migration is essential to describe the movement of salt, carbon, nitrogen and other nutrient. Although there are many studies on the soil moisture, there are less works on the mechanism of soil water migration in the unsaturated zone in Loess Plateau of arid region in China based on successive observations and isotopes. The scientific issue of this paper is sure, research idea is clear, research method is correct, and research results are reliable. From my point of view, the work is well-done and provides reliable results to soil water migration in arid region of China.

1. Adding relevant detailed introduction about soil and hydrology in the research area.

Response: Done. Page4, lines 91-111. We have revised it.

The watershed area is 8.91km^2 , which belongs to the hilly and gully region of Loess Plateau. Gully density is $3.14\text{km}/\text{km}^2$, and the ditch depth ranges from 30 to 50m. Soil type is yellow loessal soil and saline soil, and the average thickness ranges from 40 to 60m. The soil density of soil layer ranges from 1.1 to $1.4\text{ g}/\text{cm}^3$, average soil porosity is 55%. Soil structure has a vertical joint, and the nature of soil is loose and its wet collapsibility is serious. The grassland and shrubland ecosystems are the most extensively dominant ecosystems in the Anjiugou river basin. As it is a representative area of Loess Plateau area, the Anjiagou River basin is a suitable area for soil water study in semi-arid region.

The study area has broken terrain and serious soil erosion, with the terrain being loess long beam and terraces, and gully valley landscape. Geological structure is the uplift zone between the eastern part of Qilian fold system and the west Qinling fold system, at an altitude of $1700\text{ m} \sim 2580\text{ m}$, with the gully density being $3 \sim 5\text{ km}/\text{km}^2$, ditch slope being $5 \sim 10\%$, and the mountain slope being generally 20 to 50. The sunny slope is steep, while the shade slope is relatively flat.

The soil parent material is quaternary eolian loess, and the zonal soil mainly is yellow spongy soils, sierozem, which belongs to the typical semiarid loess hilly-gully region. It has soft soil, homogeneous structure, thicker soil layer, good water performance, and the widest distribution. The average thickness is 40 ~ 60 m. Clay soil is between 33.12% ~ 42.17%, organic matter content is between 0.37% ~ 1.34%, soil bulk density is 1.17 g/cm³, wilting moisture content is 7.3%, and the saturated moisture content is 21.9% at the 0 ~ 20 cm. The soil bulk density is 1.09~1.36g cm⁻³, and the porosity is 50% ~ 55% at the 2 m soil layer. The soil has vertical joint and strong collapsibility, so the soil erosion is easily happened, and the soil erosion modulus is 5000 t/km² · a.

2. As mentioned in the paper, “Soil moisture content was determined by oven drying method simultaneously” How much about the depth of soil sample? Does every depth be concluded respectively? Does the measured depth be in accordance with that of time domain reflectometry?

Response: Done. Page5, lines 124-127. We have revised it.

Soil samples in the unsaturated zone were collected at 10 cm intervals from 100 to 130 cm. Maximum depths of sampling ranged up to 130 cm, with every depth at 0-10, 10-20, 20-30, 30-40, 40-60, 60-80, 80-100and 100-130 cm being determined by oven drying method. The measured depth were accordance with that of time domain reflectometry (TDR).

3. Please explain the reason for choosing caraganakorshinshiikom as the object of study, and illustrate its characteristics in the part of research area summary.

Response: Done. The grassland and shrubland ecosystems were the most extensive dominant ecosystems in Loess Plateau. The area accounts for about 50% of the total area, and it is the main part of loess plateau surface cover. Caraganakorshinshiikom is the dominant species in the study area. It is most widely distributed. Therefore, we choose caraganakorshinshiikom as the object of the study.

4. There are many mentioned in the paper, such as “Plant Xylem” “The isotopic composition of soil layer is similar to the xylem water is confirmed by comparing D and 18O of xylem water and soil water, which reflect the signatures of soil at the depth of soil water uptake by plants.” “soil water

from the surface horizons (20-40cm) contributed to 8%-21% of plant xylem water,” plant roots. Is it the xylogen and root system of a certain kind of plant? Caraganakorshinshiikom?

In general, this is a well-written paper containing reliable results which merits publication.

Response: Done. Yes, the xylogen and root system is Caraganakorshinshiikom,

Hess-2016-555-RC5

The authors have tried to do some fundamental research based on field survey and water samples analyses. The study sampled soil profile water stable isotopes of soil, and discussed soil water-plant water recharge sources. This study provides some useful information for soil moisture dynamics and stable characteristics. So it is significant.

However, I could recommend the paper for publication after the following issues have been addressed by the authors:

P1 L19, D is used in the text for the first time, so please state this is Deuterium.

Response: Done. Page1, lines 13.We have revised it. It should be deuterium when D is used in the text for the first time.

P4, L80 “groundwater in the unsaturated zone” There is no relevant analysis about in the whole passage, so it should be deleted. P7, L52-53 P8, 167 “The measured D and ^{18}O of soil water range from -72.42‰ to -37.05‰ and -11.74‰ to -3.57‰ respectively.” Add the little “delta” before ^{18}O such as ^{18}O .

Response: Done. Page4, lines 79-81. Page8, lines 172.We have revised it. We have deleted the groundwater and added the little “delta” before ^{18}O .

p10, L 230-233 “ Most of the sampling sites in Anjiagou River basin (except 14-7-20 and 14-9-24

sites), the isotopic composition of soil water in shallow layer are enriched in D and ^{18}O due to evaporation. The isotopic composition of soil water with depth in the soil profile. ” Here the changes should be specifically explained.

Response: Done. Precipitation is infiltrating and completely mixing with free water. The vertical trend in δD and $\delta^{18}\text{O}$ profile of soil water is simple. There is an abrupt peak value in the isotopic profiles, which suggests that the older water is pushed downward by the new water and infiltrates to the deep soil.

p12, L 261-264 “the preferred flow is rarely found in the Loess Plateau, except that there are which is caused by plant root system or animal invasion, etc.” “The macrospores can be found in the 100-200cm soil layer in the study area, which are caused by plant root or animal activity.” Here “plant root or animal activity” should be changed into “plant root or animal invasion”, and exact illustration is needed.

Response: Done. Page12, lines 265, 284-286. We have revised it. The macrospores are caused by plant root in Loess Plateau.

P13, L 292 “often-used” Here “often-used” should be changed into “often used”.

Response: Done. Page13, lines 311. We have revised it into “often used”.

Soil water migration in the unsaturated zone of semi-arid region in China from isotope evidence

Yonggang Yang^{1, 2}, Bojie Fu²

¹ Institute of Loess Plateau, Shanxi University, Taiyuan, Shanxi 030006, China;

5 ² Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing, 100085, China

Correspondence to: Bojie Fu (bfu@rcees.ac.cn)

Abstract. Soil water is an important driving force of the ecosystems, especially in the semi-arid hilly and gully region of northwestern Loess Plateau in China. The mechanism of soil water migration in the reconstruction and restoration of Loess Plateau is a key scientific problem that must be solved. Isotopic tracers can provide valuable information associated with complex hydrological problems, which is difficult to obtain by other methods. In this study, the oxygen and hydrogen isotopes are used as tracers to investigate the migration processes of soil water in the unsaturated zone in the Loess Plateau of arid region in China. Samples of precipitation, soil water, plant xylems and plant roots are collected and analysed. [The conservative elements deuterium \(D\) and oxygen \(¹⁸O\) are used as tracers](#) to identify variable source and mixing processes. The mixing model is used to quantify the contribution of each end member and calculate mixing amounts. The results show that the isotopic composition of precipitation in the Anjiagou River basin is affected by isotopic fractionation due to evaporation. The isotopic compositions of soil waters are plotted between or near the local meteoric water lines, indicating that soil waters are recharged by precipitation. The soil water migration is dominated by piston-type flow in the study area, but rarely preferential flow. Water migration exhibited a transformation pathway from precipitation to soil water to plant water. $\delta^{18}\text{O}$ and δD are enrichment in the shallow (<20cm depth) soil water in most soil profiles due to evaporation. The isotopic composition of xylem water is close to that of soil water at the depth of 40-60 cm. These values reflect soil water signatures associated with *caraganakorshinshikom* uptake at the depth of 40-60 cm. Soil water from the surface soil lay (20-40cm) contributed to 6%-12% of plant xylem water, while soil water at the depth of 40-60 cm is the largest component of plant xylem water (range from 60% to 66%), soil water below 60 cm depth contributed 8%-14% to plant xylem water, and only 5%-8% is derived directly from precipitation. This study investigates the migration process of soil water, and

25 identifies the source of plant water, and finally provides a scientific basis for identification of model structure and parameter. It can provide a scientific basis for ecological water demand, ecological restoration, management of water resources and the improvement of water benefit on each scale.

Keywords: soil water; xylem water; [stable isotope](#); migration; unsaturated zone

1 Introduction

30 Water in the soil environment plays a crucial role as a carrier of dissolved and solid species and as a reservoir in the hydrological cycle. Soil water possesses a small proportion (only 0.05 %) within the hydrological cycle, but it is vital for an ecosystem and affects spatial and temporal processes at different scales (Koeniger et al., 2016; Busari et al., 2013). Understanding soil water migration in the unsaturated zone is essential to describe the movement of salt, carbon, nitrogen and other nutrient. Soil water migration plays an important role in the processes of infiltration, evaporation, 35 transpiration and percolation, hydraulic conductivity and water uptake capacity of soils in the unsaturated zone. The traditional methods have been carried out to study the movement of soil water such as hydrologic experiment, intensive observations, modeling and remote sensing (Luo et al., 2013; Yang et al., 2013; Carucci et al., 2012). However, soil water migration is a complex nonlinear and inhomogeneous flow process. It is difficult to model on the basis of Darcy's law exclusively, and is also difficult to obtain by other techniques. Climatic conditions, soil texture and soil structure, antecedent 40 moisture and vegetation cover exert influences on the movement of soil water (Bose et al., 2016; Kidron et al., 2013).

Isotopic tracers can provide valuable information on the complex hydrological problems, such as the runoff processes, residence time, runoff pathway, the origin and contribution of each runoff component (Arny et al., 2013; Ohlanders et al., 2013; Yang et al., 2012a; McInerney et al, 2011; Maurya et al., 2011; Kevin et al., 2010). The isotopes of D and ^{18}O are conservative and do not react with clay minerals and other soil materials. D and ^{18}O are widely used to investigate ecological 45 and hydrological processes. Stable isotopes can provide information about mixing, transport processes and residence time of water within a soil profile in the unsaturated zone.

Different flow mechanisms result in different isotopic profiles. Stable isotope compositions of soil water, plant xylems and precipitation can be obtained to identify soil water migration processes such as infiltration, evaporation, transpiration

and percolation (Caley et al., 2013; Yang et al., 2012b; Catherine et al., 2010; Stumpp et al., 2009). Zimmerman et al. (1967) firstly applied stable isotopes to study the soil water profile, showing that evaporation at the surface of a saturated soil column causes deuterium enrichment near the surface that decreases exponentially with depth. Robertson and Gazis (2006) studied the seasonal trends of oxygen isotope composition in soil water fluxes at two sites along the climate gradient. The hydrogen and oxygen isotopes are used to study the transforming of precipitation, soil water and groundwater of typical vegetation in Taihang mountain area (Song et al., 2009). Gazis and Feng (2004) compared the oxygen isotope compositions of precipitation and soil water from profiles at six sites with different soil textures. δD and $\delta^{18}O$ in soil water has been observed in many studies, including column experiments using sand or soil columns and field observations of unsaturated soils (Singh et al.,2013; N ávar et al.,2011; Brodersen et al.,2015; Gehrels et al., 1998).

Stable isotope is the evidence for assessing plant water sources according to their variations caused by equilibrium and kinetic isotopic fractionation mechanisms (Orlowski et al., 2016a; Haverd et al. 2011; Zhao et al. 2011; Bhatia et al., 2011). Several studies (Liu et al.,2013; Mathieu et al., 1996; Allison et al., 1985) have shown that transpiration do not cause isotopic fractionation of soil water. The isotope δD and $\delta^{18}O$ is a powerful tool to determine the water sources utilized by plants in the field, since δD and $\delta^{18}O$ values of xylem water reflect those of the water sources utilized (Gierke et al., 2016; Yang et al., 2011; Lu et al.,2011). The naturally occurring vertical gradients of δD and $\delta^{18}O$ in soil water provide similar information about plant water uptake depth from soils. A number of studies have used isotopes to characterize soil water movement in one location and thus one climatic regime (Orlowski et al., 2016b; Manzoni et al., 2013; Liu et al., 2011; Stumpp et al., 2009).

Soil moisture is an important driving force of the ecosystems, especially in the northwestern Loess Plateau. Characterized by dry climate, less precipitation, more evaporation and thicker soil layers, groundwater in this region is difficult to use due to the depth of water table. Thus, soil water is almost the only water resource in the study area, which has become the only factor controlling agricultural production and ecological restoration. It is necessary to investigate the mechanism of soil water migration in the loess plateau. Stable isotopic can provide valuable information on the mechanism of soil water migration. At present, a large number of research focuses on the soil water recharge by precipitation and its isotope variation characteristics. There is some research on the interaction of soil water, plant water and precipitation. These

studies are mostly on the individual scale in one specific region (Lin et al., 2016; Heathman et al., 2012; William et al., 2010; Ferretti et al., 2003). However, the research on the migration process of soil water, plant water, precipitation and groundwater in the unsaturated zone based on the isotopic technique is still rarely. At present, the mechanism of soil water migration in the Loess Plateau is the key scientific problem to be solved.

Therefore, this study investigates soil water migration processes in the unsaturated zone in the hilly and gully region of Loess Plateau using isotopes, integrated with sampling in the field, experimental observation and laboratory analysis. Samples include the soil water, plant xylem and root etc. The objectives of this research are: (1) to probe into the migration process and variation of plant water, soil water and precipitation; and (2) to identify each potential water source uptake by plant, and evaluate their contributions of each potential water source in the unsaturated zone. It can provide a scientific basis for ecological water demand, ecological construction, management of water resources and the improvement of water benefit on each scale.

2. Materials and methods

2.1 The experimental site

The study area is located in Anjiagou River basin in the Dingxi City of China, with latitudes 34°26'-35°35'N, and longitudes 103°52'-104°39'E. The climate is semi-arid, with an average annual temperature of 6.3°C. The annual mean precipitation is 420mm. The mean annual evaporation is 1510mm. The aridity index is 1.15. Precipitation is low and unevenly distributed in temporal and spatial. This area is a part of the typical semi-arid in the hilly and gully region of Loess Plateau, with the altitude ranging from 1900 m to 2250 m.

The watershed area is 8.91km², which belongs to the hilly and gully region of Loess Plateau. Gully density is 3.14km/km², and the ditch depth ranges from 30 to 50m. Soil type is yellow loessal soil and saline soil, and the average thickness ranges from 40 to 60m. The soil density of soil layer ranges from 1.1 to 1.4 g/cm³, average soil porosity is 55%. Soil structure has a vertical joint, and the nature of soil is loose and its wet collapsibility is serious. The grassland and shrubland ecosystems are the most extensively dominant ecosystems in the Anjiugou river basin.

The study area has broken terrain and serious soil erosion, with the terrain being loess long beam and terraces, and gully

valley landscape. Geological structure is the uplift zone between the eastern part of Qilian fold system and the west Qinling fold system, at an altitude of 1700 m ~ 2580 m, with the gully density being 3 ~ 5 km/km², ditch slope being 5 ~ 10%, and the mountain slope being generally 20 to 50. The soil parent material is quaternary eolian loess, and the zonal soil mainly is yellow spongy soils, sierozem, which belongs to the typical semiarid loess hilly-gully region. It has soft soil, homogeneous structure, thicker soil layer, good water performance, and the widest distribution. The average thickness is 40~60 m. Clay soil is between 33.12%~42.17%, organic matter content is between 0.37% ~ 1.34%, soil bulk density is 1.17 g/cm³, wilting moisture content is 7.3%, and the saturated moisture content is 21.9% at the 0~20 cm. The soil bulk density is 1.09~1.36g cm⁻³, and the porosity is 50% ~ 55% at the 2 m soil layer. The soil has vertical joint and strong collapsibility, so the soil erosion is easily happened, and the soil erosion modulus is 5000 t/km² · a.

The vegetation type belongs to arid grassland vegetation type, with less distribution of natural arbor. The grassland and shrubland ecosystems were the most extensive dominant ecosystems. Woodland area is less; with most being open forest land. The area of the crown density being greater than 0.2 are only *Caragana intermedia* and *Pinus tabulaeformis*, and species are single. The vegetation is sparse, and species are relatively poor because of long-term influence by human activities. The artificial forest vegetations mainly are *Caragana intermedia*, *Hippophae rhamnoides*, *Pinus tabulaeformis*, *Platycladus orientalis*, *Stipa bungeana*, etc.

2.2 Sample collection and field experiment

The research was carried out in Anjiagou River basin in the hilly and gully region of Loess Plateau from May 2013 to October 2015. Samples of precipitation, soil, xylem and root were collected in the study area. The locations of the sampling sites are shown in Fig.1. Precipitation was collected after each rainfall event. Precipitation was filtered and transferred to sealed glass vials prior to analysis. Plant xylem and root of *caraganakorshinshii* were collected at each sampling site, respectively. For sap water analysis, a plant twig (0.5 to 1cm in diameter and about 5 cm in length) was cut from a main branch or trunk, cortical and phloem were peeled off. All samples were kept in the 8 ml glass bottle, sealed with packaging tape to reduce evaporation. Five glass bottles were collected for plant in each sampling site. All samples were collected at

least monthly. Altogether 27 samples sites and 396 samples were collected, and sealed with parafilm. The samples were taken back to the laboratory and preserved at 4 °C, and samples of soil, plant xylem, plant stem and leaf were refrigerated at -20 °C until analysis.

125 Soil was sampled at 10 cm intervals for the first 40 cm, 20 cm intervals from 40 to 100 cm, 30 cm intervals from 100 to 130 cm. Maximum depths of sampling ranged up to 130 cm (Plant root is rarely found below 100 cm in the study area). At each sampling site, soil moisture (volumetric soil water content) was obtained with time domain reflectometry (TDR) in the field manually at 0-10, 10-20, 20-30, 30-40, 40-60, 60-80, 80-100 and 100-130 cm. Soil moisture content was determined by oven drying method simultaneously.

130 2.3 Laboratory analyses

Water was extracted from soil, xylem, root, stem and leaf by cryogenic vacuum distillation method, and the extracted water (2 to 10 ml) was trapped at liquid–nitrogen temperature. Vacuum distillation was considered as a reliable and accepted method. The moisture in the soil or plants under the condition of vacuum (vacuum below 60MT), was heated by heat set to 105 °C after evaporation. Water vapor of evaporation in -50 °C (liquid nitrogen) was collected with frozen water collecting pipe (top-down frozen, in order to increase the collecting rate), and the precision extract of δD and $\delta^{18}O$ are $\pm 3\%$ and $\pm 0.3\%$, respectively. Water samples were filtered through 0.2 μ m Millipore membrane for trace elements analyses. Isotopes $\delta^{18}O$ and δD were measured in the Key Laboratory of Ecohydrology and River Basin Science of Cold and Arid Region Environmental and Engineering Institute, Chinese Academy of Sciences. $\delta^{18}O$ of samples was analyzed by a Euro-PyrOH elemental analyzer at a temperature of 1300°C, and δD was analyzed at a temperature of 1030 °C. At both temperatures the reaction products were analyzed on a GV ISOPRIMETM continuous flow IRMS. In order to eliminate the memory effect of online and continues flow, every sample was analyzed 6 times. The analytical precision of δD determinations was $\pm 1\%$, and that of $\delta^{18}O \pm 0.2\%$. Isotopic concentration was expressed as δ -per million (‰) relative to the Vienna Standard Mean Ocean Water, according to the follow Eq. :

$$\delta^{18}O(\text{‰}) = \frac{(^{18}O/^{16}O)_{\text{sample}} - (^{18}O/^{16}O)_{\text{SMOW}}}{(^{18}O/^{16}O)_{\text{SMOW}}} \times 1000, \quad \delta D(\text{‰}) = \frac{(D/H)_{\text{sample}} - (D/H)_{\text{SMOW}}}{(D/H)_{\text{SMOW}}} \times 1000$$

When n isotope is used to determine the proportional contributions of $i+I$ sources to a mixture, standard linear mixing models can be used to mathematically solve for the unique combination of source proportions that conserves mass balance for all n isotopes (Phillips et al., 2013). In this study, the conservative elements D and ^{18}O are utilized to calculate mixing amounts of potential water source. The mixing model is used to calculate the contributions of each respective end-member.

150 The mixing model is described on the basis of the mass balance Eq.:

$$\begin{aligned}\delta\text{D}_p &= \sum_{i=1}^n f_i \delta\text{D}_i \\ \delta^{18}\text{O}_p &= \sum_{i=1}^n f_i \delta^{18}\text{O}_i \\ I &= \sum_{i=1}^n f_i\end{aligned}$$

Where δD and $\delta^{18}\text{O}$ are concentrations of the tracers; I refers to potential water source; δD_p and $\delta^{18}\text{O}_p$ refer to plant waters; f_i is the fraction of each component contributing to plant water.

155

3. Result

3.1 Isotopic composition of precipitation

Yurtsever and Gat (1981) modified Craig's global water line, and made a more accurate global precipitation linear relationship between δD and $\delta^{18}\text{O}$ of $\delta\text{D}=8.17\delta^{18}\text{O}+10.56$, which is the Global Meteoric Water Line (GMWL). Figure 2 shows the relationship between the mean $\delta^{18}\text{O}$ and mean δD in precipitation, which is defined as the local meteoric water line (LMWL) with an Eq. of $\delta\text{D}=7.41\delta^{18}\text{O}+7.22$, $R^2=0.95$. The $\delta^{18}\text{O}$ and δD compositions of the precipitation in the Anjiagou River catchment are extremely variable, ranging from -118.36‰ to -22.32‰ , and from -16.35‰ to -2.19‰ , respectively. 34 precipitation samples were collected from May 2013 to October 2015. There are significant differences in the composition of D and ^{18}O under different precipitation event. The difference reflects the extreme nature of climate and the complexity of moisture sources in the arid region. Compared with GMWL, the slope and intercept of LMWL are lower than that of GMWL (Fig.2), indicating that they are affected by the local climate and environment with less precipitation and

165

lower humidity. When the slope is low, the implication is that precipitation is subject to evaporation. The Anjiagou River basin is located in the northwestern inland region, and it is difficult for the sea water vapor to reach directly. This indicates that the significant imbalance of isotope dynamic fractionation exists, which are strongly affected by strong solar radiation and evaporation in the local environment, leading to the enrichment in $\delta^{18}\text{O}$ and δD . Therefore, the slope of LMWL is low, and intercept of LMWL decreases with the slope.

3.2 The variations of δD and $\delta^{18}\text{O}$ in soil water of vertical profiles

Stable isotope compositions of soil water are presented in Table 1. The measured δD and $\delta^{18}\text{O}$ of soil water range from -72.42‰ to -37.05‰, and -11.74‰ to -3.57‰, respectively. By comparing the isotopic composition of soil water with that of precipitation, the isotope compositions of soil water in the unsaturated zone are relatively enriched. Therefore, it could be beneficial to identify the influence of various water bodies on different precipitation event. As Fig.2 shows, isotope compositions of most soil water are plotted in the bottom of the LMWL, and are close to the LMWL. The variable trends of δD and $\delta^{18}\text{O}$ values of soil water are similar to that of precipitation. There is a strong linear relationship ($R^2 = 0.96$) for δD and $\delta^{18}\text{O}$ between precipitation and soil water, which indicates that soil water is originally recharged by precipitation. The soil water is increasingly enriched in D and ^{18}O , which are influenced by evaporation (Table 1). It has been reported that soil water is normally more enriched because precipitation enters the soil, and mixes with antecedent soil water that has been modified by evaporation.

Vertical profiles of soil water δD - ^{18}O in the Anjiagou River basin are shown in Fig.3. The surface layer (0-20 cm) showed a larger variations and higher δD - ^{18}O values than the deeper layer. δD and ^{18}O values of soil water are greatest at the surface layer (Fig.3, Table 1). δD - ^{18}O values of shallow soil water at 0-10cm are -10.29‰ to -7.36‰, -72.71‰ to -35.45‰, respectively. The δD - ^{18}O values of shallower soil water at 10-20cm are also high (ranges from -8.08‰ to -3.57‰, and ranges from -67.27‰ to -49.70‰, respectively). The net effect of evaporation is the enrichment of heavy isotopes near the soil surface. Soil profiles (Fig.3) show the increase in the D- ^{18}O with soil depths from the 0 to 20cm, suggesting the occurrence of evaporation. The large variation in isotope compositions of shallow depths (0-20cm depth) may be caused by precipitation and evapotranspiration effect. The correlation of the isotopic composition between soil water and precipitation is weak.

These have been reported by many other studies (Gazis and Feng, 2004; Robertson and Gazis, 2006).

The $\delta^{18}\text{O}$ and δD compositions of soil water at 40cm depths range from -11.35‰ to -7.09‰, and -74.21‰ to -54.06, respectively. Soil water from 20 to 40cm depths is easily influenced by precipitation infiltration and evapotranspiration, which leads to the extremely variable δD and $\delta^{18}\text{O}$ values of soil water (Fig.3). The impact of precipitation on soil water becomes weaker with increasing depths. The isotopic composition of soil water is strongly enriching in δD and $\delta^{18}\text{O}$ due to less precipitation and evaporation in the study area. The δD and $\delta^{18}\text{O}$ decrease exponentially with the increasing depths, and the variation trends are larger. Previous studies have also shown that evaporation decreases with the increasing soil depth, and its influence is generally within tens of centimeters. δD - $\delta^{18}\text{O}$ declined with depth to 60 cm but remained constant at deeper depths. When it reaches the 60cm depth, the variation trend tends to be stationary. The $\delta^{18}\text{O}$ and δD compositions of soil water at 60cm depths ranged from -12.79‰ to -9.75‰, and ranged from -96.91‰ to -57.27, respectively. There is an abrupt peak value in the plant root zone. The isotopic fluctuations quickly disappear due to the dispersive effects of the deep root system.

However, the values of δD and $\delta^{18}\text{O}$ are gradually decreasing from 60 to 80cm because the recharge of precipitation is decreasing, and the influence of antecedent isotope value of soil water is increasing. As the Fig.3 shows, D and ^{18}O values of soil water at depths of 80 cm range from -11.53‰ to -8.92‰, and range from -78.82‰ to -62.80‰, respectively. Compared with the large variation in the δD and $\delta^{18}\text{O}$ of surface soil water, soil water deeper than 80 cm is almost constant. The low value zone is at 100cm in the isotopic profiles of soil water, which is the deepest layer that recharged by precipitation. Variations of $\delta^{18}\text{O}$ and δD of soil water in the deep layer are non-significant. Evaporation is larger than precipitation in the study area. Precipitation is difficult to infiltrate to 80 cm depth in the shrubland ecosystems, except in the high flow years. Therefore, it is difficult to cause the isotope fractionation of soil water below 80cm.

3.3 Variability in soil moisture content

Values of soil moisture obtained by TDR in the Anjiagou River basin are shown in Fig.5. The variation of soil moisture content in the shallow soil layer (0-40 cm) is extremely large, ranging from 5.94 to 17.51%, which belongs to the variable greatly layer. The variation of soil moisture content at the 40-60 cm soil depth is relatively small, which belongs to the less

215 active layer. The variation of soil moisture content is relatively stable at 60-100 cm soil depth, ranging from 17.78% to 29.47%, which belongs to the relatively stable layer. Soil water content also exhibits depth variation. Variability of water content is larger in the surface horizons more than 40 cm depth in the soil profile. The shallow soil layers are impacted by evaporation and precipitation recharge more than the deeper soil layer. Therefore, high evapotranspiration rate and precipitation recharge are the main factor controlling soil moisture especially the surface horizon.

220 As the soil moisture profile shows (Fig.4), soil moisture content is low in the shallow layer (0-20 cm), soil water content is relatively high at the active layer (40-60cm), and then tends to be stable. Figure 4 shows that soil moisture content is different in different sample sites, but soil moisture content is low under 60 cm soil depths in all the sample sites. Soil moisture content generally increases with the increase of soil depth in the variable greatly layer. Soil moisture content decreases with the increase of soil depth in the active layer, and then tends to be stable.

225 As Fig.4 shows, water content increases and soil water $\delta^{18}\text{O-D}$ values increase in the shallow layer (0-40 cm), which are impacted by evapotranspiration. This is the dominant process for most of the sites. Mean value of the soil moisture content increases with depth in the soil profile (40-60cm depth), while soil water $\delta^{18}\text{O-D}$ values decrease with depth, which mixes with antecedent moisture. Low water content and more positive $^{18}\text{O-D}$ values (Fig.4) occur at the deep soil layer (below 60cm depth), which is impacted by recharge. Single precipitation is difficult to infiltrate to deeper than 80 cm, when drought lasts for a long time, especially when soil water content lower than 20%. Extremely high values of soil water content at the 230 depth of 60 cm are observed in Anjiagou River basin. The higher water content and more negative d^{18}O values are observed at the active layer (40-60 cm) (Fig.4).The variability of water content and the $^{18}\text{O-D}$ values are relatively larger for the surface horizons than deeper in the soil profile. The profile of soil moisture content is highly responsive to evaporation and precipitation. The average water content and soil water $\delta^{18}\text{O-D}$ values are representative of the trends found in most of the 235 sample sites.

4. Discussion

4.1 The migration process of soil water in the unsaturated zone

Precipitation infiltration and evapotranspiration in the vertical direction are the main processes of the water cycle, which

plays an important role in the transformation process of water cycle in the unsaturated zone. Different migration mechanisms
of soil water result in different isotopic profiles. Therefore, isotope δD and $\delta^{18}O$ of soil water and precipitation can be
240 obtained to identify soil water migration processes such as infiltration, evaporation, transpiration and percolation through the
soil waters at different depths. These different isotopic profiles may occur as a result of several processes, including
evaporation, change in isotopic composition of precipitation and mixing of new and old water. While evaporation is widely
recognized, we believe that mixing is also important factors for controlling the isotopic composition of soil water at these
245 sites. These are possibly caused by the infiltration of summer rainwater through the soil layer. Generally, there are two major
infiltration mechanisms (piston-type flow and preferential flow) that can be identified by comparing the isotope
compositions of precipitation and soil water.

4.1.1 Evidences for piston-type flow

The δD and $\delta^{18}O$ content of soil water at different depths are shown in Fig.4. By comparing the isotopic compositions of
precipitation and soil water, the mechanisms of soil water movement can be identified. Most of the sampling sites in
250 Anjiagou River basin (except 14-7-20 and 14-9-24 sites), the isotopic composition of soil water in shallow layer are enriched
in D and ^{18}O due to evaporation. The isotopic composition of soil water changes with depth in the soil profile, especially
after the continuing precipitation between August 18 and August 20, 2014. Precipitation is infiltrating and completely mixing
with free water. The vertical trend in δD and $\delta^{18}O$ profile of soil water is simple. There is an abrupt peak value in the isotopic
255 profiles, which suggests that the older water is pushed downward by the new water and infiltrates to the deep soil. That is,
while some soil water remains stationary, the mobile soil water successively displaces pre-existing mobile soil water pushing
it downward. Piston-type flow after an isotopic distinct rainfall event results in an abrupt “isotopic front” within the soil
(Gehrels et al., 1998; Song et al., 2009). This evidence for mixing, abrupt changes in the δD and $\delta^{18}O$ of soil water with depth
in the soil profile indicates that piston-flow has occurred. A relatively flat pattern is observed, slightly enriched in
260 accordance with precipitation. δD and $\delta^{18}O$ declined with depth to 60 cm but remained constant at deeper depths. The older
soil water is pushed by newer rainwater downward. Precipitation infiltration exhibits piston infiltration characteristics
(Zimmerman, 1967).

4.1.2 Evidences for preferential flow

At the 14-7-20 and 14-9-24 samples sites, the vertical trend of δD and $\delta^{18}O$ in the soil profile indicates a uniform infiltration process (Fig.5). There are two abrupt peak values in the isotopic vertical profiles of soil water. The soil water at depths between 20 cm and 40 cm decrease quickly, which is caused by the strong transpiration effect of plants roots. Abrupt change occurred at 20 cm-40 cm depth of the isotopic profile where the soil water moisture increased quickly (range from -14.97‰ to -12.78‰, range from -107.93‰ to -101.21‰, respectively). There are two continuous rainfall events on 20 July 2014 and 24 September 2014 (The rainfall event on 20 July 2014, with the precipitation of 25mm, lasted for three days. The rainfall event on 24 September 2014, with the precipitation of 16mm, lasted for two days), which infiltrating uniformly into deep depths, as shown by the reducing δD and $\delta^{18}O$ values with time after the precipitation (Fig.4). The data do not show the obvious correlation with depth at the 14-7-20 and 14-9-24 samples sites in Anjiagou River basin (Fig.4). This is mainly caused by the mixture effect occurred during the process of infiltration. That is to say, the newer water, which did not replace the old water totally during the process of infiltration, mixed with the old water stored in the soil.

The second peak of $\delta^{18}O$ and δD values of soil water is at the depth of 80 cm (range from -12.41‰ to -10.75‰, range from -98.87‰ to -82.77‰, respectively), which is caused most likely by macropore flow. The non-existence of a significant difference between the isotopic content of precipitation and soil water suggests that the infiltrating summer precipitation passes the root zone quickly and contributes to the soil water recharge. The deep soil water is recharged by precipitation in the form of preferential flow, which passes the soil porosity and quickly reaches the deep layer, and do not mix with older water. Variations of $\delta^{18}O$ and δD of deep soil water are only by mixing, and rarely by evaporation. A number of studies documenting tracer movement in the soil column have pointed out the existence of preferential fluid flow, which can be caused by cracks (in dried soils), macropores (Beven and Germann, 1982), plant roots, earthworm burrows, etc. (Vincent et al., 2001; Bronswijk et al., 1991). However, the preferred flow is rarely found in the Loess Plateau, except that there are macropores, which is caused by plant root system or animal invasion, etc. The deep soil water is quickly recharged by precipitation in the form of preferential flow through observing the excavated soil profile. The macropores can be found in the 100-200cm soil layer in the study area, which are caused by plant root. The cracks in the Loess Plateau can also provide

the important path for preferential flow. This is supported by the variations of soil water content. The soil water content at 80 cm depth increases quickly. The soil water content ranges from 34.5 to 36.8%. At corresponding depths of the isotope profile, the isotope values are relatively depleted. Significant soil water content increase in the deep soil is similar to variability of isotopic composition of soil water. The isotope profile can provide some evidence for preferential flow.

4.2 water origin and contribution of each potential water sources are uptake by plants

4.2.1 Water origin of water sources are uptake by plants

Zimmermann et al. (1967) studied the effect of transpiration with an experiment in which the root water uptake of various plant species is monitored, and no fractionation is found. Allison et al. (1985) tested a much greater fraction of root water uptake, no conclusive proof of fractionation by root water uptake is found. The isotope fractionations are not caused when soil water is uptaken by roots of plant. By comparing the δD and $\delta^{18}O$ of plant water and each water sources, the water origin and contribution of each component for xylem could be identified.

The isotopic composition of soil layer is similar to the xylem water is confirmed by comparing δD and $\delta^{18}O$ of xylem water and soil water, which reflect the signatures of soil at the depth of soil water uptake by plants. All the soil and xylem samples are taken simultaneously in Anjiagou River basin. The isotopes of δD and $\delta^{18}O$ are conservative and do not react with clay minerals and other soil materials. During the process of transpiration, the soil water content changes, but the isotopic composition of soil water remains constant. The isotopic composition of plant xylem water is a mixture of soil water at different soil depths. Through analyzing the δD and $\delta^{18}O$ in xylem water of *caraganakorshinshiikom*, and the δD and $\delta^{18}O$ in potential source such as soil water and precipitation in Anjiagou River basin (Table 2). Comparing the signatures δD and $\delta^{18}O$ of both soil water and xylem water with a simple graphical interference approach indicates a distinct difference between both isotopes. Graphical “best match” approach is suitable to illustrate water uptake depth as plants often withdraw water from more than one distinct soil depth. The δD and $\delta^{18}O$ in xylem water ranged from -11.64‰ to -7.95‰, and from -81.62‰ to -55.21‰, respectively (Fig.6, Table 2). Plant Xylem water δD and $\delta^{18}O$ for both species is similar to that of soil water at 40-60cm depth (Fig.6). It indicates that soil water at the 40-60cm depth is mainly used by *caraganakorshinshiikom*.

310 4.2.2 The contribution of each potential water sources to plants water

Hooper et al. (1990) and Christophersen and Hooper (1992) introduced the end-member mixing analysis (EMMA), which is an often used method for analyzing possible source area contributions to flow. Multiple source mixing models (Parnell et al. 2010; Phillips et al.2003) account for water uptake from more than one discrete soil layer and weigh the importance of certain layers for water uptake by incorporating e.g. soil water potentials into the calculation. In this study, the mixing model is used to identify potential source areas and mixing processes, and to quantify the contribution of each end member using mixture fractions. The conservative elements δD and $\delta^{18}O$ are utilized as tracers to calculate mixing amounts.

It is more likely that plant xylem water is a mixture of soil water from several soil depths. Multiple source mass-balance mixing models of proportional contributions of plant xylem water (%) (Parnell et al. 2010) is used to evaluate the contributions of each potential water source (Fig.6).The results show that,soil water from the surface horizons (20-40cm) contributed to 8%-21% of plant xylem water, while soil water at 40-60cm soil depth is the largest component of plant xylem water (range from 68% to 83%), soil water below 60 cm depth contributed 10%-26% to plant xylem water, and only 0%-18% is contributed directly from precipitation. Water source is dominated by soil water at a depth of 40-60 cm.

5. Conclusions

325 The following conclusions can be drawn from the present study:

(1) $\delta^{18}O$ and δD enrichment in the shallow (< 20 cm depth) soil water, observed in most soil profiles, is due to evaporation. The isotopic composition of xylem water is close to that of soil water at a depth of 40-60 cm. These values reflect soil water signatures associated with shrubland uptake at a depth of 40-60 cm.

(2) A sharp isotopic front at approximately 40cm depth observed shortly after an isotopic distinct rainfall event suggests that infiltration into soil occurred as piston-type flow with newer water pushing older water downward in the soil profile. Soil waters are recharged from precipitation. The soil water migration is dominated by piston-type flow in the study area, but rarely preferential flow, except that there are macropores in the Loess Plateau, which is caused by plant root or animal invasion, etc. Water migration exhibited a transformation pathway from precipitation to soil water to plant water.

(3) Soil water from the surface horizons (20-40cm) contributed to 8%-21% of plant xylem water, while soil water at 40-60cm soil depth is the largest component of plant xylem water (range from 68% to 83%), soil water below 60 cm depth contributed 10%-26% to plant xylem water, and only 0%-18% is contributed directly from precipitation. Water source is dominated by soil water at depth of 40-60cm.

Acknowledgements

This research is supported by National Natural Science Foundation of China (41390464;41201043), and China Postdoctoral Science Foundation Funded Project (2014M550095). The authors are grateful to experimental research station and all participants in the field for their contributions to the progress of this study. We also express our appreciation to the anonymous reviewers of the manuscript.

References

- Allison, G. B. and Barnes, C. J.: Estimation of evaporation from the normally “dry” Lake Frome in South Australia, *J. Hydrol.*, 78, 229-242, 1985.
- Arny, E., Svein, B. D., Hans, C., Steen, L., Thorsteinn, J., Sigfus, J. and Johnsen, J.: Monitoring the water vapor isotopic composition in the temperate North Atlantic, *J. Geop. Res., Abstracts*, 15, 2013-5376, 2013.
- [Bose T, Sengupta S, Chakraborty S, Borgaonkar H, Reconstruction of soil water oxygen isotope values from tree ring cellulose and its implications for paleoclimate studies , *Quaternary International*:387-398, 2016.](#)
- Beven, K. and Germann, P.: Macropores and water flow in soils, *Water Resour. Res.*, 18, 1311-1325, 1982.
- Bhatia, M. P., Das S. B. and Kujawinski, E. B.: Seasonal evolution of water contributions to discharge from a Greenland outlet glacier: insight from a new isotope-mixing model, *J. Glaciol.*, 57, 929-941, 2011.
- [Brodersen C, Pohl S, Lindenlaub M, Leibundgut C, Wilpert KV.: Influence of vegetation structure on isotope content of throughfall and soil water, *Hydrol.Process*, 14\(14\):1439-1448, 2015.](#)
- Bronswijk, J. J. B.: Magnitude, modelling and significance of swelling and shrinkage processes in clay soils, PhD, 1991.

- Busari, M. A., Salako, F. K., Tuniz, C. and Zuppi, G. M.: Estimation of soil water evaporative loss after tillage operation using the stable isotope technique, *Int. Agrophys.*, 27, 257-264, 2013.
- 360 Caley, T. and Roche, D. M.: A paleoperspective based on present-day data-model comparison for oxygen stable isotopes in carbonates, *Geo. Sci.*, 1527-1558, 2013.
- Carucci, V., Petitta, M. and Aravena, R.: Interaction between shallow and deep aquifers in the Tivoli Plain enhanced by groundwater extraction: A multi-isotope approach and geochemical modeling, *Appl. Geochem.*, 27, 266-280, 2012.
- Catherine, C. G., Alan, F. M. and Katharine, J. M.: Hydrological processes and chemical characteristics of low-alpine patterned wetlands, south-central New Zealand, *J. Hydrol.*, 385, 105-119, 2010.
- 365 Christophersen, N. and Hooper, R. P.: Multivariate analysis of stream water chemical data: the use of components analysis for the end-member mixing problem, *Water Resour. Res.*, 28, 99-107, 1992.
- Ferretti, D. F., Pendall, E., Morgan, J. A., Nelson, J. A., LeCain D. and Mosier, A. R.: Partitioning evapotranspiration fluxes from Colorado grassland using stable isotopes: Seasonal variations and ecosystem implications of elevated atmospheric CO₂, *Plant Soil*, 254, 291-303, 2003.
- 370 Gehrels, J. C., Peeters, E. M., De Vries, J. J. and Dekkers, M.: The mechanism of soil water movement as inferred from ¹⁸O stable isotope studies, *Hydrol. Sci. J.*, 43, 579-594, 1998.
- [Gierke C, Newton BT, Phillips FM.:Soil-water dynamics and tree water uptake in the Sacramento Mountains of New Mexico: a stable isotope study, *Hydrogeology Journal*, 24\(4\):1-14,2016.](#)
- Haverd, V., Cuntz, M. and Griffith, D.: Measured deuterium in water vapour concentration does not improve the constraint on the partitioning of evapotranspiration, using a soil vegetation atmosphere transfer model, *Agr. Forest Meteorol.*, 151, 645-654, 2011.
- 375 Heathman, G. C., Cosh, M. H., Merwade, V. and Han, E.: Multi-scale temporal stability analysis of surface and subsurface soil moisture within the Upper Cedar Creek Watershed, Indiana, *Catena*, 95, 91-103, 2012.
- Hooper, R. P.: Diagnostic tools for mixing models of stream water chemistry, *Water Resour. Res.*, 39, 1055, 2003.
- 380 Kevin, W. T., Brent, B. W. and Thomas, W. D. E.: Characterizing the role of hydrological processes on lake water balances in the old Crow Flats, Yukon Territory, using water isotope tracers, *J. Hydrol.*, 386, 103-117, 2010.

- Kidron, G. J. and Gutschick, V. P.: Soil moisture correlates with shrub-grass association in the Chihuahuan Desert, *Catena*, 107, 71-79, 2013.
- 385 Koeniger P, Gaj M, Beyer M, Himmelsbach T. :Review on soil water isotope-based groundwater recharge estimations, *Hydrol.Process*, 30(16), 2016.
- Komor, S. C. and Emerson, D. G.: Movements of water, solutes, and stable isotopes in the unsaturated zones of two sand plains in the upper Midwest, *Water Resour. Res.*, 30, 253-267, 1994.
- Lin, Y., Horita, J.: An experimental study on isotope fractionation in a mesoporous silica-water system with implications for vadose-zone hydrology. *Geochimica et Cosmochimica Acta*, 184: 257-271. 2016.
- 390 Liu, H., Tian, F. and Hu, H. C.: Soil moisture controls on patterns of grass green-up in Inner Mongolia: an index based approach, *Hydrol. Earth Syst. Sc.*, 17, 805-815, 2013.
- Liu, Y., Xu, Z., Duffy, R., Chen, W., An, S., Liu, S. and Liu F.: Analyzing relationships among water uptake patterns, rootlet biomass distribution and soil water content profile in a subalpine shrubland using water isotopes, *Eur. J. Soil Biol.*, 47, 380-386, 2011.
- 395 Lu, N., Chen, S.P., Wilske, B., Sun, G. and Chen, J.: Evapotranspiration and soil water relationships in a range of disturbed and undisturbed ecosystems in the semi-arid Inner Mongolia, China, *J. Plant Ecol.*, 4, 49-60, 2011.
- Luo, G. J., Kiese, R. and Wolf, B.: Effects of soil temperature and moisture on methane uptake and nitrous oxide emissions across three different ecosystem types, *Biogeosciences*, 10, 3205-3219, 2013.
- Manzoni, S., Giulia, V., Gabriel, K., Sari, P. and Robert, B. J.: Hydraulic limits on maximum plant transpiration and the emergence of the safety–efficiency trade-off, *New Phytol.*, 198, 169-178, 2013.
- 400 Mathieu, R. and Bariac, T.: An isotopic study (^2H and ^{18}O) of water movements in clayey soils under a semiarid climate, *Water Resour. Res.*, 32, 779-789, 1996.
- Maurya, A. S., Shah, M. and Deshpande, R. D.: Hydrograph separation and precipitation source identification using stable water isotopes and conductivity: River Ganga at Himalayan foothills, *Hydrol. Process.*, 25, 1521-1530, 2011.
- 405 McInerney, F. A., Helliker, B. R. and Freeman, K. H.: Hydrogen isotope ratios of leaf wax nalkanes in grasses are insensitive to transpiration, *Geochimica Et Cosmochimica Acta*, 75, 541-554, 2011.

- Návar, J.: Stemflow variation in Mexico's northeastern forest communities: Its contribution to soil moisture content and aquifer recharge, *J. Hydrol.*, 408, 35-42, 2011.
- Ohlanders, N., Rodriguez, M. and McPhee, J.: Stable water isotope variation in a central Andean watershed dominated by glacier and snowmelt, *J. Hydrol. Earth Syst. Sci.*, 17, 1035-1050, 2013.
- Orlowski N, Pratt DL, McDonnell JJ.: Intercomparison of soil pore water extraction methods for stable isotope analysis, *Hydrol. Process*, 30(19), 2016 a.
- Orlowski, N., Breuer, L., McDonnell, J.J., Critical issues with cryogenic extraction of soil water for stable isotope analysis. *Ecohydrology*, 9: 3-10. 2016b.
- Parnell, A. C., Inger, R., Bearhop, S. and Jackson, A. L.: Source Partitioning Using Stable Isotopes: Coping with Too Much Variation, *Plos. One*, 5, 9672, 2010.
- Phillips, S. J., Anderson, R. P. and Schapire, R. E.: Maximum entropy modeling of species geographic distribution, *Ecol. Model*, 19, 231-259, 2013.
- Robertson, J. A. and Gazis, C. A.: An oxygen isotope study of seasonal trends in soil water fluxes at two sites along a climate gradient in Washington state (USA), *J. Hydrol.*, 328, 375-387, 2006.
- Singh, B. P.: Isotopic composition of water in precipitation in a region or place, *J. Appl. Radiat. Isot*, 75, 22-25, 2013.
- Song, X. F., Wang, S. Q., Xiao, G. Q., Wang, Z. M., Liu, X. and Wang, P.: A study of soil water movement combining soil water potential with stable isotopes at two sites of shallow groundwater areas in the North China Plain, *Hydrol. Process.*, 23, 1376-1388, 2009.
- Vincent, M., Didon-Lescot, J. F. and Couren, M.: Investigation of the hydrological processes using chemical and isotopic tracers in a small Mediterranean forested catchment during autumn recharge, *J. Hydrol.*, 247, 215-229, 2001.
- William, T. P. and Eric, E. S.: The influence of spatial patterns of soil moisture on the grass and shrub responses to a summer rainstorm in a Desert ecotone, *Ecosystems*, 13, 511-525, 2010.
- Yang, Y. G., Xiao, H. L., Wei, Y. P. and Zou, S. B.: Hydrochemical and hydrological processes in the different landscape zones of alpine cold region in China, *Environ. Earth Sci.*, 65, 609-620, 2012a.
- Yang, Y. G., Xiao, H. L., Wei, Y. P. and Zou, S. B.: Hydrological processes in the different landscape zones of alpine cold

regions in the wet season, combining isotopic and hydrochemical tracers, *Hydrol. Process*, 25, 1457-1466, 2012b.

Yang, Y. G., Xiao, H. L., Wei, Y. P. and Zou, S. B.: Hydrologic processes in the different landscape zones in the alpine cold region during the melting period, *J. Hydrol.*, 409, 149-156, 2011.

435 Yang, Y. G., Xiao, H. L., Zou, S. B.: Hydrogen and oxygen isotopic records in monthly scales variations of hydrological characteristics in the different landscape zones, *J. Hydrol.*, 499, 124-131, 2013.

Yurtsever, Y. and Gat, J. R.: Atmospheric waters. In: *Stable Isotope Hydrology: Deuterium and Oxygen-18 in the Water Cycle*, Technical Report Series, IAEA, Vienna, 210, 103-142, 1981.

440 Zhao, L. J., Wang, L. X. and Xiao, H. L.: The effects of short-term rainfall variability on leaf isotopic traits of desert plants in sand-binding ecosystems, *Ecol. Eng.*, 60, 116-125, 2013.

Zimmerman, U., Ehhalt, D. and Mu ninnich, K. O.: Soil-water movement and evapotranspiration: changes in the isotopic composition of the water, In: *Proceedings of the IAEA Symposium on Isotopes in Hydrol.*, 567-585, 1967.