

Impact of snow deposition on major and trace element concentrations and elementary fluxes in surface waters of Western Siberian Lowland across a 1700-km latitudinal gradient

Vladimir P. Shevchenko¹, Oleg S. Pokrovsky², Sergey N. Vorobyev³, Ivan V. Krickov³, Rinat M. Manasyrov^{3,4}, Nadezhda V. Politova¹, Sergey G. Kopysov³, Olga M. Dara¹, Yves Auda², Liudmila S. Shirokova^{2,4}, Larisa G. Kolesnichenko³, Valery A. Zemtsov³, Sergey N. Kirpotin³

¹Shirshov Institute of Oceanology RAS, 36 Nakhimovsky Pr., Moscow, Russia

²Geosciences Environment Toulouse, UMR 5563 CNRS, University of Toulouse, 14 Avenue Edouard Belin 31400, Toulouse, France

³BIO-GEO-CLIM Laboratory, Tomsk State University, 36 Lenina, Tomsk, Russia

⁴N. Laverov Federal Center for Integrated Arctic Research, Russian Academy of Science, Arkhangelsk, Russia

Correspondence to: Oleg S. Pokrovsky (oleg.pokrovsky@get.omp.eu)

Abstract. Towards a better understanding of chemical composition of snow and its impact on surface water hydrochemistry in poorly studied Western Siberia Lowland (WSL), the surface layer of snow was sampled in February 2014 across a 1700-km latitudinal gradient (c.a. 56.5 to 68°N). We aimed at assessing the latitudinal effect on both dissolved and particulate forms of element in snow and quantifying the impact atmospheric input to element storage and export fluxes in inland waters of the WSL. The concentration of dissolved+colloidal (< 0.45 µm) Fe, Co, Cu, As, La increased by a factor of 2 to 5 north of 63°N compared to southern regions. The pH and dissolved Ca, Mg, Sr, Mo and U in snow water increased with the increase in concentration of particulate fraction (PF). Principal Component Analyses of major and trace element concentration in both dissolved and particulate fractions revealed 2 factors not linked to the latitude. A hierarchical cluster analysis yielded several group of elements originated from alumino-silicate mineral matrix, carbonate minerals and marine aerosols or belonging to volatile atmospheric heavy metals, labile elements from weatherable minerals and nutrients. The main sources of mineral components in PF are desert and semi-desert regions of central Asia.

The snow water concentration of DIC, Cl, SO₄, Mg, Ca, Cr, Co, Ni, Cu, Mo, Cd, Sb, Cs, W, Pb and U exceeded or were comparable with spring-time concentration in thermokarst lakes of the permafrost-affected WSL zone. The spring-time river fluxes of DIC, Cl, SO₄, Na, Mg, Ca, Rb, Cs, metals (Cr, Co, Ni, Cu, Zn, Cd, Pb), metalloids (As, Sb), Mo and U in the discontinuous to continuous permafrost zone (64-68°N) can be explained solely by melting of accumulated snow. The impact of snow deposition on riverine fluxes of elements strongly increased northward, in discontinuous and continuous permafrost zones of frozen peat bogs. This was consistent with the decrease of the impact of rock lithology on river chemical composition in the permafrost zone of WSL, relative to the permafrost-free regions. Therefore, the present study demonstrates significant and previously underestimated atmospheric input of many major and trace elements to their riverine fluxes during spring flood. A broader impact of this result is that current estimations of river water fluxes response to the climate warming in high latitudes may be unwarranted without detailed analysis of winter precipitation.

38 1 Introduction

39 The snow cover exhibits a number of properties making it **unique** natural archive and indicator of the ecosystem
40 status (Baltrėnaitė et al., 2014; Bokhorst et al., 2016; Callaghan et al., 2011; Caritat et al., 1998, 2005; Garbarino et al.,
41 2002; Guéguen et al., 2016; Kashulina et al., 2014; Lisitzin, 2002; Niu et al., 2016; Ross and Granat, 1986; Singh et al.,
42 2011; Siudek et al., 2015; Van de Velde et al., 1999; Walker et al., 2003). The snow washes out insoluble aerosols particles
43 from the atmosphere as well as soluble compounds, including various pollutants (Telmer et al., 2004; Barrie, 1986; Tranter
44 et al., 1986, 1987). Unlike rain, the snow remains at the soil surface and thus records all atmospheric input during the
45 glacial period of the year. In boreal and subarctic regions, both dissolved and particulate **fraction** of snow water reflect the
46 air chemistry in winter, when the land is covered by snow and the water surfaces are **covered by ice**. During winter, the
47 input of mineral compounds from adjacent regions is minimal and the main factor controlling chemical composition of
48 snow is long-range, **hundred and thousand km**, atmospheric transport (Franzén et al., 1994; Huang et al., 2015;
49 Shevchenko, 2003, Shevchenko et al., 2000, 2010, 2016; Welch et al., 1991; Zdanowicz et al., 1998, 2006; Krachler et al.,
50 2005; Zhang et al., 2015).

51 Several studies of major elements and some trace metals in particulate fraction of snow have been conducted in
52 western Siberia (Boyarkina et al., 1993; Ermolov et al., 2014; Kashulina et al., 2014; Moskovchenko and Babushkin, 2012;
53 Shevchenko et al., 2015; Talovskaya et al., 2014). The dissolved ($< 0.45 \mu\text{m}$ or $< 0.22 \mu\text{m}$) fraction of snow was
54 traditionally studied in European subarctic (Caritat et al., 1998; Chekushin et al., 1998; Kashulina et al., 2014; Reimann et
55 al., 1999; Reinosdotter and Viklander, 2005) but the data on trace elements in snow water collected in boreal, arctic and
56 subarctic regions are limited. This is especially true for large and geographically homogeneous territories of western
57 Siberia, presenting relatively similar **level** of snow deposition during winter seasons (i.e., from 100 mm of water in the
58 south to 140–150 mm of water in the north) without any pronounced influence of large industrial centers, mountain regions
59 and marine aerosols over the territory close to 1.5 million km² (Resources, 1972, 1973; Boyarkina et al., 2013).

60 The originality of the present study consists in *i*) sampling of substantial (~1700 km) latitudinal transect in
61 relatively pristine zones comprising forest, forest tundra and tundra within the permafrost-free, discontinuous and
62 continuous permafrost regions; *ii*) assessment of both dissolved+colloidal and particulate forms of major and trace
63 elements in snow samples. Given the scarcity of available measurements of snow chemical and particulate composition in
64 Western Siberia, we aimed at addressing the following specific issues: (1) characterizing the effect of the latitude on major
65 and trace element concentration in dissolved ($< 0.45 \mu\text{m}$) and particulate ($> 0.45 \mu\text{m}$) fractions of snow; (2) testing the link
66 between dissolved and particulate fractions of elements and the impact of particle mineralogy on snow chemical
67 composition; (3) comparing dissolved concentrations of major and trace elements in snow to those in lakes and rivers
68 across the latitudinal gradient of WSL and (4) assessing the share of snow deposition on seasonal and annual export of
69 dissolved elements by western Siberian rivers. Via addressing quantitatively the abovementioned issues using **unified**
70 methodology in quite large geographical coverage (56 to 68°N) of orographically flat low populated terrain, we anticipate
71 to enhance our knowledge of the winter atmospheric deposition in western Siberia, in the absence of direct influence of
72 marine aerosols and large industrial centers. This should eventually allow to evaluate the impact of snow deposition on
73 chemical composition and elementary fluxes of subarctic inland waters across a large latitudinal gradient of climate and
74 permafrost parameters.

75

76 2. Study site, materials and methods

77 2.1. Geographic settings

78 Western Siberia Lowland, located between the Ural mountains and the Yenisei River, extends over 2000 km
79 from south to north and presents highly homogeneous, from physico-geographical point of view, taiga, forest-tundra
80 and tundra landscapes comprising bogs and mires in the permafrost-free zone and thermokarst lakes developed on flat
81 peat bogs (palsa) in the permafrost-bearing zone. Detailed physico-geographical description, hydrology, lithology and
82 soils can be found in earlier works (Botch et al., 1995; Smith et al., 2004; Frey and Smith, 2007; Beilman et al., 2009)
83 and in our recent geochemical studies (Manasypov et al., 2014; Stepanova et al., 2015; Pokrovsky et al., 2015; Raudina
84 et al., 2017). Because of its flat orographic context, extensive vegetation cover and relative remoteness from the Arctic
85 Coast (except the north of the Gyda and the Yamal peninsulas), the atmospheric precipitates in winter are likely to bear
86 the signature of remote desert and semi-desert regions of Central Asia. The anthropogenic impact is not expected to be
87 strongly pronounced because of *i*) low population density (average 6 people/km² but only 0.5–2 people/km² in the
88 northern half of WSL) and *ii*) moderate local pollution from the gas burning in oil wells mostly in the permafrost-free
89 zone, south of the Surgut town. The part of WSL north of 64°N contains essentially gas exploration facilities (no gas
90 burning) and minimally impacts the environment. Taken together, the latitudinal profile of the WSL presents a unique
91 opportunity to study the chemistry of atmospheric deposits within highly homogeneous physico-geographical context
92 and relatively low local anthropogenic impact.

94 2.2. Snow sampling

95 The snow of the WSL was sampled along the latitudinal transect S → N, from the vicinity of the Tomsk city (zone
96 of southern taiga) to the eastern coast of the Ob estuary (tundra zone) from 19.02.2014 to 5.03.2014 (Fig. 1). The possible
97 sources of snow deposition and the pathways of aerosols transport to the WSL were reconstructed by analyzing
98 meteorological maps and by calculating back trajectories of air transport to the observation points using NOAA's
99 HYSPLIT model (Draxler and Rolf, 2003). In order to assess a snapshot of snow deposition across 1700-km latitudinal
100 profile and collect the freshest snow that was subjected to minimal transformation, we chose to sample only the upper layer
101 of the snow cover. This technique, in contrast to traditional sampling of full snow column (i.e., Guéguen et al., 2016; Niu et
102 al., 2017) allows adequate representation of the upper fresh snow layer that had minimal transformation at the soil, and
103 frequently used in remote regions (Kang et al., 2007; Zhang et al., 2013). The isotope composition of collected snow
104 proved its fresh character, not subjected to any metamorphism (Vasil'chuk et al., 2016).

105 The upper 0–5 cm of snow was sampled in 39 locations (Fig. 1). All sampling points were located more than 500
106 m from the winter road. The sampling was performed using metal-free technique, in protected environment, using pre-
107 cleaned plastic shovel and vinyl single-used gloves. Approximately 30 L of snow was collected into single-used
108 polyethylene bags. These polyethylene bags were thoroughly washed with 1 M HCl and abundant MilliQ water in the
109 clean room class A 10,000. In the laboratory, the snow was melted at ambient temperature, and filtered through pre-
110 weighted acetate cellulose filters (Millipore, 47 mm diameter) of 0.45 µm poresize. The storage of unfiltered snow
111 water samples was less than 1 h at 4°C.

113 2.3. Particle analyses

114 The sizes and morphology of particles on filters and elemental composition of individual particles were studied
115 using scanning electron microscope VEGA 3 SEM (Tescan) with a microprobe attachment INCA Energy (Oxford
116 Instruments). The mineralogical composition of particulate fraction on selected filters was studied by X-ray powder

117 diffractometric method on the D8 ADVANCE (Bruker AXS) X-ray diffractometer equipped with the LYNXEYE linear
118 detector (Lisitzin et al., 2015). The uncertainty of the relative proportion of mineral composition was 1–2% and the
119 detection limit was 1%.

120 Freshly melted snow water was filtered through pre-weighted 0.45 μm acetate cellulose (Millipore) filters. These
121 filters were placed in **the Petri** dishes, dried at 60°C in an oven and digested using microwave acid attack which comprised
122 6.5 mL concentrated HNO_3 , 3.5 mL concentrated HCl and 0.5 mL concentrated HF . HNO_3 and HCl were bi-distilled in the
123 clean room and HF was **commercial** ultra-pure quality (Fluka). The filters were reacted 30 min in ultrasonic bath prior full
124 digestion using **Mars** 5 microwave digestion system (CEM, France). For this, 10 samples of filters, 1 certified 2711a
125 Montana II Soil standard and 1 blank filter sample were loaded into Teflon reactors **subjected** to treating at 150°C during
126 20 min. After completing the digestion, the content of reactors was transferred to 30 mL Savilex vials and evaporated at
127 70°C. The residue was dissolved in 10 mL of 10% HNO_3 and diluted by 2% HNO_3 prior to the analyses. **For the analysis of**
128 **snow particles on filters, the blanks were estimated after digestion of 6 random filters. In the digestion solution, the**
129 **concentrations of all trace elements were a factor of 10 to 100 lower than that obtained from the filters with particles after**
130 **0.5-1.0 L of snow water filtration.** The concentration of major and trace elements (TE) in filter digestion products was
131 measured using an ICP-MS Agilent 7500 ce with $\sim 3 \mu\text{g/L}$ of indium and rhenium as internal standards. **4** in-house external
132 standards were analyzed every 10 samples. Necessary corrections for oxide and hydroxide ion interferences were made for
133 rare earth elements (REE) and metals (Ariés et al., 2000). Based on replicate analyses of in-house standards and certified
134 materials, the uncertainty for TE measurement ranged from 5 % at 0.1–100 $\mu\text{g/L}$ to 10 % at 0.001–0.01 $\mu\text{g/L}$. Analyses of
135 low concentrations of Hf, Ge, Cs, Ga, and W (e.g., on the order of 0.001 $\mu\text{g/L}$, comparable with detection limits) was
136 possible with a minimal estimated uncertainty of 20%.

137

138 **2.4. Melted snow analyses**

139 The pH and specific conductivity were measured on unfiltered snow water samples using Hanna portable
140 instruments. The dissolved ($< 0.45 \mu\text{m}$) fraction of snow water was obtained via filtration using a polycarbonate
141 Nalgene vacuum filter unit, and a PVC-made Mityvac hand vacuum pump. This fraction included colloidal and truly
142 dissolved (ionic) forms. Blanks of MilliQ water were also placed in polyethylene bags for the same time as melting
143 snow ($\leq 1 \text{ h}$ at 4°C) and processed via filtration similar to snow samples. The filtrates were divided into two parts; one
144 was acidified with double distilled HNO_3 acid and stored in pre-cleaned HDPE vials for ICP MS analysis, the second
145 part was stored in HDPE bottles without acidification, for dissolved organic and inorganic carbon analysis (DOC and
146 DIC), respectively, and anion analysis.

147 The major anion concentrations (Cl^- , SO_4^{2-}) in the $< 0.45 \mu\text{m}$ fraction were measured using ion chromatography
148 (HPLC, Dionex ICS 2000i), with an uncertainty of 2%, estimated from the replicate analyses of PERADE and RAIN
149 international certified materials. The DOC and DIC in this fraction were analyzed using a Carbon Total Analyzer
150 (Shimadzu TOC-VSCN) with an uncertainty of 5% and a detection limit of 0.1 and 0.05 mg/L, respectively.

151 Filtered snow water samples were analyzed with an Element XR ICP MS allowing for much better precision of
152 the analyses of highly diluted samples and avoiding many interferences compared to Agilent 7500 ce. The uncertainty of
153 the Element XR analysis was $\pm 5\%$, while its detection limit was a factor of 100 lower than the traditional (Agilent)
154 instrument. The Element XR operated in three modes depending on the elements measured: low resolution for B, Rb, Sr,
155 Zr, Mo, Cd, Sb, Cs, Ba, REEs, Hf, W, Pb, Th, U; medium resolution for Na, Mg, Al, Si, P, Ca, Ti, V, Cr, Mn, Fe, Co, Ni,
156 Cu, Zn, Ga, Sr, and high resolution for K and As. The agreement between two instruments for most elements was within 10
157 %. The international geostandards SLRS-5 (Riverine Water References Material for Trace Metals certified by the National
158 Research Council of Canada) was used to assess the validity and reproducibility of the analyses. **For all major and most**

159 trace elements in the snow water, the concentrations in the blanks were below or comparable with analytical detection
160 limits (≤ 0.1 ng/L for Cd, Ba, Y, Zr, REEs, Hf, Pb, Th, U; 1 ng/L for Ga, Ge, Rb, Sr, Sb; ~ 10 ng/L for Ti, V, Cr, Mn, Fe,
161 Co, Ni, Cu, Zn, As). These values were at least 5 times lower than the average concentration of trace elements in snow
162 samples. Most TE presented in this work exhibited $\leq 15\%$ -agreement between the certified or recommended values and our
163 measurements. The TE for which certified or recommended data were not available were considered only for the cases
164 where we obtained good analytical reproducibility (i.e., the relative standard deviation based on our standard measurements
165 was $\leq 10\%$).

168 2.5. River fluxes and snow storage

169 The mass balance calculation of the degree of snow melt influence on element fluxes in WSL rivers was
170 performed, taking into account of *i*) the water stock in snow (in mm snow water accumulated during winter), fairly well
171 known for Western Siberia (Karnatzevitch and Khruschev, 2014; Resources, 1972, 1973; Zakharova et al., 2011) and *ii*)
172 the spring-time river runoff (in mm during May and June) calculated from hydrological parameters. For water stock
173 calculation, we used the available mean multi-annual daily and monthly discharges of WSL rivers across the latitudinal
174 profile (Resources, 1972 and 1972 and recently compiled in the database R-ArcicNET (www.r-arcticnet.sr.unh.edu)).
175 The WSL territory is covered by Russian Hydrological Survey (RHS) gauging stations which allowed to calculate the
176 discharge during May-June as described elsewhere (Pokrovsky et al., 2015). The most recent complete hydrological
177 data of small and medium size rivers in permafrost – affected area of WSL (Novikov et al., 2009) were used together
178 with RHS database to calculate the spring flood fluxes of individual rivers and snow water stock for three latitudinal
179 zones, 56-60°N, 60-64° and 64-68°N. Note that a comparison between the elementary snow stock and the river
180 elementary discharge could not be performed for individual river watersheds, since no snow water chemical data are
181 available with necessary spatial resolution. Therefore, we compared the winter snow stock with riverine spring flood
182 fluxes of major and trace element for three latitudinal zones. For this, both spring flood flux of individual rivers and
183 snow water stock were averaged for each latitudinal zone.

185 2.6. Statistical methods

186 Statistical analysis of the average, median and geometric mean values and the link between element concentration
187 in suspended and dissolved fraction as well as comparison of different sampling sets (snow water and snow particles)
188 included ANOVA, H-criterion of the Kruskal–Wallis and Mann–Whitney U tests. These tests allowed evaluating the
189 difference between two sets of data separately for each TE following the approaches developed for lakes and rivers of
190 western Siberia (Manasypov et al., 2014, 2015; Pokrovsky et al., 2015, 2016a). The multiple regressions were performed
191 for quantifying the relationship between dissolved and particulate concentration of TE and the latitudinal trends of
192 concentrations and enrichment factors. More thorough statistical treatment of both log-transformed and non-transformed
193 major and TE concentration in dissolved and particulate fraction of snow samples in each location included a normed PCA
194 analysis using the ADE-4 R package (Thioulouse et al., 1997; Chessel et al., 2004) using the methods for scores and
195 variables (De la Cruz and Holmes, 2011).

196 To identify the group of elements that behaved in a similar way in snow water and snow particles, we applied a
197 complementary hierarchical cluster analysis (HCA, Hartigan, 1975; Kaufman and Rousseeuw, 2005) which is widely
198 adopted in geochemical interpretations of element concentration data (e.g. Bini et al., 2011; Levitan et al., 2015; Schot and
199 van der Wal, 1992; Moragues-Quiroga et al., 2017). We used the Ward's method (Ward, 1963) for the linkages rule,
200 following previous studies (Gourdol et al., 2013; Lin et al., 2014). The Pearson correlation distance was used for the

201 linkage distance, which is frequently used for cluster variables (Reimann et al., 2008). These choices are in agreement with
202 the group search of the PCA loadings.

203 In order to assess the degree of element fractionation in snow particles, Al-normalized TE enrichment factor
204 (EF) with respect to the average upper part of continental earth crust (Rudnick and Gao, 2003) was calculated according
205 to:

$$EF = \frac{[TE]/[Al]_{sample}}{[TE]/[Al]_{crust}}$$

206 For the assessment of element enrichment factor in snow particles, a normalization to both general upper Earth
207 crust and the local geological background (soil, peat and moss) was used. The reason for this is that, unlike in studies of the
208 local pollution tracing in the European arctic (e.g., within the Kola Ecogeochemistry project, see de Caritat et al., 1997;
209 Reimann and de Caritat, 2000; Reimann et al., 2000) or small-scale stream bed sediments or soils (N'guessan et al., 2009;
210 Moragues-Quiroga et al., 2017; Levitan et al., 2015) where the normalization to the local soil or bedrock was necessary, the
211 present study essentially deals with winter-period long-range atmospheric transport of soluble and mineral forms of
212 elements. As such, following the common practice in this field, the normalization to upper Earth crust allowed assessing
213 the true enrichment/depletion of the atmospheric aerosols. However, in order to better represent the elementary features of
214 snow particles, the concentration of elements in PF was also compared with western Siberian mineral soils, peat and moss.
215

216

217

218 3. Results

219 3.1. Soluble fraction of the snow water

220 The latitude-averaged concentrations of dissolved and particulate fraction of snow samples are listed in **Table 1**.
221 A full data set of major and TE concentration in snow water is given in the “Data availability” section. Examples of the
222 effect of latitude on dissolved (< 0.45 µm) element concentrations are shown in **Fig. 2**. Fe and Cu demonstrated a 2 to
223 5-fold increase in dissolved concentration north of 63°N (at p < 0.05). Zn and Pb did not exhibit any systematic effect of
224 latitude, and Sb, Cd and Ni demonstrated a single maximum at c.a. 63-65°N. As exhibited two maxima, at 63.5 and
225 67.5°N with overall 2 to 3-fold decreasing trend northward. All other major and trace elements demonstrated a lack of
226 systematic variation of concentrations as a function of latitude (not shown).

227 The PCA treatment of soluble fraction suggested that at least two factors are interpretable. The PC1 x PC2
228 correlation circle revealed two large groups of variables (**Fig. 3A**). The first group is composed of Al, Fe, Cr, Zr, Pb,
229 REEs corresponding to lithogenic, poorly soluble trace elements. The second large group is composed of DOC, K, Rb,
230 Cs, Mn, Co, Ba, Sb, Co, Mo, Mg, Si, Sr, Na, Ca, pH. These highly mobile elements presumably reflect the marine
231 aerosols and leaching from soluble soil minerals such as carbonates as well as plant biomass. Similar factors determine
232 chemical composition of snow water regardless of the latitude of the sampling and no specific conditions or limiting
233 factors depended on geographical location.

234 The HCA was conducted on the basis of first two factors of the PCA. The criterion of non-intersection between
235 the groups allowed partitioning the chemical elements of the dissolved part into 6 specific groups presented in **Fig. 3B**.
236 These groups characterize the elements according to their general chemical properties, ability to mobilize in aqueous
237 solution from the solid minerals, affinity to the biota or their presence in the contaminated particles of industrial
238 activity. Thus, the first two group of the dissolved fraction shown in **Fig. 3B** and encircled in **Fig. 3A** comprise low
239 mobile elements likely originated from alumino-silicate mineral matrix (Al, Cr, REE, Ti, Zr, Fe, V) as well as some
240 volatile heavy metals typically present in the solid aerosol particles (Cu, Cd, Pb). The 4th group includes major
241 constituents of carbonate or marine aerosols matrix (elevated pH, Mg, Ca and Na). The 5th group is represented by

242 typical macro- and micronutrients (K, Rb, Mn, Co, Ba). Finally, the last 6th group of elements comprises both labile
243 elements linked to weatherable minerals (Sr, Sb, Si, Ni) and nutrients such as Sr, Ni, Si, DOC and Mo. Three of these
244 elements are strongly enriched in snow particles relative to the Earth crust (Sr, Sb, Mo, see section 3.3 below), thus
245 suggesting their possible leaching from atmospheric dust into the soluble fraction of snow. We could not find a
246 straightforward explication of the common group of Zn and U in soluble snow fraction (**Fig. 3 B**)

247 The effect of particulate fraction on dissolved element composition in snow is illustrated in **Fig. 4** where the
248 value of pH (**4 A**), Sr (**4 B**), Al (**4 C**) and Pb (**4 D**) in **dissolved** fraction are plotted as a function of total particle
249 concentration in snow water. The elements of 4th and 5th group (Ca, Mg, Sr, Mn, and Co) increase their concentration in
250 snow water by ca. an order of magnitude with the increase in particle concentration by 2 orders of magnitude. The
251 insoluble hydrolysates (Fe, Al, light REEs, Zr), **Cu** and Pb belonging to 1st and 2nd HCA group decrease their
252 concentration (less than a factor 10) when the particle concentration increases by 2 orders of magnitude. Other elements
253 in < 0.45 µm fraction exhibit the variations within an order of magnitude (DOC, DIC, Na, Cl, SO₄, K, Si, Cr, V, Ni, Cu,
254 Zn, As, Sb, Rb, Cd, Cs, Ba, heavy REE and U) or two orders of magnitude (Ti, Ga, Mo, W) and do not **demonstrate** any
255 significant (at p < 0.05) link with particle concentration.

256

257 **3.2. Particle concentration and TE in **particulate** fraction of snow**

258 Concentration of particulate fraction (PF) of snow and its elementary composition are available in the “Data
259 availability” section. The mineralogical composition of most representative snow samples is given in **Table S1** of the
260 Supplement. The dominant minerals are quartz (37%), albite (13%), K-feldspar (13%), phlogopite (10%), chrysotile
261 (8%), illite (7%), and chlorite (5%). The concentration of dolomite and calcite ranges from 1 to 48 and 1 to 19%,
262 respectively. Although mineral components dominated the composition of **particulate** fraction, the PF also contained
263 organic fibers, diatom frustules, pollens and particles produced during fuel burning (fly ash and black carbon). The
264 concentration of particles in snow water ranged from 0.4 to 67 mg/L. The highest values are encountered in the vicinity
265 of **the** Tomsk city (No SF 1) and around **towns** of Surgut (No SF 54, 14), Nojabrsk (SF 36, SF 38) and Gubkinsky (SF
266 33). Although the proportion of fly ash and black carbon in these samples is significant and higher than in the rest of
267 samples as **follows** from SEM observation, the mineral particles (1-25 µm size) still dominate. Note that high content of
268 fly ash and fuel burning spheres was not linked (p > 0.05) to high particulate and dissolved elements. The lowest
269 concentrations of particles (< 5-10 mg/L) were recorded north of 65°N, the region of gas industry, and between 58 and
270 61°N corresponding to the winter road along the Ob River with very low population density.

271 The enrichment coefficient ranged from ~1–5 (Ga, REEs, Fe) to > 100 (Mo, W, As, Sb, Ni, Cu, Pb, Mg, Ca, Na)
272 as illustrated in **Fig. 5 A**. The highest enrichment (EF ≥ 1000) is observed of Sb, Zn and Cd. The variation of the
273 enrichment factor as a function of latitude is shown for elements most enriched in particulate fraction in **Fig. S1 of**
274 **Supplement**. For Mg, Ca, Sr, Ba, Fe, Mn, Co, Ni, K, Rb, Cs, V, Cr, As, Cd, W the EF exhibits a maximum around 63-
275 64.5°N. This maximum coincides with the maximum of particulate fraction concentration (not shown).

276 The majority of chemical elements are present in particulate rather than dissolved form in snow meltwater
277 samples. This is illustrated by a histogram of the ratios averaged over **full** latitudinal profile (**Fig. 5 B**). Although the
278 variations of this ratio for different snow water samples across the WSL achieve ±0.5 order of magnitude, the average
279 values shown in this figure illustrate the importance of particulate deposition of Al, Fe, Ga, REEs, Cr, V, Ti, Zr, Mo and
280 W. For other elements, particulate and dissolved inputs in the form of snow are within the same order of magnitude.
281 Some soluble elements such as Na, Cd, Ca, Sr, Ba, K, As and Zn exhibit the dominance of dissolved transport in snow.

282 Although the use of average crust for assessment of element enrichment in snow particles is justified by long-
283 range transfer of snow components, it is known since the works of **group** of Reimann and de Caritat in NW Europe that

284 the “average crust” is unlikely to represent the local background and the use of the “upper crust” average value can
285 introduce a 2 to 3 order of magnitude uncertainty to any calculated EF (de Caritat et al., 1997; Reimann and de Caritat,
286 2000; Reimann et al., 2000). As such, western Siberia moss, peat and clay/loam horizons were used to assess relative
287 enrichment of elements in snow particles. It can be assumed that the leaching of soluble forms of elements from these
288 solid phases in winter is highly unlikely. The specificity of western Siberia is that the mineral (“geological”) local
289 substrate is completely frozen, even in summer, since the active (unfrozen) layer depth does not exceed the peat
290 thickness, and in that case, the use of “organic” substrates is most relevant. All three WSL reference substances (“local”
291 moss, peat and clays) represent latitudinal-averaged values based on large (> 50) number of samples collected in
292 previous studies across the 1700-km latitudinal gradient.

293 The elementary ratios of snow particles to that in mineral soil, peat and moss of the WSL are illustrated in **Fig. 6**
294 **A, B, and C**, respectively. Given significant variation on the latitude-averaged values of element concentration in snow
295 particles, mineral, peat and moss of soil column, the deviation of the ratios from unity is significant if it exceeds a factor
296 of 2 to 3. Compared to mineral soil of WSL, the snow particles are strongly ($\geq 10\times$) enriched in Sb, Zn, Ni and Cd and
297 in a lesser degree ($\geq 5\times$) in Mg, Ca, Pb, Mo, and As (**Fig. 6 A**). Note that western Siberian soils, developed on sand and
298 clay (silt) deposits (Vasil’evskaya et al., 1986), are quite poor in Ca and Mg, especially in the permafrost-bearing zone
299 north of 62°N. The enrichment of snow particles relative to peat is observed for all elements, being particularly high (>
300 50 \times) for Ni, Cr, Pb, Cu, Zn, Mg, Na and Sb (**Fig. 6 B**). Only P, Ge and Cd, exhibiting high affinity to peat (Shotyk et
301 al., 1990, 1992), are not significantly ($p > 0.05$) higher in snow particles compared to the peat column. Finally, the
302 mosses are most depleted by all elements relative to snow PF with only biogenic elements (P, K, Rb, Mn and Cd)
303 known to be concentrated in bryophytes being non-significantly higher in snow particles relative to mosses (**Fig. 6 C**).

304 The PCA of elementary composition of particulate fraction demonstrated the F1 x F2 structure (**Fig. S2 A** of the
305 Supplement). Here, two groups can be distinguished: highly mobile elements (Na, Ca, V, Ni, Mg, Mn) and low mobile
306 elements (REE, Zr, Pb, Cd, Ga, P). Note here that Pb is present as low-mobile ferric colloids in organic and Fe-rich
307 surface waters of the WSL (Pokrovsky et al., 2016b) but it is considered as volatile in the atmospheric transport,
308 especially during fuel burning (Reimann et al., 2000). For the particulate fraction, the HCA attributed the elements to 5
309 formal groups shown in **Fig. S2 B** and encircled in **Fig. S2 A**. This distinction, however, is less certain than that of the
310 dissolved fraction and does not allow establishing a clear link between the selected groups and physico-chemical
311 properties of elements or their possible sources in the snow particles. Thus, in the 1st group, among three labile elements
312 (Mg, Na and Ca) we identified V, which may exhibit elevated mobility in the form of anion in carbonate-bearing
313 mineral particles. Divalent metals (Co, Ni, Mn) and Sr constitute the 2nd labile group of elements, yet this group also
314 comprises low-mobile Fe and Cr. The 3rd group of insoluble low mobile elements is marked by the presence of
315 phosphates (REE and P), refractory Zr, and volatile Pb. The 4th group of elements revealed by HCA of particles is
316 composed of Sb, Cu and Zn. All these elements are strongly enriched in snow particles over the soil minerals (see **Fig. 6**
317 **A**). The last group of elements in snow particles comprises both labile (Li) and biologically-important Mo, K, Rb, Ba,
318 toxic volatile elements which could bear the signature of anthropogenic pollution (As, Cd) but also low mobile Ti and
319 Ga. We could not identify the link of elements in this group to the degree of snow particles enrichment relative to main
320 “local” substrates of the WSL (moss, peat and clays), shown in **Fig. 6**.

321
322
323
324
325

3.3. Possible impact of snow deposition on major and TE in lakes and rivers.

3.3.1. Snow water in comparison to lake and river water

The concentrations of dissolved major and trace **element** in snow water fraction can be compared with those in thermokarst (thaw) lakes of western Siberia measured in 2013-2014. These lakes are shallow (0.5-1.5 m depth) water bodies representing the largest reservoir of surface waters in western Siberia, north of 62°N (Polishchuk et al., 2017). The average concentration of major and TE in thermokarst lakes of various size (Manasypov et al., 2014) can be compared with those in snow water collected in this study across the same latitudinal gradient. Because the size of thermokarst lakes of WSL ranges from few m² to several km², 4 representative ranges of lake diameters are used for this comparison (0-10, 11-100, 101-500 and > 500 m). Given that the spring-time lake concentrations across the latitudinal gradient are not available, the summer-time elementary compositions of lakes were taken as most representative for the open water period of the year. The concentrations of low-soluble elements such as Fe, Al, trivalent and tetravalent hydrolysates in lakes are 1 to 2 **order** of magnitude above their concentrations in snow (not shown). At the same time, Zn, Cu, Cd, Pb, Sb and Mo exhibited snow-water concentrations that were comparable or significantly higher ($p < 0.05$) than the concentrations in lakes.

Because the main source of water in shallow lakes of WSL in spring is melted snow (Manasypov et al., 2015), we could compare the mean concentrations of snow water with spring-period lake water concentration for one particular region of discontinuous permafrost zone (town of Nojabrsk, Khanymey site) for which high-resolution seasonal observations on lakes of various size are available. For two classes of lake size (< 0.5 km² and > 0.5 km²), the following three groups of elements could be distinguished. The concentrations of dissolved Na, Mn, Zn, As, Rb and Sr in snow water are similar (within a factor of 2) to lake water concentrations. Concentrations of DIC, Cl, SO₄, Mg, Ca, Cr, Co, Ni, Cu, Mo, Cd, Sb, Cs, W, Pb and U in snow are close or higher ($p < 0.05$) than those in lakes. And finally, concentrations of DOC, Al, Si, K, Ti, V, Fe, Ga, Zr, Ba, and REEs in snow water are significantly lower than the lakes' concentrations. There was no distinction of elements belonging to individual groups of the HCA and this classification.

The concentrations of elements in snow water could be also compared with river water concentrations measured during **spring** flood 2014 across the full latitudinal profile, since such data for rivers of different **size** are available (Pokrovsky et al., 2015, 2016a). Examples of elements whose concentrations in snow water are higher or comparable with those in rivers during spring flood are illustrated in **Fig. S3**. Generally, the effect of snow melt is mostly pronounced north of 64°N. During this period, when the rivers are essentially fed by melted snow, the atmospheric deposition exhibited comparable or higher ($p < 0.05$) concentrations of SO₄, Cr, Co, Ni, Cu, Zn, Mo, Cd, Sb, Cs, W and Pb than those in rivers. These elements belonged to 5 dominant groups of HCA treatment. The concentrations of all other elements in WSL rivers cannot be explained **by solely** snow water concentration.

Note that, by filtering the snow and the lake/river water to 0.45 μm, the dissolved fraction includes a colloidal load, which can play a crucial role in the concentration of trace elements (Pokrovsky et al., 2016b). However, with typical concentration of DOC in snow water around 1-2 mg/L, the share of colloidal forms of metals will be an order of magnitude lower than that in **river** and lakes of the WSL, which have 10 to 30 mg/L of DOC.

3.3.2. Comparison of river fluxes in spring and snow water stock

Considering the mass balance calculation of snow melt influence on element fluxes in WSL rivers, the ratios of river fluxes in May-June to snow stock can be presented in the form of histograms for 3 latitudinal zones (**Fig. 7**). These ratios systematically decrease with the increase in the latitude. In the southern, permafrost-free zone, Zn, Cd, Pb, Ga, Cs, W, Sb and Cl fluxes in rivers can be provided essentially by snow melt. The riverine fluxes of DIC, Cl⁻, SO₄²⁻, Na,

367 Mg, Ca, Sr, Rb, Cs, Zn, Cu, Cr, Ni, Cu, Pb, As, Sb, Mo, W and U are strongly (i.e., $\geq 50\%$ at $p < 0.05$) affected by snow
368 melt in the discontinuous and continuous permafrost zones, north of $60\text{-}62^\circ\text{N}$.

369 According to the evolution of the ratio [river flux] / [snow stock] with the latitude, three **group** of element can be
370 distinguished: (i) elements that steadily decrease this ratio suggesting an increase in the impact of snowmelt northward:
371 DOC, SO_4 , Al, Ti, V, Cr, Rb, Sr, Cd, Sb, Cs, La, Ce, W, Pb; (ii) elements for which this ratio decreases abruptly to $62\pm 2^\circ\text{N}$
372 and then remains constant further northward: DIC, Na, Mg, Si, K, Ca, Ni, Cu, As, Mo and U; (iii) elements exhibiting
373 non-systematic variation of the ratio with latitude but having strong ($> 50\%$) impact of snowmelt on river fluxes (Cl, Co,
374 Zn, Ga) and (iv) elements having negligible ($< 10\%$) impact of snowmelt on river fluxes (Mn, Fe, Zr and Ba). **The impact**
375 **of snow melt on river export fluxes in spring strongly increases northward for DIC, Cl⁻, SO_4^{2-} , Na, Mg, Ca, Cr, Ni, Mo, Rb,**
376 **U whereas Cd, Pb, Sb, Cu, As, W and Cs present a high impact of snow melt water on river for the three latitude zones**
377 **(Fig. 7).** Although these elements belong to all 5 major groups of **cluster** analysis (**Fig. 3 B**), they can be characterized as
378 soluble (highly labile) elements, originated either from marine aerosols or from leaching from soluble minerals such as
379 carbonates, and also include volatile constituents of the atmospheric aerosols (Cd, Pb, Sb, As).

380

381

382 4. Discussion

383 4.1. Dissolved major and trace elements in Siberian snow

384 In accord with general knowledge of the Arctic aerosol chemistry (Barrie, 1986; Barrie and Barrie, 1990; Laing
385 et al., 2014, 2015; Nguyen et al., 2013; Pacyna and Ottar, 1989; Shevchenko et al., 2003; Weinbruch et al., 2012), the
386 principal component structure of snow water chemistry identified the combination of lithogenic source (dust and soil
387 particles dissolution, providing low-mobile, insoluble elements such as Al, Fe, Cr, Zr, REEs) and marine aerosols
388 (soluble forms, providing high **concentration** of mobile elements such as Ca, Mg, Na, Mo, Ni). The latter may also
389 originate from aeolian transport of carbonate-rich soils. The biogenic component **includes** Mn, Zn, K, Rb, DOC, Si
390 whereas the anthropogenic pollution originates from coal combustion (Sb, Co) and heating systems, gas flaring at the
391 gas oil production site as well as non-ferrous metal-smelter industry (Sb, Zn, Vinogradova et al., 1993) and ground
392 transportation (Pb, Cu, Zn, Cr, Ni, As, Rossini Oliva and Fernández Espinosa, 2007; Sutherland et al., 2000).

393 The soluble, highly mobile elements such as alkali and especially alkaline-earth elements, Sb, Mo, W and U
394 demonstrated an increase in their dissolved ($< 0.45\ \mu\text{m}$) concentration with the increase in the total particulate fraction
395 (**Fig. 4 B**). We interpret this increase in concentration, also correlated with $\text{pH}_{\text{snow water}}$ increase (**Fig. 4 A**), as a result of
396 element leaching from soluble minerals such as calcite and dolomite. There was a positive ($R^2 = 0.53$, $p < 0.05$)
397 correlation between % of calcite in the particulate fraction of snow and Ca concentration in snow meltwater (not
398 shown). Therefore, we hypothesize that simultaneous mobilization of carbonate minerals and soluble elements from the
399 soil and rocks to the atmosphere occurs in southern, carbonate-rock bearing provinces where the winter aerosols are
400 generated. The generation of insoluble elements such as trivalent and tetravalent hydrolysates in dissolved fraction of
401 snow occurs independently of snow enrichment in solid particles. Indeed, the decrease, and not increase in insoluble
402 elements dissolved concentration with the increase in particle concentration (**Fig. 4 C, D**) suggests that these elements
403 are not desorbed or leached from mineral particles, either within the origin of aerosol formation or during snow melting
404 and filtration in the laboratory. **Alternatively, these elements can be linked to specific labile pools that constitute the**
405 **mineral fraction. Unfortunately, we could not run selective extractions on very small amount of solid particles in WSL**
406 **snow available in this study.**

407 Regional background concentrations of dissolved metals in snow of Quebec, **Canada** are reported to be 1.1, 1.7,
408 and $1.6\ \text{mg/L}_{\text{meltwater}}$ for Cu, Pb, and Zn, respectively (Telmer et al., 2004). The values for Cu and Pb are comparable

409 with average snow water concentration across the WSL (0.83 and 0.68, respectively) but the concentration of Zn in the
410 WSL snow is significantly higher (10.1 ± 5.0 $\mu\text{g/L}$, excluding 3 contaminated samples near the Tomsk city). Background
411 concentrations of dissolved Cu, Pb, and Zn in snow of Alaskan Arctic are much lower (0.08, 0.09 and 1.2, respectively,
412 Snyder-Conn et al., 1997). In snow from background areas of north-eastern European Russia, the concentrations of
413 dissolved Cu are near at the same level as in snow from the WSL, whereas the concentrations of dissolved Pb and Zn
414 are 2 times lower (Walker et al., 2003). Concentrations of dissolved Cu and Zn in snow of NW Finland are few times
415 lower than in snow of WSL; concentrations of dissolved Pb are at the same level (Caritat et al., 1998).

416 Significant enrichment in Ni is known for the aerosols of the Arctic Ocean (Shevchenko et al., 2003). It may be
417 linked both to Ni transport from Norilsk and Kola smelters but also with Ni fractionation at the sea surface (Duce et al.,
418 1976). Ni concentration in snow water of the northern part of WSL significantly exceeds that in the thermokarst lakes.
419 The winter snow stock of dissolved Ni is several times higher than the river export of this element during spring flood
420 in the permafrost-bearing zone of the WSL, north of 60°N , and Ni concentration in snow particles exceeds up to 2
421 orders of magnitude its concentration in moss and peat of the territory.

422 The winter-time deposition of dissolved (< 0.45 μm) metals on the surface of northern part of the WSL can be
423 calculated taking into account the mean multi-annual volume of accumulated snow during 8 winter months (in mm of
424 snow water) and the average concentration of elements in February snow collected north of 64°N . The monthly
425 depositions of selected metals ($\mu\text{g m}^{-2} \text{ month}^{-1}$) on the north of the WSL in the form of snow are equal to 2.8, 12, 15,
426 210 and 0.9 for As, Ni, Pb, Zn, and Cd which is significantly higher than the values for winter deposition of insoluble
427 aerosols into the Russian Arctic (0.22, 0.74, 2.7, 1.3 and 0.056, respectively, Shevchenko et al., 2003). Only V exhibited
428 similar values of Arctic aerosol and snow deposition (0.71 and 0.96 $\mu\text{g m}^{-2} \text{ month}$, respectively).

429
430

431 **4.2. Particulate forms of elements in the snow across the latitudinal profile: the effect of mineralogical** 432 **substrate, industrial centers, local pollution and long-range transport.**

433 The majority of elements are transported in particulate rather than dissolved fraction in the snow water (Fig. 5
434 B). This is in general agreement with the results of other studies in Scandinavia and Kola Peninsula (Reimann et al.,
435 1996), north-eastern European Russia (Walker et al., 2003) and on drifting ice in the northern Barents Sea (Gordeev and
436 Lisitzin, 2005). The enrichment of snow particulate fraction relative to the earth crust as shown by Al-normalized
437 enrichment coefficient (Figs. 5 A, S1) can be understood via taking into account the particle concentration in snow and
438 microscopic observations. We suggest that the clays supply most trace elements in the PF. The atmospheric particles are
439 known to exert significant impact on soils and ground vegetation (Kabata-Pendias and Pendias, 1984; Rasmussen, 1998;
440 Steinnes and Friedland, 2006). In the case of WSL, the elementary composition of snow particulate fraction was
441 compared with three main reservoirs of elements within the soil, sampled over significant latitudinal profile, from 55°N
442 to 68°N (Stepanova et al., 2015). These reservoirs are averaged over full latitudinal range and include *i*) mineral fraction
443 from the bottom of the peat column; *ii*) depth-averaged peat column composition, and *iii*) *Sphagnum* mosses, collected
444 in ombrotrophic bogs, which receive their constituents essentially from the atmosphere (e.g., Santelman and Gorham,
445 1988).

446 The particularity of the northern part of western Siberia lowland is that the active (seasonally unfrozen) soil layer
447 is located within the organic (moss+peat) rather than mineral horizon; the latter is represented by poorly reactive sands
448 and clays (Baulin et al., 1967; Baulin, 1985; Tyrtikov, 1973, 1979). As a result, the surface waters drain essentially
449 organic part of the column which is very poor in lithogenic elements (Pokrovsky et al., 2015, 2016a). The supply of
450 mineral particles from the snow therefore may significantly enrich the rivers and lakes in dissolved alkaline earths,

451 metal micronutrients, phosphorus and other elements given high reactivity of incoming silicate and carbonate grains in
452 acidic (pH < 3-4), organic-rich (10 < DOC < 50 mg/L) surface waters of Western Siberia. The degree to which such a
453 supply can lead to overestimation of the calculated chemical weathering export fluxes of cations in the permafrost zone
454 is **not possible** to quantify. Therefore, in view of the importance of atmospheric input of solid particles for mineral-poor,
455 peat bogs of western Siberia, the seasonal, year-round measurements of particulate atmospheric deposition in this region
456 are necessary.

457 The main source of mineral particles in the southern part of **latitudinal** profile (56–58°N) may be soils of steppe
458 and forest-steppe regions south of WSL, where the land is cultivated and the snow cover is relatively thin. The aeolian
459 transport of soil particles under these conditions may be efficient even in winter (Evseeva et al., 2003). The main source
460 of ash particles in **southern** part of the profile is the industry and transport of the city of Tomsk (Boyarkina et al., 1993;
461 Yazikov et al., 2000; Talovskaya et al., 2014). The concentration of particles in snow collected from 58°N to 61°N
462 ranged between 0.85 and 5.7 mg/L which is comparable or slightly higher than the values reported for the Arctic snow
463 cover (Darby et al., 1974; Mullen et al., 1972; Nürnberg et al., 1994; Shevchenko et al., 2002, 2010). It is **important** that
464 in this zone of low PF concentration, combustion spheres, fly ash and black carbon of few µm diameters were
465 dominating. This can explain **relatively** low concentration of all TE at low PF concentration, as carbon compounds
466 likely contain very low **proportion** of trace metals. The most important sources of fly ash and black carbon are gas
467 flaring, land transport, heating plants, residential combustion, forest fires (mainly in summer) and industrial plants
468 (Moskovchenko and Babushkin, 2012; Quinn et al., 2008; Stohl et al., 2013). Chemical pollution of **atmosphere** during
469 gas flaring associated with oil industry is known for the WSL (Raputa, 2013; Yashchenko et al., 2014). The black
470 carbon produced during gas burning is detected not only in western Siberia but in the Russian sector of the Arctic Ocean
471 **in** high latitudes (Stohl et al., 2013; **Evangelidou et al., 2017**).

472 In the zone 62–64.5°N, where some impact of oil industry is possible, the **concentration** of insoluble particles in
473 snow were above 10 mg/L, achieving the value of 66.6 mg/L in sample SF36. Backward trajectories to this site using
474 **Draxler** and Rolf (2003) approach show that, during **last** few days before sampling, the air masses arrived from south-
475 western direction. Accordingly, the particulate fraction in these samples contained mostly mineral particles 1–25 µm
476 size with some fly ash (burning spheres). It is possible that mineral particles are supplied here via long-range transport
477 from forest-steppe, steppe and semi-desert regions south and south-west from the study site. Indeed, during **winter** snow
478 coverage period, the dominant winds in this zone have S, SW and W directions (Moskovchenko and Babushkin, 2012).
479 The events of mineral dust transport over large distances are well known in the boreal zone (Lisitzin, 1978, 2011;
480 Shevchenko et al., 2010).

481 Further north of **studied** latitudinal profile, from 65 to 68°N, the concentration of snow particles ranged from 0.8
482 to 9.2 mg/L. These values are within the background in the Arctic and subarctic (Darby et al., 1974; Mullen et al., 1972;
483 Nürnberg et al., 1994; Shevchenko et al., 2002). The particulate fraction was represented by mineral debris of 1 to 15
484 µm in size, with frequent but not significant presence of spherical ash particles, biogenic strains and porous carbon
485 particles. Because the main source of mineral particles is long-range transport from southern desert and steppe regions,
486 moving to the north **decrease** the influence of these provinces.

487 We believe that the elevated concentrations of divalent metals, As and Sb in snow particles (**Fig. 5 A, S1**) should
488 not be interpreted as necessarily pollution from the industrial centers. Rather, volatile Cd, Pb, As may originate from
489 long-range transport of desert material. Therefore, we attempted to distinguish the well-known refractory, non-volatile
490 heavy metals such as Cu, Ni and Co and more volatile elements such as Pb, Cd and As (i.e., Reimann et al., 2000) based
491 on the HCA treatment. For both particulate and dissolved **fraction**, these elements are located in three or two different

492 groups but never belong to one single group of inter-correlated elements. As such the available data do not evidence
493 similar **origin** of Cu, Ni and Co, or Pb, Cd, and As in the snapshot of WSL snow sampled in this work.

494

495 **4.3. Impact of snow on hydrochemistry of inland waters and riverine elementary fluxes.**

496 Quantitative comparison of element input to the land surface with winter snow and element **concentration** and
497 fluxes in the WSL inland waters provided the assessment of minimal atmospheric contribution to lake storage and river
498 export. The concentrations of Pb, Zn, Cu, Cd, Sb and Mo in lakes are significantly lower than those in snow. However,
499 these elements belong to 4 various **group** of elements in dissolved snow fraction, identified by the HCA (**Fig. 3 B**). In
500 rivers, SO₄, Cr, Co, Ni, Cu, Zn, Mo, Cd, Sb, Cs, W and Pb are dominated by snow input. These elements also belong to
501 4 various **group** of the HCA. It thus can be concluded that there is no direct link between the **group** of elements
502 identified by **cluster** dendrogram in the snow water and the elements whose **concentration** in rivers or lakes are
503 significantly affected by snow deposition. We believe that a natural cause of this apparent inconsistency **is** different
504 mechanisms controlling the element distribution **in the aerosols and surface waters. Whereas the aerosols are influenced**
505 **by local sources of pollution, remote desert provinces, and leaching of soluble elements from mineral particles, the**
506 **inland waters chemical composition is controlled by** interaction of melted snow with upper peat and moss/lichen
507 horizons; underground feeding, release of elements from silicate river suspended matter due to abrasion in spring flood.

508 Overall, the impact of the snowmelt on chemical composition of western Siberian thermokarst lakes may be very
509 high. This will be further accentuated by **reported** increase in the proportion of meltwater that does not reach the main
510 rivers but is stored by the wetlands (i.e., from 20-30% in early 1990s to 50-60% in the mid-2000s, Zakharova et al.,
511 2011). A comparison of snow stock/river water fluxes demonstrates **that the influence of atmospheric deposition**
512 **increases northward** (**Fig. 7**). At the same time, the chemical composition of the snow water, although **subjected** to
513 significant variation, does not exhibit any systematic trend with the latitude (**Fig. 2**) as also follows from the PCA
514 (section 3.1). The reason for this difference may be **relatively** low fluxes and concentrations in rivers of the northern,
515 permafrost-affected territory of the WSL compared to the southern, permafrost-free zone (Pokrovsky et al., 2015,
516 2016a). As a result, the impact of atmospheric deposition on the riverine transport is more pronounced in the permafrost
517 zone than in the permafrost-free zone. We expect **that the contribution of atmospheric deposition into river water fluxes**
518 **should be quite strong for all flat bog tundra areas of northern Eurasia, including, in addition to northern part of western**
519 **Siberia (~400,000 km²) studied in this work, the Yamal and Gyda Peninsula (122,000 and 160,000 km², respectively),**
520 **the North-Siberian Lowland (~700,000 km²), the Kolyma Lowland (170,000 km²), and the Yana-Indigirka Lowland**
521 **(180,000km²) with an overall territory close to 1.7 million km². The impact of snow deposition on river elementary**
522 **fluxes should be much lower in permafrost-bearing mountainous terrain such as Central and Eastern Siberia, the**
523 **Alaskan slopes, north of Scandinavian shield and the Canadian High Arctic. In those territories, two factors may**
524 **decrease the contribution of snow deposition to river fluxes: 1) the impact of local mineral dust for aerosols generation**
525 **may be well pronounced and 2) the chemical weathering occurs within the mineral seasonally unfrozen layer producing**
526 **higher fluxes of inorganic components.**

527 In contrast, in the lowlands of Northern Eurasia, the rivers drain essentially organic layer (peat bog) terrain, thus
528 mineral feeding of rivers is really low. As it is demonstrated in section 3.3.2 of this study, low chemical (cationic)
529 weathering in the north of the WSL during spring suggests that total dissolved cationic and DIC fluxes in May-June in
530 this and other similar regions are essentially controlled by snowmelt, rather than by **soil** weathering. It follows that
531 during the spring period, the intensity of chemical weathering in these latitudes can be a factor of 2 (major cations) to 5
532 (TE) lower than that deduced from riverine fluxes. However, given that the shares of spring flood period (May-June) in
533 the annual export fluxes are only 5 to 10% for major cations and 10 to 20% for TE (Pokrovsky et al., 2015, 2016), the

534 overall impact of atmospheric deposits on element export fluxes will be strongly pronounced (i.e., $\geq 50\%$ of total
535 measured river flux value) only for elements which have the ratio of the spring-time river export to snow stock less than
536 0.2, i.e., SO₄, Cu, Mo, Cd, Sb, Cs, W and Pb. With further increase of winter precipitation in western Siberia (i.e.,
537 Bulygina et al., 2009), the impact of snowmelt on element transport to the Arctic Ocean by rivers may increase thus
538 enriching the surface waters in many elements such as Cd, Pb, Sb, Cr, Cu, Ni, As, Mo, Rb, U.

539
540

541 **Conclusions**

542 The chemical composition of surface layer of snow cover was studied across a 1700-km latitudinal gradient in
543 western Siberia Lowland. The particulate fraction ranged from 0.4 to 66 mg/L_{meltwater} and increased in the regions of
544 enhanced dust deposition from southern steppe and desert provenances, in the proximity of industrial centers and due to
545 fly ash production from gas burning of the oil exploration sites. There was an increase in concentration of soluble
546 elements (Ca, Mg, Sr, Mn, Co) and pH in snow water with the increase in mineral (calcite, dolomite) fraction. The
547 elementary composition of PF demonstrated its significant enrichment in most elements relative to mineral soil horizon,
548 peat and moss across **full** latitudinal profile (~1700 km) of WSL. As such, solid atmospheric aerosols may **be** important
549 factor of insoluble element delivery to the soil surface. The supply of mineral particles from the snow may also
550 significantly enrich the rivers and lakes in dissolved alkaline earths, metal micronutrients, phosphorus and other elements
551 given high reactivity of incoming silicate and carbonate grains in acidic (pH < 3-4), organic-rich (10 < DOC < 50 mg/L)
552 surface waters of Western Siberia.

553 Concentrations of Na, Mn, Zn, As, Rb and Sr in winter aerosols are similar (within a factor of 2) to lake water
554 concentrations during spring **period**. Concentrations of DIC, Cl, SO₄, Mg, Ca, Cr, Co, Ni, Cu, Mo, Cd, Sb, Cs, Pb and U
555 in filtered snow water are **close** or higher than those in lakes. In the **southern**, permafrost-free zone, only Zn, Cd, W, Pb,
556 Cs and Sb fluxes in rivers during **May**-June period can be provided by **dissolved** fraction of the snow melt. However,
557 the impact of snow melt on river export fluxes in spring strongly increases northward for DIC, Cl, SO₄, Na, Mg, Ca, Cd,
558 Pb, Sb, Cr, Cu, Ni, As, Mo, Rb, U. In the permafrost zone, $\geq 50\%$ of riverine fluxes of these elements during spring
559 **flood** can be provided by the snowmelt. The reason for **such** high sensitivity of WSL surface reservoirs to atmospheric
560 deposition **is** feeding of surface waters by essentially organic (moss, peat) soil profiles.

561
562

562 **Data availability**

563 **Full** data set of major and trace element **concentration** in snow water (< 0.45 μm) and snow particles sampled
564 across the latitudinal profile of Western Siberia Lowland is available at **the** Research Gate,
565 <https://www.researchgate.net/publication/309666956>; DOI: 10.13140/RG.2.2.12156.54408.

566

567 **Acknowledgements:**

568 This work was supported **from** the BIO-GEO-CLIM grant No 14.B25.31.0001 **of** Russian Ministry of Science and
569 Education. RM and LS acknowledge support from RSCF (RNF) grant No 15-17-10009. Supports from GDRI CAR-WET-
570 SIB, RFBR Nos 17-55-16008, 17-05-00348_a, FCP “Kolmogorov” RFMEFI58717X0036, and Program 32 of
571 Fundamental Research of **Presidium** of Russian Academy of Sciences are also acknowledged. We would like to thank
572 Academician A.P. Lisitzin for valuable recommendations, J. Prunier, M. Henry, F. Candaudap, A. Lanzaova for help in
573 analytical work. The authors acknowledge the NOAA Air Resources Laboratory (ARL) for the provision of the HYSPLIT
574 transport model and READY website (<http://www.arl.noaa.gov/ready.html>).

- 576 Ariés, S., Valladon, M., Polvé, M., and Dupré, B.: A routine method for oxide and hydroxide interference
 577 corrections in ICP-MS chemical analysis of environmental and geological samples, *Geostandards Newslett.*, 24, 19–31,
 578 doi: 10.1111/j.1751-908X.2000.tb00583.x, 2000.
- 579 Baltrėnaitė, E., Baltrėnas, P., Lietuvninkas, A., Šerevičienė, V., and Zuokaitė, E.: Integrated evaluation of aerogenic
 580 pollution by air-transported heavy metals (Pb, Cd, Ni, Zn, Mn and Cu) in the analysis of the main deposit media, *Environ.*
 581 *Sci. Pollut. Res.*, 21, 299–313, 2014.
- 582 Barrie, L. A.: Arctic air pollution: an overview of current knowledge. *Atmospheric Environment*, 20, 643–663,
 583 1986.
- 584 Barrie, L. A., Barrie, M. J.: Chemical components of lower tropospheric aerosols in the high Arctic: Six year of
 585 observation, *J. Atmos. Chem.*, 11, 211–226, 1990.
- 586 Baulin, V. V.: Permanently frozen rocks of oil and gas producing regions of the USSR. Nauka: Moscow (in
 587 Russian), 1985.
- 588 Baulin, V. V., Belopukhova, E. B., Dubikov, G. I. and Shmelev, L. M.: Geocryological conditions of western
 589 Siberia Lowland, Nauka, Moscow (in Russian), 1967.
- 590 Beilman, D. W., MacDonald, G. M., Smith, L. C., and Reimer, P. J.: Carbon accumulation in peatlands of West
 591 Siberia over the last 2000 years, *Global Biogeochem. Cycles*, 23, GB1012, doi:10.1029/2007GB003112, 2009.
- 592 Bini, C., Sartori, G., Wahsha, M., Fontana, S.: Background levels of trace elements and soil geochemistry at
 593 regional level in NE Italy. *J. Geochemical Explor.* 109, 125–133. <http://dx.doi.org/10.1016/j.gexplo.2010.07.008>, 2011.
- 594 Bokhorst, S., Pedersen, S. H., Brucker, L., Anisimov, O., Bjerke, J. W., Brown, R. D., Ehrlich, D., Essery, R. L. H.,
 595 Heilig, A., Ingvander, S., Johansson, C., Johansson, M., Jónsdóttir, I. S., Inga, N., Luoju, K., Macelloni, G., Mariash, H.,
 596 McLennan, D., Rosqvist, G. N., Sato, A., Savela, H., Schneebeli, M., Sokolov, A., Sokratov, S. A., Terzago, S., Vikhamar-
 597 Schuler, D., Williamson, S., Qiu, Y., and Callaghan, T. V.: Changing Arctic snow cover: A review of recent developments
 598 and assessment of future needs for observations, modelling, and impacts, *Ambio*, 45, 516–537, doi: 10.1007/s13280-016-
 599 0770-0, 2016.
- 600 Botch, M. S., Kobak, K. I., Vinson, T. S., and Kolchugina, T. P.: Carbon pools and accumulation in peatlands of the
 601 former Soviet Union, *Global Biogeochem. Cy.*, 9, 37–46, doi: 10.1029/94GB03156, 1995.
- 602 Boyarkina, A. P., Baikovsky, V. V., Vasiliev, N. V., Glukhov, G. G., Medvedev, M. A., Pisareva, L. F., Rezhnikov,
 603 V. I., and Shelud'ko, S. I.: Aerosols in natural tablet of Siberia, Publishing house of Tomsk State University, 157 pp. (in
 604 Russian), 1993.
- 605 Brown, J., Ferrians, O. J. J., Heginbottom, J. A. & Melnikov, E. S.: International Permafrost Association circum-
 606 Arctic map of permafrost and ground ice conditions, scale 1:10,000,000, U. S. Geol. Surv., Washington, D. C., 1997.
- 607 Bulygina, O. N., Razuvaev, V. N., and Korshunova, N. N.: Changes in snow cover over Northern Eurasia in the last
 608 few decades, *Environ. Res. Lett.*, 4, 045026, doi: 10.1088/1748-9326/4/4/045026, 2009.
- 609 Callaghan, T. V., Johansson, M., Brown, R. D., Groisman, P. Ya., Labba, N., Radionov, V., Bradley, R. S., Blangy,
 610 S., Bulygina, O. N., Christensen, T. R., Colman, J. E., Essery, R. L. H., Forbes, B. C., Forchhammer, M. C., Golubev, V.
 611 N., Honrath, R. E., Juday, G. P., Meshcherskaya, A. V., Phoenix, G. K., Pomeroy, J., Rautio, A., Robinson, D. A.,
 612 Schmidt, N. M., Serreze, M. C., Shevchenko, V. P., Shiklomanov, A. I., Shmakin, A. B., Skold, P., Sturm, M., Woo, M.,
 613 Wood, E. F.: Multiple effects of changes in Arctic snow cover, *Ambio*, 40 (S1), 32–45, 2011.
- 614 Caritat, P. de, Reimann, C., Chekushin, C., Bogatyrev, I., Niskavaara, H., Braun, J.: Mass balance between
 615 emission and deposition of airborne contaminants. *Environ. Sci. Technol.* 31, 2966–2972, 1997.
- 616 Caritat, P. de, Åyräs, M., Niskavaara, H., Chekushin, V., Bogatyrev, I., and Reimann, C.: Snow composition in eight
 617 catchments in the central Barents Euro-Arctic region, *Atmos. Environ.*, 32, 2609–2626, 1998.
- 618 Caritat, P. de, Hall, G., Gislason, S., Belsey, W., Braun, M., Goloubeva, N. I., Olsen, H. K., Scheie, J. O., and Vaive,
 619 J. E., 2005. Chemical composition of arctic snow: concentration level and regional distribution of major elements, *Sci.*
 620 *Total Environ.*, 336, 183–199.
- 621 Chekushin, V. A., Bogatyrev, I. V., Caritat, P. de, Niskavaara, H., and Reimann, C., 1998. Annual atmospheric
 622 deposition of 16 elements in eight catchments of the central Barents region, *Sci. Total Environ.*, 220, 95–114, 1998.
- 623 Chessel, D., Dufour, A. B., and Thioulouse, J.: The ade4 package - I : One-table methods. *R News* Vol. 4/1, 5–10,
 624 2004.
- 625 De la Cruz, O., and Olmes, S.: The duality diagram in data analysis: Examples of modern applications. *Annals*
 626 *Applied Stat.*, 5(4), 2266–2277, DOI:10.1214/10-AOAS408, 2011.
- 627 Darby, D. A., Burckle, L. H., and Clark, D. L.: Airborne dust on the Arctic pack ice: Its composition and fallout rate,
 628 *Earth Planet Sci. Lett.*, 24, 166–172, 1974.
- 629 Draxler, R. R., and Rolf, G. D.: HYSPLIT (Hybrid Single-Particle Lagrangian Integrated Trajectory). Silver Spring
 630 (MD): NOAA Air Resources Lab. Mod access via NOAA ARL READY Website
 631 (<http://www.arl.noaa.gov/ready/hysplit4.html>), 2003.
- 632 Duce, R. A., Hoffman, G. L., Ray, B. J., Fletcher, I. S., Wallace, G. T., Fasching, J. L., Piotrowicz, S. R., Walsh, P.
 633 R., Hoffman, E. J., Miller, J. M., and Heffter, J. L.: Trace metals in the marine atmosphere: sources and fluxes, In:
 634 Windom, H. L., Duce, R. A. (Eds.), *Marine Pollutant Transfer*. Lexington Books, Toronto, pp. 77–119, 1976.
- 635 Ermolov, Yu. V., Makhatkov, I. D., and Khudyaev, S. A.: Background concentration of chemical elements in snow
 636 cover of the typical regions of the Western Siberia. *Optika Atmosfery i Okeana* 27, 790–800 (in Russian), 2014.

637 Evangelidou, N., Shevchenko, V. P., Yttri, K.-E., Eckhardt, S., Sollum, E., Pokrovsky, O. S., Kobelev, V. O.,
638 Korobov, V. B., Lobanov, A. A., Starodymova, D. P., Vorobiev, S. N., Thompson, R. L., Stohl, A.: Origin of elemental
639 carbon in snow from Western Siberia and Northwestern European Russia during spring 2014, 2015 and 2016, *Atmospheric
640 Chemistry and Physics*, submitted, 2017.

641 Evseeva, N. C., Kvasnikova, Z. N., Romashova, T. N., and Osintzeva N. V.: Aeolian erosion of soil during cold
642 period of the year on Tom-Yaiskoe watershed divide (Western Siberia), *Geography and Natural Resources*, No 3, 101-105,
643 2003.

644 Franzén, L. G., Hjelmroos, M., Källberg, P., Brorström-Lundén, E., Junnto, S., Savolainen, A.-L.: The “yellow
645 snow” episode of northern Fennoscandia, march 1991 – a case study of long-distance transport of soil, pollen and stable
646 organic compounds, *Atm Environ.*, 28, 3587–3604, 1994.

647 Frey, K. E. and Smith, L. C.: How well do we know northern land cover? Comparison of four global vegetation and
648 wetland products with a new ground-truth database for West Siberia, *Global Biogeochem. Cy.*, 21, GB1016,
649 doi:10.1029/2006GB002706, 2007.

650 Garbarino, J. R., Snyder-Conn, E., Leiker, T. J., and Hoffman, G. L.: Contaminants in arctic snow collected over
651 northwest Alaskan sea ice, *Water, Air, Soil Poll.*, 139, 183–214, 2002.

652 Gordeev, V. V., and Lisitzin, A. P.: Heavy metals in snow and ice cover of the Barents Sea, *Oceanology* 45 (5),
653 737–744, 2005.

654 Gourdol, L., Hissler, C., Hoffmann, L., Pfister, L., 2013. On the potential for the Partial Triadic Analysis to grasp
655 the spatio-temporal variability of groundwater hydrochemistry. *Appl. Geochem.* 39, 93–107.

656 Guéguen, C., Cuss, C. W., and Cho, S.: Snowpack deposition of trace elements in the Athabasca oil sands region,
657 Canada, *Chemosphere*, 153, 447–454, doi: 10.1016/j.chemosphere.2016.03.020, 2016.

658 Hartigan, J.: *Clustering Algorithms*. John Wiley and Sons, NY, 1975.

659 Huang, Z., Huang, J., Hayasaka, T., Shanshan Wang, S., Zhou, T., and Jin, H.: Short-cut transport path for Asian
660 dust directly to the Arctic: a case study, *Environ. Res. Lett.*, 10, 114018, 2015.

661 Kabata-Pendias, A., and Pendias, H., 1984. *Trace Elements in Soils and Plants*. CRC Press, Boca Raton, Florida,
662 304 pp, 1984.

663 Kang, S., Zhang, Q., Kaspari, S., Qin, D., Cong, Z., Ren, J., and Mayewski, P. A.: Spatial and seasonal variations of
664 elemental composition in Mt. Everest (Qomolangma) snow/firn, *Atmos. Environ.*, 41, 7208–7218, 2007.

665 Karnatzevich, I. V., and Khruschev, S. A.: A computer system of mass calculation of running water balances of
666 river watersheds from poorly studied territories, Omsk, Gos Ped Institut, 174 pp, 2014.

667 Kashulina, G., de Caritat, P., and Reimann, C.: Snow and rain chemistry around the “Severonikel” industrial
668 complex, NW Russia: Current status and retrospective analysis, *Atmos. Environ.*, 89, 672–682, 2014.

669 Kaufman, L., Rousseeuw, P. J.: *Finding Groups in Data*. John Wiley and Sons Inc., NY, (368pp), 2005.

670 Khrenov V. Ya.: *Soils of cryolithozone of western Siberia: Morphology, physico-chemical properties and
671 geochemistry*, Nauka, Moscow (in Russian), 2011.

672 Krachler, M., Zheng, J., Koerner, R., Zdanowicz, C., Fisher, D., and Shotyk, W.: Increasing atmospheric antimony
673 contamination in the northern hemisphere: snow and ice evidence from Devon Island, Arctic Canada. *J. Environ. Monit.*, 7,
674 1169–1176, 2005.

675 Laing, J. R., Hopke, P. K., Hopke, E. F., Husain, L., Durkiewicz, V. A., Paatero, J., and Viisanen, Y.: Long-term
676 particle measurements in Finnish Arctic: Part I – Chemical composition and trace metal solubility, *Atmos. Environ.* 88,
677 275–284, 2014.

678 Laing, J. R., Hopke, P. K., Hopke, E. F., Husain, L., Durkiewicz, V. A., Paatero, J., and Viisanen, Y.: Positive
679 matrix factorization of 47 years of particle measurements in Finnish Arctic, *Aerosol Air Quality Res.*, 15, 188–207, 2015.

680 Levitan, D. M., Zipper, C. E., Donovan, P., Schreiber, M. E., Seal, R. R., Engle, M., Chermak, J., Bodnar, R. J.,
681 Johnson, D. K., Aylor, J. G.: Statistical analysis of soil geochemical data to identify pathfinders associated with mineral
682 deposits: an example from the Coles Hill uranium deposit, Virginia, USA. *J. Geochemical Explor.* 154, 238–251, 2015.

683 Lin, X., Wang, X., Zhang, B., Yao, W.: Multivariate analysis of regolith sediment geochemical data from the
684 Jinwozi gold field, north-western China, *J. Geochem. Explor.* 137, 48–54, 2014.

685 Lisitzin, A. P.: Arid sedimentation in the oceans and atmospheric particulate matter, *Dispersed Russian Geology
686 Geophys.*, 52(10), 1100–1133, doi: 10.1016/j.rgg.2011.09.006, 2011.

687 Lisitzin, A. P.: *Processes of Ocean Sedimentation*. Lithology and Geochemistry, Moscow, Nauka, 391 pp, 1978.

688 Lisitzin, A. P.: *Sea-ice and Iceberg Sedimentation in the Ocean: Recent and Past*, Springer, Berlin. 563 pp, doi:
689 10.1007/978-3-642-55905-1, 2002.

690 Lisitzin, A. P., Lukashin, V. N., and Dara, O. M.: Composition and fluxes of minerals in suspended particulate
691 matter from the water column of the Caspian Sea, *Doklady Earth Sci.*, 463(2), 733-737, doi:
692 10.1134/S1028334X15070120, 2015.

693 Manasypov, R. M., Pokrovsky, O. S., Kirpotin, S. N., and Shirokova, L. S.: Thermokarst lake waters across the
694 permafrost zones of western Siberia, *The Cryosphere* 8, 1177–1193, 2014.

695 Manasypov, R. M., Vorobyev, S. N., Loiko, S. V., Kritzkov, I. V., Shirokova, L. S., Shevchenko, V. P., Kirpotin, S.
696 N., Kulizhsky, S. P., Kolesnichenko, L. G., Zemtsov, V. A., Sinkinov, V. V., and Pokrovsky, O. S.: Seasonal dynamics of
697 organic carbon and metals in thermokarst lakes from the discontinuous permafrost zone of western Siberia, *Biogeosciences*
698 12, 3009–3028, 2015.

699 Moragues-Quiroga, C., Juilleret, J., Gourdol, L., Pelt, E., Perrone, T., Aubert, A., Morvan, G., Chabaux, F., Legout,
700 A., Stille, P., and Hissler, C.: Genesis and evolution of regoliths: Evidence from trace and major elements and Sr-Nd-Pb-U
701 isotopes, *Catena* 149, 185–198, 2017.

702 Moskovchenko, D. V., and Babushkin, A. G.: Peculiarities of formation of chemical composition of snow waters (on
703 example of Khanty-Mansi autonomous district), *Earth Cryosphere*, XVI, 71–81, 2012.

704 Mullen, R. E., Darby, D. A., and Clark, D. L.: Significance of atmospheric dust and ice rafting for Arctic sediment,
705 *Geol. Soc. America Bull.* 83, 205–212, 1972.

706 Nikitin, S. P. and Zemtsov, V. A.: The variability of hydrological parameters of western Siberia, Nauka,
707 Novosibirsk, 204 pp., 1986 (in Russian).

708 Niu, H., He, Y., Kang, S., Lu, X., Shi, X., and Wang, S.: Chemical composition of snow from Mt. Yulong,
709 southeastern Tibetan Plateau, *J. Earth Syst. Sci.*, 125(2), 403–416, 2016.

710 Niu, H., Kang, S., Shi X., He Y., Lu X. et al.: Water-soluble elements in snow and ice on Mt. Yulong, *Sci. Total*
711 *Environ.*, 574, 889–900, 2017.

712 Nguyen, Q. T., Skov, H., Sørensen, L. L., Jensen, B. J., Grube, A. G., Massling, A., Glasius, M., and Nøjgaard, J.
713 K.: Source appointment of particles at Station Nord, North East Greenland during 2008–2010 using COPREM and PMF
714 analysis, *Atmos. Chem. Phys.* 13, 35–49, 2013.

715 Novikov, S. M., Moskvina, Y. P., Trofimov, S. A., Usova, L. I., Batuev, V. I., Tumanovskaya, S. M., Smirnova, V.
716 P., Markov, M. L., Korotkevich, A. E., and Potapova, T. M.: Hydrology of bog territories of the permafrost zone of
717 western Siberia, *BBM publ. House, St. Petersburg*, 535 pp., 2009 (in Russian).

718 Nürnberg, D., Wollenburg, I., Dethleff, D., Eicken, H., Kassens, H., Letzig, T., Reimnitz, E., and Thiede, J.:
719 Sediments in Arctic sea ice: Implications for entrainment, transport and release, *Marine Geol.*, 119, 185–214, 1994.

720 Pacyna, J. M., and Ottar, B.: Origin of natural constituents in the Arctic aerosols, *Atmos. Environ.* 23 (4), 809–815,
721 1989.

722 Pokrovsky, O. S., Manasypov, R. M., Shirokova, L. S., Loiko, S., Krickov, I., Kopysov, S., Kolesnichenko, L. G.,
723 Zemtsov, V. A., Kulizhsky, S. P., Vorobyev, S. N., and Kirpotin, S.N.: Permafrost coverage, watershed area and season
724 control of dissolved carbon and major elements in western Siberia rivers. *Biogeosciences*, 12, 6301–6320, 2015.

725 Pokrovsky, O. S., Manasypov, R.vM., Loiko, S., Krickov, I. A., Kopysov, S. G., Kolesnichenko, L. G., Vorobyev,
726 S. N., and Kirpotin, S.N.: Trace elements transport in western Siberia rivers across a permafrost gradient, *Biogeosciences*,
727 13, 1877–1900, 2016a.

728 Pokrovsky, O. S., Manasypov, R. M., Loiko, S. V., Shirokova, L. S.: Organic and organo-mineral colloids of
729 discontinuous permafrost zone, *Geochim. Cosmochim. Acta*, 188, 1-20, 2016b.

730 Polishchuk Y. M., Bogdanov, A. N., Polishchuk, V. Y., Manasypov, R. M., Shirokova, L. S., Kirpotin, S. N.,
731 Pokrovsky, O. S.: Size distribution, surface coverage, water, carbon, and metal storage of thermokarst lakes in the
732 permafrost zone of the Western Siberia Lowland, *Water*, 9, 228; doi:10.3390/w9030228, 2017.

733 Quinn, P. K., Bates, T. S., Baum, E., Doubleday, N., Fiore, A. M., Flanner, M., Fridlind, A., Garrett, T. J., Koch, D.,
734 Menon, S., Shindell, D., Stohl, A., and Warren, S. G.: Short-lived pollutants in the Arctic: their climate impact and possible
735 mitigation strategy, *Atmos. Chem. Phys.*, 8, 1723–1735, 2008.

736 Raputa, V. F.: Experimental and modeling studies of aerosol deposition in the vicinity of gas torch burning, *Vestnik*
737 *NGY, Ser. Mathematics, mechanics and Informatics*, 13(3), 96-102, 2013 (in Russian).

738 Rasmussen, P. E.: Long-range atmospheric transport of trace metals: the need for geoscience perspectives, *Environ.*
739 *Geol.*, 33 (2/3), 96–108, 1998.

740 Raudina, T. V., Loiko, S. V., Lim, A. G., Krockov, I. V., Shirokova, L. S., Istignichev, G. I., Kuzmina, D. M.,
741 Kulizhsky, S. P., Vorobyev, S. N., and Pokrovsky, O. S.: Dissolved organic carbon and major and trace elements in peat
742 porewater of sporadic, discontinuous, and continuous permafrost zones of western Siberia, *Biogeosciences*, 14, 3561–3584,
743 <https://doi.org/10.5194/bg-14-3561-2017>, 2017.

744 Reimann, C., Halleraker, J. H., Kashulina, G., and Bogatyrev, I.: Comparison of plant and precipitation chemistry in
745 catchments with different levels of pollution on the Kola Peninsula, Russia, *Sci. Total Environ.*, 243/244, 169–191, 1991.

746 Reimann, C., Niskavaara, H., de Caritat, P., Finne T.E., Äyräs, M., Chekushin, V.: Regional variation of snowpack
747 chemistry in the vicinity of Nikel and Zapoljarnij, Russia, northern Finland and Norway, *Sci. Total Environ.*, 182, 147–
748 158, 1996.

749 Reimann, C., de Caritat, P.: Intrinsic flaws of element enrichment factors (EFs) in environmental geochemistry,
750 *Environ. Sci. Technol.* 34, 5084-5091, 2000.

751 Reimann, C., Banks, D., de Caritat, P.: Impacts of airborne contamination on regional soil and water quality: The
752 Rola Peninsula, Russia, *Environ. Sci. Technol.*, 34, 2727-2732, 2000.

753 Reimann, C., Filzmoser, P., Garrett, R. G., Dutter, R.: *Statistical Data Analysis Explained: Applied Environmental*
754 *Statistics with R*. John Wiley and Sons Ltd., NY, (343pp), 2008.

755 Reinosdotter, K. and Viklander, M.: A comparison of snow quality in two Swedish municipalities – Luleå and
756 Sundsvall, *Water, Air, Soil Poll.*, 167, 3–16, 2005.

757 Resources of Surface Waters of USSR, Vol. 15: Altai and Western Siberia, Issue 2, Middle Ob. Ed: Panina, N. A.,
758 Leningrad, *Gidrometeoizdat*, 408 pp, 1972.

759 Resources of Surface Waters of USSR, Vol. 15: Altai and Western Siberia, Issue 3, Low Irtush and Low Ob. Ed:
760 Vodogrtesky, V. E., Leningrad, *Gidrometeoizdat*, 424 pp, 1973.

761 Ross, H. B. and Granat, L.: Deposition of atmospheric trace metals in northern Sweden as measured in the
762 snowpack, *Tellus*, 38B, 27–43, 1986.

763 Rossini Oliva, S. and Fernández Espinosa, A. J.: Monitoring of heavy metals in topsoils, atmospheric particles and
764 plant leaves to identify possible contamination sources, *Microchem. J.*, 86, 131–139, 2007.

765 Rudnick, R. L. and Gao, S.: Composition of the continental crust. *Treatise on Geochemistry*, Ed. H. D. Holland, K.
766 K. Turekian. Elsevier V. 3, 1–63, 2003.

767 Santelman, M. V. and Gorham, E.: The influence of airborne road dust on the chemistry of Sphagnum mosses, *J.*
768 *Ecol.*, 76, 1219–1231, 1988.

769 Shevchenko, V.: The influence of aerosols on the oceanic sedimentation and environmental conditions in the Arctic,
770 *Berichte zur Polar- und Meeresforschung*, 464, 149 pp, 2003.

771 Shevchenko, V., Lisitzin A., Vinogradova, A. and Stein R.: Heavy metals in aerosols over the seas of the Russian
772 Arctic, *Sci. Total Environ.*, 306 (1–3), 11–25, doi: 10.1016/S0048-9697(02)00481-3, 2003.

773 Shevchenko, V. P., Korobov, V. B., Lisitzin, A. P., Aleshinskaya, A. S., Bogdanova, O. Yu., Goryunova, N. V.,
774 Grishchenko, I. V., Dara, O. M., Zavernina, N. N., Kurteeva, E. I., Novichkova, E. A., Pokrovsky, O. S., and Sapozhnikov,
775 F. V.: First data on the composition of atmospheric dust responsible for yellow snow in northern European Russia in March
776 2008, *Doklady Earth Sciences*, 431 (2), 497–501, doi: 10.1134/S1028334X10040185, 2010.

777 Shevchenko, V. P., Lisitsyn, A. P., Polyakova, E. I., Dethleff, D., Serova, V. V., and Stein, R.: Distribution and
778 composition of sedimentary material in the snow cover of arctic drift ice (Fram Strait), *Doklady Earth Sci.*, 383A, 278–
779 281, 2002.

780 Shevchenko, V. P., Lisitzin, A. P., Vinogradova, A. A., Smirnov, V. V., Serova, V. V., and Stein, R.: Arctic
781 aerosols. Results of ten-year investigations, *Atmos. Oceanic Optics*, 13, 510–533, 2000.

782 Shevchenko, V.P., Vorob'ev, S. N., Kirpotin, S. N., Kritskov, I. V., Manasyopov, R. M., Pokrovsky, O. S., and Politova,
783 N. V.: Investigations of insoluble particles in the snow cover of Western Siberia from Tomsk to the Ob estuary, *Optika*
784 *Atmosfery i Okeana*, 28(6), 499–504, doi: 10.15372/AOO20150601, 2015. (in Russian).

785 Shevchenko, V. P., Vinogradova, A. A., Lisitzin, A. P., Novigatsky, A. N., Panchenko, M. V., and Pol'kin, V. V.:
786 Aeolian and ice transport of matter (including pollutants) in the Arctic. In: Kallenborn, R. (Ed.), *Implications and Consequences*
787 *of Anthropogenic Pollution in Polar environments, From Pole to Pole*. Springer-Verlag, Berlin Heidelberg, pp. 59–73, doi:
788 10.1007/978-3-642-12315-3_5, 2016.

789 Schot, P. P., van der Wal, J.: Human impact on regional groundwater composition through intervention in natural
790 flow patterns and changes in land use, *J. Hydrol.*, 134, 297–313, 1992.

791 Shotykh, W., Nesbitt, H. W., and Fyfe, W. S.: The behaviour of major and trace elements in complete vertical peat
792 profiles from three Sphagnum bogs, *Internat. J. Coal Geol.*, 15, 163–190, 1990.

793 Shotykh, W., Nesbitt, H. W., and Fyfe, W. S.: Natural and anthropogenic enrichments of trace metals in peat profiles,
794 *Internat. J. Coal Geol.*, 20(1-2), 49–84, 1992.

795 Singh, V. P., Singh, P., and Haritashya, U. K.: *Encyclopedia of Earth Sciences Series: Encyclopedia of Snow, Ice*
796 *and Glaciers*, Springer, 1253 pp, 2011.

797 Siudek, P., Frankowski, M., and Siepak, J.: Trace element distribution in the snow cover from an urban area in
798 central Poland, *Environ. Monit. Assess.* 187, 225, 2015.

799 Smith, L. C., Macdonald, G. M., Velichko, A. A., Beilman, D. W., Borisova, O. K., Frey, K. E., Kremenetsky, K.
800 V., and Sheng, Y.: Siberian peatlands as a net carbon sink and global methane source since the early Holocene, *Science*,
801 303, 353–356, 2004.

802 Snyder-Conn, E., Garbarino, J. R., Hoffman, G. L., and Oelkers, A.: Soluble trace elements and total mercury in
803 Arctic Alaskan snow, *Arctic* 50(3), 201–215, 1997.

804 Steinnes, E. and Friedland, A. J.: Metal contamination of natural surface soils from long-range atmospheric
805 transport: Existing and missing knowledge, *Environ. Rev.* 14, 169–186, 2006.

806 Stepanova, V. M., Pokrovsky, O. S., Viers, J., Mironycheva-Tokareva, N. P. Kosykh, N. P., and Vishnyakova, E.
807 K.: Major and trace elements in peat profiles in Western Siberia: impact of the landscape context, latitude and permafrost
808 coverage, *Appl. Geochem.*, 53, 53–70, 2015.

809 Stohl, A., Klimont, Z., Eckhardt, S., Kupiainen, K., Shevchenko, V. P., Kopeikin, V. M., and Novigatsky, A. N.:
810 Black carbon in the Arctic: the underestimated role of gas flaring and residential combustion emissions, *Atmos. Chem.*
811 *Phys.*, 13, 8833–8855, doi:10.5194/acp-13-8833-2013, 2013.

812 Sutherland, R. A., Tolosa, C. A., Tack, F. M., and Verloo, M. G.: Characterization of selected element concentration
813 and enrichment ratios in background and anthropogenically impacted roadside areas, *Arch. Environ. Contam. Toxicol.*, 38
814 (4), 428–438, 2000.

815 Talovskaya, A. V., Simonenkov, D. V., Filimonenko, E. A., Belan, B. D., Yazikov, E. G., Rychkova, D. A., and
816 Il'enok, S. S.: Study of aerosol composition in Tomsk region background and urban stations (the winter period 2012/13),
817 *Optika Atmosfery i Okeana*, 27, 999–1005, 2014.

818 Telmer, K., Bonham-Karter, G. F., Kliza, D. A., and Hall, G. E. M.: The atmospheric transport and deposition of
819 smelter emissions: Evidence from the multi-element geochemistry of snow, Quebec, Canada, *Geochim. Cosmochim. Acta*,
820 68, 2961–2980, 2004.

821 Thioulouse, J., Chessel, D., Doledec, S., and Olivier, J. M.: ADE-4: a multivariate analysis and graphical display
822 software, *Statistics and Computing*, 7, 75–83, doi:10.1023/A:1018513530268, 1997.

823 Tranter, M., Brimblecombe, P., Davies, T. D., Vincent, C. E., Abrahams, P. W., and Blackwood I.: The chemical
824 composition of snowpack, snowfall and meltwater in the Scottish Highlands Highlands - evidence for preferential elution,
825 *Atmos. Environ.*, 20, 517-525, 1986.

826 Tranter, M., Davies, T.D., Abrahams, P. W., Blackwood, I., Brimblecombe, P., and Vincent, C. E.: Spatial
827 variability in the chemical composition of snowcover in a small, remote Scottish catchment, *Atmos. Environ.*, 21, 853-862,
828 1987.

829 Tyrtikov, A. P.: Thawing of soils in tundra of western Siberia, In: *Natural environment of western Siberia*. Popov,
830 A. I. (ed.) Izd-vo MGU, Moscow, 1973 (in Russian)

831 Tyrtikov, A. P.: Dynamics of vegetation coverage and permafrost development in western Siberia, Nauka, Moscow,
832 1979 (in Russian).

833 Van de Velde, K., Ferrari, C., Barbante, C., Moret, I., Bellomi, T., Hong, S., and Boutron, C.: A 200 year record of
834 atmospheric cobalt, chromium, molybdenum, and antimony in high altitude alpine firn and ice, *Environ. Sci. Technol.*, 33,
835 3495–3501, 1999.

836 Vasil'chuk, Yu. K., Shevchenko, V. P., Lisitzin, A. P., Budantseva, N. A., Vorobiov, S. N., Kirpotin, S. N., Krizkov,
837 I. V., Manasyrov, R. M., Pokrovsky, O. S., Chizhova, Ju. N.: Oxygen isotope and deuterium composition of snow cover on
838 the profile of West Siberia from Tomsk to the Ob Bay, *Doklady Earth Sciences*, 471(12), 1284-1287.

839 Vasil'evskaya, V. D., Ivanov, V. V., and Bogatyrev, L. G.: Soils of North of western Siberia, Moscow University
840 Publ. House, Moscow, 228 pp, 1986 (in Russian).

841 Vinogradova, A. A., Malkov, I. P., Polissar, A. V., Khramov, N. N.: Elemental composition of the surface
842 atmospheric aerosol in the Arctic regions of Russia, *Izvestiya, Atmospheric and Oceanic Physics* 29, 149–157, 1993.

843 Walker, T. R., Young, S. D., Crittenden, P. D., and Zhang, H.: Anthropogenic metal enrichment of snow and soil in
844 north-eastern European Russia, *Environ. Pollut.*, 121, 11–21, 2003.

845 Weinbruch, S., Wiesemann, D., Ebert, M., Schütze, K., Kallenborn, R., and Ström, J.: Chemical composition and
846 sources of aerosol particles at Zeppelin Mountain (Ny Ålesund, Svalbard): An electron microscopy study, *Atmos. Environ.*,
847 49, 142–150, 2012.

848 Welch, H. E., Muir, D. C. G., Billeck, B. N., Lockhart, W. L., Brunskill, G. J., Kling, H. J., Olson, M. P., and
849 Lemoine, R. M.: Brown snow: a long-range transport event in the Canadian Arctic, *Environ. Sci. Technol.*, 25, 280–286,
850 1991.

851 Yashchenko, I. G., Svarovskaya, L. I., and Alexeeva, M. N.: Assessment of environmental risk associated with gas
852 flaring in Western Siberia, *Optika Atmosfery i Okeana*, 27(6), 560–564, 2014 (in Russian).

853 Yeghicheyan, D., Bossy, C., Bouhnik Le Coz, M., Douchet, Ch., Granier, G., Heimburger, A., Lacan, F.,
854 Lanzanova, A., Rousseau, T. C. C., Seidel, J.-L., Tharaud, M., Candaudap, F., Chmeleff, J., Cloquet, C., Delpoux, S.,
855 Labatut, M., Losno, R., Pradoux, C., Sivry, Y., and Sonke, J. E.: A Compilation of Silicon, Rare Earth Element and
856 Twenty-One other Trace Element Concentrations in the Natural River Water Reference Material SLRS-5 (NRC-CNRC),
857 *Geostand. Geoanal. Res.*, 37, 449–467, doi:10.1111/j.1751-908X.2013.00232.x, 2013.

858 Zakharova, E. A., Kouraev, A. V., Biancamaria, S., Kolmakova, M. V., Mognard, N. M., Zemtsov, V. A., Kirpotin,
859 S. N., and Decharme, B.: Snow cover and spring flood flow in the northern part of Western Siberia (the Poluy, Nadym,
860 Pur, and Taz Rivers, *J. Hydrometeorol.*, 12, 1498-1511, 2011.

861 Zdanowicz, C., Hall, G., Vaive, J., Amelin, Y., Percival, J., Girard, I., Biscaye, P., and Bory, A.: Asian dustfall in
862 the St. Elias Monttains, Yukon, Canada, *Geochim. Cosmochim. Acta*, 70, 3493–3507, 2006.

863 Zdanowicz, C. M., Zielinski, G. A., and Wake, C. P.: Characteristics of modern atmospheric dust deposition in snow
864 on the Penny Ice Cap, Baffin Island, Arctic Canada, *Tellus*, 50B, 506–520, 1998.

865 Zhang, Y., Mahowald, N., Scanza, R.A., Journet, E., Desboeufs, K., Albani, S., Kok, J.F., Zhuang, G., Chen, Y.,
866 Cohen, D. D., Paytan, A., Patey, M. D., Achterberg, E. P., Engelbrecht, J. P., and Fomba, K. W.: Modeling the global
867 emission, transport and deposition of trace elements associated with mineral dust, *Biogeosciences* 12, 5771–5792, 2015.

868

869

870

871

872

873

874

875

876

877

878


879 **Table 1.** Minimal, maximal, median and geometric mean concentration of dissolved ($\mu\text{g L}^{-1}$ snow water), n=35, and particulate
880 ($\mu\text{g g}^{-1}$ particles), n=34 snow components. The data for upper continental crust (UPC) are from Rudnick and Gao (2003). N.A.
881 stands for non analyzed.

| Element | Dissolved | | | | Particulate | | | | UPC |
|---|-----------|--------|--------|----------------|-------------|--------|--------|----------------|-------|
| | Min | Max | Median | Geometric mean | Min | Max | Median | Geometric mean | |
| pH | 4.38 | 8.73 | 5.11 | 5.44 | N.A. | N.A. | N.A. | N.A. | N.A. |
| SC, $\mu\text{S cm}^{-1}$ | 9 | 35 | 15.5 | 16.3 | N.A. | N.A. | N.A. | N.A. | N.A. |
| DIC, mg/L | 0.26 | 2.12 | 0.37 | 0.47 | N.A. | N.A. | N.A. | N.A. | N.A. |
| DOC, mg/L | 0.46 | 1.87 | 0.84 | 0.85 | N.A. | N.A. | N.A. | N.A. | N.A. |
| Cl, mg/L | 0.07 | 2.94 | 0.51 | 0.48 | N.A. | N.A. | N.A. | N.A. | N.A. |
| SO₄, mg/L | 0.41 | 2.01 | 0.71 | 0.72 | N.A. | N.A. | N.A. | N.A. | N.A. |
| Li | N.A. | N.A. | N.A. | N.A. | 2.6 | 32.2 | 10.8 | 10.7 | 24 |
| Be | N.A. | N.A. | N.A. | N.A. | 0.12 | 2.11 | 0.59 | 0.59 | 2.1 |
| Na | 47 | 1982 | 295 | 303 | 1452 | 39156 | 6717 | 7314 | 24200 |
| Mg | 19 | 862 | 114 | 114 | 3492 | 156712 | 19089 | 21411 | 14900 |
| Al | 1.6 | 35.2 | 15.5 | 12.3 | 6444 | 138267 | 31079 | 31565 | 81500 |
| P | N.A. | N.A. | N.A. | N.A. | 70 | 1928 | 481 | 503 | 660 |
| Si | 3.5 | 180 | 64.6 | 33.2 | N.A. | N.A. | N.A. | N.A. | N.A. |
| K | 39.2 | 120 | 55.5 | 63.0 | 1682 | 38395 | 5895 | 6023 | 23200 |
| Ca | 57 | 2266 | 267 | 296 | 3944 | 159272 | 17331 | 17775 | 25600 |
| Ti | 0.001 | 0.338 | 0.032 | 0.018 | 194 | 5762 | 674 | 689 | 3800 |
| V | 0.007 | 0.221 | 0.051 | 0.049 | 23.8 | 322 | 67.4 | 69.7 | 97 |
| Cr | 0.027 | 0.340 | 0.111 | 0.117 | 43.8 | 841 | 138 | 156 | 92 |
| Mn | 0.62 | 9.54 | 3.06 | 2.99 | 180 | 1242 | 400 | 404 | 780 |
| Fe | 1.8 | 62.2 | 14.6 | 12.0 | 7206 | 41255 | 15873 | 16488 | 39100 |
| Co | 0.006 | 0.418 | 0.097 | 0.094 | 5.9 | 60.7 | 19.4 | 18.6 | 17.3 |
| Ni | 0.04 | 5.66 | 0.36 | 0.36 | 28.1 | 1067 | 149 | 145 | 47 |
| Cu | 0.16 | 2.51 | 0.57 | 0.63 | 13.1 | 273 | 63.4 | 75.1 | 28 |
| Zn | 1.7 | 31.0 | 8.3 | 8.3 | 70.7 | 3832 | 202 | 255 | 67 |
| Ga | 0.0001 | 0.0185 | 0.0023 | 0.0014 | 1.8 | 26.7 | 8.32 | 7.73 | 17.5 |
| Ge | N.A. | N.A. | N.A. | N.A. | 0.36 | 3.18 | 0.88 | 0.91 | 1.4 |
| As | 0.02 | 0.46 | 0.19 | 0.15 | 3.8 | 67.2 | 16.1 | 16.0 | 4.8 |
| Rb | 0.033 | 0.262 | 0.066 | 0.075 | 6.1 | 124 | 24.1 | 23.4 | 84 |
| Sr | 0.26 | 10.2 | 1.04 | 1.23 | 26.2 | 580 | 117 | 115 | 320 |
| Y | N.A. | N.A. | N.A. | N.A. | 1.52 | 40.2 | 7.1 | 7.5 | 21 |
| Zr | 0.0001 | 0.0403 | 0.0024 | 0.0015 | 7.7 | 383 | 36.8 | 38.5 | 193 |
| Nb | N.A. | N.A. | N.A. | N.A. | 0.80 | 25.5 | 3.54 | 3.72 | 12 |
| Mo | 0.0001 | 0.059 | 0.010 | 0.005 | 0.55 | 10.4 | 2.12 | 2.24 | 1.1 |
| Cd | 0.015 | 0.180 | 0.047 | 0.046 | 0.11 | 3.37 | 0.71 | 0.71 | 0.09 |

| | | | | | | | | | |
|-----------|---------|---------|---------|---------|-------|------|------|------|------|
| Sn | N.D. | N.D. | N.D. | N.D. | 1.13 | 29.3 | 7.76 | 7.36 | 2.1 |
| Sb | 0.009 | 0.132 | 0.036 | 0.038 | 1.67 | 27.2 | 5.96 | 6.15 | 0.4 |
| Cs | 0.0015 | 0.0105 | 0.0034 | 0.0036 | 0.32 | 4.78 | 1.35 | 1.24 | 4.9 |
| Ba | 0.74 | 13.6 | 3.35 | 3.32 | 88 | 1664 | 374 | 391 | 628 |
| La | 0.001 | 0.049 | 0.012 | 0.011 | 2.0 | 60.2 | 10.3 | 10.8 | 31 |
| Ce | 0.003 | 0.095 | 0.022 | 0.019 | 4.05 | 128 | 19.0 | 20.6 | 63 |
| Pr | 0.0001 | 0.0084 | 0.0022 | 0.0019 | 0.50 | 15.5 | 2.30 | 2.35 | 7.1 |
| Nd | 0.0013 | 0.0275 | 0.0085 | 0.0067 | 1.86 | 58.6 | 8.32 | 8.70 | 27 |
| Sm | 0.0001 | 0.0072 | 0.0020 | 0.0016 | 0.39 | 11.8 | 1.78 | 1.79 | 4.7 |
| Eu | 0.00010 | 0.00253 | 0.00096 | 0.00083 | 0.11 | 2.56 | 0.45 | 0.47 | 1.0 |
| Gd | 0.0004 | 0.0082 | 0.0022 | 0.0022 | 0.40 | 10.3 | 1.71 | 1.77 | 4.0 |
| Dy | 0.00002 | 0.0041 | 0.0016 | 0.0008 | 0.32 | 7.83 | 1.35 | 1.42 | 3.9 |
| Ho | 0.00006 | 0.00123 | 0.00061 | 0.00054 | 0.06 | 1.51 | 0.26 | 0.27 | 0.83 |
| Er | 0.0002 | 0.0029 | 0.0010 | 0.0010 | 0.18 | 4.71 | 0.77 | 0.80 | 2.3 |
| Tm | 0.00002 | 0.00088 | 0.00011 | 0.00009 | 0.03 | 0.72 | 0.11 | 0.11 | 0.3 |
| Yb | 0.00000 | 0.00289 | 0.00089 | 0.00049 | 0.16 | 4.91 | 0.73 | 0.73 | 1.96 |
| Lu | N.A. | N.A. | N.A. | N.A. | 0.024 | 0.76 | 0.11 | 0.11 | 0.31 |
| Hf | N.A. | N.A. | N.A. | N.A. | 0.25 | 13.2 | 1.10 | 1.18 | 5.3 |
| Ta | N.A. | N.A. | N.A. | N.A. | 0.18 | 4.35 | 0.62 | 0.62 | 0.9 |
| W | 0.002 | 0.108 | 0.020 | 0.017 | 2.0 | 102 | 35.9 | 28.8 | 1.9 |
| Tl | N.A. | N.A. | N.A. | N.A. | 0.04 | 0.73 | 0.23 | 0.23 | 0.90 |
| Pb | 0.02 | 3.67 | 0.51 | 0.38 | 13.2 | 703 | 71.9 | 67.9 | 17 |
| Th | N.A. | N.A. | N.A. | N.A. | 0.43 | 17.1 | 2.22 | 2.33 | 10.5 |
| U | 0.0007 | 0.0063 | 0.0031 | 0.0028 | 0.19 | 4.69 | 0.92 | 0.93 | 2.7 |

882


883



884

885 **Figure 1.** Map of the study site. with permafrost zone boundaries. The size of the sampling points reflects the
 886 concentration of particulate fraction (mg/L_{snow water}). The boundaries of permafrost zones are from Brown et al. (1997).

887



888


889

890

891 **Fig. 2.** Examples of dissolved ($< 0.45 \mu\text{m}$) metal concentrations in snow water as a function of latitude. The cause for
 892 the elevated concentrations of Ni, Cd and Sb at ca. 64-65°N is most likely industrial impact, but given relatively low
 893 number of data points around industrial centers it is hard to prove it unambiguously.

894


895
896
897
898
899
900
901
902
903
904
905
906
907
908



dissolved fraction

A

Cluster Dendrogram



dissolved fraction


B

909
910
911
912
913
914

Fig. 3 A: PCA Factorial map F1x2 of elements of a reconstructed table for the dissolved fraction. Partition of elements into 6 groups revealed by HCA is shown by a contour line. **B:** Dendrogram of a hierarchical cluster analysis (HCA) performed on variables of a reconstructed table for the dissolved fraction using the Pearson correlation as a distance measure and Ward's method for the linkage rule.

915

916



917

918

919

920

921 **Fig. 4.** pH value (A) and Sr (B), Al (C) and Pb (D) concentration in dissolved fraction of snow as a function of
922 concentration of particles. Note log X scale for Sr and Pb.

923

924

925

926

927

928


929

930


931

932

933



5 **Fig. 5 A.** The latitude-averaged Al-normalized enrichment coefficient of snow particles with respect to the earth crust.



10 **Fig. 5 B.** The average values (56 to 68°N) of the ratio of particulate to dissolved element concentration in snow water of western Siberia. Bold red line indicates statistically non-significant deviation from 1.





Fig. 6. The ratios of the average concentrations of elements in snow particles (neglecting sample SF22) to those in mineral soil (A), peat (B) and mosses (C) of WSL. The peat, moss, and underlying mineral horizons data are averaged over the latitude of 55 to 68°N as described in Stepanova et al (2015). Note normal Y scale for mineral soil (A) and log Y scale for peat and moss (B, C).



5 **Fig. 7.** The ratio of mean dissolved flux of rivers in three latitudinal zones (56-60°N, 60-64°N, and 64-68°N) of WSL to the stock of dissolved fraction of snow. For this calculation, the snow volume (in mm of water) accumulated over full winter and mean river runoff over May and June were used. **The impact of snow melt on river export fluxes in spring strongly increases northward for DIC, Cl⁻, SO₄²⁻, Na, Mg, Ca, Cr, Ni, Mo, Rb, U whereas Cd, Pb, Sb, Cu, As, W and Cs present a high impact of snow melt water on river for the three latitude zones.**

SUPPLEMENTARY INFORMATION






Fig. S1. The Al-based enrichment factor (relative to the upper part of continental earth crust) of particulate fraction of snow for Mo, Sb, Cr, Ni, Zn, Cd, Cu and Pb as a function of latitude.

5

**A**

15

**B****B**

25

30

Fig. S2 A. PCA Factorial map F1x2 of variables (elements) of a reconstructed table for the particulate fraction. Partition of elements into 5 groups revealed by HCA is reported by a contour line. **B:** Dendrogram of a hierarchical cluster analysis (HCA) performed on variables of a reconstructed table for the particulate fraction using Pearson correlation distance as distance measure and Ward's method for the linkage rule.




Fig. S3. Snow water soluble ($< 0.45 \mu\text{m}$) of Zn (A), Cu (B), Cd (C), Pb (D), Sb (E) and Mo (F) (blue asterisk) compared with actual concentrations in rivers during spring flood (May-June) of different size of the watershed (closed diamonds, squares, triangles and circles correspond to < 100 , $100\text{-}1000$, $1000\text{-}10,000$ and $> 10,000$ km² surface area, respectively) in western Siberia along the latitudinal gradient.

Table S1. Mineralogical composition of selected snow particles.

| No | Quartz | Albite | K-Fs | Calcite | Dolomite | Chlorite | Illite | Phlogopite | Amphibole | Pyroxene | Chrysotile | Magnesite | Forsterite | Talc | Magnetite |
|-------|--------|--------|------|---------|----------|----------|--------|------------|-----------|----------|------------|-----------|------------|------|-----------|
| SF-1 | 37 | 20 | 11 | 19 | < 1 | 4 | 7 | | 2 | | | | | | |
| SF-3 | 38 | 27 | 9 | 3 | | 10 | 9 | | 3 | | | | | | |
| SF-14 | 20 | 8 | 6 | 1 | 48 | 5 | 4 | | 3 | 1 | 2 | | | 1 | |
| SF-31 | 30 | 20 | 12 | | 8 | 8 | | 12 | 6 | | 3 | | | | |
| SF-33 | 35 | 16 | 9 | | 10 | 3 | | 16 | 4 | | 3 | 4 | | | |
| SF-36 | 47 | 7 | 11 | | 1 | 4 | | 5 | 3 | 3 | 12 | | 4 | 3 | |
| SF-38 | 48 | 3 | 6 | | 1 | 6 | | 4 | 3 | | 16 | | 6 | 4 | 3 |
| SF-39 | 41 | 8 | 6 | 4 | 8 | 5 | | 12 | 2 | 2 | 8 | | 2 | 2 | |
| SF-40 | 35 | 12 | 9 | 6 | 3 | 4 | | 8 | 4 | 3 | 10 | | 4 | 2 | |