Hydrol. Earth Syst. Sci. Discuss., doi:10.5194/hess-2016-626-SC2, 2017 © Author(s) 2017. CC-BY 3.0 License.



HESSD

Interactive comment

Interactive comment on "Evaluating primary productivity, ripple effect and resilience of fluvial ecosystems: a new approach to assessing environmental flow requirement" by Yui Shinozaki and Naoki Shirakawa


A.V. Pastor

amandine.pastor22@gmail.com

Received and published: 13 February 2017

The authors brought new insights into the global eco-hydrology field by considering NPP, resilience and trophic levels. These last ideas are necessary to improve eco-hydrology relationships at global scale. However, these last deserve more investigation, monitoring and testing before being upscaled to global EF methods.

I recommend to test their method with empirical case study first and second to acknowledge worked developed by (Gerten, Hoff et al. 2013, Pastor, Ludwig et al. 2014) with the development of which the VMF method which was not mentioned in this re-



Discussion paper



view, neither the Q90_Q50 and Tessman methods which shown better results than the Smakthin and Hanasaki and Tennant methods (Pastor, Ludwig et al. 2014). The VMF method was acknowledged by other global assessments including: (Gerten, Hoff et al. 2013, Boulay, Bare et al. 2015, Gaupp, Hall et al. 2015, Sadoff 2015, Steffen, Richardson et al. 2015) and was used to defined freshwater planetary boundaries in science (see references). This study is based on the Tennant method which was created for temperate case studies and which showed low performances for intermittent rivers Ppastor et al. 2014). Moreover, an explanation on the choice of method (Tennant over Smakhtin, parametric vs. non-parametric methods) is required and why the latest methods were ignored (Hoekstra and Mekonnen 2011, Pastor, Ludwig et al. 2014).

I recommend the authors to test their parameters and methods with different case studies and other global global EF methods worldwide before validation and global upscaling. It is also necessary to compare these last EF methods in different contexts (different ecoregions, flow regime types). Overall, this study has the merit to extend knowledge on the eco-hydrology field but to my point of view it should be first described the use of NPP, resilience and trophic levels for all freshwater ecoregions (Abell, Thieme et al. 2008) including the acknowledgement of the latest global eco-hydrological studies (Oberdorff, Tedesco et al. 2011, Tisseuil, Cornu et al. 2013).

I wish good luck to the authors to develop further their study,

References

Abell, R., et al. (2008). "Freshwater ecoregions of the world: a new map of biogeographic units for freshwater biodiversity conservation." BioScience 58(5): 403-414.

Boulay, A.-M., et al. (2015). "Consensus building on the development of a stressbased indicator for LCA-based impact assessment of water consumption: outcome of the expert workshops." The International Journal of Life Cycle Assessment: 1-7.

Gaupp, F., et al. (2015). "The role of storage capacity in coping with intra-and inter-

Interactive comment

Printer-friendly version

Discussion paper



annual water variability in large river basins." Environmental Research Letters 10(12): 125001.

Gerten, D., et al. (2013). "Towards a revised planetary boundary for consumptive freshwater use: role of environmental flow requirements." Current Opinion in Environmental Sustainability 5(6): 551-558.

Hoekstra, A. Y. and M. M. Mekonnen (2011). Global water scarcity: the monthly blue water footprint compared to blue water availability for the world's major river basins. 333. 222. Delft, The Netherlands, , UNESCO-IHE Institute for Water Education. Value of Water Research Report: 78pp.

Oberdorff, T., et al. (2011). "Global and Regional Patterns in Riverine Fish Species Richness: A Review." International Journal Ecology 2011, Article ID 967631: 12 pp.

Pastor, A. V., et al. (2014). "Accounting for environmental flow requirements in global water assessments." Hydrol. Earth Syst. Sci. 18(12): 5041-5059.

Sadoff, C. W. (2015). Securing Water, Sustaining Growth: Report of the GWP/OECD Task Force on Water Security and Sustainable Growth.

Steffen, W., et al. (2015). "Planetary boundaries: Guiding human development on a changing planet." Science 347(6223): 1259855.

Tisseuil, C., et al. (2013). "Global diversity patterns and crossâĂŘtaxa convergence in freshwater systems." Journal of Animal Ecology 82(2): 365-376.

Interactive comment on Hydrol. Earth Syst. Sci. Discuss., doi:10.5194/hess-2016-626, 2016.

HESSD

Interactive comment

Printer-friendly version

