

Dear Editor:

We greatly appreciate the efforts of the two reviewers and we have carefully considered and responded to all of their comments with appropriate changes to our manuscript. Our responses are shown here (in blue typeset), including we made our changes and additions. The page and line numbers we report for our changes refer to those in the edited manuscript (marked up) that was submitted. Thank you all for your diligence and we feel that the paper has been greatly improved based on these critiques. We hope that these major revisions will satisfy the reviewers and Editor.

### **Anonymous Referee #1**

This manuscript reports on rainfall, throughfall, and soil moisture measurements on a steep slope. The topic is of high relevance to HESS readers and the manuscript is clearly written. It has four important limitations: (1) poor data on throughfall intensity; (2) lack of relevance of soil moisture measurements for slope stability; and (3) overstating conclusions beyond data support:

Thank you for your review comments, these are addressed in the specific responses to your comments that follow.

1. The substantial uncertainty in throughfall measurements using the trough system necessitated large calibration correction of 50-78% (P7L14). The total corrected throughfall estimates are 10-14% less than rainfall, so calibration errors are 3-8 times larger than the difference being quantified. I admire the authors for confessing this limitation, but I don't understand why they find data from this instrument sufficient for addressing their hypotheses. I also think the calibration procedure cannot be omitted from the paper given its paramount importance. There are some apparent absurdities in the data that may be explainable by the calibration procedure as well: e.g., it appears total TF>RF for the early portions of the most intense storms in the dataset (Fig 5a-d), followed by RF>TF later in the storm during low-intensity RF; I am not aware of this pattern ever being reported. It appears that conditioning the calibration on total TF:RF may have resulted in plausible estimates of total mass balance (where they can be compared to expectations), but measurements of intensity are both the least reliable and most important data. I disagree with the "clarity" claimed P17L22.

Thank you for raising this query. After careful consideration, we chose to address this issue in a new section of the paper entitled "Limitations and recommendations":

### **6 Limitations and recommendations**

There are some important limitations to our methods. In a prior study, Ziegler et al. (2009) compared the same troughs used herein with several movable tipping bucket gauges, finding no statistical difference between the two approaches. However, in the previous study, total event precipitation was examined, not minutely changes occurring over the course of storms. We caution that the trough method may create a somewhat confusing signal because the area-integrated pattern of throughfall, in which the records are delayed as water captured in the trough flows through the trough, compared to an individual rain gauge placed above the canopy. Nevertheless, at high rates of throughfall, through flow would be more efficient. Additionally, estimates of both throughfall and rainfall have errors. Measurements by tipping-bucket rain gauges installed above a canopy are affected by turbulent exchange at this interface, and wind affects rainfall catch (e.g., Kamph and Burges, 2010). As mentioned before, the

throughfall troughs had a large associated splash error during high-intensity events. While our attempt to correct the splash error resulted in reasonable total event values, individual minutely values could still have substantial errors; these would affect the time series we compare with measured rainfall (Figure 6). However, such limitations are inherent in most all studies reported in the literature (albeit typically not articulated).

In future experiments, we urge researchers to minimize uncertainties by: (a) using troughs that are deeper to minimize splash loss; (b) collect ambient rainfall in more than one location, preferably gauges positioned just above ground level with appropriate wind shields to minimize wind effects; (c) collect more throughfall values, potentially employing other types of collection devices to help interpret the measurements; and (d) perform a rigorous assessment of splash losses to facilitate error correction (again our splash error data are few). We also encourage researchers to spend time in the forest plots during events, recording the various types of phenomena that may affect the capture of throughfall over time, to help with the interpretation of data. Here, we re-emphasize that throughfall reaching the forest floor is highly variable in space and time. Multi-stored canopies can create wet and dry zones below them, which change over the course of a storm with respect to variable wind direction, changes in rainfall phenomena (rate, drop size), and changes in canopy wetness (e.g., Konishi et al., 2006). The oddities apparent in the data of some of our recorded storms (e.g., higher throughfall than rainfall during some periods, but not others) may possibly be related to inadequacies in our error correction; however, these may be realistic. For example, they may result because wind-driven rain is captured by portions of a large tree and then channelled directly to the throughfall trough (as was documented at the prior study site; Ziegler et al., 2009).

We believe that these uncertainties do not undermine the integrity of our conclusions. While the uncertainties may prevent us from producing a high-precision budget of the portion of rainfall converted to throughfall at minutely scales, they do allow us to address the primary goals of this investigation, which are to assess whether secondary tropical canopies intercept sufficient rainwater during large storms to mitigate landslide initiation compared to open areas.

2. Hillslope hydrology is poorly constrained, so it is difficult to understand relevance of the soil moisture data to slope stability. There are two problems in the manuscript that arise because of this. (1) The instrumented slope was obviously not near failure during conditions represented in the dataset as evidenced by low soil moisture at depth, and the deep and highly weathered soils suggest this site is not prone to failure in general. It is unclear specifically how soil moisture responses in this slope is useful for understanding slope stability, but the lack of responsiveness at timescales relevant to canopy interception is not enough evidence to conclude a general lack of canopy interception effect. (2) The analysis of paired TF and soil water measurements implies a one-dimensional water balance is relevant for slope failure, but in fact hillslope- and watershed-scale hydrologic conditions are important. Depending on slope configuration, there may be little reason to expect substantial effect of local canopy interception on soil moisture at depth and thus slope stability. These conceptual problems can be addressed by modifying the discussion, but I think the conclusion linking interception to stability through soil moisture at this site (P17L26) is oversimplified.

While we agree that the slope hydrology (actually the soil physics) could have been better constrained, we emphasize this was only a supplemental part of the study – i.e., as stated in the third objective “(to) determine the effect of canopy interception on the potential for soil water increases that could trigger landslides”. True, the site was not near failure during the storms that were monitored, in spite of some of them exceeding the conservative global slope failure thresholds for intensity – duration. It is also true that for our site we were only able to assess one-dimensional (vertical) flow. But we never intended this to be an assessment of stability of

3-D hillslopes where issues like convergent topography and related flow pathways exert controls on landslide initiation. We simply wanted to address the issue of the role of canopy interception during storms of different sizes on the delivery of rainwater to and into the soil. Our inclusion of the soil moisture data was simply to show that rather homogeneous soils (without extensive macropores) could effectively buffer peak rainfall inputs that have been postulated by others to affect pore water pressure at depth and thus affect landslides. We recognize the limitations of such a 1-D assumption in the context of 3-D hillslope hydrology and now address this more clearly. The reviewer is correct in stating that “Depending on slope configuration, there may be little reason to expect substantial effect of local canopy interception on soil moisture at depth and thus slope stability.”, and we have modified the Discussion to note this more clearly. But this is clearly not recognized by a number of researchers who keep supporting the idea that canopy interception buffers rain inputs and potentially ameliorates peak pore water pressure response at depth. Additionally, to address these concerns we have modified the “Conclusions”.

See additions on pg. 12, L 20-23 and pg. 17, L 13.

3a. Intensity-duration quantification of slope stability is useful for general purposes, but limitations of the concept prevent literal application of thresholds. Obviously none of the thresholds were correct for the instrumented slope or it would have failed about 30 times in the 30 months of monitoring. So, each slope must have its own threshold, and presumably some slopes have thresholds that pass between paired TF and RF intensities in triggering storms (Fig 7). How many slopes is that? The answer to that question is the true effect of canopy interception on slope stability, and the effect of canopy interception on stability of one instrumented slope cannot be reasonably extrapolated to encompass all slopes.

Please understand that the intensity – duration ‘thresholds’ are the minimum combinations of average storm intensity and duration which have been recorded that have triggered landslides somewhere in the world. Exceedance of thresholds does not mean a landslide will occur, it means that the very minimum rainfall conditions for global landslides has been met. True, it is a very general indicator of landslide susceptibility, but in countries like Thailand where detailed geotechnical, soils, and geological measurements are not widely available, such simple rainfall – landslide relations may be useful. And, of course, we agree that each hillslope (even hillslope segment) has a unique threshold for actual slope failure based on a number of factors and predispositions, but the point here was to show that canopy interception during storms that exceeded the global thresholds did not have a big influence on rainwater delivery to the soil surface. The question about how many hillslopes ‘have unique thresholds that pass between the throughfall and rainfall intensities’ seems unclear and rhetorical, but it was never our intention to apply our findings to the catchment-scale, rather we aimed to clarify the effects of canopy interception on water delivery to the soil surface (where much speculation occurs). We have noted the reviewer’s concerns and have tried to address these by additions on pg. 8, L. 16-17 and pg. 9, L. 5-7, plus other places.

3b. The strong conclusion that there was no intensity smoothing (P15L30) is dubious and contradicted elsewhere in the manuscript (P16L27). In “large events” (Fig 7d) and in most events overall (Fig 7), storm-total TF intensity was lower intensity than RF, so in that sense there was smoothing. Later statements (P16L4-7) rightly focus on peak intensities, but are based on highly uncertain data. Blanket characterization of “no effect” is not credible.

The statement beginning on Pg. 16, L27 does not state or infer that there was intensity smoothing by the canopy. Smoothing of rainfall peaks cannot be derived from Figure 7 – these intensity values are conservative, average storm intensity (this was originally stated on pg. 8, L 14 and on pg. 18, L 1, and is now emphasized on pg. 8, L 16, pg. 9 , L5-7; pg. 14, L 11 and 27; Pg. 17, L 21). The fact that average intensity over the course of an entire storm is less in TF compared to rainfall does not necessarily imply smoothing of rainfall peaks. Nevertheless, we have modified our statement in the Discussion (see pg. 16, L 21-22, 29).

Minor points:

P3L10-18 why present a review of root reinforcement literature when this work has nothing to do with root reinforcement?

We have now reduced this information on root reinforcement. We feel it serves the purpose of putting the issues of vegetation effects on slope stability into perspective, but, as suggested, we have reduced the reference to root reinforcement to one sentence. See changes to second paragraph of the Introduction.

P6L23 can you use these field data to convert soil moisture content ( $m^3/m^3$ ) to %saturation? The Results and Discussion refer to degree of saturation (e.g., P13L9, P13L18, P16L16) but no information is presented in the figures or text on porosity or soil moisture release curves and the reader cannot link volumetric soil moisture data to pore pressure.

This information is now presented in the text, see query below. In addition, in Table 2 we now indicate the maximum wetness (relative to saturation) that occurs during an event, extending to 24-h after the event ends.

There are some problems with the figures to clean up. Fig 6a: “axis title”; Fig 6c rainfall bars are not at the same interval as the time interval labels; panels in Fig 5-7 are often different sizes and not aligned.

All the figures 5-7 have been redrawn, with special attention paid to alignment of the axis.

P14L8 editing error muddies a critical statement about the TF-RF comparison.

The sentence “The three events in which incident rainfall exceeded the threshold but throughfall did not were similar in canopy interception ( $C_i = 0.20-0.23$ ); duration (3.4-4.6 h); and event intensity (7.1- 7.7  $mm\ h^{-1}$ )”.

Was modified to:

“The three events in which incident rainfall exceeded the threshold, but throughfall did not, were similar in canopy interception ( $C_i = 0.20-0.23$ ); duration (3.4-4.6 h); and event intensity (7.1- 7.7  $mm\ h^{-1}$ )”.

## **Anonymous Reviewer #2**

### *General comment*

The manuscript deals with a topic falling within the scope of HESS, to which part of the readership will be interested in. The paper is well structured and clearly written, and the

presented experimental data are innovative, as very few examples of similar measurements can be found in the literature. Apart of this merit, however, as the focus of the paper is about the possible effects of canopy interception on the triggering of shallow landslides caused by infiltration into the soil mantle up to a depth of 2 m, the analysis of the results in view of the infiltration processes is poor, lacking important information about soil properties, and more in-depth discussion of the soil moisture dynamics should be provided.

Thank you for recognizing the uniqueness of this research. Actually the focus of the paper is not on the effects of infiltration processes into the soil mantle; as stated in the third objective “(to) determine the effect of canopy interception on the potential for soil water increases that could trigger landslides”, this was only a supplemental part of the study. We have clarified that our data only assessed one-dimensional (vertical) soil moisture fluxes (pg. 12, L 20-22) and that we did not attempt to assess the effect of slope shape on subsurface water flux (pg. 17, L 14). Our inclusion of the soil moisture data was simply to show that homogeneous soils could buffer peak rainfall inputs (with or without canopy interception) in contrast to other speculations that canopies alone can buffer pore water pressure at depth and thus affect landslides. Our primary objective was to assess the role, magnitude, and timing of tropical canopy interception during storms of different sizes on the delivery of rainwater to and into the soil. Please see our responses that follow to specific comments.

Therefore, my recommendation is that some major revisions are needed before this manuscript could be published in HESS. Some of the following detailed comments will hopefully clarify my point of view.

#### *Detailed comments*

Page 5, lines 13-14 (minor issue). Please clarify the meaning of “landslides (...) associated with road runoff”. A clear definition of the possible triggering mechanisms of landslides in the area would indeed help to better focus the discussion of the measured soil moisture responses to precipitations.

We have added text and a reference to clarify the landslide triggering mechanisms; pg. 5, L14-16.

Page 6, lines 6-10 (major issue). Providing more information about soil properties would allow a better understanding of the observed soil moisture changes. Soil porosity is not given, but in the following section 4.4 the authors state that when volumetric moisture content approaches 0.45 the soil is saturated. The provided bulk density data seem to indicate that, at least in the upper layer, the porosity should be greater (by the way, what is the moisture content corresponding to the provided values bulk density?). As the following discussion points out that the triggering of landslide is expected to occur at depths >1.0m, would it be possible to get some information about soil properties (at least porosity and  $k_{sat}$ ) at depths larger than 25cm? (indeed, the authors say that the upper 20cm are characterized by a soil horizon different from the deeper one).

We have added information on soil properties, particularly, bulk density, porosity, and saturated hydraulic conductivity as follows (pg. 6, L 9-17):

The soil at the site is an Ultisol with a thin (< 20 cm) brown A horizon underlain by a dark red B horizon that extends below a depth of two meters. Saturated hydraulic conductivity declines exponentially from the surface ( $\sim 136 \text{ mm h}^{-1}$ ) to approximately

$< 4 \text{ mm h}^{-1}$  at a depth of 25 cm; values at 1 m and 2 m are  $1\text{-}2 \text{ mm h}^{-1}$  ( $n = 3$  measurements for all depths; unpublished data, determined with a bore-hole permeameter). Bulk density does not change much over this depth range ( $1.08\text{-}1.38 \text{ g cm}^{-3}$  for the surface, 1 m. and 2 m depths). Corresponding porosity for the three depths is 0.59, 0.52, and 0.48 (based on a particle density of  $2.65 \text{ g cm}^{-3}$ ). The decrease in saturated hydraulic conductivity is typical of that in other profiles found in southeast Asia (cf. Ziegler et al., 2004; 2006). Macropores and fissures, features that could influence preferential flow through the soil, were not abundant in the subsoil.

During the storms we did not observe the occurrence of a perched water table, which implies that there is no restricting K layer at depths (at least down to 2 m). Saturation was only approached at the surface for a brief period of time (see Table 2 and Figure 6). Bulk density is reported for dry soil conditions as typical in the soils literature.

Page 6, lines 15-16 (typo). I think it should read “(stations 429, figures 1b)”.

Changed to “station 429, Figure 1b”

Page 6, line 17 (major issue). The definition of an event should be motivated in view of the expected triggering mechanism. Why the thresholds of 8mm and 4hours have been chosen?

Our objective was to examine all rainstorms that occurred during this  $\approx 3$  yr period. We wanted to show how interception varies for different size events. Later in the paper we address how bigger storms (e.g., Table 2, Figures 5 and 6 and related discussions), which are of the size that could potentially trigger landslides, affect interception and water delivery to and into the soil. Some clarification is now included on page 6, L 24-25: “As such, we included a range of monsoon storms to assess interception losses for potentially landslide-triggering events and those that were smaller.”

Page 7, lines 6-8 (minor issue). The “dynamic calibration correction” is not clear. Please provide some description of the applied correction.

The following was modified/added (with new references):

A dynamic calibration correction was then applied to account for differences in tip volume over the range of observed tipping rates (Ziegler et al., 2009). Dynamic calibration accounts for differences in tip volume over the range of observed tipping rates (Calder and Kidd, 1978; Marsalek, 1981; Humphrey et al., 1997). These differences are caused by “splash” losses occurring as the tipping mechanism is moving when rainfall is draining from the funnel. This relationship was determined by draining known volumes of water through the tipping bucket system ( $\text{mm tip}^{-1}$ ) and recording the number of tips registered.

Page 7, lines 25-30 (moderate issue: I don't know if this issue is minor or major). It is clear that using a large throughfall collector allows the integration over a relatively large area of an inherently inhomogeneous process (in space). However, in the following discussion, in some cases the authors point out that, owing to differences in canopy structure and to the effects of wind (and possibly also to the effects of rainfall intensity, I would add), the dripping of throughfall from canopy could follow different paths, leading to local concentration of drops.

How did the authors conclude that the shape, size and position of their collector are adequate? What do the authors think about using several randomly distributed ordinary rain gages? In such a case it could be possible to get information about the adequacy of the obtained spatial mean by subtracting one (or more) gages and then check if the obtained (spatially averaged) throughfall is affected or not.

In a prior work we compared the same troughs used herein with several movable tipping-bucket (round) gauges. In that work we found no statistical difference between the two. However, we were looking at “event” totals, not dynamic changes over the course of a storm. It might be that the trough method leads to a confusing signal because the integrated pattern of throughfall, which has a delay as the water flows down the trough, is being compared to an individual rain gauge placed above the canopy. In this paper we now speak of this potential problem in the new limitations section, which is introduced above in the Reviewer 1 queries.

Page 9, line 28 – page 10, line 1 (minor issue). The outliers could be an artifact due to concentration of throughfalling drops in the collector, caused by the shape and position of the adopted collector.

This comment cuts to the issue of using a combined “Results and Discussion” rather than separating these. In our original submission, we combined these sections, but were then asked to separate them. The answer to your question appears on pg. 15, L 29-31, but we now made changes to the last sentence of this paragraph which appears on pg. 10, L 9-11.

Page 11, lines 16-25, and figure 5 (moderate issue). Looking at the provided hyetographs, it seems simply that, regardless of the timing of a peak within the event, when the intensity is below 1.0-1.1 mm/min, it results  $RF > TF$ , while it is the other way around when the intensity is larger.

You make a good point, but this is not true for event #63 (and some other events as well that are not shown in Fig. 5). To clarify, we added the following sentence on pg. 12, L13-15: While in five of these six large events rainfall exceeded throughfall when intensities were  $< 1.0 - 1.1 \text{ mm min}^{-1}$  and, typically, throughfall exceeded rainfall when intensities were  $> 1.1 \text{ mm min}^{-1}$ , this pattern was not consistently found in other storms.

Section 4.4, as a whole (major issue). The whole discussion is too simplistic, and some deeper interpretation should be made. I just give some possible keys. In a soil with  $k_{sat} < 5 \text{ mm/h}$  at the depth of 25cm (and maybe further reducing with depth), it is easily expectable that it may take many hours before water reaches 2.0m depth (even if we don't know soil properties at depth larger than 25cm), so I strongly suggest to extend the time interval over which the soil moisture changes are visualized and discussed (this issue has to do also with the previously raised issue about the adopted definition of a rainfall event). The interpretation of the (clearly visible) effect of initial soil moisture on the effectiveness of a rain event on the following soil moisture changes should be linked to the degree of saturation (but we don't know soil porosity) of the soil and to its hydraulic conductivity (once saturated, the upper layer cannot retain more water, and so, if the hydraulic conductivity allows it, it is “obliged” to release the excess water to the underlying soil). In other words, there should be a maximum storable soil moisture increase, depending on initial moisture condition, over which the excess water penetrates deeper or runs off laterally (above or below surface, or both).

We examined the soil moisture response at depth (1 m and 2 m) for all events and these longer term data are now presented in Fig. 6 – i.e., we have extended the time interval on the x-axis by 2- to 10-fold in order to show where the small soil moisture increases at depths 1 and 2 m completely subside. The degree of saturation is now shown in three new columns (Wetness, 0 m, 1 m, and 2 m) introduced in the revised Table 2. Please note that during only two events (#156 and 158) was saturation reached (or nearly reached) in the surface soil; and in no events was the subsoil approaching saturation. Please see rather extensive changes and additions have been made based on these longer time assessments referenced to new data now shown in Fig. 6 and Table 2 (see pg. 13, L 17 to pg. 14, L8); however, please note that these changes do not affect our general conclusions.

Page 13, lines 5-6 (major issue). It seems to me that limiting the observation of soil moisture to the (widely variable) duration of rainfall events in many cases may be the reason why a (later) deep soil moisture change was not detected.

See response to previous comment and changes made on pg. 13, L 17 to pg. 14, L8.

Page 15, line 22 (typo). It should probably be “environmental conditions change during the storm”.

Corrected (now pg. 16, L 13)

Page 16, lines 16-21 (major issue). See my previous comment about section 4.4. As RF and TF are quite similar in the considered forest, this paragraph would mislead the reader to the conclusion that soil moisture at 2.0m would not be affected by any rainfall event.

We have addressed this issue in our revisions in section 4.4, and we also have modified the text accordingly in the Discussion section as requested: see pg. 17, L 8-15.

Page 16, lines 30-31 (major issue). It is clear that for the considered rain events canopy interception has negligible effects. But, as I already commented above, the rain events have been defined arbitrarily >8,0mm, and there is (maybe obvious) evidence that canopy interception could be larger for smaller events. Could these neglected smaller events affect the initial moisture state of the soil at the beginning of the considered larger events? And, if so, can the authors exclude that canopy interception may play a role in the establishment of such initial moisture state? I would like to read some discussion about this point, before concluding that canopy interception has no effect on landslide initiation.

We agree. Certainly interception versus rainfall differences for very small events that just precede a large storm could affect soil moisture response. However, combined evapotranspiration and infiltration would negate such effects after a day or at most a couple days (depending on weather conditions). We have added a comment to address this issue on pg. 17, L 13-15.

Page 17, section 6 as a whole (major issue). In view of the previously raised issues, some of the conclusion drawn could be different.

We have modified our “Conclusions” section accordingly.



Figure 1, caption (minor issue). It does not seem that the topography and the major stream channels are actually shown in Figure 1a.

The figure has been revised to show this information:

Figure 1: Site map of the Mae Sa experiment site in northern Thailand. Panel (a) shows the catchment location in Thailand, major contours, and the stream system. Panel (b) shows the major land covers in the Mae Sa catchment including hillslope and plantation agriculture (AG, 23%), greenhouse agriculture (GH, 7%), urbanized or peri-urban areas (U, 8%), and forest cover with various degrees of disturbance (F, 62%). Grid cell dimensions are 2 x 2 km. Rectangles demarcate hydro-meteorological measurement sites. Streamflow, total suspended solids, particulate organic carbon, and particulate organic nitrogen were measured at the stream gage station 434. Rainfall is measured at all other numbered hydro-meteorological stations (rectangles). The throughfall investigation in this paper was conducted at station 429, where rainfall, throughfall, and soil moisture were monitored.

Figure 6a (typo). The title of x-axis is missing.

This error has been fixed.

# The canopy interception–landslide initiation conundrum: insight from a tropical secondary forest in northern Thailand

Roy C. Sidle<sup>1</sup>

Sustainability Research Centre

5 University of the Sunshine Coast  
90 Sippy Downs Drive, Sippy Downs  
Queensland 4556, Australia  
Email: rsidle@usc.edu.au

10 Alan D. Ziegler  
Department of Geography  
National University of Singapore  
AS2, #03-01, 1 Arts Link, Kent Vale  
Singapore 117570  
15 Email: adz@nus.edu.au

20

---

<sup>1</sup> Corresponding author

**Abstract.** The interception and smoothing effect of forest canopies on pulses of incident rainfall and its delivery to the soil has been suggested as a factor in moderating peak pore water pressure development within soil mantles, thus reducing the risk of shallow landslides. Here we provide three years of rainfall and throughfall data in a tropical secondary dipterocarp forest characterized by few large trees in northern Thailand, along with selected soil moisture dynamics, to address this issue. Throughout the sampling period, throughfall was an estimated 88% of rainfall, varying from 86-90% in individual years. Data from 167 events demonstrate that canopy interception was only weakly associated (via a non-linear relationship) with total event rainfall, but not significantly correlated with duration, mean intensity, or antecedent 2-day precipitation ( $API_2$ ). Mean interception during small events ( $\leq 35$  mm) was 17% ( $n = 135$  events) compared with only 7% for large events ( $> 35$  mm;  $n = 32$ ). Examining small temporal intervals within the largest and highest intensity events that would potentially trigger landslides revealed complex patterns of interception. The tropical forest canopy had little ~~or no~~ smoothing effect on incident rainfall during the largest events. During events with high wind speeds and/or moderate-to-high pre-event wetting, measured throughfall was occasionally higher than rainfall during large event peaks, demonstrating limited buffering. However, in events with little wetting and low-to-moderate wind speed, early event rainfall peaks were buffered by the canopy. As rainfall continued during most large events there was little difference between rainfall and throughfall depths. Comparing both rainfall and throughfall depths to conservative mean intensity–duration thresholds for landslide initiation, throughfall exceeded the threshold in 75% of the events in which rainfall exceeded the threshold for both wet and dry conditions. Throughfall intensity for all the 11 largest events (rainfall = 65-116 mm) plotted near or above the intensity-duration threshold for landslide initiation during wet conditions; five of the events were near or above the threshold for dry conditions. Soil moisture responses during a range of rainfall conditions in large events were heavily and progressively buffered at depths of 1 to 2 m, indicating that the time-scale of any short-term smoothing of peak rainfall inputs (i.e.,  $\leq 1$  h) has little ~~influence or~~ ~~no effect~~ on peak pore water pressure at depths where landslides would initiate. Given these findings, we conclude that canopy interception would have little effect on mitigating shallow landslide initiation during the types of monsoon rainfall conditions in this and similar tropical secondary forest sites.

25

**Key Words:** throughfall, rainfall, canopy interception, soil moisture, shallow landslides, intensity-duration thresholds, tropical secondary forest, Thailand

## 1 Introduction

Mechanisms of slope failure in relatively shallow soil mantles during rain events are generally well understood. Typically a positive pore water pressure develops just above a hydrologic discontinuity in the regolith causing an abrupt decline in shear strength and resultant rapid landslide (e.g., Sidle and Swanston, 1982; Harp et al., 1990; Fernandes et al., 1994; Kuriakose et al., 2008; [Sidle and Bogaard, 2016](#)). Alternatively, landslides have been known to occur due to an increase in soil weight and reduction in soil suction as soils wet during events (Sasaki et al., 2000; Lacerda, 2007; Godt et al., 2009; Yamao et al., 2016). In contrast, some interactions amongst vegetation, site hydrology, and slope stability are not as well understood. In particular, the role of canopy interception of precipitation has drawn considerable speculation with little supporting data.

~~Root systems of woody vegetation contribute significantly to the R~~reinforcement of potentially unstable slopes ~~by~~  
~~root systems of woody vegetation is well recognized, especially when roots anchor into stable substrate~~ (Gray and  
Megahan, 1981; ~~Riesterberg and Sovonick-Dunford, 1983; Schmidt et al., 2001; Schwartz et al., 2012~~Roering et al.,  
2003; Stokes et al., 2009; ~~Schwartz et al., 2012~~) as is the effect of vegetation management on root strength, ~~or enhance~~  
~~lateral reinforcement within the soil mantle (O'Loughlin and Ziemer, 1982; Schmidt et al., 2001; Schwartz et al., 2012).~~  
15 ~~When trees are cut, the root strength declines with time (Sidle, 1991). If forests then regenerate, the period of~~  
~~significantly reduced root strength ranges from about 3 to 20 years after harvesting (Ziemer, 1981; Sidle and Wu, 1999;~~  
~~Sidle et al., 2006; Imaizumi et al., 2008; DeGraff et al., 2012; Schwartz et al., 2013).~~ ~~In contrast to~~ However, if forest  
~~sites are converted to weak, shallow rooted agricultural species or plantations, root strength remains low indefinitely~~  
~~(Sidle et al., 2006; DeGraff et al., 2012).~~ While this mechanical reinforcement of shallow soil mantles by roots, ~~is well~~  
20 ~~recognized and has been successfully modelled (Sidle, 1992; Wu and Sidle, 1995; Dhakal and Sidle, 2003; Schwartz et~~  
~~al., 2013),~~ the effects of the presence or absence of trees on the hydrological processes of transpiration, interception,  
water redistribution, and subsequent pore pressure formation in the subsurface remain a topic of controversy, especially  
when related to shallow landslide initiation (Keim and Skaugset, 2003; Reid and Lewis, 2009; Ghestem et al., 2011;  
Greco et al., 2013; Dhakal and Sullivan, 2014).

25 Deep-rooted woody vegetation extracts soil water near potential failure planes during periods of high transpiration;  
however, such effects typically are not expected to augment slope stability during extended rainy periods when soils are  
already at field capacity, especially in temperate regions (Megahan, 1983; Sidle and Ochiai, 2006). In the tropics, where  
evapotranspiration rates are sustained year-round, the potential for modification of the soil moisture regime when trees  
are removed may be greater. Nevertheless, simulations of soil moisture in a Peninsular Malaysia rain forest indicate that  
30 evapotranspiration more significantly affects soil moisture during events preceded by dry conditions than events

preceded by wet conditions; it is during these wet periods that shallow landslides are more likely to occur (Sidle, 2005; Sidle et al., 2006).

Because forests intercept and evaporate rain water back to the atmosphere, less rainfall typically reaches the forest floor when canopies are intact (e.g., Rowe et al., 1999; Crockford and Richardson, 2000; Reid and Lewis, 2009; Ziegler et al., 2009; Kato et al., 2013). However, the effects of canopy interception are complicated by antecedent precipitation, wind, rainfall intensity and duration, and canopy structure (Xiao et al., 2000; Scott et al., 2003; Pypker et al., 2005; Germer et al., 2006; Kato et al., 2013). As such, canopy interception can vary greatly from event to event at a given site (Keim et al., 2004; Ziegler et al., 2009). Most studies show that the percentage of rainfall intercepted by tree canopies is most variable and highest for small events compared to larger events (Filoso et al., 1999; Keim et al., 2004; Germer et al., 2006; Reid and Lewis, 2009; Ziegler et al., 2009; Dhakal and Sullivan, 2014). In addition to interception of rain water, forest canopies have been reported to exert a buffering effect on short-term pulses of incident rainfall (Xiao et al., 2000; Keim and Skaugset, 2003; Keim et al., 2006). Using a stochastic representation of rainfall, canopy evaporation, and rainfall transfer through the canopy, Keim et al. (2004) showed that effective rain intensity during large events was reduced more for short duration events than for long duration events; during small events such differences with storm duration were not apparent.

Based on reports of interception and intensity smoothing in forest canopies, it has been advocated that canopy removal could lead to more intense pulses of rainfall infiltrating into forest soils and subsequently higher pore water pressures in the subsurface that could exacerbate landsliding (Rowe et al., 1999; Keim and Skaugset, 2003; Keim et al., 2004; Reid and Lewis, 2007, 2009). If this is true, then in addition to the delayed effect of root decline after tree removal, there could also be an immediate negative effect due to the lack of canopy interception and subsequent potential to increase pore water pressure. While attempts have been made to include canopy interception losses into subsurface hydrology (Keim et al., 2006) and landslide models (Wilkinson et al., 2002), it is not clear what the mechanistic effects are on slope stability. Most canopy interception studies have been conducted in temperate forests, but this information is especially needed in tropical rain forests to assess the possible effects of canopy removal on landslide initiation in these regions where management pressures are rapidly increasing.

The objectives of this research are three-fold: (1) evaluate rainfall interception by a secondary tropical forest canopy for a large number of monsoon events; (2) compare throughfall and rainfall rates to intensity–duration threshold relationships established for shallow landslides; and (3) determine the effect of canopy interception on the potential for soil water increases that could trigger landslides. One major question related to slope stability is to determine if canopy interception significantly reduces incident rainfall with respect to established conservative rainfall intensity-duration thresholds for landslide initiation. Another question relates to finding evidence that canopy interception significantly

affects event soil moisture dynamics to an extent that it would influence shallow landslide initiation. These questions are addressed within the context of a three-year field investigation in a disturbed, secondary hill dipterocarp forest stand in northern Thailand.

## 2 Site description

5 The Mae ('river' in Thai language) Sa Experimental Catchment, a headwater catchment of the Ping River, is located northwest of Chiang Mai city in northern Thailand (18°54'06.8"; 98°53'14.2"; Figure 1a). The 74.2 km<sup>2</sup> catchment is mountainous, with elevations ranging from 500 to 1400 m asl. The topography is characterized by steep (some exceeding 45°) slopes and narrow valleys. The geology of the catchment includes granites and gneiss, with some marble and limestone. Soils include Ultisols, Alfisols, and Inceptisols; soil depth typically exceeds 2 m. Land cover is primarily  
10 mixed secondary forests and scrublands (together approximately 80% of the land area), with ongoing conversion to intensive agriculture, especially tree crops, floriculture, and greenhouse operations (Figure 1b). These agricultural activities, in addition to ecotourism, support the economies of several small villages. Much of the development, including the building and maintenance of major roads, is located immediately adjacent to the Sa River and its tributaries. A few recent landslides have occurred within the upper catchment (personal observations); most are  
15 triggered by runoff, either from compacted road surfaces or intercepted subsurface flow along cutslopes, that is discharged onto fillslopes or unstable slopes below the road (e.g., Sidle et al., 2006)are associated with road runoff. Given the rapid revegetation of these tropical sites, it is difficult to detect older landslides.

The catchment is the site of ongoing investigations of hydrological and land-use change (Sidle and Ziegler, 2010; Bannwarth et al., 2014a,b; Ziegler et al., 2014a,b). Associated instrumentation includes 11 spatially-distributed rain gages  
20 and one stream gaging station that monitors discharge and turbidity at the mouth of the catchment at a sub-hourly time scale (Figure 1). Mean annual rainfall in the catchment varies from 1500 to 2000 mm y<sup>-1</sup>. The vast majority of the annual precipitation is delivered as intense rainfall (often exceeding 20 mm h<sup>-1</sup>) during the monsoon season between May and November. The catchment has a mean annual runoff ratio of approximately 30% (Ziegler et al., 2014b).

The throughfall experiment was conducted at station 429 (Figure 1b), which consists of a hydrometeorological  
25 tower that measures water and energy fluxes within an upland, dipterocarp forest. Trees are typically 10-16 m tall in the forest; tree trunk diameter ranges from 2-77 cm. Tree density in the plot is moderate: 127 trees within a 350 m<sup>2</sup> plot. Leaf Area Index ranges from 1.8-3.2, as determined at 117 point locations using fisheye digital photographs and Delta-T Hemiview software ([www.delta-t.co.uk](http://www.delta-t.co.uk)). The forest is secondary and is typically burned annually, but fire only affects the understory vegetation.

A total of 34 tree species are found in the plot, along with three types of bamboo: *Bambusa tulda* Roxb (Poaceae); *Bambusa pallida* (Poaceae); and *Dendrocalamus nudus* Pilg. (Poaceae). Dominant tree species are *Shorea obtusa* Wall. Ex Bl. (Dipterocarpaceae) and *Quercus kerrii* Craib var. *kerrii* (Fagaceae). Other important species include: *Lithocarpus polystachyus* (Wall. Ex A. DC.) Rehder. (Fagaceae); *Tectona grandis* L. f. (Verbenaceae); *Craibiodendron stellatum* (Pierre) W.W. Sm. (Ericaceae); *Cratogeomys formosum* (Jack) Dyer ssp. *Pruniflorum* (Kurz) Gogel. (Guttiferae, Hypericeae); *Dipterocarpus tuberculatus* Roxb. Var. *tuberculatus* (Dipterocarpaceae); *Gardenia sootepensis* Hutch. (Rubiaceae); *Pterocarpus marocarpus* Kurz (Leguminosae, Papilionoideae); *Shorea siamensis* Miq. var. *siamensis* (Dipterocarpaceae); and *Wendlandia tinctoria* (Roxb.) DC. subsp. *Orientalis* Cowan (Rubiaceae).

The soil at the site is an Ultisol with a thin (< 20 cm) brown A horizon underlain by a dark red B horizon that extends below a depth of two meters. Saturated hydraulic conductivity declines exponentially from the surface (~136 mm h<sup>-1</sup>) to approximately < 45 mm h<sup>-1</sup> at a depth of 25 cm; values at 1 m and 2 m are 1-2 mm h<sup>-1</sup> (n = 3 measurements for all depths; unpublished data, determined with a bore-hole permeameter). (n = 3 measurements; unpublished data). Bulk density does not change much over this depth range (1.0817-1.38 g cm<sup>-3</sup> for the surface, 1 m. and 2 m depths). Corresponding porosity for the three depths is 0.59, 0.52, and 0.48 (based on a particle density of 2.65 g cm<sup>-3</sup>). The decrease in saturated hydraulic conductivity is typical of that in other profiles found in southeast Asia (cf. Ziegler et al., 2004; 2006). Macropores and fissures, features that could influence preferential flow through the soil, were not abundant in the subsoil.

### 3 Methods

#### 3.1 Measurements

To assess real-time rainfall interception in the forest stand, a tipping-bucket rain gage was mounted on a meteorological tower at a height of 18 m, about 1 m above the tallest canopy trees (820 m asl), to measure incident rainfall (stations 429, Figures 1b). For our analyses we examined all events that occurred during the period from 6 May 2005 to 21 November 2007. To be considered an event, total rainfall during a period had to be ≥ 8 mm with no precipitation break > 4 h occurring. As such, we included a range of monsoon storms to assess interception losses for potential landslide-triggering events and those that were smaller.

Soil moisture was monitored at the soil surface and depths of 1 and 2 m in the same forest patch using Campbell Scientific (Logan UT, USA) CS-615 soil moisture probes, connected to a Campbell CR23x data logger. The probes were situated less than 10 m from the throughfall collection system under the forest canopy. Soil moisture measurements were recorded at 20 min intervals (note: these are instantaneous measurements, not means). Water

content reflectometer values recorded with the CS-615 were converted to volumetric water contents via sensor-specific calibration curves determined from manual samples collected within the soil profile at the time of installation and during subsequent periods of both wet and dry seasons. During the latter periods, manual samples were collected by augering holes to a 2 m depth near the probe site. Volumetric samples were collected with an AMS bulk density sampler. The calibration curve was determined via linear regression from the reflectometer (independent variable) and paired volumetric water content (dependent variable) data. Details of this calculation are provided by the manufacturer (<https://s.campbellsci.com/documents/us/manuals/cs616.pdf>).

Throughfall was collected in a system consisting of six 4-m long gutters radiating from a central tipping-bucket device that was installed under the canopy and secured at heights of 0.5-1.0 m above the ground at a slight angle ( $\leq 6^\circ$ ) to promote rapid drainage (Figure 2a); the angle was based on prior experience (Ziegler et al., 2009). Each collection gutter of the throughfall system was 43.5 mm wide, with a triangular-shaped channel and 25 mm vertical risers to reduce rain splash loss. All gutters drained into a large tipping-bucket to measure real-time throughfall response for comparison with incident rainfall. The volume of throughfall required to produce one tip was 230-240 cm<sup>3</sup> (0.22-0.23 mm). A dynamic calibration correction was then applied to account for differences in tip volume over the range of observed tipping rates (Calder and Kidd, 1978; Marsalek, 1981; Humphrey et al., 1997; Ziegler et al., 2009). These differences are caused by “splash” losses that occur as the tipping mechanism moves when rainfall drains from the funnel. This relationship was determined by draining known volumes of water through the tipping bucket system (mm tip<sup>-1</sup>) and recording the number of tips registered.

The volume of rainfall within a given time interval was divided by the total surface collection area of the entire gutter system, corrected for the angle of inclination (area = 1.044 m<sup>2</sup>), to calculate total throughfall for the interval. A correction was also applied to account for splash error occurring during high-intensity throughfall. Based on data collected from seven events using a paired tipping bucket rain gage and a throughfall system installed in an open area (i.e., both were used to measure rainfall), we observed that substantial splash loss occurred during high-intensity events. Total event rainfall depths between the devices could be achieved when calibrated tip volumes were increased 50-78% (via linear regression) during high-intensity periods of events (i.e., for rates of 5-12 tips per minute). We used this relationship to adjust the throughfall rates for high-intensity throughfall measured in this study.

We recognize several limitations in this correction: (1) the correction is based on limited data (unpublished); (2) some splash error probably also occurred at lower intensities (< 5 tips per minute), but the data set does not allow us to quantify it; and (3) the splash error associated with open rainfall may not be the same as that for throughfall, owing to different drop sizes and drop direction (both of which vary from event to event). Nevertheless, after applying this crude correction, the total event throughfall depths were within the ranges (relative to rainfall) expected for the range



of events measured (i.e., the  $C_i$  of increasingly large events approached a value of 0; see Figure 3a). Thus, we believe any residual errors due to splash (after correction) are minor; importantly, these errors would not change our final interpretations.

The time interval used to assess both rainfall and throughfall inputs via the respective tipping bucket devices was 1 min. Although the collector was kept in the same location during the three year study, it has an advantage over using conventional, movable, tipping-bucket rain gauges because it integrates throughfall response under much of the variable canopy structure (Ziegler et al., 2009). Spatial integration of throughfall is especially important in tropical forests where multi-tiered canopies create considerable variability in throughfall (Lloyd and Marques, 1988; Dykes, 1997; Konishi et al., 2006).

## 10 3.2 Calculations

Canopy interception ratio ( $C_i$ ) is calculated as

$$C_i = (RF - TH) RF^{-1} \quad [1]$$

where, RF is the incident rainfall during an event or a portion of the event (mm) and TH is the throughfall (mm) during the same period of time. Values of  $C_i$  approaching zero indicate no canopy interception for that event or period of the event.

As a conservative predictive measure for shallow landslides, a number of regional studies have generated the lowest thresholds for slope failure based on average rainfall intensity–duration relationships (e.g., Larson and Simon, 1993; Aleotti, 2004; Guzzetti et al., 2007; Dahal and Hasegawa, 2008) based on an earlier global concept developed by Caine (1980). Sidle and Ochiai (2006) modified Caine’s global intensity–duration threshold by removing some very short and very long (> 10 days) events that misrepresented rainfall – landslide initiation data. The resulting relationship is given as:

$$I = 13.58 D^{-0.38} \quad [2]$$

where I is the mean event intensity ( $\text{mm h}^{-1}$ ) and D is the duration of the event (h). To assess the effects of antecedent rainfall on the intensity–duration relationship, all of Caine’s (1980) data that included 2-day antecedent rainfall ( $API_2$ ) together with new data were plotted separately for  $API_2 \leq 20$  mm and  $API_2 > 20$  mm (Sidle and Ochiai, 2006). Two-day antecedent rainfall was used because this parameter correlated well with maximum piezometric response in unstable hollows (Sidle, 1992). The intensity–duration relationship developed for the events preceded by dry ( $\leq 20$  mm) antecedent conditions is the following (Sidle and Ochiai, 2006):

$$I = 19.99 D^{-0.38} \quad [3]$$

where I and D are as in Eq. 2. For the events preceded by wet antecedent conditions, the following relationship is used:

$$I = 12.64 D^{-0.49} \quad [4]$$

where I and D are as above. The global modified Caine threshold (Eq. 2) is lower than the dry condition threshold (Eq. 3) for all combinations of event intensity–duration. Furthermore, because the equations are based on duration, the modified Caine threshold is higher than the wet condition threshold (Eq. 4) for events exceeding 1-h duration. It should be noted that storms that exceed any of these thresholds will not necessarily trigger a landslide, rather it means that the very minimum rainfall conditions (i.e., average intensity and duration) for global landslides has been met.

## 4 Results

### 4.1 Canopy interception for all events

10 The maximum, minimum, and mean rainfall totals for the 167 recorded events during the three year study were 116.4, 8.1, and 24.3 mm, respectively. The corresponding totals for throughfall were 114.6, 5.7, and 21.3 mm, respectively. Event mean rainfall intensity (depth/duration) ranged from 0.5 to 88.0 mm h<sup>-1</sup>. The mean event intensity of all 167 events was 9.1 mm h<sup>-1</sup>. The duration of the 167 events ranged from about 9 min to 57 h, with a mean duration of about 7 h.

15 A total of 52, 59, and 56 events were monitored in 2005, 2006, and 2007 (Table 1). Mean event size ranged from 22-28 mm. Rainfall depths for the three years varied from 1149-1678 mm; the corresponding throughfall depth range was 1037-1450 mm (Table 1). Annual estimates of throughfall (fraction of rainfall) were 0.90, 0.86, and 0.87, respectively. We can only speculate that annual variations result from minor changes in canopy characteristics (based on observations) and differences in event rainfall characteristics. Further, we believe the inherent error in the calculation  
20 of throughfall for any one event is on the order of  $\pm 6\%$ . The differences in the yearly calculations are within this tolerance.

The three-year throughfall estimate was 0.88, which is near the higher end of values reported for forests in Southeast Asia (e.g., Sinun et al., 1992; Dykes, 1997; Konishi et al., 2006; Ziegler et al., 2009; Tanaka et al., 2015). The throughfall estimate may be slightly elevated because of the following reasons: (a) the stand was a recovering secondary forest (i.e.,  
25 lacking a multi-story canopy) with low LAI (1.8-3.2); (b) the event-based estimate does not include many very small events when canopy interception is expected to be high (i.e., we only report throughfall for events representing 78, 87, and 82% of incident rainfall entering the forest in 2005, 2006, and 2007; data not shown); and (c) under-catch of rainfall

above the canopy during windy conditions. Nevertheless, we believe the method provided reasonably accurate data for this type of analysis.

For the 167 events, throughfall ranged widely from 62 to 129% of rainfall (Figure 4). When expressed as a canopy interception ratio ( $C_i$ , Eq. 1), values ranged from -0.29 to 0.38, with the mean for all 167 events being 0.15 (Figure 3).

5 Throughfall was greater than incident rainfall during 9 events (Figure 34b), but only for four events was the difference greater than 6%, a value we consider to be approximately the uncertainty in the throughfall estimate. Most of the events where throughfall was greater than rainfall occurred in 2007 ( $n = 4$ ), with one and three occurring in 2005 and 2006, respectively (Figure 4b). We consider the four exceptionally low values (ranging from -0.18 to -0.24) to be outliers in this analysis that can partly be explained by the preferential channelling of intercepted water via canopy “pour points”  
10 to the collector (Konishi et al., 2006; Ziegler et al., 2009), as we cannot fully explain the nature of the very high throughfall that generated them (see discussion that follows). Both the boundary of uncertainty and the outliers are shown in Figure 4b.

A total of 35 events had  $C_i$  values  $\geq 0.25$  (Figure 3a). Most of these events were small-to-moderate in size (range = 8.7-32.0 mm; median = 12 mm), but rainfall depth in three events exceeded 30 mm. These 35 events ranged from 0.5  
15 to 7.4 h in duration; event intensities ranged from 2 to 37 mm h<sup>-1</sup>. Three events had mean intensities  $> 20$  mm h<sup>-1</sup>. The events also ranged greatly in antecedent moisture conditions:  $API_2$  ranged from 0.0 to 60 mm.

Except for four events with anomalously low  $C_i$  values, the overall tendency was for  $C_i$  to converge towards zero as total rainfall increased, particularly beyond 60 mm (Figure 3a). However, the relationship between  $C_i$  and total rainfall was not strong (non-linear regression;  $R^2 = 0.11$ ; significant at  $\alpha = 0.05$ ; Figure 3a). No meaningful relationship existed  
20 between  $C_i$  and  $API_2$ , event duration, or mean event intensity (Figures 3b-d). However, during longer events with short periods of high intensity, mean intensity would not be a good index to compare with  $C_i$ .

## 4.2 Canopy interception during large storms

Because many shallow landslides occur after a high-intensity burst of rainfall that follows an initial period of lower  
25 intensity rain (e.g., Okuda et al., 1979; Sidle and Swanston, 1982; Sidle and Chigira, 2004; Sidle and Bogaard, 2016), we focus mainly on the largest and longest events (Table 2). Eleven of the events summarized in Table 2 have total rainfall depths  $> 65$  mm; and one had a duration  $> 55$  h (event #1; total rainfall = 40 mm). The eleven large events have low  $C_i$  values (-0.01 to 0.11), indicating that most rainfall was converted into throughfall (open orange colored circles in Figure 3). The long-duration event #1 had a relatively high  $C_i$  value (0.14; triangle in Figure 3b).

30 Events #54 and #158 had  $C_i$  values  $\leq 0$  (Table 2). Event #54 was short-duration, high intensity with moderately high wind speed, while event #158 had relatively low intensity and wind speed, but high surface soil moisture prior to

the storm. The high intensity and moderately high wind speed during event #54 likely generated non-vertical rainfall that may have been underestimated in the gauge above the canopy and may have dislodged and transferred rainfall from proximate trees to the plot canopies and throughfall collector troughs. The slightly negative  $C_i$  (-0.01) during event #158 was likely affected by wet conditions (soil moisture in surface and 1 m depths were 0.45 and 0.42 g cm<sup>-3</sup>, respectively) at the onset of the storm.

Five events had canopy interception values ranging from 0.02-0.05 (#99, #156, #48, #56, #63; Table 2), indicating very limited canopy buffering during these large (65-116 mm) events. Durations and intensities of these five events were variable, ranging from 2.4 to 38.8 h and 5.3-41.6 mm h<sup>-1</sup>, respectively. High antecedent precipitation ( $API_2 = 11-22$  mm) and maximum wind speed (2.3-4.2 m s<sup>-1</sup>) occurred prior to and during four of these events, while event #63 had the driest antecedent conditions (Table 2).

The large event with greatest interception (#1;  $C_i = 0.14$ ) had the driest antecedent conditions ( $API_2 = 0$  mm) of all large events considered and very low maximum wind speed (1.9 m s<sup>-1</sup>). Because of these conditions and the relatively small total rainfall depth (40 mm), canopy interception was likely higher than that for the other large events (Figure 3b). We included event #1 as a “large” event because of its exceptionally long duration (> 56 h).

The remaining four large events with intermediate  $C_i$  values ranging from 0.09-0.11 (#115, #96, #85, and #26) all had relatively long durations (10.6-29.7 h), moderate intensities (2.4-7.4 mm h<sup>-1</sup>), and relatively low maximum wind speeds (2.2-2.7 m s<sup>-1</sup>) (Table 2). The main difference among these events was the much higher  $API_2$  value for event #96 (52 mm) compared to the other three events (0-1 mm).

Collectively, the variation in rainfall characteristics (e.g., duration, intensity) and other hydro-climatic phenomena (e.g., maximum wind speed,  $API_2$ ) demonstrate the inherent variability in (or the measurement of) canopy interception across a range of large events. In most cases it is difficult to pinpoint the key factor or combination of factors dictating  $C_i$  because meteorological conditions (rainfall intensity, wind speed) vary across time and space scales that our methodology does not measure. Again, we consider the error in the interception estimate to be on the order of 6%.

### 4.3 Throughfall and rainfall patterns during large storms

We also examined the pattern of incident rainfall versus throughfall during the six of the largest events (Figure 5). This sub-group includes the three events (#54, #56, and #156) with the highest intensities (36.9-45.6 mm h<sup>-1</sup>), short durations (1.8-2.8 h), and low  $C_i$  values (0.0-0.3). Three other events (#63, #99, and #158) had low intensities (2.0-6.8 mm h<sup>-1</sup>), long durations (16.4-38.8 h), and a range of  $C_i$  values (-0.01 to 0.05). For events #56, #99, and #156, throughfall exceeded rainfall during early large peaks, but was generally lower than rain intensity during latter parts of these events (Figures 5a,b,c). Short-duration event #54 was characterized by an initial burst of rainfall intensity of nearly

3 mm min<sup>-1</sup> (Figure 5d), which immediately translated into substantial throughfall despite the initially dry canopy (API<sub>2</sub> = 0; SM<sub>0m</sub> = 0.17; Table 2). Event #158 produced a complicated pattern of throughfall response – during the initial rain burst, throughfall exceeded rainfall, but after about a 5-h period of little precipitation, rainfall exceeded throughfall for the rest of the event (Figure 5f). Throughout most of event #63, rainfall exceeded throughfall (Figure 5e). The three largest events (#56, #99, and #156) exhibited minor canopy storage during early, low intensity rainfall; however, during large subsequent peaks (> 100 mm h<sup>-1</sup>, minutely rates) throughfall exceeded rainfall (Figures 5a,b,c). Moderate to high maximum wind speeds occurred during these three events (2.9-4.2 m s<sup>-1</sup>). The highest wind speed was associated with the largest rainfall event #56, during which throughfall depth was similar to rainfall depth (115-116 mm; C<sub>i</sub> = 0.02). In addition, wet canopy conditions preceding these three events are supported by high API<sub>2</sub> (11-22 mm) and associated surface soil moisture (0.40-0.48 g cm<sup>-3</sup>) values. During the early peaks of large events with relatively low intensity and wind speed (#63 and #158), more rain water was likely stored in the canopy (Figure 5e,f). The findings of early storage agree with the generally accepted idea that canopies store a larger proportion of rainwater during the early stage of events (e.g., Xiao et al., 2000; Zeng et al., 2000; Iida et al., 2012). While in five of these six large events rainfall exceeded throughfall when intensities were < 1.0 – 1.1 mm min<sup>-1</sup> and, typically, throughfall exceeded rainfall when intensities were > 1.1 mm min<sup>-1</sup>, this pattern was not consistently found in other storms.

#### 4.4 Soil moisture dynamics

We examined ~~vertical~~ the soil moisture dynamics near the soil surface and at depths of 1 m and 2 m under the forest canopy to ascertain the effects of canopy buffering on water movement in the soil to depths where shallow landslides may occur. Such a 1-dimensional assessment does not capture effects of complex topography which can influence flow pathways and hydrologic response at depth, rather it gives a general perspective on the potential of relatively homogeneous soils to attenuate soil infiltration. In particular, we focus on periods during events when the canopy exerts a maximum influence on short-term incident rainfall. Such canopy interception effects have been suggested to provide benefits to slope stability during large, landslide-producing events (Rowe et al., 1999; Keim and Skaugset, 2003; Reid and Lewis, 2009). We show soil moisture dynamics for the same six events used to assess intra-storm patterns of rainfall/throughfall (Figure 6), but also consider changes in the other large events (Table 2). The six events include a range of dry and wet antecedent moisture conditions (e.g., API<sub>2</sub> ranged from 0 to 22 mm). Corresponding surface, 1-m, and 2-m initial soil moisture ranged from 0.17-0.48 m<sup>3</sup> m<sup>-3</sup>, 0.34-0.42 m<sup>3</sup> m<sup>-3</sup>, and 0.33-0.37 m<sup>3</sup> m<sup>-3</sup>, respectively (Table 2; Figure 6). These values indicate large differences in pre-event soil moisture near the soil surface, but not at depths where landslides may initiate.

Events #56, #99, #156, and #158 are representative of relatively wet antecedent conditions. Initial surface soil moisture values for these events range from 0.40-0.48 (Figure 6). Throughfall infiltration into the soil produced peaks in surface soil moisture that lagged behind throughfall peaks by typically 20-60 min (Figure 6; note that soil moisture is measured every 20-min). In some cases, surface soil moisture increased  $> 0.1 \text{ m}^3 \text{ m}^{-3}$  during the event (e.g., events #56, #156, #158). The rainfall rates triggering these increases typically exceeded  $60\text{-}100 \text{ mm h}^{-1}$  for 20 min periods (Figure 6a,c,f). During event #99, two periods of rainfall resulted in corresponding peaks in surface soil moisture. For all four events, increases in soil moisture at the 1-m depth occurred 100-180 min after the onset of rainfall, or 30-90 min following the maximum rainfall and/or throughfall rate (Figures 6a,b,c,f).

During events with drier antecedent conditions (#54 and 63;  $API_2 = 0$  and surface soil moisture  $\leq 0.34 \text{ m}^3 \text{ m}^{-3}$ ), much greater wetting occurred in the surface soil (increases of  $0.14\text{-}0.27 \text{ m}^3 \text{ m}^{-3}$ ) compared with that during wetter antecedent conditions (Figures 6d,e). Event #54 was characterized by an initial burst of rainfall on dry soil ( $0.17 \text{ m}^3 \text{ m}^{-3}$ ), which rapidly elevated surface soil moisture over the next hour (Figure 6d). Event #63 consisted of two rainfall/throughfall peaks that produced corresponding peaks in surface soil moisture with short lags (20-40 min). This event also caused a small, but abrupt lagged increase in soil moisture at the 1-m depth just after the second rainfall peak, and a lagged and very minor increase in moisture at the 2 m depth. ~~whereas~~ Subsoil moisture during event #54 was unaffected during this shorter (1.8 ~~versus 10.6~~ h) event.

Soil moisture at the 2-m depth increased incrementally about 1-2 h after the final rainfall peaks of exhibited delayed and very minor increases in response to inputs of rainfall/throughfall during events #156 and #158, but then tapered off after about 3-6 h only (Figures 6c,f). Both events occurred under some of the wettest conditions observed (surface soil moisture =  $0.45\text{-}0.48 \text{ m}^3 \text{ m}^{-3}$ ). ~~Although the events had very different durations (2.4 versus 16.4 h), the increases at 2-m occurred more than an hour after the last rainfall peak, at which time most of the event rainfall had already occurred. The short-term bursts of intensity during the two largest~~ ~~It is not clear why the largest events~~ (#56, (116 mm and #99, 106 mm) generated only a very minor and lagged response of soil moisture at the 2 m depth. ~~falling in 2.8 h on wet soil) did not affect soil moisture at the 2-m depth.~~ Three other large events summarized in Table 2 produced soil moisture changes at 2 m (#96, #85, and #48). In all cases, the maximum observed change in soil moisture at 2-m was only on the order of  $0.03 \text{ m}^3 \text{ m}^{-3}$ .

Six of the eight large ~~All~~ events that increased soil moisture at the 2-m depth had very wet surface conditions ~~saturated to nearly saturated surface soils~~ (surface soil moisture values  $\geq 0.445 \text{ m}^3 \text{ m}^{-3}$ ; Table 2;  $Wetting_{max}$ ). Together with these wet surface conditions, total depth of throughfall (or rainfall) appeared to be more important than event intensity in propagating water fluxes to the 2 m depth. In contrast, the two events that only affected surface soil moisture had initial soil moisture values of  $0.17\text{-}0.29 \leq 0.34 \text{ m}^3 \text{ m}^{-3}$  ( $API_2 = 0.0$  for ~~both~~ ~~three~~ events; Table 2).

Throughfall depths during these drier antecedent conditions ranged from 34-81 mm. Event #54 had the highest average measured event intensity (45.6 mm h<sup>-1</sup>), but produced no increases in subsoil moisture. Intermediate of these responses, some events with initial soil moisture conditions ranging from 0.30-0.405 m<sup>3</sup> m<sup>-3</sup> produced soil moisture increases down to one meter depth for a variety of throughfall inputs (64-115 mm) occurring over 2.8 to 18.9 h (Table 2). Observed changes in soil moisture at 1 m were on the order of 0.02-0.08 m<sup>3</sup> m<sup>-3</sup>. Saturated or nearly saturated conditions in surface soils occurred during only two events (#156 and 158), both of which had high levels of soil moisture (0.45-48 m<sup>3</sup> m<sup>-3</sup>) prior to the events. Even during the largest events, soil moisture contents were well below saturation at the 1 m and 2 m depths (Table 2).

#### 4.5 Rainfall duration–intensity landslide thresholds

To further assess the potential of the monitored events to initiate shallow landslides, we compared incident rainfall and throughfall to three conservative intensity-duration landslide threshold relationships (Eqs. 2, 3, and 4). Considering all 167 events, regardless of antecedent rainfall, 37 rainfall events exceeded the threshold (Eq. 2) for potential landslide initiation, while throughfall from 30 events fell on or above this threshold (Figure 7a). This difference of seven events is associated with 10 events with rainfall > throughfall (positive canopy interceptions ranging from 0.03 to 0.34) and three events with throughfall > rainfall (negative C<sub>i</sub> ranging from -0.06 to -0.24). Five of these events had rainfall depths > 25 mm and six had intensities > 10 mm h<sup>-1</sup> (not shown). Only one had an API<sub>2</sub> value > 20 mm – event #80, 29 mm of rainfall during a period of 3.3 h (9 mm h<sup>-1</sup>) with a C<sub>i</sub> = 0.14 (not shown).

We segregated the 167 events into those preceded by dry and wet conditions to compare with Eqs. 3 and 4 (Figure 7b,c). A total of 120 events were preceded by dry conditions (API<sub>2</sub> ≤ 20 mm); 47 events were preceded by wet conditions (API<sub>2</sub> > 20 mm). For dry conditions, 16 and 12 rainfall and throughfall events, respectively, plotted above the corresponding threshold (Eq. 3). For the events where both rainfall and throughfall were above the threshold, durations did not exceed 3 h three hours and intensities were greater than 26.7 mm h<sup>-1</sup> (Figure 7b). Events where rainfall exceeded the threshold but throughfall did not were characterized by low to moderate rainfall depths (12-30 mm), appreciable antecedent rainfall (API<sub>2</sub> = 7-18 mm; except for one event with 0 mm), positive C<sub>i</sub> values (0.07-0.38), and short durations (0.3-1.7 h). In contrast, 12 and 9 rainfall and throughfall events, respectively, plotted above the threshold for wet conditions (Eq. 4; Figure 7c). The nine events with throughfall above the threshold were variable in length (0.42-11.8 h) and average storm intensity (4.7-54.1 mm h<sup>-1</sup>). The three events in which incident rainfall exceeded the threshold, but throughfall did not, were similar in canopy interception (C<sub>i</sub> = 0.20-0.23); duration (3.4-4.6 h); and event intensity (7.1-7.7 mm h<sup>-1</sup>). Rainfall depth for these events varied from 24-34 mm.

Nearly all of the 12 largest events (Table 2) plot on or above the threshold for wet conditions (Eq. 4; Figure 7d). The long-duration event #1 (56.6 h) plotted well below all thresholds (Figure 7d). Throughfall for three events (#54, #56, and #156) plotted well above the threshold for dry conditions demonstrating their potential for landslide generation, despite having relatively short durations (about 2-3 h) (Table 2; Figure 7d). Collectively, these comparisons show the limited potential of canopy interception to reduce the probability of landslide initiation during large annual storms (e.g., those listed in Table 2), particularly under wet antecedent conditions.

## 5. Discussion

Our estimated 3-year interception loss from a tropical secondary forest in northern Thailand based on 167 storms with total precipitation  $\geq 8$  mm was 12%. Interception losses for all individual events ranged from -29% to 38% and from -1% to 11% during the 11 largest storms. Overall, we found that events with larger total precipitation had lower rates of interception compared to smaller events (albeit weakly correlated), which agrees with most other studies (e.g., Filoso et al., 1999; Keim et al., 2004; Germer et al., 2006; Reid and Lewis, 2009; Bäse et al., 2012). Given the spatially distributed gutter system we employed to collect throughfall under this secondary forest stand, we believe our estimates are realistic within a measurement error of about 6%.

The measured interception losses in this tropical secondary dipterocarp forest are on the low side of most ranges reported for temperate and semi-arid canopies (e.g., Xiao et al., 2000; Iida et al., 2005; Reid and Lewis, 2007; Kato et al., 2013; Allen et al., 2014; Swaffer et al., 2014; Nanko et al., 2016). However, our values and variabilities are very similar to those reported in an open tropical Brazilian rainforest with many palm trees ( $10.2 \pm 5.6\%$ ) where similar magnitudes and numbers of storms were recorded (Germer et al., 2006). Interception losses in other native and secondary Amazonian forests have been reported in the range from about 6% to 22% (Lloyd and Marques, 1988; Elsenbeer et al., 1994; Filoso et al., 1999; Tobón Marin et al., 2000; Bäse et al., 2012; Zanchi et al., 2015). Several studies in both native and plantation forests in southeast Asia that experience monsoon storms reported similarly low interception losses in the range of about 7 to 20% (Sinun et al., 1992; Dykes, 1997; Konishi et al., 2006; Ziegler et al., 2009; Tanaka et al., 2015). Given that the preponderance of the research on canopy interception has been conducted in temperate and arid or semi-arid environments, it is not surprising that this secondary tropical forest in Thailand has relatively lower canopy interception during intense monsoon events.

These slightly lower values of interception we measured may reflect a combination of factors that vary among individual events. Stable isotope differences between throughfall and rainfall during low intensity events show the complexity of pre-event storage on contributions to throughfall in a conifer forest in the Oregon Cascades, but indicate



that the release of residual water stored in canopies may be significant (Allen et al., 2014). Given the wet, humid conditions at the Thailand site, it is possible that pre-event canopy wetness may have augmented throughfall during large events as evidenced by high soil moisture and/or API<sub>2</sub>. In other large events, it appears that the higher maximum 20-min wind speed was a factor in dislodging water from within plot and surrounding canopies to restrict interception.

5 Many studies have shown that storage of water in tree canopies is reduced under windy conditions (e.g., Hutchings et al., 1988; Llorens and Gallart, 2000; Xiao et al., 2000; Keim and Skaugset, 2003; Kato et al., 2013). While wind can increase evaporation and dry the canopy during storms (e.g., Kelliher et al., 1992; Xiao et al., 2000), it may increase measured throughfall by increasing canopy drip, changing the angle of incoming rainfall, and capturing wind-blown rain from adjacent trees (Xiao et al., 2000; Ziegler et al., 2009). Non-vertical rainfall, which is common during windy

10 conditions associated with many monsoon storms, is also often under-recorded by small, standard gauges (Kamph and Burges, 2010). Finally, canopy “drip points” or “pour points” may develop, where intercepted water is channelled preferentially to the collector (Konishi et al., 2006; Ziegler et al., 2009); these points are dynamic and change as canopies develop and environmental conditions change during the storm ~~(e.g., wind) change~~. Throughfall measurements during four moderate-sized events with anomalously low C<sub>i</sub> values (indicated as outliers in Figures 3 and 4) may also have been

15 affected by a combination of these factors.

It should be noted that we did not account for losses due to interception of litter cover on the forest floor, which in some ecosystems can be significant (e.g., Kelliher et al., 1992; Gerrits et al., 2010). Nevertheless, such interception would be little affected by forest removal and subsequent regeneration, unless significant site disturbance occurred during logging. Thus, the ultimate influence of changes in litter interception on pore water pressures at soil depths that

20 could trigger landslides are expected to be minimal (Dhakal and Sullivan, 2014).

We found no strong evidence of canopy smoothing of incident rainfall on throughfall during large events; such intensity smoothing has been suggested as a factor in moderating peak pore water pressures within soil mantles, thus reducing the risk of shallow landslides (Rowe et al., 1999; Keim and Skaugset, 2003; Keim et al., 2004; Reid and Lewis, 2009). The throughfall data we present from northern Thailand are unique in terms of number of events and temporal

25 resolution, which allows us to better assess peak responses related to rainfall inputs. During six of the largest events that would potentially trigger landslides, throughfall intensity actually exceeded rainfall intensity during the largest storm peaks in five of the six events (Figure 5). In event #63, which had the lowest peak intensity of these six events, peak rainfall intensity exceeded peak throughfall intensity. Although the peak intensities of rainfall and throughfall differed amongst events, little ~~no~~ smoothing (i.e., flattening) of throughfall peaks relative to rainfall peaks was evident. While a

30 few studies have alluded to intensity smoothing by forest canopies on throughfall regime (Xiao et al., 2000; Keim et al., 2004, 2006; Nanko et al., 2016), only one presented specific evidence of intra-storm smoothing (i.e., flattening) of

rainfall peaks (Xiao et al., 2000). The only storm that Xiao et al. (2000) presented was small (13 mm) and low-intensity compared to the monsoon events in northern Thailand, and only throughfall under one oak was assessed. In some cases, the smoothing effects are derived through modelling (Keim et al., 2004, 2006), but even in these cases, the effects on pore water pressure in the substrate during events were low.

5 Furthermore, only soil moisture in the surface horizon was responsive to individual rainfall peaks during the same six large events shown in Figure 5. Soil moisture increases at the depth of 1 m were highly dampened, lagged the rainfall peak by nearly an hour or more, and never approached saturation (Figure 6). At the depth of 2 m, near where shallow landslides would typically initiate in this terrain, ~~only very only two of these large storms produced very dampened and~~ minor increases in soil moisture ~~( $\leq 0.01\text{-}0.02\text{ m}^3\text{ m}^{-3}$ ) were recorded during five of the six large events and;~~ no soil  
10 moisture response at 2 m was ~~measured~~~~recorded during event #54~~~~in the four other large events (Figure 6)~~. This dampened or lack of soil moisture response at depth, shows that even during events with higher peak rainfall versus throughfall inputs (e.g., event #63; Figure 5e), the impact on pore water pressure at the depth of a potential failure plane would be ~~small~~~~insignificant~~ (Figure 6e). We did not assess the effect of slope shape on soil moisture response. It is possible that higher canopy interception during small events just preceding (e.g., one to a few days) a large storm, could mitigate soil moisture response at depth to some extent.  
15

The comparisons of incident rainfall and throughfall to established intensity-duration landslide threshold relationships allow us to compare the potential for rainfall-initiated landslides in secondary tropical forests (throughfall measurements) versus cutover or converted sites (incident rainfall). Most large events had intensity-duration relationships that fell above global thresholds for potential landslide initiation during wet conditions. The three events  
20 that greatly exceeded the most conservative threshold (i.e., for dry conditions) were very short-duration (2-3 h), high average intensity (37-46 mm h<sup>-1</sup>) storms. Considering that peak rainfall intensity was generally not much greater than corresponding peak throughfall intensity during large events and that soil moisture response at depths where landslides may initiate ( $\geq 1$  m) did not respond rapidly to peak rainfall inputs, it appears that canopy interception would have little  
~~or no~~ influence on mitigating pore water accretion at depths where shallow landslides typically occur in this secondary  
25 tropical forest. Furthermore, similar behaviour of incident rainfall and throughfall during individual events with respect to intensity-duration thresholds for landslide initiation support our conclusion that canopy interception at the site has negligible influence on landslides for the rainfall conditions we observed.

## 6 Limitations and recommendations

30 There are some important limitations to our methods. In a prior study, Ziegler et al. (2009) compared the same troughs used herein with several movable tipping bucket gauges, finding no statistical difference between the two approaches.

5 However, in the previous study, total event precipitation was examined, not minutely changes occurring over the course of storms. We caution that the trough method may create a somewhat confusing signal because the area-integrated pattern of throughfall, in which the records are delayed as water captured in the trough flows through the trough, compared to an individual rain gauge placed above the canopy. Nevertheless, at high rates of throughfall, through flow would be more efficient. Additionally, estimates of both throughfall and rainfall have errors. Measurements by tipping-bucket rain gauges installed above a canopy are affected by turbulent exchange at this interface, and wind affects rainfall catch (e.g., Kamph and Burges, 2010). As mentioned before, the throughfall troughs had a large associated splash error during high-intensity events. While our attempt to correct the splash error resulted in reasonable total event values, individual minutely values could still have substantial errors; these would affect the time series we compare with measured rainfall (Figure 6). However, such limitations are inherent in most all studies reported in the literature (albeit typically not articulated).

15 In future experiments, we urge researchers to minimize uncertainties by: (a) using troughs that are deeper to minimize splash loss; (b) collect ambient rainfall in more than one location, preferably gauges positioned just above ground level with appropriate wind shields to minimize wind effects; (c) collect more throughfall values, potentially employing other types of collection devices to help interpret the measurements; and (d) perform a rigorous assessment of splash losses to facilitate error correction (again our splash error data are few). We also encourage researchers to spend time in the forest plots during events, recording the various types of phenomena that may affect the capture of throughfall over time, to help with the interpretation of data. Here, we re-emphasize that throughfall reaching the forest floor is highly variable in space and time. Multi-stored canopies can create wet and dry zones below them, which change over the course of a storm with respect to variable wind direction, changes in rainfall phenomena (rate, drop size), and changes in canopy wetness (e.g., Konishi et al., 2006). The oddities apparent in the data of some of our recorded storms (e.g., higher throughfall than rainfall during some periods, but not others) may possibly be related to inadequacies in our error correction; however, these may be realistic. For example, they may result because wind-driven rain is captured by portions of a large tree and then channelled directly to the throughfall trough (as was documented at the prior study site; Ziegler et al., 2009).

25 We believe that these uncertainties do not undermine the integrity of our conclusions. While the uncertainties may prevent us from producing a high-precision budget of the portion of rainfall converted to throughfall at minutely scales, they do allow us to address the primary goals of this investigation, which are to assess whether secondary tropical canopies intercept sufficient rainwater during large storms to mitigate landslide initiation compared to open areas.

## 76 Conclusions

Our examination of the effects of canopy interception in a secondary dipterocarp forest of northern Thailand on the potential for shallow landslide initiation revealed some interesting findings. Compared to temperate and semi-arid forests, throughfall in our secondary forest plot was relatively high owing to wind effects (transferring rain water from surrounding trees and causing under-catch of precipitation), wet and humid antecedent conditions, and preferential channelling of canopy drip into collection troughs. Nevertheless, our throughfall measurements are in line with many values reported from both native and secondary forests in Amazonia and elsewhere southeast Asia.

Few studies have reported intra-storm comparisons of incident rainfall and throughfall at temporal resolutions that could be used to assess effects on shallow landslide initiation (i.e.  $\leq 1$  h). Most studies that have noted smoothing effects of canopy interception on incident rain intensity (that would potentially reduce the risk of landslides) seem to focus on the lowering of throughfall relative to rainfall peaks or how the perceived flattening (smoothing) of throughfall peaks can be represented in models (Keim and Skaugset, 2003; Keim et al., 2006; Reid and Lewis, 2009; Greco et al., 2013; Nanko et al., 2016). None of these studies conducted across a range of temperate forest cover types show any physical evidence that smoothing of rainfall peaks by forest canopies during large storms lowered soil moisture or pore pressures at depths that would reduce landslide susceptibility. In contrast, the few studies that have presented throughfall versus rainfall inputs at small time intervals for large events, show little or no evidence of flattening of peaks (Iida et al., 2012; Safeeq and Fares, 2014; Dhakal and Sullivan, 2014). Only the study of Xiao et al. (2000) measured intensity smoothing effects of single oak canopy during a relatively small storm (13 mm). Although our throughfall results from many large and intense monsoon events in northern Thailand were affected by instrumental errors (common in all studies of this type), our findings indicate clearly show that these secondary tropical forest canopies have relatively small/negligible smoothing effects on incident rainfall peaks.

Although numerous studies have noted the reduction of throughfall relative to incident rainfall, there is little information on effects of the reduction of rainfall under canopies on soil moisture dynamics. Our findings show that soil moisture response is quite dampened or even non-responsive at depths where potential failure planes exist. These data coupled with our analysis of mean rain intensity – duration thresholds that have been used to estimate the lower global limit of rainfall conditions that may trigger shallow landslides, show that both rainfall and throughfall for the 11 largest events exceeded or reached thresholds for wet antecedent conditions and plotted very closely on the intensity – duration graph (Figure 6d). As such, there is little evidence that canopy interception in this secondary tropical forest has a significant any mitigating effect on shallow landslides. The possibility of intercepted rainfall during a small event prior to a large storm that would alter antecedent soil moisture, could potentially increase pore water pressure response at depth to a small degree. More likely and much better documented anthropogenic causes of landslide increases in

similar tropical environments include root strength deterioration following timber harvesting, forest conversion, or swidden (e.g., Harper, 1993; Sidle et al., 2006; DeGraff et al., 2012), roads and trails (e.g., Douglas et al., 1999; Chappell et al., 2004; Sidle and Ziegler, 2012), and possibly the effects of increased antecedent soil moisture following clearing or conversion of forest cover (Sidle et al., 2006; [Sidle and Bogaard, 20016](#)).

5

*Acknowledgements.* Gratitude is expressed to Aaron Pruitt for his assistance in data analysis. We also acknowledge Tom Giambelluca, Mike Nullet, and Jefferson Fox for their efforts in the establishment of the Mae Sa experiment site. [The constructive suggestions and critiques by two external reviewers and the Editor are much appreciated.](#)

## 10 References

- Aleotti, P.: A warning system for rainfall-induced shallow failures, *Eng. Geol.*, 73, 247-265, 2004.
- Allen, S. T., Brooks, J. R., Keim, R. F., Bond, B. J., McDonnell, J. J.: The role of pre-event canopy storage in throughfall and stemflow by using isotopic tracers, *Ecohydrology*, 7, 858-868, 2014.
- Bannwarth, M. A., Sangchan, W., Hugenschmidt, C., Lamers, M., Ingwersen, J., Ziegler, A. D., and Streck, T.:  
15 Pesticide transport simulation in a tropical catchment by SWAT, *Environ. Pollut.*, 191, 70-79, 2014a..
- Bannwarth M. A., Hugenschmidt C., Sangchan W., Lamers M., Ingwersen J., Ziegler A. D., Streck T.: Simulation of stream flow components in a mountainous catchment in northern Thailand with SWAT, using the ANSELM calibration approach, *Hydrol. Process.*, 29, 1340-1352. doi: 10.1002/hyp.10268, 2014b.
- Bäse, F., Elsenbeer, H., Neill, C., and Krusche, A.V.: Differences in throughfall and net precipitation between  
20 soybean and transitional tropical forest in the southern Amazon, Brazil, *Agric., Ecosys. And Environ.*, 159, 19-28, 2012.
- Caine, N.: Rainfall intensity-duration control of shallow landslides and debris flows, *Geograf. Ann.*, 62A, 23-27, 1980.
- Calder, I.R., Kidd, C.H.R., 1978: A note on the dynamic calibration of tipping-bucket gauges. *J. Hydrol.* 39, 383–386.
- Chappel, N.A., Douglas, I., Hanapi, J.M., and Tych, W.: Sources of suspended sediment within a tropical catchment  
25 recovering from selective logging, *Hydrol. Process.*, 18, 685-710, 2004.
- Crockford R., and Richardson, D.: Partitioning of rainfall into throughfall, stemflow and interception: Effect of forest type, ground cover and climate, *Hydrol. Process.*, 14, 2903-2920, 2000.
- Dahal, R. K., and Hasegawa, S.: Representative rainfall thresholds for landslides in the Nepal Himalaya, *Geomorphology*, 100, 429-443, 2008.

- DeGraff, J. V., Sidle, R. C., Ahmad, R., and Scatena, F. N.: Recognizing the importance of tropical forests in limiting rainfall-induced debris flows, *Environ. Earth Sci.*, 67, 1225-1235, doi: 10.1007/s12665-012-1650-8, 2012.
- Dhakal A. S., and Sullivan, K.: Shallow groundwater response to rainfall on a forested headwater catchment in northern coastal California: implications of topography, rainfall, and throughfall intensities on peak pressure head generation, *Hydrol. Process.*, 28(3), 446-463, 2014.
- ~~Dhakal A. S., and Sidle, R. C.: Long-term modeling of landslides for different forest management practices, *Earth Surf. Process. Landforms*, 28, 853-868, 2003.~~
- Douglas, I., Biden, K., Balamurugan, G., Chappell, N.A., Walsh, R.P.D., Greer, T., and Sinun, W.: The role of extreme events in the impacts of selective tropical forestry on erosion during harvesting and recovery phases at Danum Valley, Sabah, Phil. Trans. Roy. Soc. London, 354, 1749-1761, 1999.
- Dykes, A.P.: Rainfall interception from a lowland tropical rainforest in Brunei, *J. Hyrol.*, 200, 260-279, 1997.
- Elsenbeer, H., Cassel, D.K., and Zuniga, L.: Throughfall in the terra firme forest of western Amazonia, *J. Hydrol.*, 32, 30-45, 1994.
- Fernandes, N. F., Netto, A. L. C., and Lacerda, W. A.: Subsurface hydrology of layered colluvium mantles in unchanneled valleys – south-eastern Brazil, *Earth Surf. Proc. Landforms*, 19(7), 609-626, 1994.
- Filoso, S., Willimas, M.R., and Melack, J.M.: Composition and deposition of throughfall in a flooded forest archipelago (Negro River, Brazil), *Biogeochem.*, 45, 169-195, 1999.
- Germer S., Elsenbeer, H., and Moraes, J. M.: Throughfall and temporal trends of rainfall redistribution in an open tropical rainforest, south-western Amazonia (Rondônia, Brazil), *Hydrol. Earth Syst. Sci.*, 10, 383-393, 2006.
- Gerrits A.M.J., Pfister, L., and Savenije, H.H.G.: Spatial and temporal variability of canopy and forest floor interception in a beech forest, *Hydrol. Processes*, 24, 3011-3025, 2010 doi: 10.1002/hyp.7712
- Ghestem, M., Sidle, R. C., and Stokes, A.: The influence of plant root systems on subsurface flow: Implications for slope stability, *BioScience*, 61(11), 869-879, 2011.
- Godt, J. W., Baum, R.L., and Lu, N.: Landsliding in partially saturated materials, *Geophys. Res. Lett.*, 36, L02403, doi: 10.1029/2008GL035996, 2009.
- Gray, D. H., and Megahan, W. F.: Forest vegetation removal and slope stability in the Idaho Batholith, Res. Pap. INT-271, 23 p., For. Serv., U.S. Dep. of Agric., Ogden, Utah, 1981.
- Greco R., Comegna, L., Damiano, E., Guida, A., Olivares, L., and Picarelli, L.: Hydrological modelling of a slope covered with shallow pyroclastic deposits from field monitoring data, *Hydrol. Earth Syst. Sci.*, 17, 4001-4013, 2013.
- Guzzetti, F., Peruccacci, S., Rossi, M., and Stark, C. P.: The rainfall intensity-duration control of shallow landslides and debris flows: an update, *Landslides*, 5, 3-17, 2008.

- Harp, E. L., Wells II, W. G., and Sarmiento, J. G.: Pore pressure response during failure in soils. *Geol. Soc. Am. Bull.*, 102(4), 428-438, 1990.
- Harper, S.B.: Use of approximate mobility index to identify areas susceptible to landsliding by rapid mobilization to debris flows in southern Thailand. *J. Southeast Asia Earth Sci.*, 8, 587-596, 1993.
- 5 [Humphrey, M.D., Istok, J.D., Lee, J.Y., Hevesi, J.A., Flint, A.L., 1997: A new method for automated dynamic calibration of tipping-bucket rain gauge. \*J. Atmos. Oceanic Tech.\* 14, 1513–1519.](#)
- Hutchings, N.J., Milne, R., and Crowther, J.M.: Canopy storage capacity and its vertical distribution in a Sitka spruce canopy, *J. Hydrol.*, 104, 161-171, 1988.
- Iida, S., Tamai, K., Shimizu, T., Nobuhiro, T., Kabeya, N., and Shimizu, A.: Characteristics of process of rainfall  
10 interception in a Japanese cedar forest within the Tsukuba Experimental Watershed, Japan, *Kanto Forestry Res.*, 63(2), 121-124.
- Iida, S., Tanaka, T., and Sugita, M.: Change of interception process due to the succession from Japanese red pine to evergreen oak, *J. Hydrol.*, 315, 154-166, 2005 (in Japanese with English abstract).
- Imaizumi, F., Sidle, R. C., and Kamei, R.: Effects of forest harvesting on occurrence of landslides and debris flows in  
15 steep terrain of central Japan, *Earth Surf. Process. Landforms*, 33, 827-840, 2008.
- Kamph, S. K., Burges, S. J.: Quantifying the water balance in a planar hillslope plot: Effects of measurement errors on flow prediction, *J. Hydrol.*, 380, 191-202, 2010.
- Kato, H., Onda, Y., Nanko, K., Gomi, T., Yamanaka, and Kawaguchi, S.: Effect of canopy interception on spatial variability and isotopic composition of throughfall in Japanese cypress plantations, *J. Hydrol.*, 504, 1-11, 2013.
- 20 Keim, R. F, and Skaugset, A. E.: Modelling effects of forest canopies on slope stability, *Hydrol. Process.*, 17, 1457-1467, 2003.
- Keim, R. F., Skaugset, A. E., Link, T. E., and Iroumé, A.: A stochastic model of throughfall for extreme events, *Hydrol. Earth Syst. Sci.*, 8, 23-34, 2004.
- Keim, R. F., Tromp-van Meerveld, H. J., and McDonnell, J. J.: A virtual experiment on the effects of evaporation and  
25 intensity smoothing by canopy interception on subsurface stormflow generation, *J. Hydrol.*, 327, 352-364, 2006.
- Kelliher, F. M., Whitehead, D., Pollock, D. S.: Rainfall interception by trees and slash in a young *Pinus radiata* D. Don stand, *J. Hydrol.*, 131, 187-204, 1992.
- Konishi, S., Tani, M., Kosugi, Y., Takanashi, S., Mohd Md S., Abdul Rahim N., Niiyama, K., and Okuda, T.: Characteristics of spatial distribution of throughfall in a lowland tropical rainforest, Peninsular Malaysia, *Forest  
30 Ecol. Mgmt.*, 224, 19-25, 2006.

- Kuriakose, S.L., Jetten, V. G., van Westen, C. J., Sankar, G., and van Beek, L. P. H.: Pore water pressure as a trigger of shallow landslides in the Western Ghats of Kerala, India: some preliminary observations from an experimental catchment, *Physical Geog.*, 29, 374-386, 2008.
- Lacerda, W. A.: Landslide initiation in saprolite and colluvium in southern Brazil: Field and laboratory observations, *Geomorphology*, 87, 104-119, 2007.
- 5 Larsen, M. C., and Simon, A.: A rainfall intensity-duration threshold for landslides in a humid-tropical environment, Puerto Rico, *Geografiska Ann.*, 75A, 13-23, 1993.
- Llorens, P., and Gallart, F.: A simplified method for forest water storage capacity measurement, *J. Hydrol.*, 240, 131-144, 2000.
- 10 Lloyd, C. R., and de O Marques, A.: Spatial variability of throughfall and stemflow measurements in Amazonian rainforest, *Agric. For. Meteorol.*, 42, 63-73, 1988.
- ~~Marsalek, J., 1981: Calibration of the tipping-bucket raingage. *J. Hydrol.* 53, 343-354.~~
- Megahan, W. F.: Hydrologic effects of clearcutting and wildlife on steep granitic slopes in Idaho, *Water Resour. Res.*, 19(3), 811-819, 1983.
- 15 Nanko, K., Onda, Y., Kato, H., and Gomi, T.: Immediate change in throughfall spatial distribution and canopy water balance after heavy thinning in a dense mature Japanese cypress plantation, *Ecohydrology*, 9, 300-314, 2016.
- Okuda, S., Ahida, K., Gocho, Y., Okunishi, K., Sawada, T., and Yokoyama, K.: Characteristics of heavy rainfall and debris hazard, *J. Nat. Disaster Sci.*, 1(2), 41-55, 1979.
- ~~O'Loughlin, C. L., and Ziemer, R. R.: The importance of root strength and deterioration rates upon edaphic stability in steepland forests, in Carbon Uptake and Allocation in Subalpine Ecosystems as a Key to Management, 70-78, Proc. IUFRO Workshop, Oregon State Univ., Corvallis, OR, 1982.~~
- 20 Pypker, T. G., Bond, B. J., Link, T. E., Marks, D., and Unsworth, M. H.: The importance of canopy structure in controlling the interception loss of rainfall: Examples from a young and old-growth Douglas-fir forest, *Agric. Forest Meteorol.*, 130, 113-129, 2005.
- 25 Reid, L. M., and Lewis, J.: Rates and implications of rainfall interception in a coastal redwood forest, *USDA Forest Serv. Gen. Tech. Rep. PSW-GTR-194*, 107-117, 2007.
- Reid, L. M., and Lewis, J.: Rates, timing, and mechanisms of rainfall interception loss in a coastal redwood forest, *J. Hydrol.*, 375, 459-470, 2009.
- ~~Riesterberg, M. M., and Sovonick-Dunford, S.: The role of woody vegetation in stabilizing slopes in the Cincinnati area, Ohio, *Geol. Soc. Am. Bull.*, 94, 506-518, 1983.~~
- 30



- Roering, J., Schmidt, K. M., Stock, J. D., Dietrich, W. E., and Montgomery, D. R.: Shallow landsliding, root reinforcement, and the spatial distribution of tress in the Oregon Coast range, *Can. Geotech. J.*, 40, 237-253, 2003.
- Rowe, L. K., Marden, M., and Rowan, D.: Interception and throughfall in a regenerating stand of kanuka (*Kunzea ericoides* var. *ericoides*), East Coast region, North Island, New Zealand, and implications for soil conservation, *J. Hydrol. N.Z.*, 5 38, 29-48, 1999.
- Safeeq, M., and Fares, A.: Interception losses in three non-native Hawaiian forest stands, *Hydrol. Process.*, 28, 237-254, 2014.
- Sasaki, Y., Fuijii, A., and Asai, K.: Soil creep process and its role in debris slide generation – field measurements on the north side of Tsukuba Mountain in Japan, *Eng. Geol.* 56, 163-183, 2000.
- 10 Schmidt, K. M., Roering, J. J., Stock, J. D., Dietrich, W. E., Montgomery, D. R., and Schaub, T.: The variability of root cohesion as an influence on shallow landslide susceptibility in the Oregon Coast Range, *Can. Geotech. J.*, 38, 995-1024, 2001.
- Schwartz, M., Cohen, D., and Or, D.: Spatial characterization of root reinforcement at stand level: Theory and case study, *Geomorphology*, 171-172, 190-200, 2012.
- 15 Schwartz M., Giadrossich, E., and Cohen, D.: Modeling root reinforcement using a root-failure Weibull survival function, *Hydrol. Earth Syst. Sci.*, 17, 4367-4377, 2013.
- Scott R., Watts, C., Payan, J., Edwards, E., Goodrich, D. Williams, D., and Shuttleworth, W.: The understory and overstory partitioning of energy and water fluxes in an open canopy, semiarid woodland, *Agric. Forest Meteor.*, 114, 127-139, 2003.
- 20 ~~Side, R. C.: A conceptual model of changes in root cohesion in response to vegetation management, *J. Environ. Qual.*, 20(1), 43-52, 1991.~~
- Side, R. C.: A theoretical model of the effects of timber harvesting on slope stability, *Water Resour. Res.*, 28, 1897-1910, 1992.
- Side, R. C.: Influence of forest harvesting activities on debris avalanches and flows, in M. Jakob and O. Hungr (eds.) 25 *Debris Flow Hazards and Related Phenomena*, 345-367, Springer-Praxis, Heidelberg, 2005.
- Side, R. C., and Swanston, D. N.: Analysis of a small debris slide in coastal Alaska, *Can. Geotech. J.*, 19, 167-174, 1982.
- Side, R. C., and Wu, W.: Simulating the effect of timber harvesting on temporal and spatial distribution of shallow landslides, *Z. für Geomorphologie*, 43, 185-201, 1999.
- Side, R. C., and Chigira, M.: The July 20, 2003, landslides and debris flows in southern Kyushu, Japan. *Eos, Trans.,* 30 *Am. Geophys. Union* 85(15), 145-151, 2004.
- Side, R. C., and Ochiai, H.: Landslides: Processes, Prediction, and Land Use, *Am. Geophysical Union, Water Resour.*

- Monogr. No. 18, AGU, Washington, D.C., 312 p., 2006.
- Sidle R. C., and Ziegler, A. D.: Elephant trail runoff and sediment dynamics in northern Thailand, *J. Environ. Qual.*, 39, 871–881. DOI: 10.2134/jeq2009.0218, 2010.
- Sidle, R.C., and Ziegler, A.D.: The dilemma of mountain roads. *Nature Geosci.* 5: 437-438, 2012.
- 5 [Sidle, R.C., and Bogaard, T.A.: Dynamic earth system and ecological controls of rainfall-initiated landslides. \*Earth-Sci. Rev.\* 159: 275-291. doi: 10.1016/j.earscirev.2016.05.013, 2016.](#)
- Sidle R. C., Ziegler, A. D., Negishi, J. N., Abdul Rahim N., Siew, R., and Turkelboom, F.: Erosion processes in steep terrain – Truths, myths, and uncertainties related to forest management in Southeast Asia, *Forest Ecol. Mgmt.*, 224, 199-225, 2006.
- 10 Stokes A., Atger, C., Bengough, A. G., Fourcaud, T., and Sidle, R. C.: Desirable plant root traits for protecting natural and engineered slopes against landslides, *Plant Soil*, 324, 1-30, 2009.
- Sunin, W., Meng, W.W., Douglas, I., and Spencer, T.: Throughfall, stemflow, overland flow and throughflow in the Ula Segama rain forest, Sabah, Malaysia, *Phil. Trans. Royal Soc., Series B*, 335, 389-395, 1992.
- Swaffer, B.A., Holland, K.L., Doody, T.M., and Hutson, J.: Rainfall partitioning, tree form and measurement scale: a  
15 comparison of two co-occurring, morphologically distinct tree species in a semi-arid environment, *Ecohydrology*, 7, 1331-1344, 2014.
- Tanaka, N., Levia, D., Igarashi, Y., Nanko, K., Yoshifuji, N., Tanaka, K., Tantasirin, C., Suzuki, M., and Kumagai, T.: Throughfall under a teak plantation in Thailand: a multifactorial analysis on the effects of canopy phenology and meteorological conditions. *Int. J. Biometeorol.*, 59, 1145-1156, 2015.
- 20 Tobón Marin, C., Bouten, W., and Sevink, J.: Gross rainfall and its partitioning into throughfall, stemflow and evaporation on intercepted water in four forest ecosystems in western Amazonia, *J. Hydrol.* 237, 40-57, 2000.
- Wilkinson, P. L., Anderson, M. G., and Lloyd, D. M.: An integrated hydrological model for rain-induced landslide prediction, *Earth Surf. Process. Landforms*, 27, 1285-1297, 2002.
- ~~Wu, W., and Sidle, R. C.: A distributed slope stability model for steep, forested basins, *Water Resour. Res.*, 31, 2097-  
25 2110, 1995.~~
- Xiao, Q., McPherson, E. G., Ustin, S. L., Grismer, M. E., and Simpson, J. R.: Winter rainfall interception by two mature open-grown trees in Davis, California, *Hydrol. Process.* 14, 763-784, 2000.
- Yamao, M., Sidle, R. C., Gomi, T., and Imaizumi, F.: Characteristics of landslides in unwelded pyroclastic flow deposits, southern Kyushu, Japan, *Nat. Hazards Earth Syst. Sci.*, 16, 617-627, doi:10.5194/nhess-16-617-2016, 2016.
- 30 Zeng, N., Shuttleworth, W., and Gash, J. H. C.: Influence of temporal variability of rainfall on interception loss. Part I. Point analysis, *J. Hydrol.*, 228, 228-241, 2000.

- Zanchi, F.B., Waterloo, M.J., Tapia, A.P., Alvarado Barrientos, M.S., Bolson, M.A., Luizão, F.J., Manzi, A.O., and Dolman, A.J.: Water balance, nutrient and carbon export from a heath forest catchment in central Amazonia, Brazil, *Hydrol. Process.*, 29, 3633-3648, 2015.
- Ziegler, A.D., Giambelluca, T.W., Tran, L.T., Vana, T.T., Nullet, M.A., Fox, J.M., Vien, T.D., Pinthong, J., Maxwell, J.F., Evett, S. 2004. Hydrological Consequences of Landscape Fragmentation in Mountainous Northern Vietnam: Evidence of Accelerated Overland Flow Generation. *J. Hydrol.*, 287, 124–146, 2004.
- Ziegler, A.D., Negishi, J.N., RC Sidle, R.C., S Noguchi, S., Abdul Rahim Nik. Impacts of logging disturbance on saturated hydraulic conductivity in a tropical forest in peninsular Malaysia. *Catena*, 67, 89-102, 2006.
- Ziegler, A. D., Giambelluca, T. W., Nullet, M. A., Sutherland, R. A., Tantasarin, C., Vogler, J. B., and Negishi, J. N.: Throughfall in an evergreen-dominated forest stand in northern Thailand: Comparison of mobile and stationary methods, *Agric. Forest Meteorol.*, 149, 373-384, 2009.
- Ziegler, A. D., Sidle, R. C., Phang, V. X. H., and Wood, S. H.: Bedload transport in SE Asian streams – uncertainties and implications for reservoir management, *Geomorphology*, 227, 31-48. doi: 10.1016/j.geomorph.2014.01.015, 2014a.
- Ziegler, A. D., Benner, S. G., Tantasarin, C., Wood, S. H., Sutherland, R.A., Sidle, R. C, Jachowski, N. R. A., Lu, X., Snidvongs, A., Giambelluca, T.W., and Fox, J.M.: Turbidity-based sediment monitoring in northern Thailand: hysteresis, variability, and uncertainty. *J. Hydrol.*, 519, 2020-2039, doi: 10.1016/j.jhydrol.2014.09.010, 2014b.
- Ziemer, R. R.: Roots and the stability of forested slopes, in *Erosion and Sediment Transport in Pacific Rim Steeplands*, T. R. H. Davies and A. J. Pearce (eds.), 343-361, IAHS, Christchurch, New Zealand, 1981.

Figure 1: Site map of the Mae Sa experiment site in northern Thailand. Panel (a) shows the catchment location in Thailand, the topography and major stream channels. Panel (b) shows major land covers in the Mae Sa catchment including hillslope and plantation agriculture (AG, 23%), greenhouse agriculture (GH, 7%), urbanized or peri-urban areas (U, 8%), and forest cover with various degrees of disturbance (F, 62%). Grid cell dimensions are 2 x 2 km.

5 Rectangles demarcate hydro-meteorological measurement sites. Streamflow, total suspended solids, particulate organic carbon, and particulate organic nitrogen were measured at the stream gage station 434. Rainfall is measured at all other numbered hydro-meteorological stations (rectangles). The throughfall investigation in this paper was conducted at station 429, where rainfall, throughfall, and soil moisture were monitored.

Figure 2: (a) Stationary collector with six collection troughs (gutters). (b) Schematic of stationary gauge tipping bucket mechanism (located inside the collector base); the inset shows the dimensions for the collection troughs (gutters).

10

Figure 3. (a) Canopy interception ratio ( $C_i$ ) versus total event rainfall for all monitored events. The non-linear regression curve describes the tendency for  $C_i$  to approach 0 as total rainfall increases. (b)  $C_i$  plotted with respect to event duration (h). (c)  $C_i$  plotted against 2-day antecedent precipitation ( $API_2$ ). (d)  $C_i$  plotted with respect to mean event intensity ( $\text{mm h}^{-1}$ ). The open circles and triangle in all panels refer to the 11 large and one long storm investigated in detail (Table 2).

15 The four events that plot below the dashed line are considered outliers.

Figure 4. (a) Comparison of throughfall to runoff depths for the three years of study (2005-2007). (b) Throughfall fraction of rainfall for events ranging in depths of 8 to 116 mm, during the three-year study. Uncertainty of any throughfall measurement is estimated to be  $\pm 6\%$  (indicated in the figure for the case of throughfall = rainfall). The four values labelled “outliers” are exceptionally high estimates for which we cannot completely explain the possible errors.

20

Figure 5. Five-minute running means of throughfall (solid) and rainfall (hashed) for the six largest storms (Table 2).

Figure 6. Volumetric soil moisture response at three depths during the six largest events (Table 2). Events with initial surface soil moisture (0 cm)  $\geq 0.40 \text{ m}^3 \text{ m}^{-3}$  are considered to have wet antecedent conditions: #56 (116 mm of rainfall); #99 (111 mm); #156 (89 mm); and #158 (78 mm). The other two events, #54 (81 mm) and #63 (77 mm), are associated with drier antecedent moisture conditions ( $\leq 0.30 \text{ m}^3 \text{ m}^{-3}$ ). All data are plotted as 20-minute aggregated values. Soil moisture curves in all panels correspond to surface (thin line), 1-m (hashed line), and 2-m (thick line) depths (key shown in panel a). The thickness of each rainfall bar has no meaning; each represents a 20-min value.

25

Figure 7. Comparison of event rainfall (open circles) and throughfall (solid circles) intensity-duration relationships with modified Caine thresholds for shallow landslide initiation. (a) All 167 events are plotted against the general threshold

(Eq. 2). (b) Events associated with dry conditions ( $API_2 < 20\text{mm}$ ) are plotted against the threshold defined by Eq. 3. (c) Events associated with wet conditions ( $API_2 \geq 20\text{mm}$ ) are plotted against the threshold defined by Eq. 4. (d) The 12 largest storms summarized in Table 2 are plotted against all thresholds (Eqs. 2-4). Some apparently missing throughfall data points plot behind their paired corresponding rainfall value.

5

**Table 1. Number of events sampled each year and corresponding rainfall and throughfall totals.**

Year	Events	Mean depth (mm)	Rainfall (mm)	Throughfall (mm)	Throughfall (-)
2005	52	22	1149	1037	0.90
2006	59	28	1678	1450	0.86
2007	56	22	1235	1078	0.87
total	167	24	4062	3564	0.88

Mean depth refers to the mean rainfall depth of the 52, 58, or 56 events in a given year (2005, 2006, 2007). Throughfall is listed as both a depth and a fraction of total rainfall.



Table 2. Characteristics of 12 large/long rainfall events considered in this analysis.

<u>Event</u>	<u>RF</u>	<u>TF</u>	<u>C<sub>i</sub></u>	<u>D</u>	<u>I</u>	<u>API<sub>2</sub></u>	<u>Wind</u>	<u>SM (0m)</u>	<u>SM (1m)</u>	<u>SM (2m)</u>	<u>Wetness (0m)</u>	<u>Wetness (1m)</u>	<u>Wetness (2m)</u>	<u>Wetting<sub>max</sub></u>
-	(mm)	(mm)	(-)	(h)	(mm h <sup>-1</sup> )	(mm)	(m s <sup>-1</sup> )	(m <sup>3</sup> m <sup>-3</sup> )	(m <sup>3</sup> m <sup>-3</sup> )	(m <sup>3</sup> m <sup>-3</sup> )	(-)	(-)	(-)	(m)
56	116.44	114.66	0.02	2.8	41.6	11	4.2	0.40	0.34	0.33	0.89	0.81	0.70	2
99	111.14	106.07	0.05	16.4	6.8	16	3.7	0.45	0.41	0.37	0.87	0.81	0.78	2
156	88.54	85.48	0.03	2.4	36.9	22	2.9	0.48	0.41	0.37	1.02	0.83	0.82	2
85	78.04	69.43	0.11	19.3	4.1	1	2.2	0.44	0.41	0.37	0.89	0.81	0.70	2
63	77.07	75.54	0.02	38.8	2.0	0	2.3	0.34	0.40	0.36	0.82	0.82	0.79	2
96	70.95	63.65	0.10	29.7	2.4	52	2.7	0.46	0.41	0.37	0.87	0.83	0.80	2
158	69.76	70.67	-0.01	16.4	4.2	6	1.8	0.45	0.42	0.37	0.98	0.83	0.81	2
48	65.02	62.81	0.03	12.2	5.3	15	3.0	0.45	0.39	0.36	0.87	0.83	0.83	2
115	78.64	70.10	0.11	10.6	7.4	1	2.2	0.30	0.34	0.33	0.80	0.70	0.68	1
26	70.22	63.97	0.09	18.9	3.7	0	2.2	0.34	0.40	0.37	0.84	0.81	0.78	1
54	81.28	80.87	0.00	1.8	45.6	0	3.2	0.17	0.34	0.33	0.77	0.65	0.68	0
1	39.91	34.13	0.14	56.6	0.7	0	1.9	0.29	0.34	0.33	0.72	0.64	0.68	0

5 RF is rainfall; TF, throughfall; C<sub>i</sub>, canopy interception; D, duration; I, mean intensity (RF/D), API<sub>2</sub>, two-day antecedent precipitation index; Wind, maximum 20-min wind speed (recorded above the canopy); SM<sub>xm</sub>, soil moisture measured at 0-, 1- and 2-m depths; Wetness 0 and 1 m, is fraction of pore space filled with water (red highlighted values indicate the two storms where saturation occurred in the surface soil or was nearly achieved; and Wetting<sub>max</sub>, the maximum soil depth (m) at which wetting during the event was measured by the soil moisture probes. The first 11 events are largest recorded (based on depth); event #1 is the longest (ranked only 23<sup>rd</sup> in size). Events are ranked by Wetting<sub>max</sub>, then rainfall depth.



Event	RF (mm)	TF (mm)	Gi -	D (h)	I (mmh <sup>-1</sup> )	API <sub>2</sub> (mm)	Wind (ms <sup>-1</sup> )	SM(0m) (m <sup>3</sup> ·m <sup>-2</sup> )	SM(1m) (m <sup>3</sup> ·m <sup>-2</sup> )	SM (2m) (m <sup>3</sup> ·m <sup>-2</sup> )	Wetting <sub>max</sub> —(—)
156	88.54	85.48	0.03	2.4	36.9	22	2.9	0.48	0.41	0.37	2
85	78.04	69.43	0.11	19.3	4.1	1	2.2	0.44	0.41	0.37	2
96	70.95	63.65	0.10	29.7	2.4	52	2.7	0.46	0.41	0.37	2
158	69.76	70.67	-0.01	16.4	4.2	6	1.8	0.45	0.42	0.37	2
48	65.02	62.81	0.03	12.2	5.3	15	3.0	0.45	0.39	0.36	2
56	116.44	114.66	0.02	2.8	41.6	11	4.2	0.40	0.34	0.33	1
99	111.14	106.07	0.05	16.4	6.8	16	3.7	0.45	0.41	0.37	1
115	78.64	70.10	0.11	10.6	7.4	1	2.2	0.30	0.34	0.33	1
26	70.22	63.97	0.09	18.9	3.7	0	2.2	0.34	0.40	0.37	1
54	81.28	80.87	0.00	1.8	45.6	0	3.2	0.17	0.34	0.33	0
63	77.07	75.54	0.02	38.8	2.0	0	2.3	0.34	0.40	0.36	0
1	39.91	34.13	0.14	56.6	0.7	0	1.9	0.29	0.34	0.33	0





Figure 1




Figure 2




Figure 3







Figure 4


a) Event #56




b) Event #99




c) Event #156



d) Event #54



e) Event #63



f) Event #158






Figure 5


**a) Event #56**




**b) Event #99**




**c) Event #156**




**d) Event #54**



**e) Event #63**




**f) Event #158**




**Figure 6**


(a) All events



(b)  $API_2 < 20$  mm (dry)



(c)  $API_2 > 20$  mm (wet)



(d) Large events




Figure 7