

Reply to reviewers [paper hess–2017–2]

Temperature signal in suspended sediment export from an Alpine catchment

Anna Costa, Peter Molnar, Laura Stutenbecker, Maarten Bakker, Tiago A. Silva, Fritz Schlunegger, Stuart N. Lane, Jean–Luc Loizeau, Stéphanie Girardclos

We thank the reviewers and the Editor for their helpful reviews. We have analysed their suggestions and we have revised the manuscript accordingly. We report in the following our response to each specific comment.

Editor

Dear authors, following up on the reviews received for your manuscript it appears that both referees rate your contribution as highly interesting and relevant for the HESS reader community. The two referees (and I do agree) see much potential for increasing the readability and structure of your paper – please follow their quite detailed suggestions for improvement in this respect. One referee sees some need for clarification related to the conceptualization and results of the ice model (essentially due to the fact that no glacier retreat is modelled). You may also consider to develop on this more specific aspect in your revised version.

We thank the Editor for the feedbacks. We have edited the manuscript to increase its readability. We have followed the suggestions of the reviewers and we have added to the revised manuscript results related to the following four main aspects:

- (1) sensitivity analysis of our snowmelt model parameters;
- (2) quantification of the possible overestimation of icemelt rates due to fact that glacier retreat is not modelled;
- (3) analysis of the effects of discontinuous SSC sampling on trend and jumps detected during the observation period;
- (4) estimation of potential impacts of gravel mining activities on SSC rise detected in mid–1980s.

Anonymous Referee #1

General comments

This manuscript focuses on the role of climatic forcing in sediment production and transport in a large Alpine catchment. It applies a degree–day model to explain changes in suspended sediment concentrations resulting from hydro–climatic activation of sediment sources within the study catchment. This represents an interesting approach that has been implemented with consideration of processes influencing rates of sediment supply from the range of catchment sediment sources. The manuscript is well written and describes clearly the modelling approach. There is scope for potential re–structuring of the manuscript, which contains some repetition and is over–long in places. Nonetheless, such changes amount to only a minor level. Overall, this is a technically sound and interesting study that deserves publication.

We thank Referee #1 for this helpful review. We have analysed her/his suggestions and we report in the following our response.

Specific comments

1) Lines 20–35, page 2: Description of catchment sediment sources. Given the number of figures in the manuscript, I suggest the authors remove the overly simplistic schematic in Fig. 1 as it adds little beyond that which is available from the summary in the text.

- 1) We agree with Referee #1 and we removed Fig. 1 from the revised manuscript.

2) Lines 20–25, page 4: The discussion of anthropogenic impacts in the catchment mentions gravel mining operations along the main channel and tributaries. Such direct disturbance of the channel could increase suspended sediment supply. Can this effect be discounted completely as a factor contributing to the observed trend in suspended sediment concentration (Fig 7c)?

- 2) We are aware that, for short periods after river bed disturbance, gravel extraction may cause local releases of fine sediment from the river bed. However, this process is unlikely to affect the suspended sediment load and balance at the outlet of the basin over seasonal and annual timescales. This is confirmed by the volume of gravel extracted along the main Rhone River and along tributaries, available from 1989. Annual volumes of gravel extracted, expressed as the difference from the average over the period 1989–2014, do not show any clear correlation with mean annual suspended sediment concentration

(Fig. 1). In agreement with Fig. 1, the coefficient of determination between the two variables is very low ($R^2 = 0.08$). Therefore, we conclude that, although gravel mining data for comparing the periods before and after mid-1980s are not available, gravel extraction is unlikely to play a significant role in the suspended sediment concentration rise observed in mid-1980s.

In the revised manuscript, we report these comments in Sect. 6.2 where we discuss other anthropogenic factors that could have potentially affected SSC dynamics but most likely did not play a relevant role in the SSC rise occurred in mid-1980s (i.e. river channelization, hydropower reservoirs).




Fig. 1: Mean annual suspended sediment concentration (SSC) and total annual volume of gravel extracted, expressed as difference from the average over the period 1989–2014.

3) Lines 10–15, page 6: Table 1 summarises some of the information given in Section 4 Data Description. Moreover, Section 3.3 Calibration and Validation also contains some description of the datasets used. To reduce repetition, can section 4 be shortened or consolidated? Perhaps a shortened descriptive summary of the datasets could be provided alongside Table 1 before introducing the models.

3) We agree with Referee #1. We reduced Sect. 4 and we changed Table 1, which now contains the list of the analysed variables, their source and the spatial and temporal resolution adopted in this analysis.

4) Lines 30–35, page 10: Could the use of fixed interval sampling (twice per week) for suspended sediment concentrations (SSC) influence the interpretation of trends during the observation period? The absence of continuous measurements (i.e. turbidity) or stage-triggered sampling may result in under-estimation of mean annual SSC because elevated but short-duration peaks in flow and SSC are less likely to be captured by fixed interval sampling. For this reason, the potential effect of the reported increase in direct rainfall on snow-free surfaces ('effective rainfall') on SSC could be underestimated because such events are short duration compared to the longer duration ice and snowmelt effect on SSC. This deserves consideration when evaluating the relative contributions of rainfall, snow and icemelt (page 18) to observed trends in SSC.

4) We thank Referee #1 for this constructive comment. We agree that discontinuity in the sampling procedure may indeed influence the interpretation of trends in suspended sediment concentration, and we agreed that this point deserved further investigation. Therefore, we estimate this potential effect by considering total daily, basin-averaged values of the hydroclimatic variables SM, IM, ER and Q only on days corresponding to SSC-measurement days. We compare these new time series ("SSC-measurement days") with the original ones ("all days") by computing the cumulative distribution functions and by testing the equality of mean monthly and annual values before and after mid-1980s. We consider only values greater than zero when computing the cumulative distribution function, to avoid the influence of many zeros. Although extremely high and low values are missed by the sampling method, cumulative distribution functions of SM, IM, ER and Q on SSC-measurement days and on all days result very similar (Fig. 2). This indicates that, although measurements of SSC are collected at a fixed interval, the sampling is representative of the process. In addition, results of the statistical tests on mean monthly and mean annual values of all analyzed hydroclimatic variables are unchanged. This confirms that our main results are not significantly influenced by the discontinuity of the SSC sampling method.

The description of this sampling effect has been added to the revised manuscript in the discussion Section 6.3 including the new figure below.




Figure 2: Cumulative distribution functions of total daily basin-averaged SM (a), IM (b), ER (c) and Q (d), computed on all days and only on days corresponding to SSC-measurements. Only positive values of SM, IM and ER are included in the computations.

5) Sections 5 & 6: I recommend merging the Results and Discussion. The Results section contains some elements of discussion (e.g. lines 20–25, page 15, on climate patterns), while in multiple locations within the Results section the authors write ‘see discussion’. The integration of Results and Discussion could produce a more coherent paper that presents findings and their interpretation in relevant sub-sections. For example, the discussion of snow and icemelt modelling in terms of previously reported melt factors (lines 10–25, page 20) would fit logically with the presentation of the calibration results (section 5.1).

5) We agree that sections of results and discussion can be organized in a more coherent way. We moved the discussion on snowmelt and icemelt models to the results as well as the trend/jump detection analysis.

In the revised manuscript the discussion section now only has two main points:

- (1) a discussion of the potential impact of neglecting glacier dynamics, where we demonstrate and discuss why although we are not including glaciers dynamics, we are not significantly overestimating icemelt rates (Sect. 6.1); and
- (2) a discussion of other potential anthropogenic factors on our results, such as river channelization, hydropower reservoirs regulation and gravel mining activities, which we argue are not likely to have a significant role in the SSC rise detected in mid-1980s (Sect. 6.2). As possible anthropogenic impact, we consider also the discontinuity of the SSC sampling procedure following the discussion presented at point n.4 and we also discuss the climatic signal on fine sediment dynamics in the context of a heavily regulated catchment.

6) Section 6.2 (lines 15–35, page 21): The discussion of future climate change effects on the sediment regime should be shortened and focus mainly on the point about the value of a more process-based approach. The paper offers no evidence based on future change simulations, so should limit speculative discussion in this area.

6) We agree that the discussion on climate change projections in Switzerland was too extended. However, as suggested by Referee #1, we would like to focus on the importance of adopting a more process-based approach when analysing the effects of climate change on suspended sediments. Therefore, in the revised manuscript we consistently reduced the text but we kept this point present in the discussion in Sect. 6.2.

Anonymous Referee #2

In this manuscript the increase in suspended sediments observed in the Rhone river after 1987 (compared to the period 1960–198) is statistically compared to hydro-climatic factors as temperature, rainfall, discharge, snow cover, snow melt and ice melt. Interpolated meteorological products and satellite observation of snow cover were used to model snow cover, snow and ice melt for the Rhone basin with degree-day melt models. The meteorological and the modelled data were then statistically compared to measured discharge and measured sediment supply. From this comparison, the authors suggest that the observed changes in the suspended sediment concentration is mainly due on the one hand to a reduced extent and duration of snow cover leading to larger areas exposed to erosive rainfall and on the other hand to an increase in icemelt in the summer months.

This study represents a valuable contribution for understanding the impact of climatic data on sediment transport. The presented approach is interesting and innovative and the results and analysis are meaningful. The study is also well-documented. Overall the manuscript is well written but substantial improvement can be made on the structure of the paper. This would avoid many repetitions throughout the text. Especially the discussion part needs to be built more on the results of the studies itself to be less speculative (see main comments below). Furthermore, I have some concern on the assumption to not model glacier retreat in the ice model, as this can lead to overestimation of the ice melt. There is therefore a need to further discuss the impact of the simplicity of the ice model on the results (see main comments below). The issues presented in this manuscript are in the scope of HESS and as my comments concern substantial restructuration of the manuscript and further development in the analysis of the ice model results, I would support its publication after major revisions.

We thank Referee #2 for these valuable suggestions. We implemented his/her comments in the revised manuscript as described in the following.

Main comments:

Goal formulation (p. 3, l. 22–25): One of the goals of the study (aim c) is to assess the future climate changes and hydropower operation impacts. This is only discussed later in the discussion, but only in a speculative way as the analysis itself does not include any hydropower data or future climate scenario. As it is formulated in the aims of the study, I would expect future climate scenario being taken into account and hydropower operation being explicitly analyzed. Instead the observations on this impact are only speculative (see for example discussion p. 21, l. 16–33). I would therefore suggest the authors to reformulate their study aims and also to minimize the weight they give to this topic in the discussion as it is only speculative.

We agree with Referee #2. We have reformulated the aims of the study by removing point c (“(c) to discuss the results in the context of future climate change and hydropower operation impacts on sediment budgets.”). However, we partially kept the discussion on future climate change investigations to highlight the importance of applying a more process based approach when investigating the impacts of climate change on suspended sediment dynamics (Sect. 6.2).

Overall structure:

The overall structure of the paper leads to many repetitions throughout the text.

1) The main problem is that parts of the discussion can be found in the methods, results, and discussion sections. This makes the text repetitive and the information hard to find. The authors should define a clear structure for their manuscript. They should decide if they want the results and discussion together in one section, meaning a discussion following each results description, or if they want 2 distinct sections (which I personally would recommend), meaning that the result section only describe the results and all discussion and interpretation of the results is moved to discussion section.

We agree with Referee #2 that the structure of the manuscript can be improved. In the revised manuscript, we moved large part of the discussion to the results sections (Sect. 5.1, 5.2, and 5.3) and we discuss separately only two specific aspects (see response to Referee no. 1)

2) The data are described first in the method section and then in the data description section. This is redundant. Best would be to shortly describe the data before the method section.

2) We agree that data description is somewhat redundant. In the revised manuscript we describe all the hydroclimatic variables that we analyse (T, P, Q, SSC, SM, SCF, IM and ER) in Sect. 3 and we list them in Table 1. In Sect. 4 (Data Description) we present all the datasets that we use in this analysis.

3) Section 3.1, snow model:

Several parameters that play an important role in the degree day approach (as SD, Trs, Tsm) are set before calibration. A sensitivity analysis of the impacts of these parameters would be interesting to assess their impact on the results and give them more credit.

3) We agree with Referee #2. We performed a sensitivity analysis on TRS, TSM and K_{snow} as the key parameters of the snow model. We analysed parameters sensitivity in two ways:

- (1) we analyze the relative change of some of the goodness of fit measures adopted during the calibration (TSS, NS RMSE) as function of the parameters perturbation, by perturbing one single parameter at the time (Fig. 3);
- (2) we estimate the influence of these three parameters on the trends and the jumps that we identified over the period 1975–2015 for the hydroclimatic variables simulated with the model: SM, IM, ER, SCF. For this purpose, we apply statistical tests for equality of mean annual and monthly values before and after mid–1980s after perturbing one parameter at the time within physically reasonable ranges of values.

Results show that the snowmelt factor k_{snow} is the most sensitive parameter (Fig. 3). For k_{snow} between 1.6 and 5.6 mm day⁻¹ °C⁻¹, the relative reduction of TSS and NS results respectively lower than 10% and 15% (Fig. 3). Although reducing the snowmelt factor k_{snow} below 2.6 mm day⁻¹ °C⁻¹ increases RMSE by almost 40%, it results in incrementing RMSE only by 0.06 units. For T_{SM} varying within the range of $-2 \div 2$ °C, TSS and NS decrease less than 10%. While the change in RMSE results larger similarly to the case of k_{snow} (Fig. 3). The effect of T_{RS} is even more negligible, with relative changes of goodness of fit measures within 5.5% (Fig. 3).




Figure 3: Parameter perturbation analysis on: (a) snowmelt factor k_{snow} , (b) threshold temperature for the onset of melt T_{SM} , (c) rain-snow threshold temperature T_{RS} . The relative change [%] of TSS, NS and RMSE are shown for each parameter.

In agreement with the results of our analysis (Sect. 5.3), a statistically significant increase of mean annual IM and ER and a statistically significant decrease of mean annual SCF after mid-1980s are detected for all selected parameter values. Similarly, mean annual SM shows a decreasing tendency after mid-1980s for all parameters selections, and in about 50% of the parameters selections statistical tests even show a statistically significant drop in mean annual SM. The increases in mean monthly ER in June and July are identified respectively in more than 90% and in 100% of the cases. Similarly, all parameters sets show an abrupt rise in IM after mid-1980s for all spring and summer months (May–August).

This confirms that by perturbing the parameters of the snow model within reasonable ranges, the overall results of our analysis do not change. This is also depicted in Fig. 4. The differences in mean monthly values of IM and ER between the period after and before mid-1980s are shown for multiple values of each individual parameter. The confidence interval depicted in Fig. 4 is artificially built by selecting, for each month, the highest and the lowest values of the confidence interval (5% significance level) among all parameter sets. By comparing Fig. 4 with Fig. 9c and Fig. 9d of the revised manuscript, it is possible to see that by changing parameters within reasonable ranges would not change significantly the results of our analysis.

These results are presented in the Supplementary Material of the revised manuscript (“S1. Sensitivity Analysis on Snow Model parameters”).




Figure 4: Sensitivity analysis on monthly differences between the periods after and before the year-of-change (1987–2015 and 1975–1986) for IM (left) and ER (right). Box plots represent monthly differences for all selections of parameters within given ranges of values: $1.6 \text{ mm day}^{-1} \text{ }^{\circ}\text{C}^{-1} \leq k_{\text{snow}} \leq 5.6 \text{ mm day}^{-1} \text{ }^{\circ}\text{C}^{-1}$ (a,b); $-2^{\circ}\text{C} \leq T_{\text{SM}} \leq 2^{\circ}\text{C}$ (e,f); $-1^{\circ}\text{C} \leq T_{\text{RS}} \leq 3^{\circ}\text{C}$ (c,d). Grey shaded areas represent the widest confidence interval among all selections of the parameters.

4) Ice model:

In my opinion the most critical point of the method is the simplicity of the ice melt simulation. The ice model uses a simple degree day routine and do not considers changes in glacier areas, which might be consequent over a time period of ca 50 years. The authors claim “Temporal dynamics of glacier coverage are not accounted for — ice accumulation, glacier retreat and ice movement are disregarded. The reduction of Alpine glaciers for the period 1950–2000 was estimated to be within the 10 range 500– 1000 m (Hoelzle, 2003; Oerlemans, 2005), while our effective climate grid resolution is 2_2 km, i.e. the retreat is considerably lower than the grid resolution of climatic inputs. The consideration of ice dynamics would therefore add a degree of complexity that our spatial resolution cannot take advantage of” (p. 11, l. 8–12). I disagree with this argumentation. The grid size of the meteorological product is surely important for the modelling of glacier retreat, but many studies used these meteorological dataset for modelling changes in glacier mass balance in the Alps and proved that the results were accurate enough. The model resolution for the ice melt calculation is 250 m and is therefore “small” enough to be influenced by a glacier retreat in range of 500–1000 m. My concern is that if the glacier retreat is not taken into account in the modelling of such a long period (50 years, with very high rates of glacier retreat in the last decade), there is a real risk of overestimating the glacier melt by the model. The results of the analysis suggest that there is a shift in the discharge from a snow–melt dominated regime to an icemelt dominated regime (Fig. 10 and 12). I wonder if part of this increase in icemelt can be induced by the fact that glacier retreat is not taken into account. Therefore I would suggest the authors to discuss in

more details the impact of this model assumption on the results. Many glaciers in the Rhone basin are well documented and have yearly mass balance data. It would be worth to compare the modeled ice melt values with measured time series of glacier mass balance to exclude that the increase of ice melt is due to model assumptions.

This is indeed a very important point, and we agree with Referee #2 that neglecting glacier evolution may influence the estimation of icemelt rates. The Referee raises the possibility that by neglecting glacier area loss (retreat) we are possibly overestimating the icemelt contribution over our study period. To provide evidence that this is not the case, we compared our simulations with time series produced with the Global Glacier Evolution Model (GloGEM), a model accounting for both glacier mass balance and evolution. For comparison, we used total monthly runoff (snowmelt + icemelt + rainfall) generated over the glacierized surfaces of the upper Rhone basin, simulated with GloGEM (Huss and Hock, 2015) for the period 1980–2010.

GloGEM computes the mass balance for every glacier in 10-m elevation bands by estimating snow accumulation, snow and ice melt, and refreezing of rain and melt water. The response of glaciers to changes in mass balance is modelled on the basis of an empirical relation between ice thickness change and normalized elevation (Huss et al., 2010). Starting from initial ice thickness values derived by the method of Huss and Farinotti (2012), ice thickness is updated at the end of each hydrological year. GloGEM is calibrated and validated over the period 1980–2010 with estimates of glacier mass changes by Gardner et al. (2013) and in situ measurements provided by the World Glacier Monitoring Service.

Although in our hydrological model – which considers precipitation, snowmelt, icemelt at pixel scale, but integrates them to basin-average values – we do not include glacier evolution, the annual runoff volumes (snowmelt + icemelt + rainfall) from glacierized areas during the period 1980–2010 correlates very well with the results of GloGEM (Fig. 5a). Measures of performance confirm the agreement between the two models: the correlation coefficient is equal to 0.86 and the Nash–Sutcliffe efficiency is equal to 0.67. We are also capable to capture quite well the seasonal pattern of runoff generated from glacierized areas (Fig. 5b).

Perhaps most importantly, GloGEM simulations show that total annual runoff is increasing throughout the period and there is no evidence for decreasing icemelt rates. This confirms that, although glaciers of the upper Rhone basin are retreating, icemelt fluxes originating from glacierized and proglacial areas are increasing during the 1980–2010 period. As expected, total runoff from glacierized surfaces and icemelt are highly correlated (Fig. 5a, correlation coefficient = 0.95), thus indicating that the increase in total runoff is due to an increase of the icemelt component. Indeed, non-parametric Mann–Kendall tests indicate an increasing trend with 5% significant level. Trend slopes, estimated with the Theil–Sen estimator, confirm the agreement between the two models: we find a total runoff of ~ 27.65 mio $\text{m}^3 \text{ year}^{-2}$ with GloGEM and ~ 21.71 mio $\text{m}^3 \text{ year}^{-2}$ with our model. We also computed the basin-averaged mass balance for each hydrological year. The mean mass balance over the period 1980–2010 is equal to -0.78 ± 0.22 m w.e./year (Fig 6). This value is slightly greater than that found by Fischer et al. (2015) for the upper Rhone basin (-0.59 m w.e./year), but within the uncertainty of the estimate.

In summary: although we do not account for glaciers retreat, our model results agree well with a state-of-the-art glaciological model that includes glacier evolution. Both comparisons with GloGEM and our basin-averaged mass balance indicate that we are not significantly overestimating icemelt contribution during the period 1975–2015.

Clearly, if we were looking at future climate projections and at glaciers that are rapidly retreating we would need to include a glacier retreat. Under climate change, even the largest glacier in the basin, the Aletsch Glacier, is expected to shrink at a rate where its icemelt contribution would start decreasing before 2050 (Farinotti et al., 2012; FOEN, 2012; Brönnimann et al., 2014).

We present the results of this analysis in Sect. 6.1 (“Glacier retreat and ice-melt”) and we include Fig. 5 in the revised manuscript.




Figure 5: Runoff (snowmelt + icemelt + rainfall) generated from glacierized areas within the upper Rhone basin, simulated with GloGEM and with our snowmelt and Ice-melt models (degree-day) for the period 1980–2010: (a) total annual values; (b) mean monthly values. Fig. 11a also depicts the time series of total annual Ice-melt simulated with our Ice-melt model.




Fig. 6: Mass balance rate for glacierized areas of the upper Rhone basin, simulated with our temperature index model for the period 1975–2015.

5) Detailed comments:

We thank the reviewer for the following detailed comments. We modified the text accordingly.

p.2, l. 13–18: “In such...”: this sentence is too long, make 2 sentences out of it. Isn’t rainfall and liquid precipitation the same?

We divided the sentence into two sentences.

p.3, l. 25–27: “Although our results...”: move to the discussion.

Done.

p.4, Fig. 1: In my opinion Fig. 1 does not give any additional information. The aim of this figure is not very clear. I would suggest to suppress it or to develop it in a way where it is clear how the 4 sediment sources are build and how they can play a role. Maybe the authors could also link the different sources to the hydro-climatic factors (which factor influences which source).

We agree with the reviewer that Fig. 1 does not give additional information and we removed it.

p.5, l. 1–2: “However ... of our investigation”: This does not belong to the description of the study site, it belongs rather to the introduction or to the discussion.

We removed this sentence.

p.5, l. 9–11: which are the observed and the modeled data?

We specify in parentheses “observed” and “simulated” to clarify.

p.6, l. 1–3: how do the authors calculate the 250–250m daily temperature and precipitation from the MeteoSwiss gridded dataset (ca. 2x2km)? It is not clear to me if or when you use monthly average over the basin or daily gridded data. Please clarify.

We thank the reviewer for pointing out the necessity of clarifying this point. We changed the text in order: (1) to specify the interpolation method (nearest-neighbor interpolation) applied to obtain the input datasets on a 250×250 m resolution grid; (2) to clarify the time scale of the analysis (daily for the snow and icemelt model simulations, monthly and annual for the analysis of the temporal evolution). We hope it is clearer now.

p.6, Table 1: the gridded dataset from MeteoSwiss is originally in degree; therefore the grid size is ca 2x2km

Indeed the spatial resolution of the gridded dataset is approximately equal to 1.6 km×2.1 km. We agree with the reviewer that it has to be specified by adding the approximation symbol tilde “~”.

p.8, l. 4–10: this belongs to the discussion (see also comments above on the ice model). This is also repeated in p.11, l. 8–12. Regarding Sect. 3.2 (p.8, l. 4–10 of the reviewed manuscript), we changed the text according to the suggestions of the reviewer, while regarding Sect. 4.3 (p.11, l. 8–12 of the reviewed manuscript) we removed the comments because they are now discussed in more details in the discussion Sect. 6.2.

p.9, l. 6–18: Most of this Sect. belongs to the description of the study area, not to the methods section.

We agree and we partly moved this description to the description of the study site (Sect. 2).

p.10, l. 18–21: “we also verified...” And? Where are the results of this analysis? What are the conclusions? I don’t understand the sense of these sentences.

We changed this part and we tried to clarify.

p. 10, l. 23–26: this is a repetition.

We reduced the text trying to avoid repetitions.

p.11, l.20: “This value...”, it is enough to discuss it only in the discussion, delete this sentence.

We kept this comment in the results since the manuscript has been re-structured.

p.11, l.31–32: belongs to the discussion.

In the revised manuscript, comments on calibration/validation of snow and icemelt models are reported in the results section (5.1) and removed from the discussion section. We hope that this make the text clearer and reduce repetitions.

p.12, l.1: “we are confident” avoid this kind of expression (all over the text) as it gives the impression that the observation are not based on results but on feelings.

We agree with the reviewer and changed the text accordingly.

p.12, l.3–5: “Despite...”, it is enough to discuss it only in the discussion, delete this sentence.

In the revised manuscript, comments on calibration/validation of snow and icemelt models are reported in the results section (5.1) and removed from the discussion session. We hope that this make the text clearer and reduce repetitions.

Fig. 3a–b: I am not convinced that these 2 figures bring additional information. I would suggest deleting them.

We removed the figures from the revised manuscript.

p.15, l. 5–8; 19–11: Move to the discussion.

In the revised manuscript, comments on changes in hydroclimatic variables are reported in the results section (5.2 and 5.3) and removed from the discussion section.

p.15, l.16–24: In this Sect. it is confusing when annual and monthly discharge is meant.

We changed the text in order to differentiate better between mean monthly and mean annual values.

p.15, l.22: “this variability is reflected in streamflow”, where?

We deleted this sentence and we keep the discussion on the potential impact of large-scale climatic patterns both on precipitation and discharge.

p.16, l.1–5: move to the discussion.

In the revised manuscript, comments on changes in hydroclimatic variables are reported in the results section (5.2 and 5.3) and removed from the discussion section.

p.16, l.6–10: this part belongs to the discussion and is a repetition.

It was removed from the discussion.

p.16, Fig 8: _Q and _P seems to react inversely, is there an explanation? Is it due to hydropower?

While for discharge hydropower operations might possibly explain the increase in winter (as discussed in the chapter 5.2 of the revised manuscript), we cannot identify a clear connections between precipitation and releases for hydropower reservoirs, on the basis of this dataset. However, we think that this could be an interesting point to investigate more in details in the future. We thank the reviewer for pointing this out.

p.17, l.2: “The fact that sediment supply impacts suspended sediment concentration is evident...” really? Where do I see that?

The sediment rating curve shows that both transport capacity and sediment supply (i.e. sediment sources and sediment production and transport processes) influence suspended sediment concentration. Indeed, given the transport capacity (discharge), suspended sediment concentration covers a wide range of values. We changed the text in order to clarify this point.

p.17, Fig. 9: would it be possible to highlight (different colors) < 1987 and >1987?

We thank the reviewer for this suggestion. We changed the figure by differentiating data for the periods 1965–1986 and 1987–2015.

p.17, l.27: “Our simulation: : ”, it is enough to discuss it only in the discussion, delete this sentence.

In the revised manuscript, comments on changes in hydroclimatic variables are reported in the results section (5.2 and 5.3) and removed from the discussion section.

p.17, l.28–30: How did you come to this conclusion? Did you compare it with literature? These sentences belong to the discussion.

We kept this comment in the results since the manuscript has been re-structured. This argument is based on the physical processes occurring in mountainous environments. For example, snow cover influences icemelt both by insulating glacierized surfaces and so preventing melting, and by affecting albedo and therefore the energy balance.

p.18, l.2–6; 8–10: move to the discussion.

We kept this comment in the results since the manuscript has been re-structured.

p.18, l. 16: where is this shown? Cite the figure.

Done.

p.18, l.22: “a shift of snow–melt to ice–melt”: I would not call it a shift as it is only slightly visible from fig. 12. Is it statistically significant? Cite the figure where the shift can be seen (fig. 12?).

We agree with the reviewer. In addition, we changed the text in agreement with the figure, which we changed showing the relative contribution of the three hydroclimatic variables during the summer months, when the rise in suspended sediment concentration is observed.

p.19, l.1–15: Most of this belongs to the discussion.

We moved this comment into the result section.

p.20, l. 20–23: this has already been said and is only a repetition.

We removed the entire section 6.1.

p.21, l.10: “...decrease in fine sediment load...” where is this decrease in the 60s to see? The analysis start around 1965, where the sediment load is low, but there is no decrease...

We agree with this comment and we clarified that this decrease was observed and documented by Loizeau et al. (1997) on the basis of sediment cores recovered in the Rhone delta region. We changed the text to clarify this point.

p.21, l. 10–12: how do you come to the conclusion that it can be an effect of hydropower?

This is the hypothesis that has been proposed so far in the literature (Loizeau et al., 1997; Loizeau and Dominik, 2000). The analysis that we conducted in this study does not allow us to reject this hypothesis. However, we demonstrate that climate has a significant effect on suspended sediment dynamics, although the system is highly regulated. Therefore, we argue that together with hydropower operations, the colder temperature that characterizes the 1960s could have contributed to the reduction in suspended sediment concentration observed by previous studies.

p.21, l.16–34: This is only speculative and take too much importance in the discussion.

We agree and we removed it almost completely.

p.22, l.8–10: repetition

We removed it.

p.22, l.14–19: sentence too long, shorten it or separate it.

This sentence was moved to chapter 5.3 and separated into two sentences according to the reviewer.

p.22, l.21: there is no increase in rainfall intensity in the analysis (fig. 7)

Here we refer to the increase in R-factor identified by Meusburger et al. (2012). Although we do not identify any increase in mean annual and mean monthly values of total daily precipitation (Fig. 5b and 6b of the revised manuscript), the analysis of rainfall at sub-daily scale could provide different results. However, we agree that this must be specified and we changed the text accordingly.

p.23, l.18–21: There were 4 sources of sediments in fig. 1, please clarify. To me it is not clear how these sediment sources were taken into account in the analysis. You should maybe emphasis in the discussion how each factor influences each sediment source.

We agree that this needs to be clarified. We changed the term “sediment sources” with the term “sediment fluxes”, and we list the three main sediment fluxes that we are analysing, i.e. sediments originated and transported by (1) snowmelt, (2) erosive rainfall, and (3) icemelt. Because we removed Fig 1 and do not specify the sediment sources anymore we think it is not necessary anymore to extend the discussion on how each factor influences each sediment source.

References

Fischer, M., Huss, M., Hoelzle, M.: Surface elevation and mass changes of all Swiss glaciers 1980–2010, *Cryosphere* 9, 525–540, doi:10.5194/tc-9-525-2015, 2015.

Gardner, A. S., Moholdt, G., Cogley, J. G., Wouters, B., Arendt, A. A., Wahr, J.: A reconciled estimate of glacier contributions to sea level rise: 2003 to 2009, *Science*, 340, 852–857, doi:10.1126/science.1234532, 2013.

Huss, M. and Farinotti, D.: Distributed ice thickness and volume of all glaciers around the globe, *J. Geophys. Res.*, 117: F04010, doi: 10.1029/2012JF002523, 2012.

Huss, M. and Hock, R.: A new model for global glacier change and sea-level rise, *Front. Earth Sci.*, 3:54, doi: 10.3389/feart.2015.00054, 2015.

Huss, M., Juvet, G., Farinotti, D., Bauder, A.: Future high-mountain hydrology: a new parameterization of glacier retreat, *Hydrol. Earth Syst. Sci.*, 14, 815–829, doi:10.5194/hess-14-815-2010, 2010.

Loizeau, J.-L. and Dominik J., Luzzi, T., Vernet J.-P.: Sediment Core Correlation and Mapping of Sediment Accumulation Rates in Lake Geneva (Switzerland, France) Using Volume Magnetic Susceptibility, *J. Great Lakes Res.*, 23(4), 391–402, 1997.

Loizeau, J.-L. and Dominik J., Evolution of the Upper Rhône River discharge and suspended sediment load during the last 80 years and some implications for Lake Geneva, *Aquat. Sci.*, 62, 54–67, 1015–1621/00/010054–14, 2000.

Meusburger, K., Steel, A., Panagos, P., Montanarella, L., Alewell, C.: Spatial and temporal variability of rainfall erosivity factor for Switzerland, *Hydrol. Earth Syst. Sci.*, 16, 167–177, 2012.

List of all relevant changes

We performed additional analysis on the following four main aspects and we added the results to the revised manuscript:

- (1) sensitivity analysis of our snowmelt model parameters (included in the Supplementary Material, Fig. S1 and Fig. S2);
- (2) quantification of the possible overestimation of icemelt due to fact that glaciers retreat is not modelled (included in Sect. 6.1, Fig. 11a and Fig. 11b);
- (3) analysis of the effects of discontinuous SSC sampling on trend and jumps detected during the observation period (included in Sect. 6.2 and Fig. 12);
- (4) estimation of potential impacts of gravel mining activities on SSC rise detected in mid-1980s (included in Sect. 6.2).

We re-structured the manuscript by merging the previous discussion and results sections in the current results section (Sect. 5). We created a new discussion section (Sect. 6) with new contents (mainly discussion on icemelt model and on anthropogenic impacts).

We changed the text to reduce repetition and increase clarity and we changed the references accordingly.

In particular, we:

- reformulated the aim of the study;
- reduced consistently the discussion on climate change (Sect. 6.2);
- changed the text referring to the decrease in suspended sediment load in 1960s. We clarify that it was observed and documented by Loizeau et al. (1997) on the basis of sediment cores recovered in the Rhone delta region;
- specified comments on rainfall erosivity: we refer to the increase in R-factor, computed on the basis of 10 min resolution precipitation data, identified by Meusburger et al. (2012). Although we do not identify any increase in mean annual and mean monthly values of total daily precipitation (Fig. 5b and 6b of the revised manuscript), the analysis of rainfall at sub-daily scale could provide different results.
- clarified in the text the tests that we applied on the MeteoSwiss gridded dataset in order to ensure that they are not affected by non-homogeneities (due to the variability of the number of stations used in developing the dataset)
- removed previous Fig. 1 (sediment sources), Fig. 3a (K_{snow}) and Fig. 3b (K_{ice});
- changed Fig. 7 (previous Fig. 9) to highlight observations of the period 1965–1986 and 1987–2015. We estimated sediment rating curves separately for the two periods;
- changed Fig. 10 (previous Fig. 12) by depicting the relative contribution of rainfall, snow and icemelt on the sum of these three components, only in July and August, months in which SSC increased.

Temperature signal in suspended sediment export from an Alpine catchment

Anna Costa¹, Peter Molnar¹, Laura Stutenbecker², Maarten Bakker³, Tiago A. Silva⁴, Fritz Schlunegger², Stuart N. Lane³, Jean-Luc Loizeau⁴, Stéphanie Girardclos⁵

¹Institute of Environmental Engineering, ETH Zurich, 8093 Zurich, Switzerland

²Institute of Geological Sciences, University of Bern, 3012 Bern, Switzerland

³Institute of Earth Surface Dynamics, University of Lausanne, 1015 Lausanne, Switzerland

⁴Department F.-A. Forel for Environmental and Aquatic Sciences, University of Geneva, 1211 Geneva, Switzerland

⁵Department of Earth Sciences and Institute for Environmental Sciences, University of Geneva, 1205 Geneva, Switzerland

Correspondence to: Anna Costa (costa@ifu.baug.ethz.ch)

Abstract

Suspended sediment export from large Alpine catchments ($> 1000 \text{ km}^2$) over decadal timescales is sensitive to a number of factors, including long-term variations in climate, the activation–deactivation of different sediment sources (proglacial areas, hillslopes, etc.), transport through the fluvial system, and potential anthropogenic impacts on the sediment flux (e.g. through impoundments and flow regulation). Here, we report on a marked increase in suspended sediment concentrations observed near the outlet of the upper Rhône River Basin in the mid–1980s. This increase coincides with a statistically significant step–like increase in basin–wide mean air temperature. We explore the possible explanations of the suspended sediment rise in terms of changes in water discharge (transport capacity), and the activation of different potential sources of fine sediment (sediment supply) in the catchment by hydroclimatic forcing. Time series of precipitation and temperature–driven snowmelt, snow cover and icemelt simulated with a spatially distributed degree–day model, together with erosive rainfall on snow–free surfaces, are tested to explore possible reasons for the rise in suspended sediment concentration. We show that the abrupt change in air temperature reduced snow cover and the contribution of snowmelt, and enhanced ice–melt. The results of statistical tests show that the onset of increased icemelt was likely to play a dominant role in the suspended sediment concentration rise in the mid–1980s. Temperature–driven enhanced melting of glaciers, which cover about 10% of the catchment surface, can increase suspended sediment yields through increased contribution of sediment–rich meltwater, increased sediment availability due to glacier recession, and increased runoff from sediment–rich proglacial areas. The reduced extent and duration of snow cover in the catchment are also potential contributors to the rise in suspended sediment concentration through hillslope erosion by rainfall on snow free surfaces, and increased meltwater production on snow–free glacier surfaces. Despite the rise in air temperature, changes in mean discharge in the mid–1980s were not statistically significant, and their interpretation is complicated by hydropower reservoir management and the flushing operations at intakes. Overall, the results show that to explain changes in suspended sediment transport from large Alpine catchments it is necessary to include an understanding of the multitude of sediment sources involved together with the hydroclimatic conditioning of their activation (e.g. changes in precipitation, runoff, air temperature). In addition, this study points out that climate signals in suspended sediment dynamics may be visible even in highly regulated and human impacted systems. This is particularly relevant for quantifying climate change and hydropower impacts on streamflow and sediment budgets in Alpine catchments.

1. Introduction

Erosion processes and sediment dynamics in Alpine catchments are determined by geological, anthropogenic, and climatic factors. Geological forcing is one of the main drivers of sediment production and landscape development, through crustal thickening, deformation and isostatic uplift, and glacier inheritance (e.g. England and Molnar, 1990; Schlunegger and Hinderer, 2001; Vernon et al., 2008). Almost continuous glacier recession in the European Alps since the late 19th Century

(Paul, 2004; 2007; Haeberli, 2007) has maintained large parts of the landscape in early stages of the paraglacial phase where unstable or metastable sediment sources (Ballantyne, 2002; Hornung et al., 2010) can maintain high sediment supply rates. Glacier inheritance influences sediment production and transport as demonstrated by a strong spatial association between sediment yield and past and current glacial cover (Hinderer et al., 2013; Delunel et al., 2014). Anthropogenic impacts on sediment yields are more recent, and on a global scale largely related to land cover change through intensified agriculture and the trapping of sediment in reservoirs (e.g. Syvitski et al., 2005). Land use changes impact mainly fine sediment production (e.g. Foster et al. 2003; Wick et al., 2003), while river channelization, flow regulation, water abstraction, and sediment extraction have caused a general reduction in sediment yield and consequently led to sediment-starved rivers world-wide (Kondolf et al., 2014). In Alpine catchments, in addition to trapping in reservoirs, sediment transfer is also disturbed by flow abstraction at many hydropower intakes. The reduction of sediment transport capacity downstream of intakes and the periodic flushing of locally trapped sediment has severe impacts on the sediment budget (e.g. Anselmetti et al., 2007) and downstream river ecology (e.g. Gabbud and Lane, 2016).

In this paper we focus on the dominant role of climate in sediment production and transfer in Alpine environments (e.g. Huggel et al., 2012; Zerathe et al., 2014; Micheletti et al., 2015; Palazon and Navas, 2016; Wood et al., 2016). **In such environments, intense rainfall triggers erosion and mass movements on hillslopes (landslides, debris flows, etc.), air temperature and (solid, liquid) precipitation drive the hydrological processes that generate runoff (e.g. Quinton and Carey, 2008), and so determine the capacity of the rivers to transport sediment. Snow cover accumulation and melt, snow avalanching, as well as the seasonal dynamics of ice-melt, also directly affect sediment supply (e.g. Moore et al., 2013).**

To assess the relative contribution of climatic variables to the sediment production and yield of an Alpine catchment we identify four main sediment sources: glacial erosion, hillslope erosion, channel bed/bank erosion and mass wasting events (e.g. rockfalls, debris flows). Climatic conditions, specifically precipitation and air temperature, contribute to the activation of these four sediment sources through different processes and at different rates. Erosive processes of abrasion, bed-rock fracturing and plucking at the base of glaciers provide proglacial areas with large amounts of sediment (Boulton, 1974). Due to glacial erosion, discharge from subglacial channels has high suspended sediment concentrations (e.g. Aas and Bogen, 1988). Temperature-driven snow and icemelt in spring and summer, as well as intense rainfall on snow-free surfaces, may lead to entrainment from proglacial areas provided they are connected to the river network (Lane et al., 2017). Hillslope erosion driven by overland flow and rainfall erosivity (Wischmeier, 1959) may be exacerbated in Alpine catchments by permanently or partially frozen ground (Quinton and Carey, 2008). In summer and autumn, when Alpine catchments are largely free of snow, intense rainfall may erode large amounts of sediment and transport it in rills and gullies to the river network. Intense rainfall is also responsible for triggering mass wasting events, such as debris flows and landslides, where a large mass of sediment is delivered to the channel network instantaneously (e.g. Bennett et al., 2012). Flow conditions (e.g. shear stress, stream power) then determine the sediment transport capacity and in-stream sediment mobilization along rivers, and hence its transfer to downstream locations.

The close link between precipitation, air temperature, runoff and the activation-deactivation of sediment sources in Alpine catchments becomes critical in the context of climate change. Alpine regions represent a very sensitive environment in relation to current rapid warming. In Switzerland, substantial glacier retreat has been observed since 1850, and considerably higher ice loss rates were found after the mid-1980s (Paul, 2004; 2007; Haeberli, 2007). A reduction in snow cover duration and mean snow depth has also been observed during the last thirty years (e.g. Beniston, 1997; Laternser and Schneebeli, 2003; Scherrer et al., 2004; Marty, 2008; Scherrer et al., 2006). Although current effects of climate change are much less clear for precipitation (Brönnimann et al., 2014) than for temperature, a sharp reduction in the number of snowfall days has been observed at many Swiss meteorological stations (Serquet et al. 2011).

The premise behind this paper is that to explain impacts of changes in climate on Alpine catchment suspended sediment yield, it is necessary to consider at least (a) the sediment sources involved; and (b) the hydroclimatic conditioning of their

activation (e.g. precipitation, runoff and air temperature). The traditional rating–curve analysis, which relates suspended sediment concentration only to discharge (e.g. Campbell and Bauder, 1940; Walling, 1974; 1977; Asselman, 1999; 2000; Lenzi and Marchi, 2000; Horowitz, 2003; Mao and Carrillo, 2015), is insufficient for analyzing detailed changes in sediment yield because sediment supply is not explicitly accounted for (e.g. Walling, 2005; De Vente et al., 2006; Mao and Carrillo, 2015). In this study we look at specific sediment sources and their activation with a process–based perspective with the aim to infer the possible effects of changes in hydroclimate, such as increases in temperature and/or precipitation intensity, on suspended sediment dynamics.

The upper Rhône River Basin draining into Lake Geneva in Switzerland is at the center of our investigation. The basin has experienced a rise in air temperature that coincided with a rise in suspended sediment concentrations in the mid–1980s. Our main objective is to explore the presence of the signal of a warmer climate in the suspended sediment dynamics of this regulated and human–impacted Alpine catchment. In this work, we refer to fine sediment as the sediment transported in suspension. We investigate the potential causes of the observed increase in suspended sediment concentration by focusing on two factors: transport capacity and sediment supply. For transport capacity we analyze daily streamflow at the catchment outlet, while for sediment supply we conceptualize the upper Rhône Basin as a series of spatially distributed sediment sources that are activated or deactivated throughout the hydrological year by hydroclimatic forcing. Hydroclimatic forcing is given by daily basin–wide observed precipitation, and simulated snow cover, snowmelt and ice–melt. Our aims are: (a) to estimate daily hydroclimatic variables over the Rhône Basin for the last 40 years, which explicitly address the activation of the main sediment sources; and (b) to provide statistical evidence for possible reasons for the rise in suspended sediment concentrations in the mid–1980s.

2. Study Site Description

The upper Rhône Basin is located in the southwestern part of Switzerland, in the Central Swiss Alps (Fig. 1). It has a total surface area of 5338 km², and an altitudinal range of 372 to 4 634 m a.s.l. About 10% of the surface is covered by glaciers, which are mostly located in the eastern and southeastern part of the catchment (Stutenbecker et al., 2016). The Rhône River originates at the Rhône Glacier and flows for about 160 km through the Rhône valley before entering Lake Geneva, a few kilometers downstream of the gauging station located at la Porte–du–Scex (Fig. 1). Basin–wide mean annual precipitation is about 1400 mm yr^{−1} and shows strong spatial variability driven mostly by orography and the orientation of the main valley. The hydrological regime of the catchment, typical of Alpine environments, is strongly influenced by snow and icemelt with highest discharge in summer and lowest in winter. Mean annual discharge is 180 m³ s^{−1}, which corresponds to about 1060 mm yr^{−1} and an annual runoff coefficient of 75%.

The catchment has been strongly affected by anthropogenic impacts during the last century. The main course of the Rhône River has been extensively channelized for the purposes of flood protection: levees were constructed and the channel was narrowed and deepened in the periods 1863–1894 and 1930–1960 (First and Second Rhône Corrections). Due to the residual flood risk that affects the main valley, a third project was started in 2009 with the main objectives to increase channel conveyance capacity and river ecological rehabilitation (Oliver et al., 2009). In addition, significant gravel mining operations are carried out along the main channel and many tributaries. Since the 1960s, several large hydropower dams have been built in the main tributaries of the Rhône River. The total storage capacity of these reservoirs corresponds to about 20% of the mean annual streamflow (Loizeau and Dominik, 2000). Flow impoundment, water abstraction and diversion through complex networks of intakes, tunnels and pumping stations, have significantly impacted the flow and sediment regime of the catchment. Flow regulation due to hydropower production has resulted in a considerable decrease of discharge in summer and increase in winter (Loizeau and Dominik, 2000). Impacts upon sediment supply are more complicated because flow abstraction and diversion tends to leave a proportion of sediment behind in the associated streams (Gabbud and Lane, 2016).

That said, the construction of dams and start of hydropower operation has coincided with a drop in the suspended sediment load of the main Rhône River measured at la Porte-du-Scex in the 1960s (Loizeau et al., 1997; Loizeau and Dominik, 2000). Two sub-catchments of the Upper Rhône basin are used for the calibration/validation of the icemelt model: the Massa and the Lonza. The Massa (Fig. 1) is a medium-sized basin (195 km^2) with a mean elevation of 2945 m a.s.l. More than 60% of the surface is glacierized, and the remaining surface is classified mostly as rock and firn (Boscarello et al., 2014). The basin includes the Aletsch Glacier, which is the largest glacier in the European Alps with a length of around 23.2 km and a surface area of approximately 86 km^2 (Haeberli and Holzhauser, 2003). The Lonza is a relatively small basin located to the west of the Massa (Fig. 1) with an average elevation of 2630 m a.s.l. It has a total drainage area of 77.8 km^2 and its surface consists of 36% of glacier cover. Daily discharge measurements are available for the Massa and for the Lonza respectively at the gauging stations of Blatten bei Naters and Blatten.

3. Methods

Our objective is to explore the potential effect of climate on suspended sediment dynamics of the upper Rhône Basin in the period 1975–2015. To this end we use observed and simulated hydroclimatic and sediment transport variables, as listed in Table 1: temperature T , precipitation P , discharge Q , suspended sediment concentration SSC (observed), snow cover fraction SCF, snowmelt SM, icemelt IM, and effective rainfall ER (simulated).

This selection of variables is based on the consideration that three main hydroclimatic factors (ice-melt, snowmelt and rainfall) are responsible for activating sediment sources with the following processes in mind: (a) icemelt runoff evacuates accumulated fine sediment, the product of glacial erosion, through subglacial channels (e.g. Swift et al., 2005); (b) snow cover dynamics influence ice-melting onset, impact icemelt efficiency through albedo and may result in more rapid erosion and sediment production through an increased glacier basal velocity (e.g. Herman et al., 2015); (c) snowmelt runoff from snow-covered catchment areas may generate downstream hillslope erosion and channel erosion (e.g. Lenzi et al., 2003); and (d) rainfall on snow-free surfaces leads to hillslope erosion, mass wasting and, due to high discharge, also channel erosion (e.g. Bennet et al., 2012; Meusburger and Alewell, 2014).

Some variables originate from observations or interpolation of observations (T , P , Q , SSC), others from simulations by spatially distributed snow and icemelt models (SCF, SM, IM), or from a combination thereof (ER). The snowmelt model is described in Sect. 3.1, the icemelt model in Sect. 3.2, and their calibration in Sect. 3.3. We first interpolate the input datasets of precipitation and temperature on a $250 \times 250 \text{ m}$ grid resolution by the nearest-neighbor interpolation method. Second we run the snow and icemelt model at the daily timescale over the period 1975–2015. In a third step, we analyze the variables (Table 1) as mean monthly and mean annual values averaged over the basin area. To quantify changes in the hydroclimatic variables and in suspended sediment concentration, we apply standard statistical tests for change detection described in Sect. 3.4. A description of all observations used in this analysis is reported in more details in Sect. 4.

3.1 Snowmelt Model

We use a snowmelt model to predict SM and SCF over the entire basin, because snow station measurements are sparsely and irregularly distributed and a physical consistency between precipitation and air temperature as climatic driving forces and snowmelt and snow cover as response variables are needed. The spatially distributed temperature index method (degree-day model) was used due to its simplicity, low data requirements, and demonstrated success at daily temporal scales over large basins (e.g. Hock, 2003; Boscarello et al., 2014). The degree-day approach also matches the coarse spatial ($250 \times 250 \text{ m}$) and temporal (daily) resolution of our analysis and the integration at the basin scale. Models based on energy balance, or enhancements of the degree-day approach, represent physical processes better and could be used when higher spatial and temporal resolution and accuracy is needed (e.g. Pellicciotti et al., 2005).

The snowmelt model includes snow accumulation and melt. At the grid scale, precipitation P [mm day⁻¹] is first partitioned into solid and liquid form based on (a) daily minimum T_{\min} [°C] and maximum air temperature T_{\max} [°C] and (b) a rain–snow threshold temperature T_{RS} [°C]. If minimum air temperature T_{\min} is above the threshold temperature T_{RS} , all precipitation falls as rainfall R ; if the maximum air temperature T_{\max} is below the threshold temperature T_{RS} , all precipitation falls as snow S ; otherwise precipitation is a mixture of liquid and solid form, partitioned proportionally to the temperature differences:

$$\begin{cases} R = c_p P \\ S = (1 - c_p) P \end{cases}, \quad (1)$$

where

$$\begin{cases} c_p = 1 & T_{\min} > T_{RS} \\ c_p = 0 & T_{\max} \leq T_{RS} \\ c_p = \frac{T_{\max} - T_{RS}}{T_{\max} - T_{\min}} & T_{\min} \leq T_{RS} < T_{\max} \end{cases}, \quad (2)$$

The daily snowmelt rate M_{snow} [mm day⁻¹] is estimated from a linear relation with air temperature:

$$\begin{cases} M_{\text{snow}} = k_{\text{snow}}(T_{\text{mean}} - T_{SM}) & T_{\text{mean}} > T_{SM} \\ M_{\text{snow}} = 0 & T_{\text{mean}} \leq T_{SM} \end{cases}, \quad (3)$$

where T_{mean} [°C] is the mean daily air temperature, T_{SM} [°C] is a threshold temperature for the onset of melt, and k_{snow} is a melt factor [mm day⁻¹ °C⁻¹]. Snow depth (SD), in mm snow water equivalent, for time t is then simulated from a balance between accumulation and melt at every grid cell i :

$$SD_i(t) = SD_i(t-1) + S_i(t) - M_{\text{snow}_i}(t). \quad (4)$$

The snow cover fraction SCF for a chosen area containing $i = 1, \dots, N$ grids, is:

$$SCF(t) = \frac{1}{N} \sum_{i=1}^N H[SD_i(t)], \quad (5)$$

where H is a unit step function: $H = 0$ when $SD = 0$ and $H = 1$ when $SD > 0$. The area of integration N can be the entire catchment, sub-basins, elevation bands, etc. For the entire catchment, we estimate mean daily snowmelt SM [mm day⁻¹] as the arithmetic average over all grid melt rates:

$$SM(t) = \frac{1}{N} \sum_{i=1}^N M_{\text{snow}_i}(t). \quad (6)$$

The threshold temperatures for defining the precipitation type T_{RS} and the onset of melt T_{SM} depend on many factors such as atmospheric boundary layer conditions, temperature, humidity, cloud types, among others. Different parametrizations and temperature values are available in the literature (Wen et al., 2013). Depending on region, altitude and modelling approach, rain–snow temperature thresholds show a wide range of variability from -5 °C (Collins et al., 2004) to more than 6 °C (Auer, 1974). For the upper Rhône Basin we assume a constant rain–snow temperature threshold $T_{RS} = 1^\circ\text{C}$, resulting from a calibration and validation of the physically–based fully distributed hydrological model Topkapi–ETH in the catchment (Fatichi et al., 2015). To reduce degrees of freedom, the threshold temperature for the onset of melt T_{SM} is set equal to 0°C and the calibration of the snowmelt model consists of estimating the melt factor k_{snow} with methods described in Sect. 3.3.

3.2 Icemelt Model

Similar to snowmelt, icemelt is also simulated with a temperature index (degree–day) model on grid cells that are identified as glacier–covered from remote sensing data. The daily icemelt M_{ice} [mm day⁻¹] on glacier surfaces that are snow–free is estimated as:

$$\begin{cases} M_{\text{ice}} = k_{\text{ice}}(T_{\text{mean}} - T_{IM}) & T_{\text{mean}} > T_{IM} \\ M_{\text{ice}} = 0 & T_{\text{mean}} \leq T_{IM} \end{cases}, \quad (7)$$

where T_{mean} [°C] is mean daily air temperature, T_{IM} [°C] is a threshold temperature for the onset of ice–melt, and k_{ice} [mm day⁻¹ °C⁻¹] is the icemelt factor. For the entire catchment, we estimate mean daily icemelt IM [mm day⁻¹] as the arithmetic average over all ice–covered grid cells:

$$IM(t) = \frac{1}{N} \sum_{i=1}^N M_{ice_i}(t) . \quad (8)$$

The threshold temperature for glacier melting T_{IM} is set equal to 0 °C. Icemelt occurs only if the glacier cell is snow-free. The snow cover simulated by the snowmelt model in Sect. 3.1 is thus essential for estimating ice-melt. The calibration of the icemelt model consists of estimating the melt factor k_{ice} with data from two highly glacierized sub-basins in the Rhone: Massa and Lonza (Fig. 1). The calibration method is described in Sect. 3.3.

3.3 Calibration and Validation

We perform the calibration and validation of the snow and icemelt model parameters in sequence, since the snow-covered surface is required for icemelt estimation on glaciers. The snowmelt factor k_{snow} is calibrated based on comparisons with snow cover data derived from satellite images (MODIS). Snow cover observations are split into two periods: 1 October 2000 – 30 September 2005 for calibration and 1 October 2005 – 31 December 2008 for validation. MODIS maps of snow cover are filtered to reduce the impacts of clouds on SCF estimations. The resulting number of calibration and validation days is equal to 217 and 143 respectively. Snow cover maps at 500×500 m resolution are distributed by proximal interpolation to the snowmelt model 250×250 m computational grid. Snow depth maps simulated with Eq. (4) and transformed into a simulated snow cover fraction SCF^{sim} in Eq. (5) are compared with the MODIS maps SCF^{obs} .

The objective function for calibration is based on a combination of mean absolute error and true skill statistic. The mean absolute error MAE is estimated as:

$$MAE = \frac{1}{n} \sum_{j=1}^n |SCF_j^{obs} - SCF_j^{sim}| , \quad (9)$$

where n is the number of MODIS image maps, and it captures the overall ability of the model to reproduce the snow cover fraction accurately. The true skill statistic TSS is a spatial statistic that measures the grid-to-grid performance of the model in capturing snow-no snow presence. It is computed as the sum of sensitivity SE (correct snow predictions) and specificity SP (correct no-snow predictions) computed from contingency tables (e.g. Wilks, 1995; Mason and Graham, 1999; Corbari, 2009) in each image j and averaged over the n MODIS maps in the simulation period:

$$TSS = \frac{1}{n} \sum_{j=1}^n TSS_j = \frac{1}{n} \sum_{j=1}^n SE_j + SP_j - 1 \quad (10)$$

Because TSS includes both sensitivity and specificity, it captures both predictions of snow-covered and snow-free areas. It takes on values between 0 and 1, where 1 indicates perfect performance. TSS is a widely applied metric for assessing spatial model performance (e.g. Begueria, 2006; Allouche et al., 2006). We combine both goodness-of-fit measures (MAE and TSS) into an objective function OF, by giving more weight to MAE. Finally, we evaluate the objective function OF over $k = 5$ different elevation bands in order to better capture the topographic gradients in snowmelt distribution in the Rhône Basin:

$$OF = \sum_{k=1}^5 OF_k = \sum_{k=1}^5 -0.6 MAE_k + 0.4 TSS_k . \quad (11)$$

This objective function is maximized in calibration. The rationale of using both MAE and TSS in evaluating performance is to give weight to both basin-integrated snow cover as well as to grid-based predictions. Indeed, the same value of snow cover fraction can result in two different spatial arrangements of snow-covered pixels, and a correct spatial distribution of snow-covered and snow-free areas is relevant for this analysis insofar as it affects the activation and deactivation of specific sediment sources. The weights assigned to MAE and TSS in Eq. (11) are the outcome of sensitivity tests with the model. After calibration, we also estimate the Nash–Sutcliffe efficiency NS (Nash and Sutcliffe, 1970) and the mean square error MSE to quantify the performance of the model:

$$NS = 1 - \frac{\sum_{j=1}^n (SCF_j^{obs} - SCF_j^{sim})^2}{\sum_{j=1}^n (SCF_j^{obs} - \overline{SCF})^2} , \quad (12)$$

$$MSE = \frac{1}{n} \sum_{j=1}^n (SCF_j^{obs} - SCF_j^{sim})^2 , \quad (13)$$

where \overline{SCF} is the average observed snow cover fraction during the calibration-validation period.

We calibrate the icemelt factor k_{ice} on the sub-basin of the river Massa (Fig. 1), on the basis of daily discharge measurements, and focusing only on months when the icemelt contribution is not negligible (June–October). The gauging station is located upstream of the Gebidem dam, therefore discharge is not influenced by reservoir regulation and represents undisturbed natural flow. Calibration is performed on the period 1 January 1975 -- 31 December 2005, while validation covers the remaining ten years of available data, i.e. the period 1 January 2006 -- 31 December 2015. We then validate the model on the Lonza sub-basin with the same procedures and goodness of fit measures.

The optimal value of k_{ice} is found by minimizing the mass balance error MBE_s computed for the period June–October:

$$MBE_s = 100 \frac{\sum_{i=1}^{ny} (V_i^{obs} - V_i^{sim})}{\sum_{i=1}^{ny} V_i^{obs}}, \quad (14)$$

where ny is the number of calibration years, V_i^{obs} and V_i^{sim} [mm year⁻¹] are the observed and simulated discharge volumes per unit area reaching the outlet of the catchment during the period June–October of each calibration year i :

$$V^{obs} = \sum_{j=1}^{nd} Q_j^{obs}, \quad (15)$$

$$V^{sim} = \sum_{j=1}^{nd} Q_j^{sim} = \sum_{j=1}^{nd} (R_j + SM_j + IM_j). \quad (16)$$

Here, nd is the number of observation days from June to October, Q_j^{obs} [mm day⁻¹] is the daily discharge per unit area observed at Blatten Bei Naters (Blatten), R_j , SM_j , IM_j are respectively the total daily rainfall, snowmelt and icemelt aggregated over the Massa (Lonza) basin. Rainfall (R) and snowmelt (SM) are simulated with the snow accumulation and melt model in Sect. 3.1, while icemelt (IM) is simulated with the icemelt model in Sect. 3.2.

3.4 Statistical Testing for Change

We use the non-parametric Pettitt test (Pettitt, 1979) for the detection of the time of change (year-of-change) in the air temperature data. We then test the other variables (SSC, P, Q, SM, SCF, IM and ER) for changes in the mean (and variance) by splitting the time series into two periods before and after the identified year-of-change, and by applying two-sample two-sided t -tests for the equality of the means (and variances). The null hypothesis of no change is tested at the 5% significance level. The t -test is a parametric test commonly used in hydrology and atmospheric science to assess the validity of the null hypothesis of two samples having equal means and unknown unequal variances. We apply the t -test to all hydroclimatic variables averaged at the annual and monthly timescales with the same year-of-change to determine which hydroclimatic variables, and therefore the activation or deactivation of which sediment sources, are possibly responsible for the observed changes in suspended sediment concentration.

4. Data Description

4.1 Precipitation and Air Temperature

For precipitation and air temperature we use spatially distributed datasets provided by the Swiss Federal Office of Meteorology and Climatology (MeteoSwiss). Total daily precipitation, mean, minimum and maximum daily air temperature are available on a $\sim 2 \times 2$ km resolution grid for Switzerland (MeteoSwiss, 2013). All four datasets are developed by spatial interpolation of quality-checked data collected at MeteoSwiss meteorological stations (Frei et al., 2006; Frei, 2014). We apply the statistical analysis of change to basin-averaged values of precipitation and temperature and not to individual grid point values, which might be potentially affected by substantial interpolation errors. Moreover, the variability in time of the number of stations involved in the spatial interpolation may induce non-homogeneities in the datasets. This is particularly relevant when analyzing long-term changes as in the case of this study. Therefore, we verify the effects of potential non-homogeneities by using an experimental dataset developed by MeteoSwiss specifically for this research, based on a constant

number of stations (294 for precipitation and 48 for temperature) for the period 1971–2013. Results confirm the validity of the original datasets for analyzing long-term changes.

4.2 Discharge and Suspended Sediment Concentration

We use daily discharge data measured by the Swiss Federal Office for the Environment at three gauging stations: la Porte-du-Scex, Blatten Bei Naters and Blatten (Fig. 1). For suspended sediment concentration, two in-stream samples per week collected by the Swiss Federal Office for the Environment at la Porte-du-Scex are available since October 1964 (Grasso et al., 2012).

In this work, with the term fine sediment we refer to sediment transported in suspension. Previous analysis on the grain size distribution of suspended sediment at the outlet of the upper Rhône River report a bimodal distribution, with mode diameters equal to 13.7 μm (silt) for the finer fraction and 39.6 μm (silt) for the coarser grains (Santiago et al., 1992). Grain sizes cover a wide range of values, including clay (16.9%), silt (64.7%) and sand (18.4%). The mean suspended sediment size is reported to be equal to 17.7 μm (silt), and the largest grains transported in suspension during summer high flow conditions are in the range of coarse sand ($> 500 \mu\text{m}$) (Santiago et al., 1992).

4.3 Snow Cover and Glacier Data

We use snow cover maps derived from satellite imagery for the upper Rhône Basin over the period 2000–2008 processed in previous research (Fatichi et al., 2015). We use the 8-day snow cover product MOD10A2 retrieved from the Moderate Resolution Imaging Spectroradiometer (MODIS) (Dedieu et al., 2010) for the calibration and validation of the snowmelt model. MOD10A2 is provided at a 500×500 m spatial resolution, where cells are classified as snow-covered, snow-free, inland water or cloud-covered. In order to reduce the impacts of clouds in estimating snow cover fraction, maps with cloud cover greater than 30% are excluded from the dataset, resulting in a total number of usable images equal to 360, i.e. on the average 40 days per year.

The surface covered by glaciers is assigned based on the GLIMS Glacier Database (Fig. 1). Ice-covered cells identified based on the GLIMS data of 1991 give more than 10% of the upper Rhône Basin as covered by ice with a total glacier surface of almost 620 km^2 .

4.4 Digital Terrain Model

We use a digital terrain model (DTM) with 250×250 m resolution (85409 cells in total, Fig. 1), obtained by resampling a finer model (25×25 m) provided by SwissTopo in the ETH geodata portal (GeoVITe). The DTM is used as a mask for extracting climatic inputs and for elevation information in the snowmelt modelling.

5. Results

5.1 Calibration of Snowmelt and Icemelt Models

The snowmelt factor, calibrated for the upper Rhône Basin following the procedure described in Sect. 3.3, is $k_{\text{snow}} = 3.6 \text{ mm day}^{-1} \text{ }^{\circ}\text{C}^{-1}$. This value is in agreement with previous studies carried out in this region. In the upper Rhône Basin, Boscarello et al. (2014) found a snowmelt factor equal to 4.3 $\text{mm day}^{-1} \text{ }^{\circ}\text{C}^{-1}$ based on previous studies on the Toce Basin in Italy (Corbari et al., 2009). Calibration of a semi-lumped conceptual model for the three tributary catchments of the upper Rhône Basin – Lonza, Drance and Rhône at Gletsch – led to snowmelt factors equal to 6.1, 4.5 and 6.6 $\text{mm day}^{-1} \text{ }^{\circ}\text{C}^{-1}$, respectively (Schaepli et al. 2005). Differences in k_{snow} between this and previous studies are attributable to different

temporal resolution of models, lengths of calibration datasets, type and thresholds of precipitation partitioning, climatic inputs, threshold temperature for melt, and others.

The calibrated snowmelt model reproduces well the seasonal fluctuations of snow cover fraction (SCF) in the basin, with Nash–Sutcliffe efficiencies (NS) close to 0.90 and low mean square errors (MSE). The model maintains good performances also in the validation period showing very slight reduction in the goodness of fit measures (Table 2). The temporal variability of SCF is also well simulated at the basin scale. Although the comparison between observed and simulated SCF is affected by the discontinuous nature of the MODIS data (8–day resolution), Fig. 2 shows that the model with a single constant k_{snow} for the entire catchment reproduces the snow cover dynamics reasonably well for all of the studied elevation bands. At lower elevations, the model tends to slightly underestimate SCF in autumn and overestimate it in winter. This is likely related to errors in partitioning precipitation into solid and liquid form. The model performs better at higher elevation bands, even at the very highest elevations with permanent snow cover (Fig. 2 bottom). Good snow cover simulations at the highest elevation bands, where most of the glaciers are located, are a prerequisite for successful icemelt estimation.

One of the main problems of degree–day models is related to their poor performance in reproducing the spatial distribution of snow accumulation and melt in complex topography. The temperature–index approach does not take into account features that affect melting, such as topographic slope, aspect, surface roughness and albedo (Pellicciotti et al., 2005). In our case, the spatial distribution of snow cover is satisfactory, with average values of sensitivity and specificity greater than 0.7 (Table 2). Goodness of fit measures indicate that, on average, more than 70% of snow–covered and snow–free pixels are correctly identified. However, sensitivity and specificity are characterized by a strong seasonal signal. In summer, when a large part of the basin is snow–free, it is much easier for the model to capture snow–free pixels correctly than snow–covered pixels. In winter, when the basin is largely snow–covered, the situation is reversed. The true skill score, which combines both metrics, results in values around 0.5 (Table 2). Nonetheless, snow cover duration maps averaged over the period 2000–2008 for MODIS observations and simulations show a good spatial coherence (Fig. 3). In summary, we conclude that the snowmelt model represents the spatial and temporal dynamics of snow cover in the Rhône Basin satisfactorily.

The icemelt factor, calibrated following the procedure described in Sect. 3.3, is $k_{\text{ice}} = 6.1 \text{ mm day}^{-1} \text{ }^{\circ}\text{C}^{-1}$. Despite the large regional and temporal variability that characterizes melt factors, comparison with previous studies confirms that this value is reasonable for the Alpine environment (e.g. Schaepli et al., 2005; Boscarello et al., 2014). A range from 5 to 20 $\text{mm day}^{-1} \text{ }^{\circ}\text{C}^{-1}$ has been reported in the literature (e.g. Hock, 2003; Schaepli, 2005).

Calibration and validation results are summarized in Table 2. In Figure 4 it is shown the seasonal pattern of the three main components of the hydrological cycle which contribute to runoff in glacierized high Alpine sub–basins (IM, SM, R), obtained with the calibrated k_{ice} . The fit to the observed discharge is very good, with mass balance errors about 7% for the Massa and 8% for the Lonza. It should be noted that the simulated fluxes shown in Fig. 4 are upper estimates of runoff, because we are neglecting evaporation (evapotranspiration) in these high Alpine catchments. Evaporation plays indeed a secondary role in the long–term water balance in Alpine environments compared to precipitation and snowmelt (Braun et al., 1994; Huss et al., 2008). Considering that the aim of this study is to evaluate long–term changes in hydroclimatology and sediment dynamics of the upper Rhône Basin and not the short–term variability of icemelt at the daily scale, we consider the Snowmelt and Icemelt model performances as satisfactory. However, in applications which require spatial distributions of snow and icemelt we recommend to use approaches based on the energy balance (e.g. Pellicciotti et al., 2005).

5.2 Temperature, Precipitation, and SSC in the Rhône Basin

Mean annual air temperature shows a clear and statistically significant increase in 1987 ($p\text{-value} < 0.01$). A two–sample t –test for equal means ($p\text{-value} < 0.01$) confirms an increase in mean daily temperature greater than 1 $^{\circ}\text{C}$ (Fig. 5a). Such dramatic jumps rather than a gradual change in air temperature have been observed globally (e.g. Jones and Moberg, 2003;

Rebetez and Reinhard, 2008). Observations indicate that Switzerland has experienced two main rapid warming periods in the past, with the 1940s and 1980s being the warmest decades of the last century (Beniston et al., 1994; Beniston and Rebetez, 1996). Statistical tests on monthly means reveal that the 1987 temperature jump is mainly in spring and summer months from March to August, while changes in the autumn and winter months are not statistically significant (Fig. 6a). For the period March–August, mean monthly temperatures have risen by about 1.2 °C on the average.

The change in air temperature around 1987 coincides with statistically significant changes in mean annual suspended sediment concentration, which has increased by about 70 mg l⁻¹ (Fig 5c). This change can be ascribed to statistically significant (p -value < 0.01) increases in summer (July–August) concentrations (Fig 6c). After the abrupt warming, mean annual suspended sediment concentrations are roughly 40% larger than before, with average values rising from 172 ± 6.86 mg l⁻¹ before 1987 up to 242 ± 14.45 mg l⁻¹ after 1987, where the ranges express the standard error of the mean. The simultaneous increase in temperature and suspended sediment concentration indicates that changes in climatic conditions may effectively impact sediment dynamics, especially in Alpine environments where temperature-driven processes, like snow and ice-melt, have a strong influence on the basin hydrology. Suspended sediment concentration is also characterized by much larger inter-annual variability after 1987 than before. A statistically significant change in the variance supports the finding that processes related to fine sediment regime of the upper Rhône Basin have been altered by changing climatic conditions, resulting in greater concentrations and higher variability of suspended sediment reaching the outlet of the basin.

While the upper Rhône Basin underwent an abrupt warming around 1987, mean annual precipitation (Fig. 5b) and mean monthly precipitation (Fig. 6b) do not change significantly in time. Likewise, mean annual discharge does not show any statistically significant change over time (Fig 5d). Mean monthly discharge (Fig. 6d) is characterized by a small statistically significant increase in winter (November–February) runoff, most likely due to increased snowmelt and possibly changes in hydropower generation. Periodic decadal-scale oscillations appear to be present in both mean annual precipitation and mean annual runoff. These may be caused by large-scale climatic patterns, e.g. North Atlantic Oscillation (NAO), especially during winter months (Hurrell, 1995; Hurrell et al., 2003; Casty et al., 2005). Positive phases of NAO are associated with increased moist and warm air over Western Europe and consequent enhanced winter precipitation in Scandinavia and lower precipitation in Southern Europe (Hurrell, 1995; Hurrell et al., 2003; Bartolini et al., 2009). Although the Alpine region is recognized to be in a transition region that weakens the effect of NAO on climatic conditions (Casty et al., 2005; Bartolini et al., 2009), a correlation between decadal-frequency oscillations patterns and climatic features has been demonstrated in some Swiss locations (Beniston and Jungo, 2002).

In summary, differences in precipitation before and after 1987 are within the 95% confidence interval and are not statistically significant. Differences in discharge are also not statistically significant except in winter, when the suspended sediment concentration doesn't show changes. Therefore, it is very unlikely that the abrupt increase in suspended sediment concentration around mid-1980s in July and August is caused by changes in mean precipitation and/or discharge, i.e. transport capacity.

5.3 Hydroclimatic Activation of Sediment Sources

The concentration of suspended sediment transported at the outlet of the catchment does depend both on the transport capacity (discharge) and on the sediment supply. Indeed, given the same discharge (transport capacity), actual suspended sediment concentration covers a wide range of values (Fig. 7). Sediment supply depends on many factors, most importantly the spatial location of sediment sources (e.g. different lithology, distance to outlet, connectivity) and the specific processes of sediment production (e.g. hillslope erosion, glacial erosion, release of subglacially stored sediment, channel bed and bank erosion, mass wasting events) and transport (e.g., hysteresis). All these factors contribute to the variability around sediment rating curves (Fig. 7). Since we demonstrate that discharge has not changed significantly, the increase in suspended sediment concentration observed at the outlet of the Rhône Basin during mid-1980s is more likely related to altered sediment supply

conditions than to a greater transport capacity. Therefore, to investigate the link between the rise in mean air temperature and suspended sediment concentration, we consider here the three main sediment production and transfer processes typical of Alpine environments: (1) the continuous effect of snowmelt runoff on hillslope and channel erosion, (2) the intermittent but potentially considerable contribution of hillslope, channel bed and bank erosion, and mass wasting triggered by rainfall events, (3) the sediment-rich flux coming from glacierized areas during the icemelt season. We analyse the time series of simulated snowmelt, snow cover fraction, icemelt and effective rainfall at annual (Fig. 8) and at monthly (Fig. 9) timescales, to identify possible changes in those variables around 1987. Results are described in the following.

First, mean annual simulated snowmelt (SM) shows a decreasing tendency during the last thirty years (Fig. 8a). The reduction in snowmelt after 1987 occurs mostly in summer and early autumn (Fig. 9a) mainly due to poor snow cover. However, except July and September, the changes in all months are within the 95% confidence interval. The increase of snowmelt in March and April is due to warmer temperatures in spring. Results are coherent with the temporal evolution of simulated snow cover fraction, which is also gradually decreasing (Fig. 8b), especially in spring and summer (Fig. 9b). Statistical analysis reveals a step-like reduction of more than 10% for mean annual values of snow cover fraction in 1987 (p -value < 0.01).

Our simulations are in agreement with snow observations across Switzerland. The decreasing tendency in snow cover after mid or late 1980s has been demonstrated for the Swiss Alps (Beniston, 1997; Laternser and Schneebeli, 2003; Scherrer et al., 2004; Marty, 2008; Scherrer et al., 2006). Snow depth, number of snowfall days, and snow cover show similar patterns during the last century: a gradual increase until the early 1980s, interrupted in late 1950s and early 1970s, and a statistically significant decrease afterwards (Beniston, 1997; Laternser and Schneebeli, 2003). Previous analyses also state that the reduction in snow cover after mid-1980s is characterized more by an abrupt shift than by a gradual decrease (Marty, 2008), in agreement with our simulations. The reduction in snow cover duration, which is observed to be stronger at lower and mid altitudes than at higher elevations, is mainly the result of earlier snow melting in spring due to warmer temperatures (Beniston, 1997; Laternser and Schneebeli, 2003; Marty, 2008). Moreover, by analysing 76 meteorological stations in Switzerland, Serquet et al. (2011) demonstrated a sharp decline in snowfall days relative to precipitation days, both for winter and early spring, showing the impact of higher temperature on reduced snowfall, independently of variability in precipitation frequency and intensity. Therefore, despite the high complexity that characterizes snow dynamics in the Alps (Scherrer et al., 2006; 2013), the dominant effect of temperature rise on snow cover decline after late 1980s has been clearly shown (Beniston, 1997; Marty, 2008; Serquet et al. 2011; Scherrer et al., 2004; 2006).

Second, although mean annual and monthly precipitation were shown not to change significantly in the mid-1980s, effective rainfall (ER) on snow free areas has increased, especially in early summer (Fig. 8d, Fig. 9d). The direct impact of rainfall on sediment detachment and transport depends on precipitation form, soil and land cover, as well as snow cover which protects the soil from the erosive effect of rainfall. Therefore, ER provides a metric for evaluating changes in the erosive power of rainfall during last forty years. The effective rainfall increases in conjunction with decreases in snow cover fraction, and a statistically significant jump is identified in 1987 (p -value < 0.01) (Fig. 8d). However, although snow cover fraction is significantly lower throughout the entire melting season, only June and especially July show statistically significant increases in ER after 1987 (Fig. 9d).

The increase in potentially erosive rainfall, is partially confirmed by recent observations. Rainfall erosivity, expressed by the R-factor of the Revised Soil Loss Equation (Wischmeier and Smith, 1978; Brown and Foster, 1987), computed on the basis of 10 min resolution precipitation data, was recently analysed for Switzerland. Although, the upper Rhône Basin together with the Eastern part of Switzerland was found to have relatively low rainfall erosivity (low R-factor) compared to the rest of the country due to a lower frequency of thunderstorms and convective events (Schmidt et al., 2016), there is evidence of an increasing trend for the R-factor from May to October during the last 22 years (1989–2010) (Meusburger et al., 2012).

This reinforces our argument that the increase in effective rainfall on snow-free surfaces may have contributed to suspended

sediment concentration rise, through a combination of reduced snow cover fraction, increased rainfall–snowfall ratio and possible increases in rainfall intensity at sub-daily scale. However, simulations show a statistically significant jump in effective rainfall in June and July, while SSC is significantly larger in July and August. Therefore, we argue that erosive rainfall alone is unlikely to explain the abrupt jump in suspended sediment concentration observed around mid-1980s.

Third, our results show that temporal evolution of icemelt is consistent with suspended sediment concentration rise. Although the change is rather gradual at the annual scale (Fig. 8c), the step-like increase in icemelt is evident in the ice-melting season (May–September) and reaches highest magnitudes in July and August (Fig. 9c) in conjunction with rises in suspended sediment concentration in those months. The simultaneous increase in icemelt and decrease in snowmelt suggests that the abrupt warming has led to important alterations of the hydrological regime. To evaluate this alteration, we compute the relative contribution of rainfall, snow and icemelt on the sum of these three components in July and August. The average relative contribution of icemelt has almost doubled after 1987, while the relative contribution of snowmelt has reduced by more than 30%. This indicates the substantial effect of the sharp temperature rise on the basin hydrology.

Enhanced icemelt is coherent with the observed acceleration of Alpine glacier retreat after mid-1980s. Ground-based and satellite observations, combined with mass balance analysis, reveal that current rates of glacier retreat are consistently greater than long-term averages (Paul, 2004; 2007; Haeberli, 2007). Estimations of glacier area reduction rates indicate a loss rate for the period 1985–1999, which is seven times greater than the decadal loss rate for the period 1850–1973 (Paul, 2004). Investigations with satellite data and in-situ observations suggest that the volume loss of Alpine glaciers during the last thirty years is more attributable to a remarkable down-wasting rather than to a dynamic response to changed climatic conditions (Paul, 2004; 2007). Haeberli et al. (2007) estimated that glaciers in the European Alps lost about half of their total volume (roughly $0.5\% \text{ year}^{-1}$) between 1850–1975, another 25% ($1\% \text{ year}^{-1}$) between 1975–2000, and an additional 10–15% ($2\text{--}3\% \text{ year}^{-1}$) in the period 2001–2005. The appearance of proglacial lakes and rock outcrops with lower albedo and high thermal inertia, separation of glaciers from the accumulation area, and general albedo lowering in European Alps (Paul, 2005), are among the main positive feedbacks that accelerate glacier disintegration and make it unlikely to stop in the near future (Paul, 2007). Although glacier dynamics are quite complex and involve many variables and feedbacks, the dominant role played by temperature rise in glacier wasting has been clearly demonstrated (e.g. Oerlemans, 2000). The major volume loss in the recent past in Swiss Alpine glaciers is attributable to negative mass balances during the ablation season rather than to a lower accumulation by precipitation (Huss, 2008). For small high altitude Alpine glaciers, Micheletti and Lane (2016) showed negligible ice melt contributions to runoff between the mid-1960s and mid-1980s, after which contributions increased markedly.

Fluxes coming from glaciers are notoriously rich in sediments. Very fine silt-sized sediment resulting from glacier erosion is transported in suspension most often as wash load (Aas and Bogen, 1988). Proglacial areas generally represent rich sources of sediment due to very active glacier erosive processes of abrasion, bed-rock fracturing and plucking (Boulton, 1974; Hallet et al., 1996). Glacier retreat discloses large amount of sediments available to be transported by proglacial streams. Moreover, change in climatic conditions and specifically temperature-driven glacier recession and permafrost wasting may initiate specific erosional processes that consequently enhance sediment supply in proglacial environments (Micheletti et al., 2015; Micheletti and Lane, 2016; Lane et al. 2017). We conclude that the significant increase in icemelt detected in the mid-1980s (Fig. 9c, 11c, 12) is likely to be the main cause of the sharp rise in suspended sediment concentration entering Lake Geneva, through a combination of: (1) increased discharge originated in proglacial environments, which implies higher suspended sediment concentration; (2) larger relative contribution of sediment-rich icemelt compared to snowmelt and precipitation fluxes; and (3) intensified sediment production and augmented sediment supply in proglacial areas due to rapid ice recession. In summary, while the rise in temperature can be detected for all spring and summer months, SSC has increased significantly only in July and August, which coincides with the greatest increases in simulated ice-melt. This reinforces the argument for enhanced ice-melting being the main reason for greater fine sediment concentrations at the outlet of the catchment.

6. Discussion

6.1 Glacier Retreat and Ice-melt

It should be noted that in this study we do not consider glacier evolution, i.e. changes in ice thickness due to accumulation and melt, as well as glacier ice flow. Neglecting glacier retreat rises the possibility that we overestimate the icemelt contribution over the study period. To quantify the potential effect of glacier retreat, we compare our simulations with time series produced from the Global Glacier Evolution Model (GloGEM), a model accounting for both the mass balance and glacier evolution (Huss and Hock, 2015). For comparison, we use total monthly runoff generated from glacierized surfaces of the upper Rhône basin, simulated with GloGEM for the period 1980–2010. GloGEM computes the mass balance for every 10-m elevation band of each glacier, by estimating snow accumulation, snow and ice melt, and refreezing of rain and melt water. The response of glaciers to changes in mass balance is modelled on the basis of an empirical equation between ice thickness changes and normalized elevation range parametrized as proposed by Huss (2010). Normalized surface elevation changes Δh_r are derived for each elevation band from mass balance changes (mass conservation). Starting from initial values derived by the method of Huss and Farinotti (2012), ice thickness is updated at the end of each hydrological year by applying the relation between normalized elevation range h_r and normalized surface elevation change Δh_r . The area of each glacier is finally adjusted by a parabolic cross-sectional shape of the glacier bed (Huss and Hock, 2015). GloGEM is calibrated and validated over the period 1980–2010 with estimates of glacier mass changes by Gardner et al. (2013) and in situ measurements provided by the World Glacier Monitoring Service.

Although in our hydrological model we do not include glacier evolution, the annual runoff volumes (snowmelt + icemelt + rainfall) from glacierized areas during the period 1980–2010 correlates very well with the results of GloGEM (Fig. 11a). Measures of performance confirm the agreement between the two models: the correlation coefficient is equal to 0.86 and the Nash–Sutcliffe efficiency is equal to 0.67. We are also capable to capture quite well the seasonal pattern of runoff generated from glacierized areas (Fig. 11b). Perhaps most importantly, GloGEM simulations show that total annual runoff is increasing throughout the period and there is no evidence for decreasing icemelt rates. This confirms that, although glaciers of the upper Rhône basin are retreating, melt-water discharges from glacierized and proglacial areas are increasing during the 1980–2010 period. As expected, total runoff from glacierized surfaces and icemelt are highly correlated (Fig. 11a, correlation coefficient = 0.95), thus indicating that the increase in total runoff is due to an increase of the icemelt component. Indeed, non-parametric Mann–Kendall tests indicate an increasing trend with 5% significant level. Trend slopes, estimated with the Theil–Sen estimator, confirm the agreement between the two models: we find a total runoff of $\sim 27.65 \text{ mio m}^3 \text{ year}^{-2}$ with GloGEM and $\sim 21.71 \text{ mio m}^3 \text{ year}^{-2}$ with our model. We also computed the basin-averaged mass balance accounting for snow accumulation and snow and icemelt for each hydrological year. The mean mass balance over the period 1980–2010 is equal to $-0.78 \pm 0.22 \text{ m w.e. year}^{-1}$ which is within the uncertainty range of recent studies (Fischer et al., 2015). In summary: although we do not account for glaciers retreat, our model results agree well with a state-of-the art glaciological model that includes glacier evolution. Both comparisons with GloGEM and our basin-averaged mass balance indicate that we are not significantly overestimating icemelt contribution during the period 1975–2015. We also compute the basin-averaged mass balance accounting for snow accumulation and snow and icemelt for each hydrological year. The mean mass balance rate over the period 1980–2010 is equal to $-0.78 \pm 0.22 \text{ m w.e. year}^{-1}$ which is within the uncertainty range of recent studies (Fischer et al., 2015). We therefore argue that although we do not account for glacier retreat, our model results agree with a state-of-the art glaciological model that includes glacier evolution. Both comparisons with GloGEM and our basin-averaged mass balance indicate that we are unlikely to be overestimating the icemelt contribution during the period 1975–2015. Clearly, if we were looking at future climate projections and at glaciers that are rapidly retreating we would need to include a glacier retreat. Under climate change, even the largest glacier in the basin, the Aletsch Glacier, is expected to

shrink at a rate where its icemelt contribution would start decreasing before 2050 (Farinotti et al., 2012; FOEN, 2012; Brönnimann et al., 2014).

6.2 Anthropogenic Factors and Climate Signals

Another potential explanation of increases in suspended sediment concentration may be connected to anthropogenic drivers and changes in the mid-1980s. Three main anthropogenic activities may have potentially influenced the suspended sediment regime of the upper Rhône basin: river channelization, construction of reservoirs and hydropower operations, and gravel extraction along the main stream and tributaries. The second and last large channelization project was completed in 1960 (Oliver et al., 2009), much earlier than the observed increase in SSC. Likewise, the largest reservoirs in the catchment have been in operation since 1975 (Loizeau and Dominik, 2000). Therefore, it is unlikely that these two anthropogenic factors have contributed to the SSC rise detected in mid-1980s. The same holds for mining activities. Annual volumes of gravel extracted from the Rhone, provided by the Cantonal Authorities as differences from the average over the period 1989–2014, do not show any significant correlation with mean annual suspended sediment concentration ($R^2 = 0.08$). Although gravel mining operation may perturb SSC for short periods after river bed disturbance by causing local pulses of fine sediments, this process does not affect significantly the suspended sediment load at the outlet of the basin over seasonal and annual timescales. A possibility still remains that changes in the hydropower operation itself, i.e. the distribution of flow responding to electricity demand, and the flushing of dams have increased SSC concentrations. We currently do not have any evidence for such changes and it is very unlikely that they would have long-term effects on SSC.

Finally, a possible anthropogenic effect on the results may come for the intermittent sampling of SSC (twice per week) which may affect the changes found in the observation period. We estimate this potential effect by considering total daily, basin-averaged values of the hydroclimatic variables SM, IM, ER and Q only on days corresponding to SSC-measurement days. We compare these new time series (“SSC-measurement days”) with the original ones (“all days”) by estimating the cumulative distribution functions of the variables and by testing the equality of mean monthly and annual values before and after mid-1980s. In this analysis we considered only the positive (non-zero) part of the distributions. Although extremely high and low values may indeed be missed by the sampling method, cumulative distributions of SM, IM, ER and Q on “SSC-measurement days” and on “all days” are very similar (Fig. 12). This indicates that, although SSC is measured at a fixed interval, the sampling captures accurately the process variability. In addition, results of the statistical tests on mean monthly and mean annual values of all analyzed hydroclimatic variables are unchanged. We therefore conclude that our results are not significantly influenced by the discontinuous nature of the SSC sampling.

Our results show that even in highly human impacted and regulated catchments such as the Rhône Basin, a strong climatic signal in hydrological and sediment dynamics can persist. This also suggests that the decrease in fine sediment load at the outlet of the upper Rhône Basin observed in the 1960s on the basis of sediment cores recovered in the Rhône delta region and reported by Loizeau et al. (1997), could be the result of a combined effect of hydropower system development, as it has been hypothesized (Loizeau et al., 1997; Loizeau and Dominik, 2000), but also reduced icemelt loads due to colder temperatures at the time. The cooling period, which occurred between 1950s and late 1970s (e.g. Beniston et al., 1994) was characterized by colder and snowy winters (e.g. Laternser and Schneebeli, 2003) and has been accompanied by reduced icemelt rates, glacier advance and positive glacier mass balances (Zemp et al., 2008; VAW-ETH, 2015).

The climate signal in sediment dynamics takes on particular importance in the context of climate change projections into the future. Despite the large uncertainty, future projections under different climate change scenarios show a common tendency for Switzerland, characterized by a shift from snow-dominated to rain-dominated hydrological regime, reduced summer discharge, increased winter discharge, reduced snow cover, and enhanced glacier retreat (Bavay, 2009; Juvet et al., 2011; Brönnimann et al., 2014; Fatichi et al., 2015; Huss, 2016). In contrast to these hydrological predictions, changes in sediment

fluxes are highly uncertain due to the complexity and feedbacks of the processes involved, inherent stochasticity in sediment mobilization and transport, and large regional variability in sediment connectivity across the Alpine landscape (Cavalli et al., 2013; Heckmann and Schwanghart, 2013; Bracken et al., 2015; Lane et al., 2017). Although at this stage we cannot reliably conclude in which direction sediment fluxes will change in the future, our paper clearly shows that a more process-based approach based on the connections between hydrological change and the activation of sediment sources will provide us with a better framework for analysing and attributing changes in sediment yields in Alpine catchments in the future.

7. Conclusions

The aim of this research was to analyze changes in the hydroclimatic and suspended sediment regimes of the upper Rhône Basin during the period 1975–2015. We show an abrupt increase in basin-wide mean air temperature in the mid-1980s. The simultaneous step-like increase in suspended sediment concentration at the outlet of the catchment, detected in July and August, suggests a causal link between fine sediment dynamics and climatic conditions. Two main factors link warmer climate and enhanced SSC: increased transport capacity and increased sediment supply resulting from spatial and/or temporal activation–deactivation of sediment sources. Our results show that transport capacity, through discharge, is not likely to explain the increases in SSC, because no statistically significant changes in the mid-1980s are present in Rhône Basin discharge, neither at the annual nor monthly timescales. The suggestion is that the impact of warmer climatic conditions acts on fine sediment dynamics through the activation and deactivation of different sediment sources and different sediment production and transport processes.

To understand sediment supply conditions we analyze the temporal evolution of three main sediment fluxes: (1) sediments sourced and transported by snowmelt along hillslopes and channels; (2) sediments entrained and transported by erosive rainfall events over snow-free surfaces, including hillslope and channel bank erosion, and mass wasting events; (3) fine sediment fluxes generated by glacier ice-melt. The fluxes of snow and icemelt together with snow cover fraction and rainfall are analyzed to detect changes in time and their coherence with changes in SSC.

Our results show that while mean annual precipitation does not show any evident change between the periods before and after the SSC jump in mid-1980s, potentially erosive rainfall clearly increases over time especially in June and July, but not in August. On the other hand, icemelt has significantly increased due to temperature-driven enhanced ablation. Statistically significant shifts in icemelt were identified for summer, with highest increases in July and August, in accordance with the rise in SSC. Concurrently to the temperature and SSC rise, the relative contribution of icemelt to total annual runoff (sum of rainfall, snow and ice-melt) presents a significant increase in mid-1980s, substantially altering the hydrological regime of the Rhône Basin. Based on these results we propose that climate has an effect on fine sediment dynamics by altering the three main fluxes of suspended sediment in the Rhône Basin, and that icemelt plays a dominant role in the suspended sediment concentration rise in the mid-1980s through: (1) increased flow derived from sediment-rich subglacial and proglacial areas; (2) larger relative contribution of sediment-rich icemelt compared to snowmelt and precipitation; and (3) increased sediment supply in hydrologically connected proglacial areas due to glacier recession. While snowmelt has decreased, the reduced extent and duration of snow cover may also have contributed to the suspended sediment concentration rise through enhanced erosion by heavy rainfall events over snow free surfaces.

Because changes in SSC are not consistent with changes in discharge and transport capacity, our work emphasizes how the inclusion of sediment sources and their activation through different processes of production and transport is necessary for attributing change. This analysis also demonstrates that climate-driven changes of suspended sediment dynamics are significantly strong even in highly regulated and human impacted catchments such as the Upper Rhône basin, where sediment fluxes are affected by flow regulation due to hydropower production and by grain-size dependent trapping in

reservoirs. This has consequences for climate change impact assessments and projections for Alpine catchments with hydropower systems, where climate change signals are sometimes thought to be secondary to human regulation.

Author contribution

A. Costa and P. Molnar designed the methodology. A. Costa developed the code and carried out simulations and computations. A. Costa prepared the manuscript with contributions from all co–authors. The authors declare that they have no conflict of interest.

Acknowledgements

We thank Christoph Frei (Federal Office of Meteorology and Climatology MeteoSwiss) for providing us with experimental temperature and precipitation datasets and for suggestions on the right use of MeteoSwiss gridded data and the application of statistical tests. We also thank Daniel Farinotti (Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Department of Civil, Environmental and Geomatic Engineering ETH Zurich) for providing us with GloGEM simulations, for the fruitful discussion on glaciers retreat and glacier dynamics and for kindly revising the manuscript. The Federal Office of the Environment (FOEN) provided discharge and suspended sediment concentration data. We thank Alessandro Grasso (FOEN) for the explanation on the SSC data collection procedures. Finally, we would like to thank the Valais Cantonal Authorities for supplying information on gravel mining extraction. This research was supported by the Swiss National Science Foundation Sinergia grant 147689 (SEDFATE).

References

- Aas, E. and Bogen, J.: Colors of Glacier Water, *Water Resour. Res.*, 24(4), 561–565, 1988.
- Allouche, O., Tsoar, A., Kadmon, R.: Assessing the accuracy of species distribution models: prevalence, kappa and the true skill statistic (TSS), *J. Appl. Ecol.*, 43, 1223–1232, 2006.
- Anselmetti, F.S., Bühler, R., Finger, D., Girardclos, S., Lancini, A., Rellstab, C., Sturm, M.: Effects of Alpine hydropower dams on particle transport and lacustrine sedimentation. *Aquat. Sci.*, 69, 179–198, 2007.
- Asselman, N. E. M.: Suspended sediment dynamics in a large drainage basin: the River Rhine, *Hydrol. Process.*, 13, 1437–1450, 1999.
- Asselman, N. E. M.: Fitting and interpretation of sediment rating curves, *J. Hydro.*, 234, 228–248, 2000.
- Auer, A. H.: The rain versus snow threshold temperatures, *Weatherwise*, 27, 67–67, doi:10.1080/00431672.1974.9931684, 1974.
- Ballantyne, C.K.: A general model of paraglacial landscape response, *The Holocene*, 12, 371–6, 2002.
- Bartolini, E., Claps, P., D’Odorico, P.: Interannual variability of winter precipitation in the European Alps: relations with the North Atlantic Oscillation, *Hydrol. Earth Syst. Sci.*, 13, 17–25, 2009.
- Bavay, M., Lehning, M., Jonas, T., Löwe, H.: Simulations of future snow cover and discharge in Alpine headwater catchments, *Hydrol. Process.*, 23, 95–108, 2009.
- Beguiria, S.: Validation and Evaluation of Predictive Models in Hazard Assessment and Risk Management, *Nat. Hazards*, 37, 315–329, DOI 10.1007/s11069-005-5182-6, 2006.
- Beniston, M.: Variations of snow depth and duration in the Swiss Alps over the last 50 years: Links to changes in large-scale climatic forcings, *Climatic Change*, 36, 281–300, 1997.
- Beniston, M., Junco P.: Shifts in the distribution of pressure, temperature and moisture and changes in the typical weather patterns in the Alpine region in response to the behavior of the North Atlantic Oscillation, *Theor. Appl. Climatol.*, 71, 29–42, 2002.
- Beniston, M. and Rebetez, M.: Regional behavior of minimum temperatures in Switzerland for the period 1979–1993, *Theor. Appl. Climatol.*, 53, 231–243, 1996.
- Beniston, M., Rebetez, M., Giorgi, F., Marinucci, R.: An analysis of regional climate change in Switzerland, *Theor. Appl. Climatol.*, 49, 135–159, 1994.
- Bennett, G., Molnar, P., Eisenbeiss, H., and McArdell, B. W.: Erosional power in the Swiss Alps: characterization of slope failure in the Illgraben, *Earth Surf. Proc. Land.*, 37, 1627–1640, doi:10.1002/esp.3263, 2012.
- Boscarello, L., Ravazzani, G., Rabuffetti, D., Mancini, M.: Integrating glaciers raster-based modelling in large catchments hydrological balance: the Rhône case study, *Hydrol. Process.*, 28, 496–508, doi: 10.1002/hyp.9588, 2014.
- Boulton, G.S.: Processes and patterns of glacial erosion, (In Coates, ed.), *Glacial Geomorphology*, New York State University, 41–87, 1974.
- Bracken, L. J., Turnbull, L., Wainwright, J., Bogaart, P.: Sediment connectivity: a framework for understanding sediment transfer at multiple scales, *Earth Surf. Process. Landforms* 40, 177–188, 2015.
- Braun, L. N., Aellen, M., Funk, M., Hock, R., Rohrer, M. B., Steinegger, U., Kappenberger, G., Müller-Lemans, H.: Measurement and simulation of high alpine water balance components in the Linth–Limmern head watershed (north-eastern Switzerland), *Zeitschrift für Gletscherkunde und Glazialgeologie*, 30, 161–185, 1994.
- Brown, L. C. and Foster, G. R.: Storm erosivity using idealized intensity distributions, *Transactions of the Asae*, 30, 379–386, 1987.
- Brönnimann, S., Appenzeller, C., Croci-Maspoli, M., Fuhrer, J., Grosjean, M., Hohmann, R., Ingold, K., Knutti, R., Liniger, M. A., Raible, C. C., Röthlisberger, R., Schär, C., Scherrer, S. C., Strassmann, K., Thalman, P.: Climate change in

- Switzerland: a review of physical, institutional, and political aspects, *WIREs Clim Change* 2014. doi: 10.1002/wcc.280, 2014.
- Campbell, F. B., and Bauder, H. A.: A rating curve method for determining silt discharge of streams, *Am. Geophys. Union Trans.*, part 2, 603–607, 1940.
- Casty, C., Wanner, H., Luterbacher, J., Esper, J., Böhm, R.: Temperature and precipitation variability in the European Alps since 1500, *Int. J. Climatol.* 25, 1855–1880, DOI: 10.1002/joc.1216, 2005.
- Cavalli, M., Trevisani, S., Comiti, F., Marchi, L.: Geomorphometric assessment of spatial sediment connectivity in small Alpine catchments, *Geomorphology*, DOI: 10.1016/j.geomorph.2012.05.007, 2013.
- Collins, W. D., Rasch, P. J., Boville, B. A., Hack, J. J., McCaa, J. R., Williamson, D. L., Kiehl, J. T., Briegleb, B.: Description of the NCAR community atmosphere model (CAM3), Tech. Rep. NCAR/TN– 464+STR, 226, 2004.
- Corbari, C., Ravazzani, G., Martinelli, J., Mancini, M.: Elevation based correction of snow coverage retrieved from satellite images to improve model calibration, *Hydrol. Earth Syst. Sci.*, 13, 639–649, 2009.
- Dedieu, J.–P., Boos, A., Kiage, W., Pellegrini, M.: Snow cover retrieval over Rhône and Po river basins from MODIS optical satellite data (2000–2009), *Geophys. Res. Abstracts*, 12, 5532, EGU General Assembly, 2010.
- Delunel, R., van der Beek, P., Bourlès, D., Carcaillet J., Schlunegger, F.: Transient sediment supply in a high–altitude Alpine environment evidenced through a ^{10}Be budget of the Etages catchment (French Western Alps). *Earth Surf. Proc. Land.*, 39, 890–899. DOI:10.1002/esp.3494, 2014.
- De Vente, J., Poesen, J., Bazzoffi, P., Van Rompaey, A., Verstraeten, G.: Predicting catchment sediment yield in Mediterranean environments: the importance of sediment sources and connectivity in Italian drainage basins, *Earth Surf. Process. Landforms*, 31, 1017–1034, 2006.
- England, P. and Molnar, P.: Surface uplift, uplift of rocks, and exhumation of rocks, *Geology*, 18, 1173–1177, 1990.
- Farinotti, D., Usselman, S., Huss, M., Bauder, A., Funk M.: Runoff evolution in the Swiss Alps: projections for selected high-alpine catchments based on ENSEMBLES scenarios, *Hydrol. Process.*, 26, 1909–1924, 2012.
- Faticchi, S., Rimkus, S., Burlando, P., Bordoy, R., Molnar, P.: High–resolution distributed analysis of climate and anthropogenic changes on the hydrology of an Alpine catchment, *J. Hydrol.*, 525, 362–382, 2015.
- FOEN: Auswirkungen der Klimaänderung auf Wasserressourcen und Gewässer. Synthesebericht zum Projekt “Klimaänderung und Hydrologie in der Schweiz” (CCHydro). Bundesamt für Umwelt, Bern. Umwelt–Wissen, Nr. 1217, 76, 2012.
- Fischer, M., Huss, M., Hoelzle, M.: Surface elevation and mass changes of all Swiss glaciers 1980–2010, *Cryosphere*, 9, 525–540, doi:10.5194/tc-9-525- 2015, 2015.
- Foster, G.C., Dearing, R.A., Jones, R.T., Crook, D.S., Siddle, D.J., Harvey, A.M., James, P.A., Appleby, P.G., Thompson, R., Nicholson, J., Loizeau, J.–L.: Meteorological and land use controls on past and present hydro–geomorphic processes in the pre–alpine environment: an integrated lake–catchment study at the Petit Lac d'Annecy, France. *Hydrol. Process.*, 17, 3287–3305, 2003.
- Frei, C.: Interpolation of temperature in a mountainous region using nonlinear profiles and non–Euclidean distances, *Int. J. Climatol.*, 34, 1585–1605, 2014.
- Frei, C., R. Schöll, S. Fukutome, J. Schmidli and P.L. Vidale: Future change of precipitation extremes in Europe: An intercomparison of scenarios from regional climate models. *J. Geophys. Res.*, 111, D06105, doi:10.1029/2005JD005965, 2006.
- Gabbud, C. and Lane, S.N.: Ecosystem impacts of Alpine water intakes for hydropower: the challenge of sediment management. *WIREs Water*, 3(1), 41–61. Doi:10.1002/wat2.1124, 2016.
- Gardner, A. S., Moholdt, G., Cogley, J. G., Wouters, B., Arendt, A. A., Wahr, J.: A reconciled estimate of glacier contributions to sea level rise: 2003 to 2009, *Science*, 340, 852–857, doi:10.1126/science.1234532, 2013.

- Grasso, A., Bérod, D., Hodel, H.: Messung und Analyse der Verteilung von Schwebstoffkonzentrationen im Querprofil von Fliessgewässern, «Wasser Energie Luft» – 104, Jahrgang, 2012, Heft 1, CH–5401 Baden, 2012.
- Haeberli, W. and Holzhauser, H.: Alpine glacier mass changes during the past two millennia, *PAGES News*, 11, 1, 13–15, 2003.
- Haeberli, W., Hoelzle, M., Paul, F., Zemp, M.: Integrated monitoring of mountain glaciers as key indicators of global climate change: the European Alps, *Ann. Glaciol.*, 46(1), 150–160, 2007.
- Hallett, B., Hunter, L., Bogen, J.: Rates of erosion and sediment evacuation by glaciers: A review of field data and their implications, *Global and Planetary Change*, 12, 213–235, 1996.
- Heckmann, T. and Schwanghart, W.: Geomorphic coupling and sediment connectivity in an alpine catchment — Exploring sediment cascades using graph theory, *Geomorphology*, 182, 89–103, 2013.
- Herman, F., Beyssac, O., Brughelli, M., Lane, S.N., Leprince, S., Adatte, T., Lin, J.Y.Y. and Avouac, J.P.: Erosion by an Alpine glacier. *Science*, 350, 193–5, 2015.
- Hinderer, M., Kastowski, M., Kamelger, A., Bartolini, C., Schlunegger, F.: River loads and modern denudation of the Alps – A review. *Earth–Science Reviews*, 118, 11–44, 2013.
- Hock, R.: Temperature index melt modelling in mountain areas, *J. Hydrol.*, 282, 104–115, 2003.
- Hornung, J., Pflanz, D., Hechler, A., Beer, A., Hinderer, M., Maisch, M. and Bieg, U.: 3–D architecture, depositional patterns and climate triggered sediment fluxes of an alpine alluvial fan (Samedan, Switzerland), *Geomorphology*, 115, 202–14, 2010.
- Horowitz, A. J.: An evaluation of sediment rating curves for estimating suspended sediment concentrations for subsequent flux calculations, *Hydrol. Process.*, 17, 3387–3409, 2003.
- Huggel, C., Clague, J. J. and Korup, O.: Is climate change responsible for changing landslide activity in high mountains? *Earth Surf. Proc. Land.*, 37, 77–91, 2012.
- Hurrell, J. W.: Decadal trends in the North Atlantic Oscillation and relationship to regional temperature and precipitation, *Science*, 269, 676–679, 1995.
- Hurrell J. W., Kushnir Y., Ottersen G., Visbeck M.: The North Atlantic Oscillation: Climate Significance and Environmental Impact, *Geophysical Monograph Series*, 134, American Geophysical Union., 10.1029/134GM01, 2003.
- Huss, M. and Farinotti, D.: Distributed ice thickness and volume of all glaciers around the globe, *J. Geophys. Res.*, 117: F04010, doi: 10.1029/2012JF002523, 2012.
- Huss, M. and Hock, R.: A new model for global glacier change and sea–level rise, *Front. Earth Sci.*, 3:54, doi: 10.3389/feart.2015.00054, 2015.
- Huss, M., Bauder, A., Funk, M., Hock, R.: Determination of the seasonal mass balance of four Alpine glaciers since 1865, *J. Geophys. Res.–Atmos.*, 113, F01015, doi:10.1029/2007JF000803, 2008.
- Huss, M., Farinotti, D., Bauder, A., Funk, M.: Modelling runoff from highly glacierized alpine drainage basins in a changing climate, *Hydrol. Process.*, 22, 3888–3902, 2008.
- Huss, M., Juvet, G., Farinotti, D., Bauder, A.: Future high–mountain hydrology: a new parameterization of glacier retreat, *Hydrol. Earth Syst. Sci.*, 14, 815–829, doi:10.5194/hess-14-815-2010, 2010.
- Huss, M. and Fischer, M.: Sensitivity of Very Small Glaciers in the Swiss Alps to Future Climate Change, *Front. Earth Sci.*, 4–34, doi: 10.3389/feart.2016.00034, 2016.
- Jones, P.D. and Moberg, A.: Hemispheric and large–scale surface air temperature variations: an extensive revision and an update to 2001, *J. Climate*, 16, 206–223, 2003.
- Juvet, G., Huss, M., Funk, M., Blatter, H.: Modelling the retreat of Grosser Aletschgletscher, Switzerland, in a changing climate, *J. Glaciol.*, 57(206), 1033–1045, 2011.

- Kondolf, G. M. et al.: Sustainable sediment management in reservoirs and regulated rivers: Experiences from five continents, *Earth's Future*, 2, 256–280, doi:10.1002/2013EF000184, 2014.
- Lane, S.N., Bakker, M., Gabbud, C., Micheletti, N., Saugy, J. N: Sediment export, transient landscape response and catchment–scale connectivity following rapid climate warming and Alpine glacier recession, *Geomorphology*, <http://dx.doi.org/10.1016/j.geomorph.2016.02.015>, 2016.
- Laternser, M. and Schneebeli, M.: Long-term Snow Climate Trends of the Swiss Alps (1931–99), *Int. J. Climatol.* 23, 733–750, 2003.
- Lenzi, M. A., Mao, L., Comiti, F.: Interannual variation of suspended sediment load and sediment yield in an alpine catchment, *Hydrolog. Sci. J.*, 48:6, 899–915, DOI: 10.1623/hysj.48.6.899.51425, 2003.
- Lenzi, M.A. and Marchi, L.: Suspended sediment load during floods in a small stream of the Dolomites (northeastern Italy). *Catena* 39, 267–282, 2000.
- Loizeau, J.–L. and Dominik J., Luzzi, T., Vernet J.–P.: Sediment Core Correlation and Mapping of Sediment Accumulation Rates in Lake Geneva (Switzerland, France) Using Volume Magnetic Susceptibility, *J. Great Lakes Res.*, 23(4), 391–402, 1997.
- Loizeau, J.–L. and Dominik J., Evolution of the Upper Rhône River discharge and suspended sediment load during the last 80 years and some implications for Lake Geneva, *Aquat. Sci.*, 62, 54–67, 1015–1621/00/010054–14, 2000.
- Mao, L. and Carrillo, R.: Temporal dynamics of suspended sediment transport in a glacierized Andean basin, *Geomorphology*, doi:10.1016/j.geomorph.2016.02.003, 2016.
- Marty, C.: Regime shift of snow days in Switzerland. *Geophys. Res. Lett.*, 35, L12501. DOI: 10.1029/2008GL033998, 2008.
- Mason, S. J. and Graham, N. E.: Conditional Probabilities, Relative Operating Characteristics, and Relative Operating Levels, *Weather Forecast.*, 14(5), 713–725, 1999.
- Meteoswiss, Federal Office of Meteorology and Climatology: Documentation of MeteoSwiss Grid–Data Products Daily Precipitation (final analysis): RhiresD, http://www.meteoswiss.admin.ch/content/dam/meteoswiss/de/service-und-publikationen/produkt/raeumliche-daten-niederschlag/doc/ProdDoc_RhiresD.pdf, 2013.
- Meteoswiss, Federal Office of Meteorology and Climatology: Documentation of MeteoSwiss Grid–Data Products Daily Mean, Minimum and Maximum Temperature: TabsD, TminD, TmaxD, <https://www.ethz.ch/content/dam/ethz/special-interest/baug/ifu/hydrology-dam/documents/research-data/ifu-hydrologie-data-proddoctabsd.pdf>, 2013.
- Meusburger K., Alewell C.: Soil Erosion in the Alps. Experience gained from case studies (2006–2013). Federal Office for the Environment, Bern. Environmental studies no. 1408: 116 pp, 2014.
- Meusburger, K., Steel, A., Panagos, P., Montanarella, L., Alewell, C.: Spatial and temporal variability of rainfall erosivity factor for Switzerland, *Hydrol. Earth Syst. Sci.*, 16, 167–177, 2012.
- Micheletti, N., Lambiel, C., Lane, S.N.: Investigating decadal–scale geomorphic dynamics in an alpine mountain setting, *J. Geophys. Res. Earth Surf.*, 120, 2155–2175, doi:10.1002/2015JF003656, 2015.
- Micheletti, N. and Lane, S. N.: Water yield and sediment export in small, partially glacierized Alpine watersheds in a warming climate, *Water Resour. Res.*, 52(6), 4924–4943, doi: 10.1002/2016WR018774, 2016.
- Moore, J. R., Egloff, J., Nagelisen, J., Hunziker, M., Aerne, U. and Christen, M.: Sediment Transport and Bedrock Erosion by Wet Snow Avalanches in the Guggigraben, Matter Valley, Switzerland, *Arct. Antarct. Alp. Res.*, 45, 350–62, 2013.
- Nash, J. E. and Sutcliffe, J. V.: River Flow Forecasting Through Conceptual Models Part 1 – A Discussion of Principles, *J. Hydrol.*, 10, 282–290, 1970.
- Oerlemans, J. and Reichert, B.K.: Relating glacier mass balance to meteorological data using a Seasonal Sensitivity Characteristic (SSC). *J. Glaciol.*, 46(152), 1–6, 2000.
- Oliver, J.–M., Carrel, G., Lamouroux, N., Dole–Oliver, M.–J., Malard, F., Bravard, J.–P., Amoros, C.: The Rhône River Basin, In *Rivers of Europe*, Chpt. 7, pp 247–295, Academic Press, London, 2009.

- Palazón, L., Navas, A.: Land use sediment production response under different climatic conditions in an alpine–prealpine catchment, *Catena*, 137, 244–255, 2016.
- Paul, F., Kääb, A., Maisch, M., Kellenberger, T., Haeberli, W.: Rapid disintegration of Alpine glaciers observed with satellite data, *Geophys. Res. Lett.*, 31 (L21), L21402, (10.1029/2004GL020816.), 2004.
- Paul, F., Machguth, H., Kääb, A.: On the impact of glacier albedo under conditions of extreme glacier melt: the summer of 2003 in the Alps, *EARSeL eProceedings* 4(2), 139–149, 2005.
- Paul, F., Kääb, A. and Haeberli, W.: Recent glacier changes in the Alps observed from satellite: Consequences for future monitoring strategies, *Global Planet. Change*, 56(1/2), 111–122, 2007.
- Pellicciotti, F., Brock, B., Strasser, U., Burlando, P., Funk, M., Corripio, J.: An enhanced temperature–index melt model including the shortwave radiation balance: development and testing for Haut Glacier d’Arolla, Switzerland, *J. Glaciol.*, 51(175), 573–587, 2005.
- Pettitt, A. N.: A Non–parametric Approach to the Change–point Problem, *Appl. Statist.*, 28(2), 126–135, 1979.
- Quinton, W. L. and Carey, S. K.: Towards an energy–based runoff generation theory for tundra landscapes, *Hydrol. Process.*, 22, 4649–4653, 2008.
- Rebetez, M. and Reinhard, M.: Monthly air temperature trends in Switzerland 1901–2000 and 1975–2004, *Theor. Appl. Climatol.*, 91, 27, doi 10.1007/s00704–007–0296–2, 2008.
- Santiago, S., Thomas, R. L., McCarthy, L., Loizeau, J. L., Larbaigt, G., Corvi, C., Rossel, D., Tarradellas, J., Vernet, J. P.: Particle Size Characteristics of Suspended and Bed Sediments in The Rhône River, *Hydrol. Process.*, Vol. 6, 227–240, 1992.
- Schaepli, B., Hingray, B., Niggli, M., Musy, A.: A conceptual glaciohydrological model for high mountainous catchments. *Hydrol. Earth Syst. Sc.*, 9, 95–109, 2005.
- Scherrer, S.C., Appenzeller, C.: Swiss Alpine snow pack variability: major patterns and links to local climate and large–scale flow. *Climate Res.*, 32(3), 187–199, 2006.
- Scherrer, S.C., Appenzeller C., Laternser, M.: Trends in Swiss Alpine snow days: the role of local– and large–scale climate variability, *Geophys. Res. Lett.*, 31, L13215. DOI: 10.1029/2004GL020255, 2004.
- Scherrer, S. C., Wüthrich, C., Croci–Maspoli, M., Weingartner, R., Appenzeller, C.: Snow variability in the Swiss Alps 1864–2009, *Int. J. Climatol.*, 33, 3162–3173, doi: 10.1002/joc.3653, 2013.
- Schlunegger, F. and Hinderer, M.: Crustal uplift in the Alps: why the drainage pattern matters, *Terra Nova*, 13, 425–432, 2001.
- Schmidt, S., Alewell, C., Panagos, P., Meusburger, K.: Seasonal Dynamics of Rainfall Erosivity in Switzerland, *Hydrol. Earth Syst. Sci. Discuss.*, doi:10.5194/hess–2016–208, 2016.
- Serquet, G., Christoph, M., Dulex, J. P., Rebetez, M.: Seasonal trends and temperature dependence of the snowfall/precipitation–day ratio in Switzerland, *Geophys. Res. Lett.*, 38, L07703, doi:10.1029/2011GL046976, 2011.
- Stutenbecker, L., Costa, A., Schlunegger, F.: Lithological control on the landscape form of the upper Rhône Basin, Central Swiss Alps, *Earth Surf. Dynam.*, 4, 253–272, 2016.
- Swift, D.A., Nienow, P.W. and Hoey, T.B.: Basal sediment evacuation by subglacial meltwater: suspended sediment transport from Haut Glacier d’Arolla, Switzerland, *Earth Surf. Proc. Land.*, 30(7), 867–883, 2005.
- Syvitski, J.P.M., Vörösmarty, C.J., Kettner, A.J., Green, P.: Impact of Humans on the Flux of Terrestrial Sediment to the Global Coastal Ocean, *Science*, 308, 2005.
- VAW-ETH, Glaciological reports (1881–2016). "The Swiss Glaciers", Yearbooks of the Cryospheric Commission of the Swiss Academy of Sciences (SCNAT) published since 1964 by the Laboratory of Hydraulics, Hydrology and Glaciology (VAW) of ETH Zürich. No. 1–134, (<http://glaciology.ethz.ch/swiss-glaciers/>).

- Vernon, A.J., Van der Beek, P.A., Sinclair, H.D., Rahn, M.K.: Increase in Late Neogene denudation of the European Alps confirmed by analysis of a fission-track thermochronology database, *Earth Planetary Sciences Letters*, 270(3–4), 316–329, 2008.
- Walling, D. E.: Suspended sediment production and building activity in a small British basin, *IAHS–AISH, Spec. Publ.*, 113, 137–144, 1974.
- Walling, D. E.: Assessing the accuracy of suspended sediment rating curves for a small basin, *Water Resour. Res.*, 13, 531–538, 1977.
- Walling, D. E.: Tracing suspended sediment sources in catchments and river systems, *Sci. Total Environ.*, 344, 159–184, 2005.
- Wen, L. J., Nagabhatla, N., Lü, S. H., Wang, S.Y.: Impact of rain snow threshold temperature on snow depth simulation in land surface and regional atmospheric model, *Adv. Atmos. Sci.*, 30, 1449–1460, 2013.
- Wick, L., Van Leeuwen, J. F. N., Van der Knaap, W. O., Lotter, A. F.: Holocene vegetation development in the catchment of Sägistalsee (1935 m asl), a small lake in the Swiss Alps. *J. Paleolimnol.*, 30, 261–272, 2003.
- Wilks, D. S.: *Statistical Methods in the Atmospheric Sciences*, Academic Press, 467, 1995.
- Wischmeier, W. H.: Rainfall erosion index for universal soil loss equation, *Soil Sci. Soc. Am. Proc.*, 23, 246–249, 1959.
- Wischmeier, W. H. and Smith, D. D.: *Predicting Rainfall Erosion Losses – A Guide to Conservation Planning*, Agric. Handbook, No. 537, Washington D.C., 58, 1978.
- Wood, J. L., Harrison, S., Turkington, T. A. R. and Reinhardt, L.: Landslides and synoptic weather trends in the European Alps, *Climatic Change*, 136, 297–308, 2016.
- Zemp, M., Paul, F., Hoelzle, M., Haeberli, W.: Glacier fluctuations in the European Alps, 1850–2000: an overview and spatio-temporal analysis of available data, in Orlove, B [et al.], *Darkening Peaks: Glacier Retreat, Science, and Society*, Berkeley, US, 152–167, 2008.
- Zerathe, S., Lebourg, T., Braucher, R. and Bourles, D.: Mid-Holocene cluster of large-scale landslides revealed in the Southwestern Alps by ³⁶Cl dating. Insight on an Alpine-scale landslide activity, *Quaternary Sci. Rev.*, 90, 106–27, 2014.

Table 1: List of the variables analysed: observed SSC, and hydroclimatic variables originating from measurements or interpolation of measurements (T, P, Q), from simulations of the snow and icemelt model (SCF, SM, IM), or a combination thereof (ER). Information on the source and the spatial and temporal resolution are reported for each variable.

Variable	Data Source	Resolution
T	Daily mean temperature [$^{\circ}\text{C}$] on a $\sim 2 \times 2$ km grid provided by MeteoSwiss	basin-averaged, daily, 1975–2015
P	Daily total precipitation [mm day^{-1}] on a $\sim 2 \times 2$ km grid provided by MeteoSwiss	basin-averaged, daily, 1975–2015
Q	Daily mean discharge [$\text{m}^3 \text{s}^{-1}$] at three stations (Porte du Scex, Blatten, Blatten bei Naters) provided by FOEN	daily, 1975–2015
SSC	Suspended sediment concentration [mg l^{-1}] at la Porte du Scex provided by FOEN	2 times per week, 1975–2012
SCF	Snow cover fraction [0–1] simulated by the Snowmelt Model on a 250×250 m grid, and calibrated with MODIS satellite data for the period 2000–2009	basin-averaged, daily, 1975–2012
SM	Snowmelt rate [mm day^{-1}] simulated by the Snowmelt model on a 250×250 m grid	basin-averaged, daily, 1975–2012
IM	Icemelt rate [mm day^{-1}] simulated by the Icemelt model on a 250×250 m grid, and calibrated at Blatten and Blatten bei Naters	basin-averaged, daily, 1975–2012
ER	Effective rainfall [mm day^{-1}] (rainfall on snow-free pixels), estimated from P, T and SCF on a 250×250 m grid	basin-averaged, daily, 1975–2012

Table 2: (top) Calibrated snowmelt factor k_{snow} and goodness of fit measures for validation and calibration period: Nash–Sutcliffe efficiency (NS), mean square error (MSE), true skill statistic (TSS), sensitivity (SE) and specificity (SP) for the entire upper Rhône Basin; (bottom) calibrated icemelt factor k_{ice} and goodness of fit measures: mass balance error computed on June–October months (MBE_S) and on the entire year (MBE_A) for Massa and Lonza sub-basins.

$k_{\text{snow}} = 3.6 \text{ mm day}^{-1} \text{ }^{\circ}\text{C}^{-1}$		
	Calibration	Validation
NS	0.88	0.86
MSE	0.01	0.01
TSS	0.54	0.46
SE	0.77	0.76
SP	0.73	0.70

$k_{\text{ice}} = 6.1 \text{ mm day}^{-1} \text{ }^{\circ}\text{C}^{-1}$		
	MBE_S [%]	MBE_A [%]
Calibration Massa	6.10	7.22
Validation Massa	6.77	9.19
Validation Lonza	11.35	10.09




Figure 1: Map of the upper Rhône Basin with topography, glacierized areas and river network. Inset shows the position of the upper Rhône Basin in Europe (blue). Locations of gauging stations used in this analysis are shown as triangles. Massa and Lonza sub-basins used in the calibration and validation of the ice melting component are highlighted.




Figure 2: Comparison between observed (red circles) and simulated (light blue lines) snow cover fraction (SCF) of the upper Rhône Basin for five different elevation bands. Simulations are computed with calibrated snowmelt factor $k_{\text{snow}} = 3.6 \text{ mm day}^{-1} \text{ }^{\circ}\text{C}^{-1}$.




Figure 3: Map of average snow permanence over the period 2000–2008, expressed as fraction of time in which pixels are snow-covered (SCDF [0 – 1]): (a) observations (MODIS) and (b) simulations.




Figure 4: Comparison of mean monthly observed (dark blue) and simulated discharge (light blue) for the period 1975–2015: (a) Massa basin and (b) Lonza basin. Simulated discharge is the sum of three components: icemelt (IM), snowmelt (SM) and rainfall (R).




Figure 5: Observations for the period 1965–2015 of: (a) basin-averaged air temperature; (b) basin-averaged daily precipitation; (c) suspended sediment concentration measured at the outlet of the basin; (d) daily discharge per unit area measured at the outlet of the basin. Mean annual values are shown in grey and the 5-years moving average is shown with a bold line.




Figure 6: Monthly differences between the period after and before the year-of-change (1965–1986 and 1987–2015) for: (a) basin-averaged air temperature; (b) basin-averaged daily precipitation; (c) mean suspended sediment concentration measured at the outlet of the basin; (d) daily discharge per unit area measured at the outlet of the basin. Points outside the confidence interval (grey shaded area) represent statistically significant (5% significance level) changes in the monthly mean.




Figure 7: Suspended sediment concentration and discharge data, measured at la Porte-du-Scex, the outlet of the upper Rhône Basin, during the period 1965–1986 (red circles) and 1987–2015 (blue triangles). Linear regression models are fitted to the logarithm of the data.




Figure 8: Simulations for the period 1975–2015: (a) mean annual snowmelt SM; (b) mean annual snow cover fraction SCF; (c) mean annual icemelt IM; (d) mean annual effective rainfall ER computed as rain falling over snow-free pixels. Mean annual values are shown in grey and a 5-year moving average is shown with a thick line.




Figure 9: Monthly differences between the period after and before the year-of-change (1975–1986 and 1987–2015): (a) mean snowmelt SM; (b) mean snow cover fraction SCF; (c) mean icemelt IM; (d) mean effective rainfall ER computed as rain falling over snow-free pixels. Points outside the confidence interval (grey shaded area) represent statistically significant (5% significance level) changes in the monthly mean.




Figure 10: Relative contribution of total annual snowmelt (SM), rainfall (R), and icemelt (IM) computed as the ratio between each component and their sum. Rainfall is extracted from observed precipitation by using a rain–snow temperature threshold, snow and icemelt are simulated with spatially distributed temperature–index models.




Figure 11: Runoff (snowmelt + icemelt + rainfall) generated from glacierized areas within the upper Rhône basin, simulated with GloGEM and with our Snowmelt and Ice-melt models (degree-day) for the period 1980–2010: (a) total annual values; (b) mean monthly values. Fig. 11a also depicts the time series of total annual Ice-melt simulated with our Ice-melt model.




Figure 12: Empirical cumulative distribution functions of total daily basin-averaged SM (a), IM (b), ER (c) and Q (d), computed on all days and only on days corresponding to SSC-measurements. Only positive values of SM, IM and ER are included.

Supplementary Material: Temperature signal in suspended sediment export from an Alpine catchment

Costa, A., Molnar, P., Stutenbecker, L., Bakker, M., Silva, T. A., Schlunegger, F., Lane, S. N., Loizeau, J.-L., Girardclos, S.

S1. Sensitivity Analysis on Snow Model parameters

We apply a sensitivity analysis on the three main parameters of our Snowmelt model: snowmelt factor k_{snow} [$\text{mm day}^{-1} \text{ }^{\circ}\text{C}^{-1}$], threshold temperature for the onset of snow melt T_{SM} [$^{\circ}\text{C}$], and the rain–snow threshold temperature T_{RS} [$^{\circ}\text{C}$]. We analyze the impact of the parameters on the model results by perturbing one single parameter at the time. We compute some of the goodness of fit measures adopted during the calibration (TSS, NS RMSE) and we analyze their relative change as function of the parameter perturbation (Fig. S1).

As expected, the snowmelt factor k_{snow} is the most sensitive parameter (Fig. S1). For k_{snow} between 1.6 and 5.6 $\text{mm day}^{-1} \text{ }^{\circ}\text{C}^{-1}$, the relative reduction of TSS and NS results respectively lower than 10% and 15% (Fig. S1). Although reducing the snowmelt factor k_{snow} below 2.6 $\text{mm day}^{-1} \text{ }^{\circ}\text{C}^{-1}$ increases RMSE by almost 40%, it results in incrementing RMSE only by 0.06 units (SCF [0–1]). For T_{SM} varying within the range of $-2 \div 2 \text{ }^{\circ}\text{C}$, TSS and NS decrease less than 10%. While the change in RMSE results larger similarly to the case of k_{snow} (Fig. S1). The effect of T_{RS} is even more negligible, with relative changes of goodness of fit measures within 5.5% (Fig. S1).




Figure S1: Parameter perturbation analysis on: (a) snowmelt factor k_{snow} , (b) threshold temperature for the onset of melt T_{SM} , (c) rain–snow threshold temperature T_{RS} . The relative change [%] of TSS, NS and RMSE are shown for each parameter.

We also estimate the influence of these three parameters on the trends and the jumps that we identified over the period 1975–2015 for the hydroclimatic variables simulated with the model: SM, IM, ER, SCF. For this purpose, we apply statistical tests for equality of mean annual and monthly values before and after mid–1980s after perturbing one parameter at the time within physically reasonable ranges of values.

In agreement with the results of our analysis (Sect. 5.3), a statistically significant increase of mean annual IM and ER and a statistically significant decrease of mean annual SCF after mid–1980s are detected for all selected parameter values. Similarly, mean annual SM shows a decreasing tendency after mid–1980s for all parameter selections, and in about 50% of all the parameters selections statistical tests even show a statistically significant drop in mean annual SM. The increases in mean monthly ER in June and July are identified respectively in more than 90% and in 100% of the cases. Similarly, all

parameters sets show an abrupt rise of IM after mid-1980s for all spring and summer months (May–August). This confirms that by perturbing the parameters of the snow model within reasonable ranges, the overall results of our analysis do not change. This is also depicted in Fig. S2. The differences in mean monthly values of IM and ER between the period after and before mid-1980s are shown for multiple values of each of the three parameters. The confidence interval depicted in Fig. S2 is artificially built by selecting, for each month, the highest and the lowest values of the confidence interval (5% significance level) among all parameter sets. The comparison between Fig. S2 and Fig. 9c and Fig. 9d, shows that changing parameters within reasonable ranges would not substantially change the results of our analysis.




Figure S2: Sensitivity analysis on monthly differences between the periods after and before the year-of-change (1987–2015 and 1975–1986) for IM (left) and ER (right). Box plots represent monthly differences for all selections of parameters within given ranges of values: $1.6 \text{ mm day}^{-1} \text{ }^{\circ}\text{C}^{-1} \leq k_{\text{snow}} \leq 5.6 \text{ mm day}^{-1} \text{ }^{\circ}\text{C}^{-1}$ (a,b); $-2^{\circ}\text{C} \leq T_{\text{SM}} \leq 2^{\circ}\text{C}$ (e,f); $-1^{\circ}\text{C} \leq T_{\text{RS}} \leq 3^{\circ}\text{C}$ (c,d). Grey shaded areas represent the widest confidence interval among all selections of the parameters.