

Dear Editor,
Dear Referees,

We would like to thank you for the detailed comments on our discussion paper. According to the suggestions of the referees we shortened some sections of the manuscript, added some further remarks to the introduction and the abstract, and addressed all of the specific comments of the two referees. Please find below our response to the referees' comments (as published in the open discussion) and the revised version of the manuscript in track changes mode.

Kind regards,
Florian Hanzer
(on behalf of all co-authors)

Reply to Reviewer #1

The authors present a very interesting and thorough study of the effects of a changing climate on cryospheric and hydrologic processes in the Austrian Alps over the next century. The authors use a well-established physically based model, which employs a full energy balance approach to simulate snow cover evolution, and a comprehensive future climate data set to look in detail at several processes in the Ötztal Alps. Additionally and commendably, the modelling method used in the study considers the change of glaciers in terms of ice thickness and areal extent throughout the modelling period. Results are presented for the development of the glaciers in the study area, the changing depth and length of the snow covers, and for the amount and timing of the runoff in the area.

While comparable studies have been carried out in other parts of the European Alps (particularly Switzerland) this seems to be the first comprehensive study in the Eastern (Austrian Alps). This along with the already mentioned fact that the current study employs an energy balance model and includes glacier evolution in their analysis make this study very interesting for a larger readership. Overall, the paper is very well written. In total, it seems to be a bit on the long side. Maybe the authors could go over the paper again, trying to find some sections that could be shortened a bit. Especially the Method section seems to have some potential for shortening (in my opinion). Furthermore, some of the individual sentences seem to rather long which makes it a bit hard to read at times. It would probably be better to simply subdivide these sentences into two or three separate ones. Again I would encourage the authors to have one more look at the manuscript with this problem in mind.

The Abstract seems to me to be very concise while addressing all the major important points of the chosen method and the most significant results of the study. The introduction provides an adequate overview of previous studies in this field. The actual goals of this study are mentioned in one (rather short paragraph) at the end of the introduction. Here I would encourage the authors to maybe expand this part a little, explaining why this study area and this method was chosen and what sets this study apart from others before. The study site and data presentation are good and complete. The "Methods" section is very thorough, maybe even, as mentioned, a little on the long side with some potential for shortening. The "Results and Discussion" has a very logical setup and presents all relevant results clearly and concisely. Also included are appropriate comparisons of the results of the current studies to previous related studies. I especially applaud the authors for including an extensive section about the uncertainties in the modelling. Especially in modelling studies of the impact of future climate scenarios, such a section is highly valuable. Finally the "Conclusions" present the major results of the study in a concise form. The conclusions are well based on the results and give the reader a good summary. The Tables and figures are adequate and present the information in an easily understandable form.

Overall, this is in my opinion a well thought out and well presented study. It covers a topic that is one of the most pressing questions of the coming decades not only in a scientific sense, but also for society as a whole (winter tourism, freshwater availability, etc.). The chosen geographic location, which to this date has not been studied in this context and the use of a completely physically based hydro-climatological model along with an algorithm tracking the evolution of the glaciers throughout the model period present, in my opinion, a significant new addition to the overall scientific knowledge. The paper's topic falls well within the scope of the journal and is of interest to other scientists, but also to the general public

and political decision makers. I would therefore recommend publication after minor revisions. As noted above, these revisions could address the overall length of the paper and the sometimes excessively long sentences, as well as the following minor specific comments.

We would like to thank the reviewer for their thorough evaluation of our manuscript and the very helpful and constructive comments, which are very much appreciated. With regard to your general comments, we agree that the paper is comparatively long and could benefit from some shortening. As a result of some of your comments and those of reviewer #2, we will remove some paragraphs in the revised version of the manuscript. In addition, we will expand on the goals and novelty of the study in the introduction. Please find below our replies to the individual comments.

I'm guessing that the layout of the article is not the final version, but as it is now, some of the Figures are quite far away from where they are discussed in the text. This makes it somewhat hard to follow the discussion. I would make sure that the Figures are placed closer to the text discussion of them in the final version.

We definitely agree that the placement of the figures in the discussion paper is unfortunate, and will make sure that this is improved for the final version.

p.1 line 7-9: I would cut this sentence into two by putting a period after "century" and describing the situation below 1500m asl in a second sentence.

We will change this sentence to: "Results show generally declining snow amounts with moderate decreases (0–20 % depending on the emission scenario) of mean annual snow water equivalent in high elevations (> 2500 m a.s.l.) until the end of the century. The largest decreases, amounting to up to 25–80 %, are projected to occur in elevations below 1500 m a.s.l."

p.2 line 7-9: This sentence ("These general ... ") reads very awkward. Please try to rephrase.

We will change this sentence to: "Since these projected impacts are mainly triggered by increasing temperatures (shift from snowfall to rainfall, earlier onset of snowmelt), the likelihood of occurrence is very high. In lower-elevated catchments on the other hand, projected changes in precipitation exhibit a larger impact on changes in runoff (Horton et al., 2006)."

p.4: The authors mention three climate scenarios. However, the scenarios are not explained any further until in the results section. Maybe you could add a small paragraph explaining these scenarios i.e. what assumptions they make, how they are situated in the overall "scenario ensemble" (high, medium or low change assumptions)

We will add a short paragraph explaining the main characteristics of the three RCPs to the respective section of the manuscript: "For the scenario simulations until 2100, we used the EURO-CORDEX climate change projections (Jacob et al., 2013) as climatic forcing, while considering the scenarios RCP2.6, RCP4.5, and RCP8.5. The latter is a scenario assuming no implementation of climate mitigation policies, resulting in considerably and steadily increasing emissions and concentrations of greenhouse gases over time. RCP4.5 is an intermediate scenario consistent with peaking emissions around the mid-century and a strong decline afterwards, resulting in stabilizing concentrations by the end of the century. Finally, the intervention scenario RCP2.6 is at the very low end of the spectrum in terms of

future emissions and radiative forcing, assuming a peak in CO₂ concentrations in the middle of the century and a slow decline afterwards along with negative emissions toward the end of the century."

p.6 line 15: Was terrain orientation and slope included in the calculation of the incoming solar radiation or were the model grid cells assumed to be "flat"?

Yes, in the radiation model for each grid cell a unit vector normal to the surface is calculated which is the basis for the subsequent calculations. We will add this information to the manuscript and change P6 L15f. to "(...) while taking into account terrain slope and orientation, hill shading, transmission losses and gains due to scattering, absorption, and reflections (...)"

p.6 p.32 Could you add one more sentence as to what modifications for climate conditions under forest canopies were applied?

We will change the sentence to: "In forested areas the interpolated meteorological fields are modified in order to capture sub-canopy conditions, resulting in reduced shortwave radiation, precipitation, and wind speed, increased longwave radiation and humidity, and an attenuation of the diurnal temperature cycle. Additionally the effects of the forest snow processes of interception, sublimation, and melt unload are accounted for (Strasser et al., 2011)."

p.7. Line 1: How was ground heat flux considered? Did you have ground temperatures (either measured or modelled) or was a constant value used?

Ground heat flux was assumed to be constant at 2 W/m².

p.8 line 20 and following: Is this paragraph really needed? It mainly discusses a method that is not needed for the current study. Maybe you could just explain what you did and omit the rest.

We agree. We will remove this paragraph from the manuscript.

p. 13 line 19: "This is partly explainable from the fact" reads very awkward. Maybe replace with "This can be attributed to the fact"

Done.

Results Section: Generally, comparable studies start out by showing that the chosen model "system" start out by showing that the model system was able to simulate the current state (here 1997 to 2006) adequately. There is a Table (Table 3) much later, where some model efficiency values for the modelling of the current state are shown. However the focus of this Table is on showing original versus disaggregated values. I would welcome a short section showing how well the employed model system does for the historic phase at the start of the results section.

The validation of the model setup for historical conditions has been described in detail in a previous paper (Hanzer et al., 2016), which is why we generally refer to this reference and only concisely discuss the changes in model performance regarding the few changes in model setup (reduced spatial and temporal resolution, temporal disaggregation of forcing

data) in section 4.5. We agree that this might not be immediately clear from the article and will add a short paragraph for clarification to the start of the results section.

p.13 and 14: It is not entirely clear to me why observed data was used for the 1997 to 2006 period of the “regular” model runs, while RCM data was used for the 1971 – 2005 “snow cover model runs”. Is this due to a lack of observed data from 1971 to 1997 or what is the reason for this procedure? Please explain.

Yes, the reason for using the historical RCM data instead of observed station data was to avoid introducing inhomogeneities in the simulation results due to the highly varying temporal coverage of station observations in the past. By using the historical RCM simulations and applying the same downscaling and bias correction methodology as for the future period, homogeneous and gap-free time series for the period 1971–2100 are obtained for all stations and variables. The model runs using observed data on the other hand were used (i) for validation, (ii) for deriving the ice thickness distribution for 2006 used for initializing the scenario runs, and (iii) as reference for calculating the future changes in runoff (figs. 15–17). We will clarify this also in the manuscript.

p. 15 line 17: “Comparing” should be “Compared”

Done.

p. 15 line 30: End sentence after “century”.

Done.

p.15 line 32: “However, also for the” is not a correct English sentence. Please rephrase.

We will change this sentence to: "For the other two scenarios comparatively small decreases in average SWE are simulated in this period, amounting to 427 mm (–18 %) for RCP4.5 and 357 mm (–31 %) for RCP8.5."

p. 15 line 35: For “the” station Obergurgl

Done.

p. 17 line29: “occur already” should be “already occur”

Done.

p.24 line 13: End sentence after (Huss et al 2010)

Done.

References

Hanzer, F., Helfricht, K., Marke, T., & Strasser, U. (2016). Multilevel spatiotemporal validation of snow/ice mass balance and runoff modeling in glacierized catchments. *The Cryosphere*, 10(4), 1859–1881. <http://doi.org/10.5194/tc-10-1859-2016>

Jacob, D., Petersen, J., Eggert, B., Alias, A., Christensen, O. B., Bouwer, L. M., et al. (2013). EURO-CORDEX: new high-resolution climate change projections for European impact research. *Regional Environmental Change*, 14(2), 563–578. <http://doi.org/10.1007/s10113-013-0499-2>

Reply to Reviewer #2

PREMISE

The way I got to review this manuscript tells me one more time that the current peer-review system is close to collapse: I declined the invitation two times, and had no longer the courage of declining a third time, since I assumed that - at that stage - the desperation of the handling editor must have been larger than mine. Too many papers for too few reviewers seems to be at the heart of the problem. Here is a suggestion for a simple fix: No decision shall be taken on any manuscript before the submitting group of authors has themselves provided three reviews (the typical number of reviews for a manuscript) to some other articles. In the ideal world, these three reviews could be provided to any journal with decent quality, but since different publishers seem not to have any interest in collaborating between each other, the rule will probably be implementable only at the level of individual publishers. I may have overlooked some pitfalls that would go with such a suggestion, still, I'm convinced that it is worth a thought.

SUMMARY

Hanzer and colleagues present an analysis of future runoff evolution in a high- mountain catchment in the Ötztal Alps, Austria. Their analyses are based on the latest climate projections, i.e. in the results provided in the frame of the EURO-CORDEX initiative, and the physically-based model AMUNDSEN. The used methodology is sound, the paper is well written, the figure are illustrative, and the overall quality certainly adheres to HESS's standards. I have only series of minor remarks that the authors may or may not find useful to improve their work. Other than that, I look forward to see this contribution published soon.

First of all, we wish to express that we share the reviewer's concerns regarding the current state of the peer review system. While it will likely not be an easy task to fix this problem, we wish to thank you for your valuable thoughts that we will take over as good ideas to be thoroughly considered in the community. Most importantly, we are thankful that you nevertheless took the time to review our manuscript and for the many helpful and constructive comments. Please find below our replies to the individual comments.

P1 L7ff: The abstract is well written. I was wondering, however, if the authors would be able to add half-a-sentence or so to better highlight the novelty in their contribution.

We will add the following sentence to the abstract to highlight the novelties in our work: "The high level of process representation within the model, the high spatial and temporal model resolution, and the large number and range of considered climate model runs make these findings a novel contribution to the possible impacts of future climate change in the Ötztal Alps in particular and in high-elevation Alpine catchments in general."

P3 L12ff: The wording seem to suggest that the methodology used by the authors is "better" than what has been used so far. Asked provocatively: How can the authors proof that? If such a proof is provided later in the manuscript, the authors may want to "announce" that here already.

We certainly did not intend to imply that our methodology is "better" than other approaches. While we state in the introduction that more physically based models are potentially better suited for the application under changing (climatic) conditions than simple conceptual models, in the study we emphasize also the limitations of our methodology and attempt to quantify the uncertainties in the results. To follow up on this, in a future study we will compare our results with those obtained from applying a semi-distributed conceptual model

in the same region and using the same forcing data, which will allow to investigate the uncertainties induced by different modeling approaches in more detail.

P6 L14-15: Here, it was not clear to me how the authors will handle temperature lapse rate in simulations for the future (since it is said that the lapse rates are “calculated from point measurements”). One sentence of clarification could be helpful.

For the application in this study, temperature lapse rates were not calculated dynamically but rather prescribed in the form of static monthly values which do not change in the future. In the case of dynamic calculation, the lapse rates are calculated separately for each time step by regressing the point data (which can be either actual measurements or e.g. downscaled climate model data) against elevation. We agree that the term "point measurements" in P6 L15 is too specific and will replace it by "point data".

P6 21ff: This is one of the very few points that I found conceptually problematic: The authors use a “snow correction factor (SCF)” that, basically, increases the modelled amount of precipitation when precipitation is in its solid form. What does that mean for future simulations, i.e. for simulations in a warmer climate? I see the danger for the changes in precipitation to be overestimated: Since more precipitation will be liquid in future than it was in the past, less of the precipitation will be affected by the correction factor. With a correction factor chosen to be 1.15 (L198), my suspicion is that the effect could be as large as 15

We agree that this is a valid concern. The increase in snowfall amounts by 15 % (which is applied additionally to the wind speed and temperature-dependent precipitation adjustment) is an empirical correction, however this value has been derived using a very thorough validation procedure taking into account various validation data sets such as areal precipitation, point-based snow depths, lidar-derived snow depth maps, and multi-year glacier mass balances (Hanzer et al., 2016). Although it has been shown that it is absolutely necessary to correct the observed precipitation amounts for undercatch in the study region, we agree that this fixed value might change given that climate conditions change as well. Utilizing the RCM-simulated snowfall fractions and precipitation fields instead of downscaling to point locations could reduce this uncertainty. Promising advances in downscaling methods such as quasi-dynamical approaches as e.g. in the ICAR model (Gutmann et al., 2016) could bridge this gap between statistical and dynamical methods and allow deriving more realistic small-scale precipitation fields. While this was not feasible for the present study, the revised version of the manuscript will include a short discussion on the possible uncertainties in model results induced by the precipitation downscaling and snow correction approach.

P7 L6ff: I think that the “SCF” (see above) should also be flagged as a parameter “requiring site-specific calibration”.

You are right. We will change this sentence to: "Apart from the parameters of this linear reservoir model which have to be calibrated individually for each catchment, most parameters in the model have a physical meaning, and in general no site-specific calibration beyond the parameters for the runoff module and the precipitation correction method is performed."

P10 L3ff: I might have missed it later in the discussion, but here I thought that having at least one sentence addressing the limitation of quantile mapping methods would be appropriate. I refer to the limitations in handling extremes in particular.

We agree. We will briefly expand on the limitations of QM in the respective section of the article.

P10 L9: Here I'm not sure to understand the wording "to retain intervariable relations". Maybe the authors can rephrase?

We will change the wording to: "(...) and can preserve intervariable dependency structures".

P10 L30: Several options for the number of considered grid-cells are named (1x1, 2x2, etc.). Which one was used at the end?

The number of grid cells used for the averaging is not statically defined, but rather determined dynamically for each model, variable, and station using the methodology described in section 3.3 (P11 L1–6). As seen in fig. 4 of the manuscript, a value of 1x1 is the most common one, however all values of up to 10x10 are occurring.

P12 L2-3: Here, the authors seems to additionally downscale the temporal resolution of the EURO-CORDEX results. Can a sentence be provided that explains why this is necessary?

We will change the first sentence of section 3.4 to: "For the calculation of the snow and ice surface energy balance in AMUNDSEN, 1–3-hourly meteorological input time series are required in order to capture the diurnal variability of the contributing energy fluxes. As the EURO-CORDEX simulations were however only available in daily temporal resolution, an additional processing step was necessary."

P 12 L8-9: Still related to the above downscaling step: I found it rather problematic that the step apparently does not preserve the daily average temperature. Can the authors give a hint on how large the introduced deviations are, and whether these deviations are systematic? If so, an additional de-biasing step would seem appropriate.

We agree that ideally the daily mean temperature should be preserved, however given daily values of T_{min}, T_{max}, and T_{mean}, with the implemented disaggregation schemes (either assuming a sinusoidal temperature course or an "average" temperature course derived from hourly observations) it is possible to preserve only two of them (either T_{min} and T_{max}, or T_{mean} and (T_{max} – T_{min})). Preserving T_{min} and T_{max} in combination with the other chosen disaggregation methods yielded better results with respect to the multilevel validation of the AMUNDSEN model results for the past, hence this method was used. The mean absolute errors of hourly observed vs. disaggregated temperature values are similar for both methods (1.12 °C for preserving T_{mean} and 1.19 °C for preserving T_{min} and T_{max} (values are averages over all stations)), however preserving T_{min} and T_{max} leads to a general tendency towards underestimated temperatures (0.38 °C on average).

P13 L6-7: I have difficulty in understanding the author's wording. Maybe they can rephrase?

We will rephrase P13 L5-7 to: "For relative humidity, an additional disaggregation method based on Waichler and Wigmosta (2003) was implemented. Hourly humidity values are generated using [month, hour, dry/wet day] categorical mean values with the additional option to preserve the daily mean humidity."

P15 L17-21: The result that the Oetztaler Alps are projected to warm significantly less than the rest of the Alps seems an important one to me. Can the authors comment on whether this is likely to be a robust results, or whether it may just be caused by the comparison

between studies using different methodologies? If the former (= robust result), a speculation on the causes could be very insightful.

Thank you for pointing this out. In fact, further analyses show that this result can likely be attributed to the different methodologies used in the individual studies (comparison of downscaled and bias-corrected RCM data interpolated to a 100 m grid (our study) vs. raw RCM results (the other studies)). A robust comparison of future climate change in our study region vs. the Alps would need to be conducted using data sets generated with the same or comparable methodology. As this paragraph is not essential for our study, we will remove it in the revised version of the manuscript.

P20 L11: The wording seems somewhat unfortunate to me: I wouldn't call a 54

Unfortunately, parts of your comment seem to be missing, however likely you argue that a 54 % reduction in glacier area cannot be called "moderate", to which we agree. We will remove "comparatively moderate".

P20L20ff: Different reactions in terms of runoff evolutions are noted for different sub-catchments. It would be useful if the authors could add some explanatory sentences for why this is the case.

The differences are mostly due to differences in glacierization and total ice volume/average ice thickness in the respective catchments. We will add some explanatory remarks to the respective section of the manuscript.

P20 L34ff: The reported changes in winter runoff appear to be very large since they are expressed in

Indeed, the relative changes in winter runoff are very large however still corresponding to very low absolute values, which we also emphasize in the manuscript. Unfortunately, here again parts of your comment seem to have been cut off, hence we would politely ask to resend it if you suggest making changes to this part of the manuscript.

P23 L21ff: I'm not entirely convinced about the "fairness" of the analysis investigating the effect of spatial and temporal resolution: Obviously, changing the resolution without recalibrating the model will impact on model performance. The question for me would rather be about the changes in model performance once the model has been recalibrated. But maybe I simply misunderstood the authors' intentions.

In the model, it is generally aimed to reduce the need for calibration. As stated above, the only parameters that have to be calibrated are the linear reservoir coefficients and possibly the snow correction factors. The former were in fact recalibrated for the assessment of the changes in model performance due to the changes in resolution and forcing data, however in the case of the latter the SCF of 15 % used in the original model setup (in combination with the temperature and wind speed-based correction and the openness-based snow redistribution) still yielded the best overall results with respect to the multilevel validation procedure. Hence, we believe our conclusion that model performance is not majorly affected by the changes in temporal and spatial resolution is appropriate.

P28 L5ff: Here (last part of the conclusions), I would have appreciated if some quantitative statements would have been included as well. Maybe, however, is just a matter of preferences ...

We agree that some quantitative statements would be appropriate here. We will replace the respective section of the conclusions by: "While some uncertainty in the results is due to the model configuration, the largest uncertainty can be traced back to the climate projections. This leads to a considerable range in the projected snow coverage, glacier extents and hydrological regimes. Snow cover is projected to decrease by up to 80 % in elevations below 1500 m a.s.l., while only comparatively moderate decreases (up to 25 %) are found for high-elevated areas (> 2500 m a.s.l.) due to strongly increasing winter precipitation which partly compensates for the increased warming. Glaciers will continue to recede strongly throughout the century. Until 2050, glacier volume will decline by approx. 60–65 % largely independent of the emission scenario, whereas by the end of the century 80–96 % of the original ice volume will be lost. Consequently, glacier runoff will diminish proportionally and summer runoff will strongly decrease in all investigated catchments by up to 55 %, resulting in a shift of the annual runoff peak from July towards June. Winter runoff volumes on the other hand will increase, however to still low absolute values. While the total annual runoff volumes stay approximately constant during the early 21st century compared to present-day levels, they gradually decrease throughout the rest of the century. Only for some catchments and scenarios runoff volumes slightly exceed present-day levels, indicating that the peak water period of maximum runoff is currently under way or has already passed in this region."

Figure 4: I have difficulties in understanding why the median deviations for "G" and "WS" (including the full name of the variables in the figure caption would be very helpful!) are only positive. To me, this is an indication that the model debiasing is not working correctly (the mean deviation should be "zero" in that case).

Global radiation is the variable with the least amount of available stations (3). For two of these stations, the mean deviation (MD) of bias-corrected vs. observed values is (absolutely) less than 0.025 W/m² for all models, whereas for one high-altitude station (Vernagtbach, 2640 m a.s.l.) it amounts to up to 1.1 W/m² (mean: 0.80 W/m²), which however corresponds to a deviation of only approx. 0.5 percent in relative terms. Similarly, bias-corrected wind speed values partly are very slightly positively biased, however amounting to a maximum MD of 0.08 m/s over all models and stations. Hence, we believe these very small remaining biases are negligible for our subsequent analyses.

References

- Gutmann, E., Barstad, I., Clark, M., Arnold, J., & Rasmussen, R. (2016). The Intermediate Complexity Atmospheric Research Model (ICAR). *Journal of Hydrometeorology*, 17(3), 957–973. <http://doi.org/10.1175/JHM-D-15-0155.1>
- Hanzer, F., Helfricht, K., Marke, T., & Strasser, U. (2016). Multilevel spatiotemporal validation of snow/ice mass balance and runoff modeling in glacierized catchments. *The Cryosphere*, 10(4), 1859–1881. <http://doi.org/10.5194/tc-10-1859-2016>
- Waichler, S. R., & Wigmosta, M. S. (2003). Development of Hourly Meteorological Values From Daily Data and Significance to Hydrological Modeling at H. J. Andrews Experimental Forest. *Journal of Hydrometeorology*, 4(2), 251–263. [http://doi.org/10.1175/1525-7541\(2003\)4<251:DOHMFV>2.0.CO;2](http://doi.org/10.1175/1525-7541(2003)4<251:DOHMFV>2.0.CO;2)

Projected cryospheric and hydrological impacts of 21st century climate change in the Ötztal Alps (Austria) simulated using a physically based approach

Florian Hanzer^{1,2}, Kristian Förster^{1,3}, Johanna Nemec⁴, and Ulrich Strasser¹

¹Institute of Geography, University of Innsbruck, Austria

²Wegener Center for Climate and Global Change, University of Graz, Austria

³Institute of Hydrology and Water Resources Management, Leibniz Universität Hannover, Germany

⁴ENVEO IT GmbH, Innsbruck, Austria

Correspondence to: Florian Hanzer (florian.hanzer@uibk.ac.at)

Abstract. A physically based hydroclimatological model (AMUNDSEN) is used to assess future climate change impacts on the cryosphere and hydrology of the Ötztal Alps (Austria) until 2100. The model is run in 100 m spatial and 3 h temporal resolution using in total 31 downscaled, bias-corrected, and temporally disaggregated EURO-CORDEX climate projections for the RCP2.6, RCP4.5, and RCP8.5 scenarios as forcing data, [making this – to date – the most detailed study for this region](#) 5 [in terms of process representation and range of considered climate projections](#). Changes in snow coverage, glacierization, and hydrological regimes are discussed both for a larger area encompassing the Ötztal Alps (1850 km², 862–3770 m a.s.l.) as well as for seven catchments in the area with varying size (11–165 km²) and glacierization (24–77 %).

Results show generally declining snow amounts with moderate decreases (0–20 % depending on the emission scenario) of mean annual snow water equivalent in high elevations (> 2500 m a.s.l.) until the end of the century, ~~however decreases of~~. [The](#) 10 [largest decreases, amounting to up to 25–80 %, are projected to occur](#) in elevations below 1500 m a.s.l. Glaciers in the region will continue to retreat strongly, leaving only 4–20 % of the initial (as of 2006) ice volume left by 2100. Total and summer (JJA) runoff will change little during the early 21st century (2011–2040) with simulated decreases (compared to 1997–2006) of up to 11 % (total) and 13 % (summer) depending on catchment and scenario, whereas runoff volumes decrease by up to 39 % (total) and 47 % (summer) towards the end of the century (2071–2100), accompanied by a shift in peak flows from July 15 towards June.

1 Introduction

Current and future climate change is expected to significantly alter the mountain cryosphere of the European Alps. Rising temperatures are expected to result in more precipitation falling as rain rather than snow, a delayed onset of the snow-covered period, and an earlier onset of snowmelt, regardless of the considered emission scenarios (e.g., Frei et al., 2017; Gobiet et al., 2014; Marty et al. 20 [\(e.g., Frei et al., 2018; Gobiet et al., 2014; Marty et al., 2017\)](#)). Possibly increasing winter precipitation amounts are expected to partly compensate for these temperature-induced effects of reduced average snowpack depths only in very high-elevated regions (e.g., Schmucki et al., 2015b).

The European mountain glaciers have already lost substantial parts of their volume and area during the past decades. Due to their delayed response to changed climatic conditions, they are still out of balance with the current climate and would continue to recede throughout the century even without any further climatic changes (e. g., Marzeion et al., 2014; Jouvet et al., 2011; Zekollari et al., 2014). Further rising temperatures will only amplify this process, resulting in a total glacier volume reduction
5 in the European Alps of 65–100 % by 2100 according to several global-scale studies (Bliss et al., 2014; Huss and Hock, 2015; Marzeion et al., 2012; Radić et al., 2013).

Consequently, the runoff regimes of snow- and ice melt-dominated Alpine catchments will also undergo significant changes. Meltwater contribution of the seasonal snow cover will be reduced, resulting in increased winter low flows, declining summer runoff, and shifts of the peak flows towards earlier periods of the year (Barnett et al., 2005; Horton et al., 2006; Stewart,
10 2009). ~~These general projected patterns of change are of high confidence, as they are~~ Since these projected impacts are mainly triggered by increasing temperatures (shift from snowfall to rainfall, earlier onset of snowmelt), ~~whereas in the likelihood of occurrence is very high.~~ In lower-elevated catchments on the other hand, projected changes in precipitation exhibit a larger impact on changes in runoff (Horton et al., 2006). Besides a projected future increase in winter precipitation on which most current climate projections agree on, future changes in precipitation over the Alpine region are however highly uncertain
15 (Gobiet et al., 2014; Smiatek et al., 2016). Consequently, future trends in total annual streamflow volume for purely snowmelt-dominated catchments are also uncertain, as changes in precipitation might (over)compensate increased evaporation rates due to higher temperatures. While in glacierized catchments on the other hand runoff volumes will eventually decrease due to the retreating glaciers and subsequently reduced ice melt runoff volumes, prolonged periods of glacier mass loss can lead to increased glacier runoff volumes in the short- to midterm, depending on if increased melt rates are able to overcompensate the
20 loss in glacier area (e. g., Jansson et al., 2003; Beniston, 2003; Collins, 2008; Bliss et al., 2014). Detecting the occurrence of this moment of peak water is of high interest, e. g., for hydropower production and planning (Schaepli, 2015).

While there are numerous studies on climate change impacts on the future of glaciers and hydrology for various parts of the European Alps, particularly Switzerland (e. g., Addor et al., 2014; Bosshard et al., 2013; Farinotti et al., 2011; Fatichi et al., 2015; Finger et al., 2012; Horton et al., 2006; Huss et al., 2008, 2014; Kobierska et al., 2013; Milano et al., 2015), few such
25 studies exist for Austria. Kuhn and Batlogg (1998) used hypothetical temperature change scenarios to simulate future runoff for nine Austrian catchments with varying glacierization using a simple conceptual water balance model while assuming constant glacier areas. The same model was applied by Kuhn and Olefs (2007) for three catchments in the Ötztal Alps while accounting for changed glacier areas using an approach taking into account observed mean annual ice thickness changes with an additional constant surface lowering per degree of temperature increase. Tecklenburg et al. (2012) investigated climate change impacts on
30 the Ötztaler Ache catchment using the conceptual semidistributed model HBV-D REG and one realization of the A1B climate scenario, however without accounting for glacier geometry changes but rather investigating only the two extremes of either constant glacier areas or entirely ice-free areas through the entire simulation period, respectively. Weber et al. (2010) used the fully distributed physically based model PROMET with the subgrid-scale extension SURGES for simulating glacier processes (Prasch et al., 2011) to calculate future hydrological changes in the Upper Danube basin using a single RCM realization based
35 on the A1B scenario, and Wijngaard et al. (2016) applied the two conceptual hydrological models HBV and HQsim for the

simulation of future hydrology in two catchments of the Ötztal Alps while updating glacier extents in 10-year intervals using precalculated ice thickness distributions.

Most of the cited studies rely on air temperature and precipitation as meteorological forcing data, applying simple temperature index methods for calculating snow and ice melt. However, the degree-day factors are calibrated for past conditions and their transferability in space and time is uncertain. Several studies hence have pointed out that more physical methods should be favored over classical temperature index melt calculations in climate change impact studies (e. g., Farinotti et al., 2011; Huss et al., 2009; Radić et al., 2013; Viviroli et al., 2011). Some studies have for example applied enhanced temperature index methods that also take solar radiation into account for melt calculation (e. g., Addor et al., 2014; Bosshard et al., 2013; Fatichi et al., 2015; Finger et al., 2012), addressing the fact that glacier melt rates are especially sensitive to variations in solar radiation (e. g., Huss et al., 2009; Ohmura et al., 2007). Only very few studies (e. g., Kobierska et al., 2013; Weber et al., 2010) however have applied full energy balance melt models for climate change impact assessment. While their superiority to more empirical methods is undisputed under the premise of in situ recordings of the required meteorological variables at the point scale, it remains challenging to provide adequate meteorological forcing data for their application in distributed mode. Nevertheless, due to their physical basis energy balance models are in principle better suited to account for changed climatic conditions than conceptual models (e. g., Klemeš, 1990; Walter et al., 2005; Pomeroy et al., 2007).

In this study we apply the fully distributed energy and mass balance model AMUNDSEN (Strasser, 2008) to assess future climate change impacts on the cryosphere and hydrology of the Ötztal Alps (Austria). ~~Downscaled, bias-corrected and temporally disaggregated~~ This region has a long tradition of hydrological and glaciological research, however – as outlined above – only scattered information about future climate change impacts exists to date, with most studies applying simplified modeling approaches and lacking a comprehensive consideration of state of the art climate scenarios. The aim of this study is to expand on this knowledge by applying a model with a comparatively high level of process representation and considering the entire set of the currently available EURO-CORDEX climate scenario simulations for the RCP2.6, RCP4.5, and RCP8.5 scenarios ~~are used as climatic forcing data~~. We analyze the simulation results with regard to changes in snow cover, glacier extent and volume, and hydrology, and discuss the associated uncertainties.

25 **2 Study site and data**

The study site (Fig. 1) is situated in the Ötztal Alps (Austria), a heavily glacierized Central-Alpine mountain range stretching east-west at the main ridge, with Austria in the north and Italy in the south. The Ötztal Alps are characterized by a warm and dry climate, with average annual precipitation sums being as low as 660 mm at station Vent (1890 m a.s.l.). In the study, we focus on the headwaters of the three north-south trending valleys Kaunertal, Pitztal, and Ötztal (located from west to east in Fig. 1), which are tributaries to the Inn river in the north. Elevations in the study site range from 862 to 3770 m a.s.l., with a mean elevation of approx. 2400 m a.s.l. For the analysis of the cryospheric impacts in terms of changes in snow and glacierization, we take the entire area shown in Fig. 1 into account (1850 km²), while the hydrological investigations are carried out for the seven gauged catchments highlighted in the figure and listed in Table 1, with a total area of 379 km². The runoff regimes are

characterized by a strong seasonality due to being fed mostly by snow and ice melt (glacial to nival regime types). Several of the catchments contribute to the Gepatsch hydropower reservoir situated in the south of the Kaunertal valley, whose natural catchment area of 106 km² is extended by a further 171 km² by diversion from the adjacent Pitztal and Radurschltal valleys. In the year 1997, 206 glaciers with a combined area of 150.7 km² were located in the area as determined from the second
5 Austrian glacier inventory (Kuhn et al., 2015).

For the model simulations, a digital elevation model (DEM) resampled to 100 m resolution was used. Initial ice thickness distributions for all glaciers in the region were calculated based on the glacier outlines and surface elevations of the year 1997 using the methodology by Huss and Farinotti (2012). The resulting ice thicknesses as shown in Fig. 1 were validated against ground-penetrating radar measurements for eleven glaciers in the region, with deviations ranging between -12 and 26 % and
10 an average deviation of -3.2 % (Seiser et al., 2012).

Meteorological stations with long-term data records availability in the region were used to drive the model in 3-hourly temporal resolution for the historical simulations. For the scenario simulations until 2100, we used the EURO-CORDEX climate change projections (Jacob et al., 2013) as climatic forcing ~~-, while considering the scenarios RCP2.6, RCP4.5, and RCP8.5. The latter is a scenario assuming no implementation of climate mitigation policies, resulting in considerably and~~
15 ~~steadily increasing emissions and concentrations of greenhouse gases over time. RCP4.5 is an intermediate scenario consistent with peaking emissions around the mid-century and a strong decline afterwards, resulting in stabilizing concentrations by the end of the century. Finally, the intervention scenario RCP2.6 is at the very low end of the spectrum in terms of future emissions and radiative forcing, assuming a peak in CO₂ concentrations in the middle of the century and a slow decline afterwards along with negative emissions toward the end of the century.~~

The entire range of (as of July 2016) publicly available EURO-CORDEX simulations for the the high-resolution (0.11° ≈ 12.5 km) EUR-11 domain that contained all required meteorological variables (minimum, maximum and mean 2 m air temperature, precipitation, relative or specific humidity, global radiation, and wind speed) in daily temporal resolution was selected, amounting to a total of 14 GCM-RCM combinations (Table 2). All of these 14 model combinations include simulations for both the RCP4.5 and RCP8.5 scenarios. ~~In contrast to many other impact studies, we also included the intervention scenario~~
25 ~~RCP2.6, for which however, whereas~~ only three realizations ~~among the EURO-CORDEX pool for the RCP2.6 scenario~~ were available. Hence, in total 31 different sets of climate projections were available for the glaciohydrological simulations. The scenario period of the models is 2006–2100, with the exception of the three HadGEM-driven models which terminate in November 2099.

3 Methods

30 3.1 Model

For the glaciohydrological simulations in this study, we used the fully distributed hydroclimatological model AMUNDSEN (Strasser, 2008). AMUNDSEN has specifically been designed as a scenario-capable model for the application in high-mountain regions, and has been set up and extensively validated for historical conditions in the study site in a recent study (Hanzer et al.,




Figure 1. Location and topography of the study site including forested areas, catchment boundaries, the locations of the meteorological stations and runoff gauges, and the calculated initial ice thickness distribution for the year 1997. Selected meteorological stations and glaciers explicitly mentioned in the article are labeled in red and blue, respectively. Glacier abbreviations refer to Hintereisferner (HEF), Kesselwandferner (KWF), Vernagterferner (VF), Taschachferner (TF), and Gepatschferner (GF).

Table 1. Area, elevation, and glacierization (as of 1997) of the investigated catchments. The catchment IDs correspond to the labels in Fig. 1. Indentations indicate subcatchments.

ID	Catchment	Area [km ²]	z_{\min}	z_{\max}	z_{mean}	Glacierization 1997 [%]
			[m a.s.l.]	[m a.s.l.]	[m a.s.l.]	
1	Venter Ache	165.3	1880	3762	2887	35.4
2	— Rofenache	98.6	1893	3762	2889	38.3
3	—— Vernagtbach	11.5	2638	3622	3120	77.0
4	Gurgler Ache	72.4	1885	3533	2785	31.8
5	Gepatschalm	53.9	1898	3524	2824	39.7
6	Taschachbach	60.5	1791	3761	2753	24.0
7	Pitze	27.0	1812	3548	2835	48.2
Total		379.1	1791	3762	2833	34.4

Table 2. EURO-CORDEX scenario simulations used in this study.

ID	RCM	GCM	RCPs
1	CCLM4-8-17	CNRM-CM5	4.5, 8.5
2	CCLM4-8-17	EC-EARTH	4.5, 8.5
3	CCLM4-8-17	HadGEM2-ES	4.5, 8.5
4	CCLM4-8-17	MPI-ESM-LR	4.5, 8.5
5	HIRHAM5	EC-EARTH	2.6, 4.5, 8.5
6	RACMO22E	EC-EARTH	4.5, 8.5
7	RACMO22E	HadGEM2-ES	4.5, 8.5
8	RCA4	CNRM-CM5	4.5, 8.5
9	RCA4	EC-EARTH	2.6, 4.5, 8.5
10	RCA4	CM5A-MR	4.5, 8.5
11	RCA4	HadGEM2-ES	4.5, 8.5
12	RCA4	MPI-ESM-LR	4.5, 8.5
13	REMO2009	MPI-ESM-LR	2.6, 4.5, 8.5
14	WRF331F	CM5A-MR	4.5, 8.5

2016). In the following, the most important model components are briefly discussed – for a more detailed model description, we refer to, e. g., Hanzer et al. (2014, 2016), Marke et al. (2015), Pellicciotti et al. (2005), Strasser (2004, 2008), and Strasser et al. (2008).

The model is capable of operating in temporal resolutions of 1–3 h, while the spatial resolution is generally arbitrary but typically chosen in the order of 10–100 m for the application in high-mountain regions in order to accurately capture the complex topography and the underlying processes in these regions. As meteorological variables, AMUNDSEN requires point measurements or gridded inputs of air temperature, precipitation, relative humidity, global radiation, and wind speed. For the application using point data, the model includes a meteorological preprocessor for the spatial interpolation of scattered point measurements using a combination of lapse rates and inverse distance weighting. Lapse rates are either calculated from the point [measurements data](#) in each time step or are presupplied as average monthly values. A radiation model based on the method by Corripio (2002) is used to calculate potential clear-sky solar radiation while taking into account [terrain slope and orientation](#), hill shading, transmission losses and gains due to scattering, absorption, and reflections, and uses measured point values of global radiation to derive cloud factor fields and subsequently actual global radiation fields. Similarly, incoming longwave radiation is calculated using parameterizations for radiation received from the clear sky, clouds, and surrounding terrain. The precipitation phase (rain or snow) is determined as a function of the wet-bulb temperature. For solid precipitation, different correction methods are implemented in order to account for the undercatch of precipitation gauges when measuring snow accumulation. Hanzer et al. (2016) showed that a combination of a station-based snow correction factor (SCF) taking

into account wind speed and air temperature with a subsequent constant post-interpolation SCF of 1.15 yielded plausible long-term precipitation amounts for the study area, while an additional redistribution of the interpolated snowfall fields using an approach based on topographic openness (Yokoyama et al., 2002) distinctly improved the spatial snow accumulation patterns. Accumulated snow on the surface is subdivided into three layers called new snow, old snow, and firn. Transitions between new snow and old snow occur depending on snow density (which is calculated based on approaches by Anderson (1976) and Jordan (1991)), while remaining snow amounts at the end of the ablation season (September 30) are transferred to the firn layer. In glacierized catchments, an additional layer is used to track the evolution of ice amounts. A linear densification of the firn layer is employed, while firn is converted to ice once reaching a threshold density of 900 kg m^{-3} . Snow albedo is parameterized using an aging curve approach with exponential decay down to a specified minimum value, while firn and ice albedo is kept constant for this application (with values of 0.4 and 0.2, respectively). In forested areas the interpolated meteorological fields are modified in order to capture sub-canopy conditions, ~~while additionally the~~ resulting in reduced shortwave radiation, precipitation, and wind speed, increased longwave radiation and humidity, and an attenuation of the diurnal temperature cycle. Additionally the effects of the forest snow processes of interception, sublimation, and melt unload are accounted for (Strasser et al., 2011). Snow and ice melt is calculated using an energy balance approach taking into account short- and longwave radiation fluxes, latent and sensible heat fluxes, advective energy, as well as the ground heat flux. The delay of the onset of snowmelt in cold snowpacks is accounted for by a parameterization of the cold content of the snowpack, whereas melting snow may persist in the snowpack in the form of liquid water up to a certain amount (and possibly refreeze again) before actual outflow occurs (Hanzer et al., 2016). Evapotranspiration in snow-free areas is calculated following Allen et al. (1998). Finally, runoff at predefined catchment outlets is calculated using a linear reservoir approach for snow on glaciers, firn, bare ice, snow outside of glaciers, and soil (Hanzer et al., 2016). Apart from the parameters of this linear reservoir model which have to be calibrated individually for each catchment, most parameters in the model have a physical meaning, and in general no site-specific calibration beyond the parameters for the runoff module and the precipitation correction method is performed.

The setup and extensive validation of the model for the study region under historical conditions has been described in Hanzer et al. (2016). Essentially the same model setup was used for this study, aside from (i) the newly implemented glacier retreat parameterization as described in the following section, (ii) reducing the spatial and temporal model resolution from 50 m and 1 h to 100 m and 3 h due to performance reasons, and (iii) the fact that only a subset of the meteorological stations that were utilized in the simulations presented in the aforementioned study was available for this study due to the constraints of a sufficiently long measurement period required for the bias correction of the RCM data (see section 3.3).

3.2 Glacier geometry change

The model setup as described in Hanzer et al. (2016) already incorporated spatially distributed glacier thicknesses, however only accounted for the climatic forcing on the glaciers without any adjustment of glacier geometry. Due to the generally small vertical ice flow contribution for the majority of the glaciers in the study region (Helfricht et al., 2014) this is a reasonable approach for shorter simulation periods, ~~however~~. However, when performing simulations on multidecadal scales, ~~the~~ effect of ice flow dynamics must be considered in glaciohydrological models. Otherwise, systematic errors in the simulated ice

distributions (and subsequently in glacier runoff) are introduced as the glaciers thicken in their accumulation areas and retreat too quickly in the tongue parts.

For this study, we implemented the Δh method (Huss et al., 2010) to periodically update the simulated glacier geometries. This approach is particularly suited for spatially distributed models operating on the regional scale, as it does not necessarily require glacier-specific parameterizations ~~but rather~~. Rather, it uses simple assumptions to translate the climatic forcing (i. e., the surface mass balance as computed by the mass balance model) into a geometric response in terms of the glacier thickness distribution.

Essentially, the Δh parameterization scales the spatial distribution of the annual glacier surface mass balance such that the changes in glacier surface elevation match patterns observed in the past. This is accomplished by applying a prescribed function to each glacier in regular time intervals, which adjusts the simulated surface elevation change as a function of the normalized glacier elevation (assumed to be a proxy for the central flowline). To derive the Δh function, two glacier surface DEMs are required, preferably covering multidecadal periods to reduce possible errors due to DEM uncertainty.

In principle, a separate Δh function could be derived for each glacier, however Huss et al. (2010) showed that glaciers with similar characteristics show similar geometric responses, and derived three Δh functions for glaciers of different size classes: large valley glaciers ($A \geq 20 \text{ km}^2$), medium valley glaciers ($5 \text{ km}^2 \leq A < 20 \text{ km}^2$), and small glaciers ($A < 5 \text{ km}^2$):

$$\Delta h = \begin{cases} (h_r - 0.02)^6 + 0.12(h_r - 0.02) & A \geq 20 \text{ km}^2 \\ (h_r - 0.05)^4 + 0.19(h_r - 0.05) + 0.01 & 5 \text{ km}^2 \leq A < 20 \text{ km}^2 \\ (h_r - 0.30)^2 + 0.60(h_r - 0.30) + 0.09 & A < 5 \text{ km}^2, \end{cases} \quad (1)$$

where h_r corresponds to the normalized surface elevation between 0 and 1.

While these parameterizations have been derived for the Swiss Alps, they are assumed to be applicable for all mountain glaciers (Huss et al., 2010), and have already been applied in other regions of the world (e. g., Huss and Hock, 2015; Ragetti et al., 2013). While for our study region DEMs and glacier outlines exist for the years 1997 and 2006, we chose to adopt these generalized parameterizations, as (i) they have been derived over much longer time periods, hence likely being more robust, and (ii) match the average observed Δh patterns for our study area in the period 1997–2006 well (Fig. 2).

In AMUNDSEN, the glacier geometry update is performed at the end of each glaciological year (September 30): for each glacier, the total volume change in terms of the surface mass balance is used to scale the thickness change prescribed by the Δh function (determined by Eq. 1) which is applied to each glacier pixel and the respective normalized elevations. Following Huss et al. (2010), the maximum surface lowering is limited to the surface mass balance lowering at the glacier terminus, and the glacier borders (pixels with ice thicknesses $< 10 \text{ m}$) are not modified by the geometry update but rather change their thickness only due to the respective surface mass balance change.

Figure 3 demonstrates the effect of the glacier geometry update as implemented in AMUNDSEN exemplarily for the Hintereisferner and Taschachferner glaciers. While the average change in surface elevation between 1997 and 2006 is reproduced reasonably well for both glaciers in the run without updating the glacier geometries, the increase in surface elevation at the




Figure 2. Normalized observed ice thickness change for 42 small ($A < 5 \text{ km}^2$, thin blue lines) and 8 medium-sized ($5 \text{ km}^2 \leq A < 20 \text{ km}^2$, thin red lines) glaciers in the Ötztal Alps during the period 1997–2006, and the corresponding parameterizations from Huss et al. (2010) (thick lines).

uppermost glacier parts is severely overestimated for both glaciers, whereas mass losses in the middle to lower parts of the glaciers are too high. Adding the glacier retreat module to the model distinctly improves the spatial distribution of mass gain or loss for both glaciers.

~~One limitation of the Δh approach is that it allows only for geometric changes in terms of glacier retreat, essentially limiting its applicability for periods with largely negative mass balances. Huss and Hoek (2015) present an extension of the method that enables to also account for advancing glaciers, however using a semidistributed model operating in elevation bands. A possible, comparatively simple extension of the method when applied in a fully distributed way could be to allow the glaciers to grow within their historical extents, similar to the approach applied by Stahl et al. (2008). In this study, however, we apply the original retreat-only version of the Δh parameterization, which is seen a feasible approach given that future climate scenarios indicate climatic conditions causing intensive melt and, hence, glacier shrinkage.~~

3.3 Spatial downscaling of RCM data

While state-of-the-art RCM simulations already feature comparatively high spatial resolutions, usually the performance of the raw RCM data is still inadequate for directly using it in hydrological impact studies, both due to still-present mismatches in space as well as due to systematic model errors (biases) introduced by the RCMs. A common approach for impact studies is hence to use ensembles of GCM-RCM chains in combination with statistical bias correction methods (Teutschbein and Seibert, 2012).

In this study, we use quantile mapping (QM) (e. g., Déqué, 2007; Themeßl et al., 2011a), an empirical bias correction method that corrects the distribution functions of the RCM variables to fit the distribution functions of the observations. QM has reg-




Figure 3. Observed and simulated glacier surface elevation change during the period 1997–2006 as a function of elevation for the glaciers Hintereisferner and Taschachferner (see Fig. 1 for their geographical setting). “Without geometry update” refers to the simulated surface mass balances without accounting for ice flow, while “with geometry update” refers to the simulation results obtained after implementing the Δh parameterization.

ularly shown to outperform other statistical bias correction methods (e. g., Teutschbein and Seibert, 2012, 2013), is applicable in mountain regions (e. g., Finger et al., 2012; Themeßl et al., 2011b, a) (e. g., Finger et al., 2012; Themeßl et al., 2011a, b), and allows for correcting climate variables other than temperature and precipitation (e. g., Finger et al., 2012; Wilcke et al., 2013). Like Limitations of QM are, among others, that it is generally not trend-preserving with respect to both means and

5 extremes and that – like in most statistical bias correction methods – QM does not explicitly account for – spatial, temporal, or intervariable correlations, however it and intervariable correlations are not explicitly accounted for (Cannon et al., 2015; Maurer and Pierce). However, QM has shown to perform well under changed climatic conditions (Teutschbein and Seibert, 2013) and to retain intervariable relations can preserve intervariable dependency structures (Wilcke et al., 2013).

In the study, we used the QM methodology by Gudmundsson et al. (2012) as implemented in the R package *qmap*. Due to strong season-dependent biases in the RCM data, we performed the bias correction separately for DJF, MAM, JJA, and SON. Only stations and variables with a minimum amount of 20 years of data within the period 1971–2005 were considered for the bias correction procedure, with the exception of global radiation where two stations with only approx. 10 years of data were also included due to an otherwise insufficient number of stations. Given these constraints, in total 16 precipitation time series were available, as well as 13 for air temperature, 6 for wind speed, 5 for relative humidity, and 3 for global radiation.

In order to avoid the possibility of unrealistic temperature values in terms of corrected $T_{\min} > T_{\max}$, following Thrasher et al. (2012) we did not correct T_{\min} and T_{\max} independently, but rather corrected T_{\max} and the diurnal temperature range (DTR) ($T_{\max} - T_{\min}$) with a subsequent calculation of corrected T_{\min} .

For the EURO-CORDEX realizations for which only specific rather than relative humidity was available (IDs 1–4 and 13 in Table 2), mean daily specific humidity q [kg kg^{-1}] was first converted to relative humidity RH [%] as-

$$\text{RH} = 100 \frac{e}{e_s},$$

with e being the vapor pressure e calculated as-

$$e = \frac{qp}{0.622 + 0.378q},$$

with p being air pressure. Saturated vapor pressure e_s was calculated following Sonntag (1990) as-

$$e_s = \begin{cases} 6.112 \exp\left(\frac{17.62T}{243.12 + T}\right) & T \geq 0 \text{ } ^\circ\text{C} \\ 6.112 \exp\left(\frac{22.46T}{272.62 + T}\right) & T < 0 \text{ } ^\circ\text{C} \end{cases}$$

with $T = T_{\text{mean}}$ formulations by Sonntag (1990).

With regard to the selection of RCM grid points for the downscaling to the point scale, we followed the approach by Hofer et al. (2017) (who however used linear regressions rather than QM) to find the *optimum scale* (OS) for each station and target variable: For each station and variable, spatial averages of the closest $1 \times 1, 2 \times 2, \dots, 10 \times 10$ RCM grid points were calculated and subsequently used for bias correction. The OS for a given station and variable was then defined as the spatial window which minimizes the deviations between the cumulative distribution functions of the corrected and observed data in terms of the mean absolute error (MAE). Histograms of the resulting OS values for the different variables are shown in Fig. 4. While for all variables an OS of 1 (corresponding to using only the closest RCM grid point without spatial averaging) is the most common value, for all variables except global radiation all OS values from 1 to 10 are found – for precipitation, the maximum possible OS of 10 is even the second most common value.

RCM outputs from models using fixed 360- or 365-day calendars were linearly rescaled to Gregorian calendar dates (e. g., for a 360-day calendar each day from 1–360 was mapped to a respective day from 1–365 (or 366) in the Gregorian calendar). Resulting days with missing values were filled by duplicating the values from the preceding day.

Figure 5 shows the mean deviation (MD) and the standard deviation ratio (SDR) of the bias-corrected RCM data vs. the daily observations for the stations in the study area in the historical period 1971–2005. The model IDs (x-axis) correspond to the IDs in Table 2. While some models tend to perform slightly better than others on average, both measures are in general close to their optimum values (0 and 1, respectively) for all realizations, indicating that the corrected RCM outputs adequately represent the observed climate in terms of mean and variability. The higher deviations in the variability of relative humidity for some models are not a result of the bias correction per se, but rather of the required conversion of specific to relative humidity for these models.




Figure 4. Histograms of the optimum scales for bias correction found using the approach by Hofer et al. (2017) for the variables minimum temperature (T_{\min}), maximum temperature (T_{\max}), precipitation (P), relative humidity (RH), global radiation (G), and wind speed (WS). Histograms are calculated using the derived optimum scales for all available stations for a given variable and all 14 GCM-RCM combinations (Table 2).




Figure 5. Mean deviation (MD) and standard deviation ratio (SDR) of simulated (i. e., downscaled and bias-corrected RCM data) vs. observed meteorological data for the historical period 1971–2005. The individual boxes represent the downscaled station meteorology for a given GCM/RCM combination (model IDs correspond to the ones in Table 2). Units given for the different variables apply only to MD (SDR is unitless).

3.4 Temporal downscaling of RCM data

For the calculation of the snow and ice surface energy balance in AMUNDSEN, 1–3-hourly meteorological input time series are required in order to capture the diurnal variability of the contributing energy fluxes. As the EURO-CORDEX simulations were however only available in daily temporal resolution, an additional processing step was necessary ~~to derive the required sub-daily~~ ~~(1–3-hourly) data~~. This step was performed with the open-source temporal disaggregation tool MELODIST (Förster et al., 2016). MELODIST provides simple, empirical disaggregation routines for the sub-daily disaggregation of daily point-scale data for the variables air temperature, precipitation, relative humidity, solar radiation, and wind speed. For the application in this study several new disaggregation methods were added to MELODIST ~~;~~, now published as version 0.1.2 (Hanzer et al., 2017):

1. For air temperature, a disaggregation method conserving the daily mean temperature and the DTR ($T_{\max} - T_{\min}$) while applying a sinusoidal temperature course in the same way as the default method (which preserves T_{\min} and T_{\max} but not T_{mean}) was implemented. However, most stations showed no clear sinusoidal temperature course, hence the application of this method still resulted in a (positive) temperature bias for most stations. Therefore, a new method based on mean values depending on the calendar month and hour of the day derived from hourly recordings was implemented. In a preprocessing step, for each month and station an average temperature course (normalized to a $[0, 1]$ range) is calculated. Then, daily values are disaggregated on the basis of these temperature courses again by either preserving T_{\min} and T_{\max} , or T_{mean} and the DTR ($T_{\max} - T_{\min}$).
2. With regard to solar radiation, a similar method was implemented, which uses monthly varying average diurnal radiation courses derived from observations to scale the daily mean radiation. In addition, the methods for deriving daily radiation values from sunshine duration or the diurnal temperature range have been updated to allow monthly or seasonally varying conversion factors.
3. For relative humidity, an additional disaggregation method based on Waichler and Wigmosta (2003) was implemented. Hourly humidity values are generated using [month, hour, dry/wet day] categorical mean values ~~was implemented following Waichler and Wigmosta (2003). If daily humidity values are available, they can be used to scale the thereby derived values with the additional option~~ to preserve the daily mean humidity.

Figure 6 shows the density functions for the observed hourly values (black lines) as well as for the various disaggregation methods exemplarily for station Obergurgl (1938 m a.s.l.) and the variables air temperature, relative humidity, global radiation, and wind speed. Based on these comparisons and the results of the multilevel validation of the glaciohydrological simulation results when driven with the disaggregated values using multiple configurations, we decided for the following disaggregation methods: for all stations, air temperature was disaggregated using the “mean course” method while preserving daily minimum and maximum temperatures, daily precipitation amounts were distributed uniformly across the day, relative humidity was disaggregated using [month, hour, dry/wet day] categorical mean values while preserving daily mean ~~radiation~~humidity, solar radiation was disaggregated using the “mean course” method while preserving daily mean radiation, and wind speed was distributed uniformly over the day.




Figure 6. Density functions of the observed hourly values (black lines) for station Obergurgl (period 1999–2013) and the respective aggregated-and-disaggregated values using different disaggregation methods for the variables temperature, relative humidity, global radiation, and wind speed. Numbers in parentheses indicate the R^2 of the regression against the observations.

Not all of these disaggregation methods performed best with regard to the reproduction of hourly values at the station scale, however this method combination yielded the best overall glaciological model performance according to the multilevel validation procedure described in Hanzer et al. (2016). This ~~is partly explainable from~~ can be attributed to the fact that the disaggregation is performed independently for each station and variable.

5 4 Results and discussion

In the following sections, first we discuss the future climatic changes in the study area as projected by the EURO-CORDEX scenarios. Then, we analyze the subsequent changes in snow, glaciers, and hydrology according to the AMUNDSEN simulations for the scenario period. These simulations were carried out by initializing the model in 1997 (using the glacier inventory and ice thickness data) and running it until September 2006 using the observed meteorological data. Afterwards, we switched to the EURO-CORDEX scenario period and carried out the scenario simulations until 2100 (with the exception of the three HadGEM-driven models for which the simulations could only be carried out until the year 2098).

We did not perform the entire range of glaciological simulations for a longer historical period due to restrictions of the current model setup: (i) the initial ice thickness distributions were calculated for the year 1997 and are hence not applicable

for earlier periods, and (ii) the generally negative mass balances of the glaciers in the Ötztal Alps during the last decades were interrupted by a short period of glacier advance between the mid-1970s and the early 1980s (Abermann et al., 2009), which cannot adequately be accounted for using the currently implemented Δh parameterization for the glacier geometry update. However, for discussing solely the changes in snow conditions, we also carried out model runs for the period 1971–2005 (hence obtaining transient runs from 1971 to 2100) using the historical RCM simulations as forcing, while neglecting the effect of glaciers in this case.

The validation of the model setup for the study area has been described in detail in Hanzer et al. (2016). As essentially the same model setup was used for the present study, in this section only the results of the scenario simulations are discussed. A brief discussion on how model performance is impacted by the few modifications in model setup can be found in the discussion of uncertainties (section 4.5).

4.1 Future climate

Figure 7 shows the range of the projected seasonal and overall changes (spatial averages, relative to the baseline period 1971–2000) of the five meteorological variables for the study area as provided by the climate simulations. These changes are calculated based on areal means of the respective variables as calculated by the meteorological preprocessor in AMUNDSEN (i. e., after bias correction). While in the figure the changes are plotted for three time slices throughout the 21st century, for the following short discussion of the changes we focus on the period 2071–2100. If not otherwise stated, values refer to multi-model means in the following.

Temperature is projected to increase by all models and for all seasons, with average values of 1.1 (RCP2.6) to 3.8 °C (RCP8.5) in the annual mean, and a maximum spread of 2.2 °C between individual models for a given scenario. The highest warming is projected for the winter season, with up to 4.5 °C in the RCP8.5 scenario, while the smallest increases are projected for spring.

For precipitation, no clear general trend can be discerned with regard to annual sums. Both decreases (up to –14 %) and increases (up to 24 %) are projected by individual models, while the multi-model averages are close to zero for all three scenarios. However, with the exception of some outliers, there is a general consensus between the model towards a precipitation shift from summer towards winter. Projected multi-model average increases in winter precipitation are between 8 % (RCP2.6) and 21 % (RCP8.5), with individual models projecting increases of up to 57 %.

Changes in relative humidity largely follow the trend of precipitation changes, with decreases in summer (up to –1.5 %) and smaller increases in winter (up to 0.7 %). At least for the RCP8.5 scenario slight decreases in the annual mean are projected.

For global radiation, the disagreement between the individual models is considerable. The multi-model means indicate a decrease in overall global radiation for all RCPs, with values between -1.2 W m^{-2} for RCP4.5 and -2.4 W m^{-2} for RCP2.6, however with individual model results ranging between -20 W m^{-2} and 4 W m^{-2} . The spread between the models is even larger when looking at seasonal changes. While all models agree on a decrease of global radiation in winter and (with one exception) spring, projections for summer are uncertain. Here, most models tend towards an increase in radiation for the RCP4.5 and RCP8.5 scenarios, however individual results range between -29 W m^{-2} and 26 W m^{-2} .

Finally, wind speed projections also show quite a large spread between individual models. Multi-model means indicate a very small overall decrease in wind speed, with seasonal increases of up to 0.12 m s^{-1} during winter and decreases of up to -0.12 m s^{-1} during summer.

~~Comparing to other studies, the projected changes over the Ötztal Alps and their seasonal patterns are similar to the average changes projected for the entire Alpine region (Gobiet et al., 2014; Jacob et al., 2013; Smiatek et al., 2016). Notable differences are found for temperature, where the multi-model median change for 2071–2100 compared to 1971–2000 is 0.3 lower than in Jacob et al. (2013) for RCP4.5, and 1.0 lower for RCP8.5. Seasonal temperature changes for the RCP4.5 scenario are between 0.1 and 0.4 lower than in Smiatek et al. (2016).~~

4.2 Changes in snow

Changes in the amount and duration of the simulated seasonal snow cover in the study area were analyzed both temporally and spatially. As a reference for computing the changes, we performed AMUNDSEN simulation runs using the historical periods of the EURO-CORDEX simulations for the period 1970–2005. Through combination with the scenario runs we hence obtained transient simulations for the period 1970–2100.

Figure 8 shows the evolution of the simulated mean annual snow water equivalent (SWE) for three meteorological stations covering an approx. 2000 m elevation range: Prutz (870 m a.s.l.), Obergurgl (1938 m a.s.l.), and Pitztaler Gletscher (2864 m a.s.l.). At the highest-elevated station, Pitztaler Gletscher, the mean snow amounts remain relatively unchanged for all three RCPs until the middle of the century, ~~whereas only afterwards stronger decreases can be seen for the RCP4.5 and RCP8.5 scenarios.~~ For RCP2.6, snow amounts are then slowly increasing during the last third of the century, reaching an average level of 517 mm (multi-model mean), almost as high as the 530 mm in the historical period 1971–2000. ~~However, also for the RCP4.5 and RCP8.5 scenarios, only~~ For the other two scenarios comparatively small decreases in average SWE are simulated in this period, amounting to 427 mm (–18 %) for RCP4.5 and 357 mm (–31 %) ~~, respectively. For for RCP8.5. For the~~ station Obergurgl, similarly no strong changes in average snow amounts are simulated within the first half of the century, whereas afterwards the curves of the three scenarios begin to diverge more strongly than for Pitztaler Gletscher. For RCP2.6 again a general trend of increasing SWE after 2050 is projected, while snow amounts in the RCP4.5 scenario decrease to 94 mm (–31 %) in the period 2071–2100, and 59 mm (–57 %) for the RCP8.5 scenario. The largest relative changes are found for the lowest-elevated station, Prutz, where the curves for the three scenarios begin to divert by around 2020. RCP2.6 snow amounts then stay relatively constant during the remainder of the century at 6 mm (–31 %), while RCP4.5 and RCP8.5 snow amounts continue to decrease strongly, amounting to –63 % and –80 %, respectively, at the end of the century.

Similar results of stronger declining snow amounts for lower elevations can be seen in Fig. 9, where the multi-model mean change in SWE relative to the baseline period 1971–2000 is plotted against elevation. For RCP2.6, with the exception of the very lowest-elevated parts of the study area, the strongest decreases in snow amounts generally are projected for the period 2041–2070, whereas the largest average snow amounts are found at the end of the century. Generally, however, the projected decreases are comparatively small, with maximum values of approx. 12 % for elevations above 2500 m a.s.l., up to 25 % for elevations between 1500 and 2500 m a.s.l., and slightly stronger below. For both RCP4.5 and RCP8.5, average snow amounts




Figure 7. Projected seasonal changes for temperature (T), precipitation (P), relative humidity (RH), global radiation (G), and wind speed (WS) according to the selected EURO-CORDEX scenario simulations (spatial averages over the study area; changes calculated relative to 1971–2000). Small dots represent the individual GCM-RCM realizations.

stay virtually constant (even slightly increasing for RCP4.5) during the period 2011–2040 in elevations above 2500 m a.s.l., whereas for the lower-elevated parts decreases during this period amount to up to 37 %. However, for both scenarios strong decreases are projected for the remainder of the century, with the largest changes in the lowest-elevated areas, amounting to up to –64 % and –83 % for RCP4.5 and RCP8.5, respectively.




Figure 8. Evolution of mean annual SWE for the stations Prutz, Obergurgl, and Pitztaler Gletscher as simulated using the EURO-CORDEX scenarios for the RCPs 2.6, 4.5, and 8.5. Thin lines and error bars indicate the multi-model mean and standard deviation, respectively, while the thick lines show the 5-year Gaussian low-pass filtered multi-model mean.

The analysis of seasonal changes (Fig. 10) reveals that the strongest relative changes are projected for the summer months, while simulated changes for winter are comparatively small, amounting to maximum decreases of approx. 30 % for RCP8.5. For all three scenarios, a gradual shift in the timing of peak SWE amounts from April towards March is simulated.

These projected changes in snow coverage generally show similar patterns to studies for other Alpine sites (e. g., Beniston et al., 2003; Marty et al., 2017; Schmucki et al., 2015a; Steger et al., 2012), however with a tendency to lower relative decreases in average snow amounts. This is likely due to the comparatively strong increases in winter precipitation especially for the warmest (RCP8.5) scenario (Fig. 7). A comprehensive study recently done for Switzerland (Marty et al., 2017) for example projected more dramatic changes in snow amounts especially in high elevations (> 3000 m a.s.l.) until the end of the century for the A2 scenario (similar warming as in the RCP8.5 scenario), however projecting only very minor winter precipitation increases. Especially in high elevations however, increases in precipitation amounts are more likely to compensate for temperature increases, as temperatures in these elevations are usually still below melting conditions for most parts of the year.

4.3 Changes in glaciers

According to the simulation results, all glaciers in the study region will continue to retreat significantly throughout the 21st century. This is illustrated in Fig. 11, where the evolution of glacier volume and area (relative to the respective values of 2006) are plotted for all 31 model combinations as well as the respective multi-model means. In general, the evolution of glacier area and volume is very similar, with only a slightly stronger decline of glacier volume compared to the area. Looking at the multi-




Figure 9. Elevation-dependent projected change in SWE (multi-model means, shadings indicate \pm one standard deviation) relative to the period 1971–2000 for the RCPs 2.6, 4.5, and 8.5.




Figure 10. Mean monthly SWE (multi-model means, shadings indicate \pm one standard deviation) over the entire study area (Fig. 1) for the three RCPs (top) and relative changes compared to 1971–2000 (bottom).

model mean values for the three emission scenarios, the largest changes in glacierization ~~occur already~~ already occur within the first half of the century, where – relatively independent of the emission scenario – the glaciers lose between approx. 60 % (RCP2.6) and 65 % (RCP8.5) of their volume from the beginning of the century. After 2050, the slope of the curves decreases and differences between the three scenarios become more prominent. In the RCP2.6 scenario, glacier volume decreases to approx. 25 % until 2080, whereas the glaciers stabilize afterwards and only slightly further retreat until 2100. In the RCP8.5

scenario, glaciers decrease more rapidly than in the RCP4.5 case between 2050 and 2080, whereas afterwards the two curves again follow a similar course. For both scenarios, the glaciers have mostly vanished by the end of the century, with only small remains of approx. 10 % (RCP4.5) and 4 % (RCP8.5) still left in terms of their initial volume. Results for the evolution of glacier area are similar.

5 Looking beyond the multi-model mean, it becomes clear that there is a remarkable spread between the individual model runs, most strikingly for the RCP2.6 scenario, where the glacier volume decrease is between 58 % and 97 % until 2100 for the three model runs. Also for the RCP4.5 and RCP8.5 scenarios the range between the individual model realizations is considerable, with individual realizations resulting in practically ice-free conditions as early as 2070, while in other realizations up to 33 % of glacier volume are preserved until 2100.

10 The comparatively small influence of the different emission scenarios on the other side is visualized in Fig. 12, where the ice thickness evolution of a single glacier (Hintereisferner) along its centerline is plotted in 10-year intervals for a single GCM-RCM combination (REMO2009 driven by MPI-ESM-LR) and the RCPs 2.6, 4.5, and 8.5. Here again a strong loss of glacier volume can be seen in the first decades of the century. Regardless of the emission scenario, by 2040 the glacier retreats by approx. 1.4 km in length and loses approx. half of its volume (49–54 %). Only after 2060, the RCP8.5 scenario generates a
15 markedly stronger loss in ice volume compared to the other two scenarios, resulting in an almost complete disappearance of the glacier by 2100. The results for RCP2.6 and RCP4.5 largely follow the same evolution until 2090. Only in the last decade of the century, the RCP4.5 scenario produces an accelerated retreat of the glacier, whereas the RCP2.6 glacier surface stays approximately constant. However, also for these two less extreme scenarios, only 25 % (RCP2.6) and 13 % (RCP4.5) of the original ice volume is left by the end of the century.

20 Figure 13 shows the evolution of the absolute and relative glacierization as well as the absolute glacier volume and mean ice thickness for each catchment in the RCP4.5 scenario, illustrating that different catchments partly show different responses to the climatic forcing. For some of them (most notably Taschachbach) the average ice thickness stays approximately the same during the entire simulation period due to the similar rates of volume reduction and area reduction. For the Gepatschalm catchment, which includes the largest and thickest glacier in the study area (Gepatschferner), this is also the case for the first
25 third of the century, whereas afterwards the rate of volume decline increases. In other catchments such as Vernagtbach and Gurgler Ache the reverse effect is observed during the last part of the century: glacier area retreats more quickly than the volume decreases (visible by the increase in mean ice thickness in this period).

The spatial evolution of glacier coverage within the study area is shown in Fig. 14, where the simulated multi-model mean glacier coverage is displayed in 10-year intervals for the three emission scenarios. The rate of glacier retreat is mainly dependent
30 on elevation and glacier thickness. In all three scenarios, most of today's small glaciers have disappeared by 2050 according to the model. However, for the largest glaciers considerable differences resulting from the different emission scenarios appear. Gepatschferner, for example, loses **comparatively moderate** 54 % of its 1997 area (17.1 km²) by 2100 in the RCP2.6 scenario, while in the RCP4.5 and RCP8.5 scenarios 71 % and 95 %, respectively, of the glacier area disappears.




Figure 11. Simulated total glacier volume (top) and area (bottom, both shown as changes relative to the year 2006) for all glaciers in the study area and all 31 available climate scenarios. Thin lines indicate individual model results, thick lines show the multi-model mean. The small spikes in the multi-model means for RCP4.5 and RCP8.5 at the very end of the century are due to the fact that three models could only be run until 2098.

4.4 Changes in hydrology

Figure 15 exemplarily shows the simulated seasonal runoff cycle for present-day conditions as well as the projected values for the future periods for four of the study catchments. While the reaction of the catchments are slightly different depending on their characteristics, the general pattern of change is the same for all catchments and emission scenarios: summer runoff strongly decreases with simultaneously increasing spring runoff, indicating a shift from glacial/glacio-nival to nivo-glacial runoff regimes. While in the RCP2.6 scenario the month of peak runoff remains unchanged, in the RCP4.5 and RCP8.5 scenarios the peak gradually shifts from July towards June for all catchments, with the exception of Vernagtbach in the RCP4.5 scenario. For the Rofenache, Gepatschalm, and Pitze catchments, this shift already occurs in the period 2041–2070 in both scenarios. Only for Vernagtbach, the most glacierized catchment, in the RCP8.5 scenario the shift occurs only towards the end of the century, as in the prior period the catchment is still glacierized to a considerable degree. Monthly peak runoff slightly




Figure 12. Bedrock topography (thick black lines) and simulated glacier thickness (thin colored lines) along the centerline of Hintereisferner as simulated for a single GCM-RCM combination (REMO2009 driven by MPI-ESM-LR) and the three RCPs, shown in 10-year intervals. Dashed black lines indicate the initial ice thickness as of the year 1997.




Figure 13. Evolution of the absolute (top left) and relative (top right) glacierized area for each catchment in the RCP4.5 scenario, as well as the absolute glacier volume (bottom left) and the mean ice thickness (bottom right). Note that the latter is calculated only over the glacierized fraction of the catchment and not the entire catchment area (i. e., it is the result of dividing the bottom left plot by the top left plot). Lines and shadings represent multi-model means \pm one standard deviation.

increases for the Pitze catchment in the period 2011–2040, however total annual runoff volumes stay approximately constant due to lower August runoff volumes. In the other three catchments, both monthly peak runoff and total annual runoff volumes do




Figure 14. Spatially distributed simulated glacier coverage (multi-model mean) shown in 10-year intervals for the three RCPs.

not exceed the levels simulated for the historical period in all scenarios. With regard to ice runoff (dashed lines), Gepatschalm is the only catchment where ice runoff does not decrease monotonically over time, but rather increases in the period 2041–2070 compared to 2011–2040 before it strongly decreases towards the end of the century. [This mid-century increase in glacier runoff is likely due to the by far largest average ice thickness for this catchment and the slower decline in glacierization compared to the other catchments.](#)

A more detailed analysis of the changes in monthly runoff is shown in Fig. 16, where the average monthly changes in runoff relative to the historical period 1998–2013 for the catchments in the study region throughout the 21st century are displayed. With respect to the different emission scenarios, again the differences between them are very small within the first considered period (2011–2040). In this period, also no clear trend of changes in the seasonality of the runoff regimes can be discerned, apart for slight increases during the winter months and slight decreases during summer. In the following period, 2041–2071, a clearer pattern begins to emerge. While May to June runoff stays approximately unchanged, runoff strongly decreases during August and September in all three scenarios, amounting to up to –55 %. From October to November, a transition period between decreasing and increasing runoff occurs, while during winter and spring increases of up to 50–125 % (depending on emission scenario) are projected (although still amounting to very small absolute values in comparison to summer as can be seen in Fig. 15). During the last part of the century (2071–2100), summer runoff continues to decrease, although not as strongly as between the first two periods. Winter runoff also increases, most notably for the RCP8.5 scenario, where November to March runoff is at least approximately doubled for all catchments compared to the reference period. In this period, also different catchment characteristics can be more clearly distinguished. The largest changes are simulated for Vernagtbach, the smallest, highest-elevated, and (initially) most glacierized catchment. Here, increases between 43 % and 225 % for November to May runoff are simulated for the RCP8.5 scenario, while the corresponding runoff decreases by 70–75 % during August and September due to the by then almost completely melted glacier (Vernagtferner). However, while in the figure the relative (positive) changes during winter appear similar or larger than the (negative) changes during summer for all catchments, it has to be emphasized that this still corresponds to a strong **relative** decrease in total annual runoff considering the respective




Figure 15. Average monthly runoff (multi-model mean \pm one standard deviation indicated as shaded bands) as simulated for the early, middle, and late 21st century for four catchments and the three emission scenarios. Dashed lines indicate bare ice melt runoff.

absolute values (for example as seen in Fig. 15, the absolute JJA runoff volumes are approx. 2 orders of magnitude larger than the respective DJF values).

The simulated future streamflow composition of the components ice melt, snowmelt and rainfall at the end of the scenario period (2071–2100) is shown in Fig. 17. While the ice melt fractions at the end of the century are negligible for all but the most glacierized catchments, snowmelt is still the major contributor to total runoff for all seasons. Rainfall runoff contributions are comparatively low in general, which is however partly due to the model structure – as rain falling on snow contributes to the liquid water storage of the snowpack up to a certain amount, these rainfall amounts are part of the snowmelt contribution in the runoff concentration scheme. Only rain falling on bare ground, ice, firn or already saturated snow is part of the rainfall runoff contribution in this case.

10 4.5 Uncertainty

Huss et al. (2014) assessed the influence of various model assumptions in projections of glacier evolution and runoff in high-mountain catchments. Their results indicate that major uncertainty sources are especially (i) the winter snow accumulation in terms of both volume and spatial distribution, (ii) the approach to account for glacier geometry changes, (iii) the initial glacier ice volume distribution, and (iv) the individual climate projections.

15 While the model has shown to capture (i) the amount and distribution of winter snow accumulation well for the study area in a recent study (Hanzer et al., 2016), the model setup was slightly altered for the present study. The most substantial changes result




Figure 16. Average changes in monthly runoff relative to the historical period 1998–2013 over the course of the 21st century for the study catchments as simulated for the three emission scenarios (lines are multi-model means; note the differently scaled y-axes for RCP8.5).




Figure 17. Average monthly runoff and composition (multi-model mean) for the period 2071–2100. The black lines show the average runoff in the historical period 1997–2013.

from the requirement of a sufficiently long historical measurement period of the meteorological stations. As a consequence of this, only a subset of the stations utilized in the aforementioned study could be used, and from these stations only daily instead of hourly recordings were available, requiring an additional processing step in terms of temporal disaggregation. In addition, at several of these stations the daily and (sub-)hourly measurements are performed with different instruments, resulting in considerable differences in the recorded meteorological variables as illustrated in Fig. 18 for three stations. Especially the latter impacted model performance, resulting in a tendency to slightly overestimate winter snow accumulation and spring runoff volumes. Reducing the spatial and temporal model resolution from 50 m and 1 h to 100 m and 3 h on the other hand only resulted in minor impacts on model performance. Figure 19 shows the average observed monthly runoff for the seven study catchments and the period 1998–2006, as well as the corresponding simulation results for the 50 m/1 h run using original hourly meteorological recordings and the 100 m/3 h run using disaggregated daily data, highlighting the tendency to overestimate runoff particularly during spring for the latter run. Table 3 shows the corresponding skill scores derived from these two simulation runs: NSE, the benchmark efficiency (BE) (Schaeffli and Gupta, 2007), and the percent bias (PBIAS). However, as for the scenario simulations mainly changes rather than absolute values are analyzed, these partial model biases likely do not affect the main conclusions of our study. Related to this issue is the question whether the implemented precipitation correction methods are still applicable in the future, especially concerning the empirically derived fixed snow correction factor of 15 % as a threshold-based adjustment. Utilizing the RCM-simulated snowfall fractions and precipitation fields instead of downscaling to point locations could reduce this uncertainty. While this was beyond the scope of this study, in the future promising advances in downscaling methods such as quasi-dynamical approaches (e. g., Gutmann et al., 2016) could help to bridge this gap between statistical and dynamical methods and allow deriving more realistic small-scale precipitation fields.

Overestimations of winter snow accumulation and, subsequently, glacier mass balances might partly also have consequences for (ii) the approach to account for glacier geometry changes. The Δh parameterization as implemented is able to account only for conditions of glacier retreat, while in case of positive mass balances no update of the glacier extents is performed. As Fig. 20 shows, 14–24 % of all simulated specific mass balances are positive in our simulations. However, as in previous studies previous studies have shown that the Δh parameterization performed significantly better than alternative approaches such as accumulation area ratio-based methods and similarly well as complex ice flow models (Huss et al., 2010). Hence, we believe that this approach is an adequate tradeoff for the application on the regional scale where the application of of process-based ice flow models is not feasible.

We also assessed the influence of (iii) the initial glacier volume distribution by scaling the original initial ice thickness distribution by factors of 0.7 and 1.3, respectively (i. e., 30 % decrease/increase) and re-running the entire set of RCP4.5 simulations. However, as Fig. 21 (b–c) exemplarily shows for selected catchments, this does not significantly change the model results with regard to the evolution of glaciers and runoff volume. Similar results are obtained also for all other catchments.

With regard to (iv), many studies (e. g., Addor et al., 2014; Bosshard et al., 2013; Horton et al., 2006; Huss et al., 2014) have shown that the largest uncertainty in hydrological impact studies usually is due to the chosen climate models, and that the spread between individual climate model simulations is often larger than the spread between different emission scenarios within a single climate model. In the case of GCM-RCM chains, GCMs tend to have a larger impact on the hydrological




Figure 18. Scatter plots of daily vs. aggregated hourly recordings of air temperature (T), precipitation (P), relative humidity (RH), and wind speed (WS) for three stations in the study area.

model results than the RCMs. We addressed these uncertainties by utilizing the entire range of available EURO-CORDEX simulations, resulting in a total of 14 GCM-RCM chains with five different GCMs (each driving 2–4 RCMs) and six different RCMs for the RCP4.5 and RCP8.5 scenarios. The results for the RCP2.6 scenario have to be interpreted with care, as for this scenario only three GCM-RCM combinations (using two different GCMs) were available. Hence, especially when looking at multi-model averages, the RCP2.6 results are not directly comparable to the other two scenarios.

Besides the climate models themselves, also the spatial and temporal downscaling approaches from the coarse daily-resolution grids to the 3-hourly resolution point-scale time series add an uncertainty component to the results, as they are statistically derived from past conditions for which it is uncertain if they will persist in the future.

The influence of the climate models on the hydrological results is illustrated in Fig. 21 (a). Here, average monthly runoff for the period 2071–2100 as simulated by all 31 individual GCM-RCM realizations is shown exemplarily for the Pitze catchment. While the influence of the emission scenarios is visible to a part, there is significant overlap between the realizations of the different emission scenarios. For example, the realizations resulting in both the lowest and the highest August streamflow volumes are both driven by the RCP8.5 scenario. Similar results were already shown for the simulated glacier volume and area in Fig. 11.

5 Conclusions

In this study, we have forced a fully distributed physically based hydroclimatological model with the most state-of-the-art climate projections available. This is the most detailed study on cryospheric-hydrological climate change impacts in the Ötztal Alps to date, and to our knowledge also in high-elevation glacierized catchments in Austria in general.




Figure 19. Seasonal cycle of observed (black lines) and simulated monthly runoff for the study catchments and the period 1998–2006. Blue lines correspond to the simulation results for the “reference run” (50 m spatial resolution, 1 h temporal resolution, original hourly meteorological data as forcing), while the red lines correspond to the simulation run using 100 m spatial and 3 h temporal resolution as well as disaggregated daily data as forcing.




Figure 20. Histograms of the specific mass balance values as simulated for all (initially) 206 study area glaciers, all simulation years (2006–2100), and all GCM-RCM combinations. Dashed vertical lines refer to the median mass balance, and percentages indicate the total fractions of positive mass balances.

While some uncertainty in the results is due to the model configuration, the largest uncertainty can be traced back to the climate projections. This leads to a considerable range in the projected snow coverage, glacier extents and hydrological regimes. ~~However, some common results can be found for all model runs.~~ Snow cover is projected to decrease ~~, however not as dramatic as presented in some other studies for Alpine regions due to the high elevation of the study site and~~ by up to 80 % in elevations below 1500 m a.s.l., while only comparatively moderate decreases (up to 25 %) are found for high-elevated areas (> 2500 m a.s.l.) due to strongly increasing winter precipitation ~~, which partly compensates for the increased warming.~~ Glaciers will continue to recede strongly throughout the century, ~~and by 2100 most glaciers in the Ötztal Alps might have disappeared (depending on the considered emission scenario).~~ Resulting total glacierized area and ~~Until 2050, glacier volume will likely amount to less than a quarter of today’s state even for the RCP2.6 scenario~~ decline by approx. 60–65 % largely




Figure 21. a) Average monthly runoff (period 2071–2100) for the Pitze catchment as simulated for all 31 individual GCM-RCM realizations, b) evolution of the total glacierized area for the entire Ötztal Alps study site as simulated by altering the initial ice thickness distribution by $\pm 30\%$ (multi-model mean values for the RCP4.5 scenario), and c) corresponding runoff evolution (10-year Gaussian low-pass filtered) for the Rofenache and Gepatschalm catchments.

Table 3. NSE, BE, and PBIAS of observed vs. simulated runoff in the period 1998–2006 for the model runs using original hourly meteorological data and disaggregated daily data, respectively.

ID	Catchment	Original			Disaggregated		
		NSE	BE	PBIAS [%]	NSE	BE	PBIAS [%]
1	Venter Ache	0.92	0.60	8.71	0.79	-0.09	21.44
2	Rofenache	0.93	0.70	8.27	0.78	0.12	23.54
3	Vernagtbach	0.92	0.70	-1.12	0.88	0.57	13.19
4	Gurgler Ache	0.91	0.59	1.96	0.83	0.22	18.87
5	Gepatschalm	0.94	0.68	3.09	0.88	0.41	11.95
6	Taschachbach	0.92	0.54	0.01	0.84	0.08	19.19
7	Pitze	0.94	0.72	2.30	0.88	0.45	16.46

independent of the emission scenario, whereas by the end of the century 80–96 % of the original ice volume will be lost. Consequently, glacier runoff will diminish proportionally and summer runoff will strongly decrease in all investigated catchments by up to 55 %, resulting in a shift of the annual runoff peak from July towards June. Winter runoff volumes on the other hand will increase, however to still low absolute values. While the total annual runoff volumes stay approximately constant during the early 21st century compared to present-day levels, they gradually decrease throughout the rest of the century. Only for some catchments and scenarios runoff volumes slightly exceed present-day levels, indicating that the peak water period of maximum runoff is currently under way or has already passed in this region.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. This work was carried out within the framework of the projects “W01 MUSICALS II – Multiscale Snow/Ice Melt Discharge Simulation into Alpine Reservoirs”, carried out in the research programme of alpS – Centre for Climate Change Adaptation in Innsbruck, and “HydroGeM3”, financed by the Austrian Academy of Sciences. The computational results presented have been achieved
5 in part using the Vienna Scientific Cluster (VSC). The authors want to thank the COMET research programme of the Austrian Research Promotion Agency (FFG), the TIWAG – Tiroler Wasserkraft AG, and the Austrian Academy of Sciences. Meteorological and hydrological data were provided by the Zentralanstalt für Meteorologie und Geodynamik (ZAMG), the Hydrographic Service of Tyrol, the TIWAG – Tiroler Wasserkraft AG, the Commission for Glaciology of the Bavarian Academy of Sciences and Humanities, and the Autonomous Province of Bozen/Bolzano. We also would like to thank Matthias Huttenlau for managing the MUSICALS II project and many helpful discussions,
10 and Johannes Schöber for valuable comments during the preparation of [the manuscript. Finally, we wish to thank two anonymous reviewers for their helpful comments that helped to improve](#) the manuscript.

References

- Abermann, J., Lambrecht, A., Fischer, A., and Kuhn, M.: Quantifying changes and trends in glacier area and volume in the Austrian Ötztal Alps (1969-1997-2006), *The Cryosphere*, 3, 205–215, 2009.
- Addor, N., Rössler, O., Köplin, N., Huss, M., Weingartner, R., and Seibert, J.: Robust changes and sources of uncertainty in the projected hydrological regimes of Swiss catchments, *Water Resources Research*, 50, 7541–7562, 2014.
- Allen, R. G., Pereira, L. S., Raes, D., and Smith, M.: Crop evapotranspiration - Guidelines for computing crop water requirements, Tech. rep., 1998.
- Anderson, E. A.: A point energy and mass balance model of a snow cover, Tech. Rep. NWS 19, NOAA, 1976.
- Barnett, T. P., Adam, J. C., and Lettenmaier, D. P.: Potential impacts of a warming climate on water availability in snow-dominated regions, *Nature*, 438, 303–309, 2005.
- Beniston, M.: Climatic Change in Mountain Regions: A Review of Possible Impacts, *Climatic Change*, 59, 5–31, 2003.
- Beniston, M., Keller, F., Koffi, B., and Goyette, S.: Estimates of snow accumulation and volume in the Swiss Alps under changing climatic conditions, *Theoretical and Applied Climatology*, 76, 125–140, 2003.
- Bliss, A., Hock, R., and Radić, V.: Global response of glacier runoff to twenty-first century climate change, *Journal of Geophysical Research: Earth Surface*, 119, 717–730, 2014.
- Bosshard, T., Carambia, M., Goergen, K., Kotlarski, S., Krahe, P., Zappa, M., and Schär, C.: Quantifying uncertainty sources in an ensemble of hydrological climate-impact projections, *Water Resources Research*, 49, 1523–1536, 2013.
- Cannon, A. J., Sobie, S. R., Murdock, T. Q., Cannon, A. J., Sobie, S. R., and Murdock, T. Q.: Bias Correction of GCM Precipitation by Quantile Mapping: How Well Do Methods Preserve Changes in Quantiles and Extremes?, *dx.doi.org*, 28, 6938–6959, doi:10.1175/JCLI-D-14-00754.1, 2015.
- Collins, D. N.: Climatic warming, glacier recession and runoff from Alpine basins after the Little Ice Age maximum, *Annals of Glaciology*, 48, 119–124, 2008.
- Corripio, J.: Modelling the energy balance of high altitude glacierised basins in the Central Andes, Ph.D. thesis, University of Edinburgh, 2002.
- Déqué, M.: Frequency of precipitation and temperature extremes over France in an anthropogenic scenario: Model results and statistical correction according to observed values, *Global and Planetary Change*, 57, 16–26, 2007.
- Farinotti, D., Usselman, S., Huss, M., Bauder, A., and Funk, M.: Runoff evolution in the Swiss Alps: projections for selected high-alpine catchments based on ENSEMBLES scenarios, *Hydrological Processes*, 26, 1909–1924, 2011.
- Fatichi, S., Rimkus, S., Burlando, P., Bordoy, R., and Molnar, P.: High-resolution distributed analysis of climate and anthropogenic changes on the hydrology of an Alpine catchment, *Journal of Hydrology*, 525, 362–382, 2015.
- Finger, D., Heinrich, G., Gobiet, A., and Bauder, A.: Projections of future water resources and their uncertainty in a glacierized catchment in the Swiss Alps and the subsequent effects on hydropower production during the 21st century, *Water Resources Research*, 48, n/a–n/a, 2012.
- Förster, K., Hanzer, F., Winter, B., Marke, T., and Strasser, U.: An open-source MEteoroLOGical observation time series DISaggregation Tool (MELODIST v0.1.1), *Geoscientific Model Development*, 9, 2315–2333, 2016.
- Frei, P., Kotlarski, S., Liniger, M. A., and Schär, C.: Snowfall in the Alps: Evaluation and projections based on the EURO-CORDEX regional climate models, *The Cryosphere*, pp. 1–38, 2017.

- Frei, P., Kotlarski, S., Liniger, M. A., and Schär, C.: Future snowfall in the Alps: projections based on the EURO-CORDEX regional climate models, *The Cryosphere*, 12, 1–24, doi:10.5194/tc-12-1-2018, 2018.
- Gobiet, A., Kotlarski, S., Beniston, M., and Heinrich, G.: 21st century climate change in the European Alps—A review, *Science of the Total Environment*, 493, 1138–1151, 2014.
- 5 Gudmundsson, L., Bremnes, J. B., Haugen, J. E., and Engen-Skaugen, T.: Technical Note: Downscaling RCM precipitation to the station scale using statistical transformations – a comparison of methods, *Hydrology and Earth System Sciences*, 16, 3383–3390, 2012.
- Gutmann, E., Barstad, I., Clark, M., Arnold, J., and Rasmussen, R.: The Intermediate Complexity Atmospheric Research Model (ICAR), *Journal of Hydrometeorology*, 17, 957–973, doi:10.1175/JHM-D-15-0155.1, 2016.
- Hanzer, F., Marke, T., and Strasser, U.: Distributed, explicit modeling of technical snow production for a ski area in the Schladming region
10 (Austrian Alps), *Cold Regions Science and Technology*, 108, 113–124, 2014.
- Hanzer, F., Helfricht, K., Marke, T., and Strasser, U.: Multilevel spatiotemporal validation of snow/ice mass balance and runoff modeling in glacierized catchments, *The Cryosphere*, 10, 1859–1881, 2016.
- Hanzer, F., Förster, K., Winter, B., and Marke, T.: melodist: MELODIST 0.1.2, doi:10.5281/zenodo.1123277, 2017.
- Helfricht, K., Kuhn, M., Keuschnig, M., and Heilig, A.: Lidar snow cover studies on glaciers in the Ötztal Alps (Austria): comparison with
15 snow depths calculated from GPR measurements, *The Cryosphere*, 8, 41–57, 2014.
- Hofer, M., Nemeč, J., Cullen, N. J., and Weber, M.: Evaluating predictor strategies for regression-based downscaling with a focus on glacierized mountain environments, *Journal of Applied Meteorology and Climatology*, pp. JAMC–D–16–0215.1, 2017.
- Horton, P., Schaefli, B., Mezghani, A., Hingray, B., and Musy, A.: Assessment of climate-change impacts on alpine discharge regimes with climate model uncertainty, *Hydrological Processes*, 20, 2091–2109, 2006.
- 20 Huss, M. and Farinotti, D.: Distributed ice thickness and volume of all glaciers around the globe, *Journal of Geophysical Research: Earth Surface* (2003–2012), 117, F04 010, 2012.
- Huss, M. and Hock, R.: A new model for global glacier change and sea-level rise, *Frontiers in Earth Science*, 3, 22, 2015.
- Huss, M., Farinotti, D., Bauder, A., and Funk, M.: Modelling runoff from highly glacierized alpine drainage basins in a changing climate, *Hydrological Processes*, 22, 3888–3902, 2008.
- 25 Huss, M., Funk, M., and Ohmura, A.: Strong Alpine glacier melt in the 1940s due to enhanced solar radiation, *Geophysical Research Letters*, 36, 17, 2009.
- Huss, M., Juvet, G., Farinotti, D., and Bauder, A.: Future high-mountain hydrology: a new parameterization of glacier retreat, *Hydrology and Earth System Sciences*, 14, 815–829, 2010.
- Huss, M., Zemp, M., Joerg, P. C., and Salzmann, N.: High uncertainty in 21st century runoff projections from glacierized basins, *Journal of
30 Hydrology*, 510, 35–48, 2014.
- Jacob, D., Petersen, J., Eggert, B., Alias, A., Christensen, O. B., Bouwer, L. M., Braun, A., Colette, A., Déqué, M., Georgievski, G., Georgopoulou, E., Gobiet, A., Menut, L., Nikulin, G., Haensler, A., Hempelmann, N., Jones, C., Keuler, K., Kovats, S., Kröner, N., Kotlarski, S., Kriegsmann, A., Martin, E., van Meijgaard, E., Moseley, C., Pfeifer, S., Preuschmann, S., Radermacher, C., Radtke, K., Rechid, D., Rounsevell, M., Samuelsson, P., Somot, S., Soussana, J.-F., Teichmann, C., Valentini, R., Vautard, R., Weber, B., and Yiou, P.: EURO-CORDEX: new high-resolution climate change projections for European impact research, *Regional Environmental Change*, 14, 563–578,
35 2013.
- Jansson, P., Hock, R., and Schneider, T.: The concept of glacier storage: a review, *Journal of Hydrology*, 282, 116–129, 2003.

- Jordan, R.: A one-dimensional temperature model for a snow cover: Technical documentation for SNTHERM. 89., Tech. rep., Hanover, NH, 1991.
- Jouvet, G., Huss, M., Funk, M., and Blatter, H.: Modelling the retreat of Grosser Aletschgletscher, Switzerland, in a changing climate, *Journal of Glaciology*, 57, 1033–1045, 2011.
- 5 Klemeš, V.: The modelling of mountain hydrology: the ultimate challenge, IAHS Publ, 1990.
- Kobierska, F., Jonas, T., Zappa, M., Bavay, M., Magnusson, J., and Bernasconi, S. M.: Future runoff from a partly glacierized watershed in Central Switzerland: A two-model approach, *Advances in Water Resources*, 55, 204–214, 2013.
- Kuhn, M. and Batlogg, N.: Glacier Runoff In Alpine Headwaters In A Changing Climate, IAHS PUBLICATION no. 248, p. 79, 1998.
- Kuhn, M. and Olefs, M.: Auswirkung von Klimaänderungen auf das Abflussverhalten von vergletscherten Einzugsgebieten im Hinblick auf
10 Speicherkraftwerke, Tech. rep., 2007.
- Kuhn, M., Lambrecht, A., and Abermann, J.: The Austrian glacier inventory GI 2, 1998, in ArcGIS (shapefile) format, PANGAEA, 2015.
- Marke, T., Strasser, U., Hanzer, F., Stötter, J., Wilcke, R. A. I., and Gobiet, A.: Scenarios of Future Snow Conditions in Styria (Austrian Alps), *Journal of Hydrometeorology*, 16, 261–277, 2015.
- Marty, C., Schlögl, S., Bavay, M., and Lehning, M.: How much can we save? Impact of different emission scenarios on future snow cover in
15 the Alps, *The Cryosphere*, 11, 517–529, 2017.
- Marzeion, B., Jarosch, A. H., and Hofer, M.: Past and future sea-level change from the surface mass balance of glaciers, *The Cryosphere*, 6, 1295–1322, 2012.
- Marzeion, B., Jarosch, A. H., and Gregory, J. M.: Feedbacks and mechanisms affecting the global sensitivity of glaciers to climate change, *The Cryosphere*, 8, 59–71, 2014.
- 20 Maurer, E. P. and Pierce, D. W.: Bias correction can modify climate model simulated precipitation changes without adverse effect on the ensemble mean, *Hydrology and Earth System Sciences*, 18, 915–925, 2014.
- Milano, M., Reynard, E., Bosshard, N., and Weingartner, R.: Simulating future trends in hydrological regimes in Western Switzerland, *Journal of Hydrology: Regional Studies*, 4, 748–761, 2015.
- Ohmura, A., Bauder, A., Müller, H., and Kappenberger, G.: Long-term change of mass balance and the role of radiation, *Annals of Glaciology*,
25 46, 367–374, 2007.
- Pellicciotti, F., Brock, B., Strasser, U., Burlando, P., Funk, M., and Corripio, J.: An enhanced temperature-index glacier melt model including the shortwave radiation balance: Development and testing for Haut Glacier d’Arolla, Switzerland, *Journal of Glaciology*, 51, 573–587, 2005.
- Pomeroy, J. W., Gray, D. M., Brown, T., Hedstrom, N. R., Quinton, W. L., Granger, R. J., and Carey, S. K.: The cold regions hydrological
30 model: a platform for basing process representation and model structure on physical evidence, *Hydrological Processes*, 21, 2650–2667, 2007.
- Prasch, M., Marke, T., Strasser, U., and Mauser, W.: Large scale integrated hydrological modelling of the impact of climate change on the water balance with DANUBIA, *Advances in Science and Research*, 7, 61–70, 2011.
- Radić, V., Bliss, A., Beedlow, A. C., Hock, R., Miles, E., and Cogley, J. G.: Regional and global projections of twenty-first century glacier
35 mass changes in response to climate scenarios from global climate models, *Climate Dynamics*, 42, 37–58, 2013.
- Ragettli, S., Pellicciotti, F., Bordoy, R., and Immerzeel, W. W.: Sources of uncertainty in modeling the glaciohydrological response of a Karakoram watershed to climate change, *Water Resources Research*, 49, 6048–6066, 2013.

- Schaefli, B.: Projecting hydropower production under future climates: a guide for decision-makers and modelers to interpret and design climate change impact assessments, *Wiley Interdisciplinary Reviews: Water*, 2, 271–289, 2015.
- Schaefli, B. and Gupta, H. V.: Do Nash values have value?, *Hydrological Processes*, 21, 2075–2080, 2007.
- Schmucki, E., Marty, C., Fierz, C., and Lehning, M.: Simulations of 21st century snow response to climate change in Switzerland from a set of RCMs, *International Journal of Climatology*, 35, 3262–3273, 2015a.
- Schmucki, E., Marty, C., Fierz, C., Weingartner, R., and Lehning, M.: Impact of climate change in Switzerland on socioeconomic snow indices, *Theoretical and Applied Climatology*, 127, 875–889, 2015b.
- Seiser, B., Helfricht, K., Huss, M., Fischer, A., Vuillet, E., Schönlaub, H., Kuhn, M., and Strasser, U.: MUSICALS - (MULTiscale Snow/ICemelt Discharge Simulation into ALpine ReservoirS): Validation of a method to estimate ice volume and ice thickness distribution in the Ötztal Alps, Tyrol, Austria, *Geophysical Research Abstracts*, 2012.
- Smiatek, G., Kunstmann, H., and Senatore, A.: EURO-CORDEX regional climate model analysis for the Greater Alpine Region: Performance and expected future change, *Journal of Geophysical Research: Atmospheres*, 121, 7710–7728, 2016.
- Sonntag, D.: Important new values of the physical constants of 1986, vapour pressure formulations based on the ITS-90, and psychrometer formulae, *Zeitschrift für Meteorologie*, 70, 340–344, 1990.
- Stahl, K., Moore, R. D., Shea, J. M., Hutchinson, D., and Cannon, A. J.: Coupled modelling of glacier and streamflow response to future climate scenarios, *Water Resources Research*, 44, n/a–n/a, 2008.
- Steger, C., Kotlarski, S., Jonas, T., and Schär, C.: Alpine snow cover in a changing climate: a regional climate model perspective, *Climate Dynamics*, 41, 735–754, 2012.
- Stewart, I. T.: Changes in snowpack and snowmelt runoff for key mountain regions, *Hydrological Processes*, 23, 78–94, 2009.
- Strasser, U.: Spatial and temporal variability of meteorological variables at Haut Glacier d’Arolla (Switzerland) during the ablation season 2001: Measurements and simulations, *Journal of Geophysical Research*, 109, D03 103, 2004.
- Strasser, U.: Modelling of the mountain snow cover in the Berchtesgaden National Park, Tech. Rep. 55, Berchtesgaden, 2008.
- Strasser, U., Bernhardt, M., Weber, M., Liston, G. E., and Mauser, W.: Is snow sublimation important in the alpine water balance?, *The Cryosphere*, 2, 53–66, 2008.
- Strasser, U., Warscher, M., and Liston, G. E.: Modeling Snow-Canopy Processes on an Idealized Mountain, *Journal of Hydrometeorology*, 12, 663–677, 2011.
- Tecklenburg, C., Francke, T., Kormann, C., and Bronstert, A.: Modeling of water balance response to an extreme future scenario in the Ötztal catchment, Austria, *Advances in Geosciences*, 32, 63–68, 2012.
- Teutschbein, C. and Seibert, J.: Bias correction of regional climate model simulations for hydrological climate-change impact studies: Review and evaluation of different methods, *Journal of Hydrology*, 456–457, 12–29, 2012.
- Teutschbein, C. and Seibert, J.: Is bias correction of regional climate model (RCM) simulations possible for non-stationary conditions?, *Hydrology and Earth System . . .*, 2013.
- Thiemeßl, M., Gobiet, A., and Leuprecht, A.: Empirical-statistical downscaling and error correction of daily precipitation from regional climate models, *International Journal of Climatology*, 31, 1530–1544, 2011a.
- Thiemeßl, M. J., Gobiet, A., and Heinrich, G.: Empirical-statistical downscaling and error correction of regional climate models and its impact on the climate change signal, *Climatic Change*, 112, 449–468, 2011b.
- Thrasher, B., Maurer, E. P., McKellar, C., and Duffy, P. B.: Technical Note: Bias correcting climate model simulated daily temperature extremes with quantile mapping, *Hydrology and Earth System Sciences*, 16, 3309–3314, 2012.

- Viviroli, D., Archer, D. R., Buytaert, W., Fowler, H. J., Greenwood, G. B., Hamlet, A. F., Huang, Y., Koboltschnig, G., Litaor, M. I., López-Moreno, J. I., Lorentz, S., Schädler, B., Schreier, H., Schwaiger, K., Vuille, M., and Woods, R.: Climate change and mountain water resources: overview and recommendations for research, management and policy, *Hydrology and Earth System Sciences*, 15, 471–504, 2011.
- 5 Waichler, S. R. and Wigmosta, M. S.: Development of Hourly Meteorological Values From Daily Data and Significance to Hydrological Modeling at H. J. Andrews Experimental Forest, *Journal of Hydrometeorology*, 4, 251–263, 2003.
- Walter, M. T., Brooks, E. S., McCool, D. K., King, L. G., Molnau, M., and Boll, J.: Process-based snowmelt modeling: does it require more input data than temperature-index modeling?, *Journal of Hydrology*, 300, 65–75, 2005.
- Weber, M., Braun, L., Mauser, W., and Prasch, M.: Contribution of rain, snow-and icemelt in the Upper Danube discharge today and in the
10 future, *Geogr Fis Dinam Quat*, 2010.
- Wijngaard, R. R., Helfricht, K., Schneeberger, K., Huttenlau, M., Schneider, K., and Bierkens, M. F. P.: Hydrological response of the Ötztal glacierized catchments to climate change, *Nordic Hydrology*, 47, 979–995, 2016.
- Wilcke, R. A. I., Mendlik, T., and Gobiet, A.: Multi-variable error correction of regional climate models, *Climatic Change*, 120, 871–887, 2013.
- 15 Yokoyama, R., Shirasawa, M., and Pike, R. J.: Visualizing topography by openness: a new application of image processing to digital elevation models, *Photogrammetric engineering and remote sensing*, 68, 257–266, 2002.
- Zekollari, H., Fürst, J. J., and Huybrechts, P.: Modelling the evolution of Vadret da Morteratsch, Switzerland, since the Little Ice Age and into the future, *Journal of Glaciology*, 60, 1155–1168, 2014.