

Response to reviewer 1 (Prof. Gil Bohrer, Ohio State University)

General Comments – the paper uses LES to determine the mechanisms responsible for increased heat flux from forests with lower albedo than their surroundings. This “convector effect” was previously described by observations. Nonetheless, this manuscript represents a very elegant approach to determine the theory behind this observed effect. They also provide a revised approach to parameterize this effect.

We sincerely thank Prof. Bohrer for the generous comments and the constructive suggestions.

Specific comments: I would be happy if there was an explanation in simple terms of your proposed mechanism behind the convector effect. You are using very technical terms such as “atmospheric coupling”, “aerodynamic resistance to heat transfer” and “aerodynamic roughness” but should spend a few lines in section 2.1 explaining this in layman terms, Will help to expand the readership of this, as it is relevant and interesting for forest ecologists, managers and planers, and not only to forest meteorologists. If I got it correctly, the high leaf area of the forest reduces albedo, which leaves more of the incoming energy in the surface. However, the organization of these dark leaf surfaces is such that they are spread over a relatively thick canopy depth (relative to grassland of shrub land where all leaves are condensed in a much thinner layer). Because canopy in dry forests is sparse, wind can easily penetrate it (i.e., they have lower roughness length and displacement height) and can easily exchange heat with the leaf surfaces. Therefore, forests would have intrinsically lower aerodynamic resistance to heat transfer than shorter biomes (with the same leaf area) or other surfaces (with the same albedo). Given eq 1, that would mean higher heat flux.

We thank Prof. Bohrer for pointing this out. This a valuable suggestion. We have added the following text in section 2.1:

“Therefore to summarize canopy convector effect in simpler terms it can be mentioned that the darker and colder canopy surface reduces albedo, which leaves more of the incoming energy on the canopy surface. However, the organization of these dark leaf surfaces is such that they are spread over a relatively thick canopy depth (relative to grassland or shrub land where all leaves are condensed in a much thinner layer). Because canopy in dry forests is sparse, wind can easily penetrate it and can easily exchange heat with the leaf surfaces. Therefore, forests would have intrinsically lower aerodynamic resistance to heat transfer than shorter biomes because of the higher roughness.

Moreover, the same forest (with the same *physical roughness*) could have higher *aerodynamic roughness* and consequently lower aerodynamic resistance to heat transfer for more heat stressed conditions. Given eq 1, that would mean higher heat flux. Thus while CCE would always be present in a forest compared to a grass or shrubland because of the obvious roughness difference, we establish that CCE can also be present within the same forest for different conditions of heat stress- which is a more subtle point and will be further discussed in the following sections by using large eddy simulations (LES).”

Eq1 – by this equation, you are assuming the big-leaf equivalency. The reality is more complicated, as you identify later (P3 L5). I would restrict the statement in P2 line 26 “it is important to recall that, when adopting the simplified big-leaf representation of the forest as a single surface”.

Agreed and added this sentence: “it is important to recall that, when adopting the simplified big-leaf representation of the forest as a single surface”

Furthermore, I would call r_H the “apparent canopy aerodynamic resistance to heat transfer” (line 29) to indicate that this property is a construct of the formulation and not a direct physical property.

Agreed and modified accordingly. Also added the text: “the word apparent is used to indicate that this property is a construct of the formulation and not a direct physical property”

The entire roughness length formulation (eq 2-8 and table 1) is based on different variants of analytic approximation approaches to reduce the complexity of flow in and above the forest canopy to a 2-D-surface equivalent. It is widely accepted that as you go near the canopy, the MOST approach is not valid. It was proposed (and rather widely accepted) that a mixing length-driven approach can be applied (see Harman and Finnigan 2007, 2008 BLM). Nonetheless, large-scale models, which cannot vertically resolve the canopies, still use MOST and it has been demonstrated to be relatively accurate. My point here is that this section of the manuscript (eq1-8) should not be mixed with the notion that it explains the physics, but state very clearly that it revisits the current leading approach for simplification of the physics in a parameterized way that can be used by coarse-resolution models

Agreed. The following text is added to the end of the section 2.2:

“Before moving on to the usage of LES, it warrants mentioning that the entire roughness length formulation (equation 2 - 8 and table 1) is based on different variants of analytic

approximation approaches to reduce the complexity of flow in and above the forest canopy to a 2-D-surface equivalent. It is widely accepted that the MOST approach is not completely accurate close to the canopy (Foken, 2006). It was proposed that a mixing length-driven approach can be applied (Harman and Finnigan, 2007). Nonetheless, large-scale models, which cannot vertically resolve the canopies, still use MOST and it has been demonstrated to be relatively accurate. Thus from an operational perspective, the present formulation revisits the current leading approach for simplification of the physics in a parameterized way that can be used by coarse-resolution models.”

I think it will be beneficial for the manuscript if you emphasize the point that you are using an LES with an explicit 3-D canopy, where the surface assumptions are not needed to develop a revised approximation approach for the surface-equivalence that account to the forest density effects and parameterize for its outcomes in a way that will allow resolving the heat convector effect even in large-scale models.

Agreed and the following text is added in section 3: “It is worth highlighting again here that the large eddy simulations have been conducted with an explicit 3-D canopy. This means that the surface assumptions are not needed to develop a revised approximation approach for the surface-equivalence that accounts for the forest density effects. Only the outcomes of the LES are parameterized in a way that will allow resolving the canopy convector effect even in large-scale models”.

P3 L26 “d is the zero-plane displacement height taken as $2/3h_c$. . .” will be more correct to say: d is the zero-plane displacement height, often approximated as $2/3h_c$. . . (and see study of roughness length and displacement height in a forest stand and their best approximation – Maurer et al Biogeosciences, 12, 2533–2548, 2015, and Maurer et al Agricultural and Forest Meteorology 177 (2013) 24– 34)

Agreed and modified accordingly. The references are added as well.

Technical corrections: I recommend setting an acronym for Rotenberg and Yakir (2010) - RY10, after the first use. It is listed so many times that it gets rather tedious.

Agreed and changed accordingly.

Pg 2, L 32: “This ‘canopy convector effect’ is sufficiently efficient. . . Word placement is confusing. Can change “sufficiently efficient” to ‘adequate enough’, ‘suitably efficient’

Agreed and changed to ‘adequate enough’.

Pg 6, L 26: “while the output of first 6400 s. . .” change to “while the output of the first 6400 s”

Changed to “while the output of the first 6400 s”

Table 2 Please include the meaning of the Stability class (e.g. “Near-Neutral”, Weakly Unstable”. . .) as the first column of the table. Will make it easier to remember, and look up.

Added the meanings of the stability classes to the table.

Pg 9, L 6: “This description refers to a more general phenomenon as opposed to the the description. . .”. Remove “the”

Removed the extra “the”

Pg 15, L21: Change to “These assumptions also lead to a less nonlinear height variation”.

Changed to “These assumptions also lead to a less nonlinear height variation”.

Response to reviewer 2.

This manuscript is motivated by the work from Rotenberg and Yakir (2010), which observed a decrease in aerodynamic resistance to heat transfer (r_H) over a forest when compared to a shrubland region under similar conditions, an effect that is caused by an increase in surface roughness and is accompanied by an increase in atmospheric instability. This effect was called “canopy convective effect” (CCE) by Rotenberg and Yakir (2010). In this manuscript, the authors investigate the occurrence of CCE above the canopy using Large Eddy Simulation (LES). After observing a decrease in r_H with increase in unstable conditions in the simulations (used as evidence of CCE), the authors compare different models of r_H as a function of height (above the canopy) and atmospheric stability with the simulation results, and conclude that some models can not capture the correct trend of CCE at all (because they present an increase in r_H with instability), and only two (out of eight) models display the signature of CCE (decrease in r_H with instability). The authors proposed an improved parameterization of one of the r_H models by using a value of momentum roughness length scale z_{0m} that vary with atmospheric stability, improving the agreement between model and LES results. The authors conclude that CCE is a generic feature of canopy turbulence. Because the value of r_H is needed for a wide range of applications, the investigation of the behavior of different r_H parameterizations above the canopy is useful, and the use of LES for this purpose is appropriate, therefore this manuscript deals with an interesting topic. However, as described in more details below, I believe the manuscript needs an alternative motivation, better description of the simulations and models and better interpretation of the results.

We thank the reviewer for the nice summary and the constructive comments. We have attempted to respond to all the points raised by the reviewer.

Major specific comments:

I'm not sure I agree with the authors' interpretation of what CCE represents. The authors defined CCE as a decrease in aerodynamic resistance above the canopy, which can be accomplished by an increase in atmospheric instability. In my opinion, it is already well-accepted that there is an increase in turbulent transport (estimated by eddy diffusivity parameters, for example) and consequent decrease in r_H with increasing instability. This should be valid over a canopy and over bare soil. The differences between the canopy and the bare soil cases are the type of the turbulent flow and level of penetration of the transporting eddies across the heat source layer in the canopy case (compared to the no-penetration condition over bare soil), which makes the

turbulent transport different in the canopy case compared to the bare soil case, even if all other factors are the same. In Equation (1) this difference is accounted by reducing the aerodynamic resistance in the canopy case, and I think this is what Rotenberg and Yakir (2010) meant in the definition of CCE. Therefore, a study that wants to better describe the CCE phenomenon should focus on comparing turbulent transport characteristics across different canopies and bare soil, probably for different stabilities, but not only the stability difference in one canopy, as this difference is already expected. Therefore, I believe that the CCE should not be the motivation of this manuscript

It is agreed that the earlier version of the manuscript was not very clear on this issue. We have added the following text to make it more clear:



““Therefore to summarize canopy convective effect in simpler terms it can be mentioned that the darker and colder canopy surface reduces albedo, which leaves more of the incoming energy on the canopy surface. However, the organization of these dark leaf surfaces is such that they are spread over a relatively thick canopy depth (relative to grassland or shrub land where all leaves are condensed in a much thinner layer). Because canopy in dry forests is sparse, wind can easily penetrate it and can easily exchange heat with the leaf surfaces. Therefore, forests would have intrinsically lower aerodynamic resistance to heat transfer than shorter biomes because of the higher roughness. Moreover, the same forest (with the same physical roughness) could have higher aerodynamic roughness and consequently lower aerodynamic resistance to heat transfer for more heat stressed conditions. Given eq 1, that would mean higher heat flux. Thus while CCE would always be present in a forest compared to a grass or shrubland because of the obvious roughness difference, we establish that CCE can also be present within the same forest for different conditions of heat stress- which is a more subtle point and will be further discussed in the following sections by using large eddy simulations (LES).”

I'm surprised with the results of increased r_H with increase negative R_{iB} (increase instability) for the non-MOST models. I believe all models try to replicate the overall idea that turbulent transport increases with instability, and after a quick look on the equations and original manuscripts, it seems to me that r_H should decrease with instability in all models, therefore I'm confused about the results shown in Figures 3 and 4.

This is a good point. The Ri_B at each level has been calculated by the following equation (eq 9 in the paper)


$$Ri_B = \frac{g}{T_a} \frac{(T_a - T_s)(z - d)}{U_{\parallel}^2}$$

where the LES profiles have been used at each level. Thus Ri_B has typical profiles for two instability cases shown in the following figures (1) and (2). As observed, the negative Ri_B at one particular height is higher for the strongly unstable case compared to the weakly unstable case. However, the empirical formulations are highly nonlinear and the effects of the height variations of the profiles of the other parameters (such as U) are manifested in the results in figure 3 and 4. This result is not obvious and that is what motivated our study partly.



If all other parameters are fixed, we would see a variation of rH reducing with increasing negative Ri_B (as You point out correctly) as shown in the following figure (3) which used the formulation of Xie (1998) as an example. Here all parameters are fixed and only Ri_B is varied. So while Your suggestion is correct, there is no conflict with the results in the paper. The formulations are used correctly (we checked them again) as they can reproduce the correct variations as suggested in the corresponding references if all other parameters are

held constant- it is the nonlinear height variations of all other profiles that creates the reported variations in this paper.



The description of the temperature field simulated with LES is not complete. Dias-Junior et al. (2015) simulated only a near neutral case, and Patton et al. (2016) included a source profile term in the temperature equation which comes from the land-surface model, which is not present in the simulations presented here, therefore they cannot be used as references for some of the details of the simulations performed here. In Table 2, the value of $\overline{w'T'_s}$ is defined as being at the ground, where the same values used in Patton et al. (2016) were defined at canopy top. When looking into Figure 1, it is not clear where the imposed heat flux value is, as nowhere in the profile there is a match with the imposed values. If the heat source is applied in the SGS part of the model, and Figure 1(e) shows only the resolved part, maybe the resolved + SGS part of the heat flux should be presented instead. Also, the final profiles of temperature have a peak at canopy top, also different from the results obtained by Patton et al. (2016). Although this may not affect the final conclusions, the equations, sources and boundary conditions used in the temperature field of LES need to be clarified

Although the final conclusions are not affected, we admit that we have been unclear here and thank the reviewer for pointing this out. We have a heat flux imposed at the top of the canopy

and within the canopy there is an exponential decay of the incoming energy due to the absorption and reflection by the leaves. The prescribed heat flux value at the ground surface only becomes effective for grid points that do not have a canopy layer above. Therefore for gridpoints with canopy our setup indeed differs from Patton et al due to the strong absorption of solar radiation within the canopy, which also explains the peak in the temperature profile at the top of the canopy in our case. The reviewer is also correct that Fig 1e) presents only the resolved part of the flux. We have corrected this and with the subgrid part added, the output value at of the top of the canopy equals the value imposed in the simulation.

To quote the PALM manual:

“The heat source distribution is calculated by a decaying exponential function of the downward cumulative leaf area index (integral of the leaf area density), assuming that the foliage inside the plant canopy is heated by solar radiation penetrating the canopy layers according to the distribution of net radiation as suggested by Brown & Covey (1966; Agric. Meteorol. 3, 73-96)). This approach has been applied e.g. by Shaw & Schumann (1992; Bound.-Layer Meteorol. 61, 47-64).”

We have clarified this in the text and also added the necessary equations and boundary conditions in the appendix:

B1 surface heat flux formulation and boundary conditions

The ground surface heat flux for gridpoints with a canopy layer is given by:

$$\overline{w'T'_s} = \overline{w'T'_{toc}} \times \exp\left(-\epsilon_c \int_0^{h_c} LAD(z) dz\right), \quad (\text{B1})$$

with $\epsilon_c = 0.6$ the extinction coefficient of light within the canopy. Within the canopy the plant-canopy heating rate is calculated as the vertical divergence of the canopy heat fluxes:

$$\overline{w'T'_{toc}} \frac{d}{dz} \exp\left(\epsilon_c \int_z^{h_c} LAD(z') dz'\right) \quad (\text{B2})$$

The bottom boundary condition for potential temperature is a Neumann condition, the boundary condition at the top of the domain is such that the initial temperature gradient is maintained at the top of the domain.

Minor specific comments:

Introduction: include a paragraph describing why better estimations of r_H are needed, even though this is a parameter poorly defined for atmospheric transport. I believe the conclusion has some of the information that could be in the intro.

Agreed. The following text has been added:

“The canopy aerodynamic resistance is a concept borrowed from the evapotranspiration literature where it represents the resistance between the idealized ‘big-leaf’ (a reduced order representation of the fully heterogeneous three dimensional canopy) and the atmosphere for heat or vapor transfer (Monteith, 1973; Foken et al., 1995; Alves et al., 1998; Monteith and Unsworth, 2007). The Penman-Monteith equation to calculate evapotranspiration requires parameterization of the aerodynamic resistance which require information on roughness lengths for heat and momentum and stability (Penman, 1948; Allen et al., 1998; Cleverly et al., 2013). r_H parameterizations are also used in global climate models to describe the canopy-atmosphere interaction at the canopy surface layer (Walko et al., 2000). Thus better parameterizations of r_H are of fundamental importance in modeling canopy level fluxes of heat and water vapor which can be used in assessing impacts of climate change, disturbance effects such as vegetation thinning, forest fires etc., as well as for developing forest management strategies.”

Section 2.2: emphasize here that these models were developed for conditions different from canopy sublayer, again this is in the conclusion but should be discussed earlier in the manuscript. This can be a major cause of discrepancies between the models and the simulation, which could be tested by performing simulations without canopy and comparing with the models. After a quick look, I could not find such a test in the literature.

This is a valid point and also has been pointed out by the other reviewer. The following text has been added to section 2.2:

“Before moving on to the usage of LES, it warrants mentioning that the entire roughness length formulation (equation 2 - 8 and table 1) is based on different variants of analytic approximation approaches to reduce the complexity of flow in and above the forest canopy to

a 2-D-surface equivalent. It is widely accepted that the MOST approach is not completely accurate close to the canopy (Foken, 2006). It was proposed that a mixing length-driven approach can be applied (Harman and Finnigan, 2007). Nonetheless, large-scale models, which cannot vertically resolve the canopies, still use MOST and it has been demonstrated to be relatively accurate. Thus from an operational perspective, the present formulation revisits the current leading approach for simplification of the physics in a parameterized way that can be used by coarse-resolution models.”

Section 4.1: if possible, when describing the figures in the text, give some justification of the result encountered, for example, if the variations with instability observed makes physical sense.

The following text has been added.

“These results are physical consistent. The near neutral case is dominated by mechanical shear driven turbulence - given by the highest mean velocity. The free convection case is fully buoyancy driven and the motion is fully upwards-as evident by the near zero mean horizontal velocity. For the same reasons, the turbulent intensity and friction velocity follow the same pattern. The strongly unstable cases have highest heat fluxes, which is also physically consistent.”

Page 7, line 21: mention how the eddy diffusivities were estimated.

We have added this in an appendix:

B2 Eddy diffusivity formulation

The computation for the eddy diffusivities in PALM follows the standard procedure for 1.5 order turbulence closure. Thus they are computed from the subgrid-scale turbulent kinetic energy, more precisely equations (13-14) from Maronga et al. (2015).

Eddy diffusivity for momentum:

$$K_m = c_m l \sqrt{e} \tag{B3}$$

Eddy diffusivity for heat:

$$K_h = \left(1 + \frac{2l}{\Delta}\right) K_m \tag{B4}$$

With e the subgrid-scale turbulent kinetic energy (a prognostic variable), $c_m = 0.1$, and Δ the geometric mean of the grid spacings in x,y and z. Finally, l is the subgrid-scale mixing length depending on Δ , stability, and distance from the topography elements or ground surface.

Page 10, line 6: describe how these profiles were estimated. Which values came from LES, which are constant, which are a function of height, for example?

The following text has been added: “To compute r_H variations, the LES generated profiles of mean velocity u , sensible heat flux, air temperature, Prandtl number (thus the diffusivities) are used where all of them have z variations. The friction velocity u_ and the roughness lengths are fixed.”*

Page 11, line 5: not clear what “for weaker cases” mean. Do you mean for weaker instabilities?

Changed to weaker instabilities.

Technical corrections and minor suggestions:

• Why “(in)stability”?

Removed the brackets.

Figure 3: I suggest to use z/z_i (where z_i is the top of the ABL) instead of z/h , emphasizing that the entire plot is above the canopy. It can help to discuss the region where MOST (and therefore some of the models) is valid (surface layer).

We have retained the z/h scaling since the convective effect is most prominent close to the canopy. In canopy turbulence studies where the canopy sub layer is discussed, the convention is using h , since the canopy sublayer is described as the region between $h-5h$. The plots start from $z/h=1$ to 10, which clearly means they are above the canopy- which would not be as clear if z_i is used. Moreover, the reason why h is used for CSL is that is the most dominant length scales among a number of length scales in the canopy sub layer -so the choice is not arbitrary and has a physical basis.

Figure 4: I believe that the blue captions below the figures are wrong.

Thanks for pointing out. The blue captions are removed.

Explaining the convector effect in canopy turbulence by means of large-eddy simulation

Tirtha Banerjee^{1,2}, Frederik De Roo¹, and Matthias Mauder¹

¹Karlsruhe Institute of Technology (KIT) Institute of Meteorology and Climate Research, Atmospheric Environmental Research (IMKIFU), D-82467 Garmisch-Partenkirchen, Germany

²Present address: Earth and Environmental Sciences Division, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA

Correspondence to: Tirtha Banerjee (tirtha.banerjee@kit.edu, banerjeetirtha10@gmail.com)

Abstract. Semi-arid forests are found to sustain a massive sensible heat flux in spite of having a low surface to air temperature difference by lowering the aerodynamic resistance to heat transfer (r_H) – a property called ‘canopy convector effect’ (CCE). In this work large-eddy simulations are used to demonstrate that CCE appears more generally in canopy turbulence. It is indeed a generic feature of canopy turbulence: r_H of a canopy is found to reduce with increasing unstable stratification, which effectively increases the aerodynamic roughness for the same physical roughness of the canopy. This relation offers a sufficient condition to construct a general description of CCE. In addition, we review existing parameterizations for r_H from the evapotranspiration literature and test to what extent they are able to capture the CCE, thereby exploring the possibility of an improved parameterization.

1 Introduction

Understanding the role of turbulence in interactions between vegetation canopies and the atmosphere is crucial to interpret momentum and scalar fluxes above vegetation. This is relevant for a number of practical applications such as regional and global weather and climate modeling, energy balance closure studies, developing forest management strategies etc. Measurement campaign networks such as FLUXNET monitor carbon, water and energy fluxes on a long term basis for this same reason (Baldocchi et al., 2001) to study how different ecosystems interact with the atmosphere and influence local and global weather and climate. One such measurement campaign (Rotenberg and Yakir, 2011) focused on semi-arid ecosystems, specifically the Yatir forest situated in the Negev desert in Israel, to study the survival and productivity of the pine forest in spite of the high radiation load and suppressed latent heat flux. An important outcome of this campaign was the concept of ‘canopy convector effect’ (CCE) introduced by Rotenberg and Yakir (2010), [thereafter called RY10](#). To quote [them \(Rotenberg and Yakir, 2010\)RY10](#), “With suppressed latent heat flux (LE) because of lack of water, the forest is transformed into an effective ‘convector’ that exploits the low tree density and open canopy and, consequently, high canopy-atmosphere aerodynamic coupling.” [Rotenberg and Yakir \(2010\)RY10](#) ascribed the origin of the CCE to the roughness difference between desert and forest. However, in the present work, we demonstrate that the canopy convector effect appears more generally in canopy turbulence. In fact, we show that the CCE is also a generic artifact of homogeneous canopy turbulence by using

large-eddy simulations (LES). In doing so, the canopy aerodynamic resistance to heat transfer (r_H) is revisited. The canopy aerodynamic resistance is a concept borrowed from the evapotranspiration literature where it represents the resistance between the idealized ‘big-leaf’ (a reduced order representation of the fully heterogeneous three dimensional canopy) and the atmosphere for heat or vapor transfer (Monteith, 1973; Foken et al., 1995; Alves et al., 1998; Monteith and Unsworth, 2007).

5 [The Penman-Monteith equation to calculate evapotranspiration requires parameterization of the aerodynamic resistance which require information on roughness lengths for heat and momentum and stability \(Penman, 1948; Allen et al., 1998; Cleverly et al., 2013\).](#)
 [\$r_H\$ parameterizations are also used in global climate models to describe the canopy-atmosphere interaction at the canopy surface layer \(Walko et al., 2000\). Thus better parameterizations of \$r_H\$ are of fundamental importance in modeling canopy level fluxes of heat and water vapor which can be used in assessing impacts of climate change, disturbance effects such as](#)
10 [vegetation thinning, forest fires etc., as well as for developing forest management strategies.](#)

We investigate if the existing parameterizations of the canopy aerodynamic resistance exhibit CCE and we identify uncertainties in their application. As the CCE is the crucial mechanism that ensures the survival of the Yatir forest, an improved physical understanding of the CCE is of primordial importance when considering large-scale afforestation in semi-arid regions.

2 Background and theory

15 2.1 The canopy convector effect and aerodynamic resistance

As mentioned earlier, the canopy convector effect was introduced by Rotenberg and Yakir (2010, 2011) while studying the interaction of vegetation cover with the surface radiation balance for the Yatir forest. The annual average incoming solar radiation in the Yatir forest is about 238 W m^{-2} comparable to that in the Sahara desert but the net radiation (R_n) is about 35% higher than the Sahara (Rotenberg and Yakir, 2010) (RY10) due to the lower albedo of the forest. However, both remote
20 sensing and local measurements indicated that the surface temperature of the forest canopy in Yatir is lower than the surface temperature of the nearby non-forested area – on annual average by about 5K. This is striking, as firstly, the lower albedo (by 0.1) of the forest than that of the surrounding shrubland translates into an approximate 24 W m^{-2} increase of radiation load on the forest canopy. Secondly, the cooler canopy surface suppresses the upwelling longwave radiation, resulting in an additional increase of radiation load by about 25 W m^{-2} . The combined annual increase of radiation load by about 50 W m^{-2} associated
25 with the Yatir afforestation in the Negev is quite high and is comparable to net annual radiation difference between the Sahara desert and Denmark, for example (Rotenberg and Yakir, 2010) (RY10). Thirdly, the latent heat flux of evapotranspiration (LE), the obvious cooling and energy dissipation mechanism in temperate forests, is not an option since water is virtually unavailable for about 7 months a year. Thus sensible heat flux (H) is the only major heat dissipation route, translating into a Bowen ratio (H/LE) as high as 20 or more — unlike temperate forests with Bowen ratio ≈ 1 . In the Yatir forest, the entire net solar
30 radiation flux (up to 800 W m^{-2}) is equilibrated by a massive sensible heat flux (H) of similar magnitude. Note that this high H cannot be explained by the difference between surface and air temperature ($\Delta T = T_s - T_a$) as the canopy surface is cooler than the surrounding desert surface in this case, but the air temperatures above desert and forest canopy are similar. To expound this apparent contradiction of larger sensible heat flux for smaller ΔT , it is important to recall that [when adopting the simplified](#)

big-leaf representation of the forest as a single surface

$$H = -\rho C_p \frac{T_a - T_s}{r_H}, \quad (1)$$

where ρ and C_p are the density and specific heat capacity of air respectively, T_a is air temperature, T_s is canopy surface temperature and r_H is the apparent canopy aerodynamic resistance to heat transfer (the word apparent is used to indicate that this property is a construct of the formulation and not a direct physical property). Hence the large H is not explained by the temperature difference (Δ_T) but by a decreased r_H . Thus the semi-arid forest with its low tree density and large surface area becomes an efficient low aerodynamic resistance ‘convector’ that is well coupled to the atmosphere above (Rotenberg and Yakir, 2010, 2011). This ‘canopy convector effect’ (CCE) is sufficiently efficient-adequate enough to support the massive sensible heat flux larger than the surrounding Negev desert, still maintaining a relatively cooler (than the desert) surface temperature (of the canopy top). It is worth noting here that equation 1 offers a very simplistic description of the complex mixing process in the surface layer, however, it should be interpreted as a zeroth order representation of the corresponding processes. Rotenberg and Yakir (2010)-RY10 identified the difference of roughness between the desert and forest as the underlying mechanism of CCE by arguing that $r_H \propto 1/PAI$ where PAI denotes the plant area index. However, in this work, we attempt to identify a more fundamental mechanism behind CCE which is more strongly connected to the feature of canopy turbulence. Therefore we hypothesize that even with the same physical roughness, variation of the aerodynamic roughness is a sufficient condition for displaying CCE. This difference of aerodynamic roughness for the same physical roughness (of the same vegetation canopy) can be generated by changing the intensity of atmospheric stratification (Zilitinkevich et al., 2008). Thus observing the variation of the canopy aerodynamic resistance to heat transfer (r_H) with atmospheric instability is a sufficient condition to demonstrate the generality of the CCE. To be more precise, if r_H is found to decrease with increasing unstable stratification, that would exhibit the fact that canopy turbulence effectively reduces the aerodynamic resistance to cope with heat stressed environments, i.e the canopy convector effect would manifest itself.

Therefore to summarize canopy convector effect in simpler terms it can be mentioned that the darker and colder canopy surface reduces albedo, which leaves more of the incoming energy on the canopy surface. However, the organization of these dark leaf surfaces is such that they are spread over a relatively thick canopy depth (relative to grassland or shrub land where all leaves are condensed in a much thinner layer). Because canopy in dry forests is sparse, wind can easily penetrate it and can easily exchange heat with the leaf surfaces. Therefore, forests would have intrinsically lower aerodynamic resistance to heat transfer than shorter biomes because of the higher roughness. Moreover, the same forest (with the same physical roughness) could have higher aerodynamic roughness and consequently lower aerodynamic resistance to heat transfer for more heat stressed conditions. Given equation 1, that would mean higher heat flux. Thus while CCE would always be present in a forest compared to a grass or shrubland because of the obvious roughness difference, we establish that CCE can also be present within the same forest for different conditions of heat stress- which is a more subtle point and will be further discussed in the following sections by using large eddy simulations (LES).

LES provides a useful and meanwhile standard tool for studying canopy turbulence under different conditions of atmospheric stratification. A recent publication by Patton et al. (2015) studied the influence of different atmospheric (in)stability-instability

classes on coupled boundary layer-canopy turbulence. In this work, those same [\(in\)stability-instability](#) classes are simulated to put our hypothesis to the test.

2.2 Parameterizations for canopy aerodynamic resistance to heat transfer

Apart from the LES outcomes, it is also important to study if the existing parameterizations of r_H can exhibit CCE. Parameterizations of r_H in the literature use Monin-Obukhov similarity theory (MOST) extensively. MOST can provide corrections for the vertical profile of the mean longitudinal velocity u and potential temperature ($T_a - T_s$) under thermal stratification, which deviates from the traditional log-law under neutral conditions. Thus under MOST, with the assumption that the vegetation is low, dense and horizontally homogeneous,

$$u = \frac{u_*}{\kappa} \left[\ln \left(\frac{z-d}{z_{0m}} \right) - \psi_m(\zeta, \zeta_{0m}) \right], \quad (2)$$

10 and

$$T_a - T_s = P_{r0} \frac{T_*}{\kappa} \left[\ln \left(\frac{z-d}{z_{0h}} \right) - \psi_h(\zeta, \zeta_{0h}) \right], \quad (3)$$

where u_* is the friction velocity, κ is von Kármán constant, z is height from the ground, d is the zero-plane displacement height [taken, often approximated](#) as $(2/3)h_c$ as per literature [\(Seginer, 1974; Shuttleworth and Gurney, 1990; Alves et al., 1998\)](#) [\(Seginer, 1974\)](#) and $\zeta = (z-d)/L$ is called the stability parameter. L is the Obukhov length, computed as

$$15 \quad L = -\frac{u_*^3 T_a}{\kappa g \overline{w'T'}}, \quad (4)$$

where $g = 9.81 \text{ms}^{-2}$, the gravitational acceleration. $\overline{w'T'}$ is the sensible heat flux — assumed to be constant in the surface layer (Foken, 2006). Negative ζ indicates unstable stratification and thus ζ decreases with increasing instability. z_{0m} and z_{0h} are the characteristic roughness lengths for momentum and heat transfer respectively. $\zeta_{0m} = z_{0m}/L$ and $\zeta_{0h} = z_{0h}/L$ are the stability parameters associated with roughness lengths. $P_{r0} = K_m/K_h$ is the turbulent Prandtl number where K_m and K_h are eddy diffusivities of momentum and heat, respectively. T_* is a characteristic temperature scale, obtained from H and the characteristic velocity scale, i.e.,

$$H = -\rho C_p u_* T_*. \quad (5)$$

Combining equations 1, 2, 3 and 5, one can write

$$r_H = \frac{P_{r0}}{\kappa^2 u} \left[\ln \left(\frac{z-d}{z_{0m}} \right) - \psi_m(\zeta, \zeta_{0m}) \right] \left[\ln \left(\frac{z-d}{z_{0h}} \right) - \psi_h(\zeta, \zeta_{0h}) \right], \quad (6)$$

25 where ψ_m and ψ_h are the integral stability correction functions for momentum and heat, respectively. Following Liu et al. (2007), they can be parameterized for unstable conditions as (Dyer and Hicks, 1970; Paulson, 1970; Dyer, 1974; Garratt, 1977; Webb, 1982)

$$\psi_m(\zeta, \zeta_{0m}) = 2 \ln \left(\frac{1+x}{1+x_0} \right) + \ln \left(\frac{1+x^2}{1+x_0^2} \right) - 2 \tan^{-1} x + 2 \tan^{-1} x_0, \quad (7)$$

$$\psi_h(\zeta, \zeta_{0h}) = 2 \ln \left(\frac{1+y}{1+y_0} \right); \quad (8)$$

where $x = (1 - \gamma_m \zeta)^{1/4}$, $x_0 = (1 - \gamma_m \zeta_{0m})^{1/4}$, $y = (1 - \gamma_h \zeta)^{1/2}$ and $y_0 = (1 - \gamma_h \zeta_{0h})^{1/2}$. Different values for the parameters γ_m and γ_h are reported in the literature, and the ones suggested by Paulson (1970) are used, i.e., $\gamma_m = \gamma_h = 16$. This formulation for r_H given by equation 6 with some approximations ($\zeta_{0m} = \zeta_{0h} = 0$) was first used by Thom (1975) and is called the ‘reference parameterization’ (Liu et al., 2007). The full form of equation 6 was used by Yang et al. (2001) with their only approximation being $P_{r_0} = 1$. Several other studies also used semi-empirical and empirical parameterizations and included the bulk Richardson number Ri_B (Monteith, 1973) given by

$$Ri_B = \frac{g}{T_a} \frac{(T_a - T_s)(z - d)}{U_{\parallel}^2}, \quad (9)$$

with U_{\parallel} the horizontal wind speed at the height that corresponds to the T_a measurement.

Liu et al. (2007) compiled different parameterizations of r_H which we will test in the context of the canopy convector effect against our LES output. Table 1 lists the details of the different parameterizations as compiled by Liu et al. (2007). These parameterizations based on MOST (Thom, 1975; Yang et al., 2001), empirical (E) (Verma et al., 1976; Hatfield et al., 1983; Mahrt and Ek, 1984; Xie, 1988) and semi-empirical (SE) (Choudhury et al., 1986; Viney, 1991) assumptions can be classified into two categories. Formulations by Thom (1975), Choudhury et al. (1986), Yang et al. (2001) and Viney (1991) have assumed $z_{0m} \neq z_{0h}$, which should be a more realistic assumption. On the other hand, formulations by Verma et al. (1976), Hatfield et al. (1983), Mahrt and Ek (1984) and Xie (1988) assumed $z_{0m} = z_{0h}$. Different parameters used in the empirical formulations are also listed in table 1.

One important point to notice is that only the formulation by Yang et al. (2001) uses the stability parameters associated with the roughness lengths ζ_{0m} and ζ_{0h} . Also note that all parameterizations assume a turbulent Prandtl number of unity, i.e., the diffusivities for momentum and heat are assumed to be the same. We shall later discuss the consequence of letting this parameter vary. Another important approximation necessary to evaluate all formulations in table 1 is a prescription for the roughness lengths z_{0m} and z_{0h} . Effects of different roughness lengths will be investigated in the following section. However, a relation between the two roughness lengths ($\kappa B^{-1} = \ln(z_{m0}/z_{0h})$) was proposed by Owen and Thomson (1963) and Chamberlain (1968), where κB^{-1} is called an ‘excess resistance parameter’. Yang et al. (2001) suggested an average value of $\kappa B^{-1} = 2.0$ (Liu et al., 2007), which will be used throughout this work.

Before moving on to the usage of LES, it warrants mentioning that the entire roughness length formulation (equation 2 - 8 and table 1) is based on different variants of analytic approximation approaches to reduce the complexity of flow in and above the forest canopy to a 2-D-surface equivalent. It is widely accepted that the MOST approach is not completely accurate close to the canopy (Foken, 2006). It was proposed that a mixing length-driven approach can be applied (Harman and Finnigan, 2007). Nonetheless, large-scale models, which cannot vertically resolve the canopies, still use MOST and it has been demonstrated to be relatively accurate. Thus from an operational perspective, the present formulation revisits the current leading approach for simplification of the physics in a parameterized way that can be used by coarse-resolution models.

Table 1. Different parameterizations of r_H as compiled by Liu et al. (2007).

Source	Parameterization of r_H	Coefficients	Assumption
Thom (1975)	$r_H = \frac{1}{\kappa^2 u} \left[\ln \left(\frac{z-d}{z_{0m}} \right) - \psi_m(\zeta) \right] \left[\ln \left(\frac{z-d}{z_{0h}} \right) - \psi_h(\zeta) \right]$	$\zeta_{0m} = \zeta_{0h} = 0$	$z_{0m} \neq z_{0h}$, MOST
Yang et al. (2001)	$r_H = \frac{1}{\kappa^2 u} \left[\ln \left(\frac{z-d}{z_{0m}} \right) - \psi_m(\zeta, \zeta_{0m}) \right] \left[\ln \left(\frac{z-d}{z_{0h}} \right) - \psi_h(\zeta, \zeta_{0h}) \right]$	NA	$z_{0m} \neq z_{0h}$, MOST
Choudhury et al. (1986)	$r_H = \frac{1}{\kappa^2 u} \left[\ln \left(\frac{z-d}{z_{0m}} \right) \right] \left[\ln \left(\frac{z-d}{z_{0h}} \right) \right] (1 - \beta Ri_B)^{-3/4}$	$\beta = 5$	$z_{0m} \neq z_{0h}$, SE
Viney (1991)	$r_H = \frac{1}{\kappa^2 u} \left[\ln \left(\frac{z-d}{z_{0m}} \right) \right] \left[\ln \left(\frac{z-d}{z_{0h}} \right) \right] [a + b(-Ri_B)^c]^{-1}$ $a = 1.0591 - 0.0552 \ln \left(1.72 + \left[4.03 - \ln \left(\frac{z-d}{z_{0m}} \right) \right]^2 \right)$ $b = 1.9117 - 0.2237 \ln \left(1.86 + \left[2.12 - \ln \left(\frac{z-d}{z_{0m}} \right) \right]^2 \right)$ $c = 0.8437 - 0.1243 \ln \left(3.49 + \left[2.79 - \ln \left(\frac{z-d}{z_{0m}} \right) \right]^2 \right)$	$a, b, c = f((z-d)/z_{0m})$	$z_{0m} \neq z_{0h}$, SE
Verma et al. (1976)	$r_H = \frac{1}{\kappa^2 u} \left[\ln \left(\frac{z-d}{z_{0m}} \right) \right]^2 (1 - 16 Ri_B)^{-1/4}$	NA	$z_{0m} = z_{0h}$, E
Hatfield et al. (1983)	$r_H = \frac{1}{\kappa^2 u} \left[\ln \left(\frac{z-d}{z_{0m}} \right) \right]^2 (1 + \beta Ri_B)$	$\beta = 5$	$z_{0m} = z_{0h}$, E
Mahrt and Ek (1984)	$r_H = \frac{1}{\kappa^2 u} \left[\ln \left(\frac{z-d}{z_{0m}} \right) \right]^2 \left[\frac{1 + c(-Ri_B)^{1/2}}{1 + c(-Ri_B)^{1/2} - 15 Ri_B} \right]$	$c = \frac{75 \kappa^2 \left(\frac{z-d+z_{0m}}{z_{0m}} \right)^{1/2}}{\left[\ln \left(\frac{z-d+z_{0m}}{z_{0m}} \right) \right]^2}$	$z_{0m} = z_{0h}$, E
Xie (1988)	$r_H = \frac{1}{\kappa^2 u} \left[\ln \left(\frac{z-d}{z_{0m}} \right) \right]^2 \left[1 + \frac{\left[1 - 16 Ri_B \ln \left(\frac{z-d}{z_{0m}} \right) \right]^{-1/2}}{\ln \left(\frac{z-d}{z_{0m}} \right)} \right]$	NA	$z_{0m} = z_{0h}$, E

3 Methodology

The PALM large-eddy simulation model (Raasch and Schröter, 2001; Maronga et al., 2015) is used to investigate this generic nature of canopy convective effect. The representation of the canopy in the LES follows the standard distributed drag parameterization (Shaw and Schumann, 1992; Watanabe, 2004; Patton et al., 2015) by adding an additional term in the momentum budget equations as $F_{d_i} = -C_d a |\mathbf{u}| u_i$ where a is a one-sided frontal plant area density (PAD), C_d is a dimensionless drag coefficient assumed to be 0.3 (Katul et al., 2004; Banerjee et al., 2013), $|\mathbf{u}|$ is the wind speed and u_i is the corresponding velocity component ($i = 1, 2, 3$, i.e. u , v and w). The effect of the canopy on the subgrid scale (SGS) turbulence is accounted for by adding a sink term to the prognostic equation for the SGS turbulent kinetic energy (e) as $F_\epsilon = -2C_d a |\mathbf{u}| e$. For closure of the SGS covariance terms, PALM uses the 1.5 order closure developed by Deardorff (1980) as modified by Moeng and Wyngaard (1988) and Saiki et al. (2000), which assumes a gradient-diffusion parameterization. The diffusivities associated with this gradient-diffusion are parameterized using the subgrid-scale turbulent kinetic energy (SGS-TKE) and includes a prognostic equation for the SGS-TKE. This SGS-TKE scheme after Deardorff (1980) is deemed to be an improvement over the more traditional Smagorinsky (1963) parameterization since the SGS-TKE allows for a much better estimation for the velocity scale corresponding to the subgrid-scale fluctuations (Maronga et al., 2015). Further details of the LES model can be found in the literature and are not discussed here (Shaw and Schumann, 1992; Watanabe, 2004; Maronga et al., 2015; Patton et al., 2015). For our simulation the number of grid points in the x , y and z directions are 320, 320 and 640 respectively, with grid resolution of 3.91 m, 3.91 m and 1.95 m in the respective direction. Each simulation has a simulated time of 10000 s with a time step of 0.1 s, while the output of the first 6400 s are discarded before achieving computational quasi-equilibrium. The canopy height (h_c) is taken as 35.0 m with a plant area index (PAI) of 5.0. It is important to note that Rotenberg and Yakir (2011) reported an effective PAI of about 5–6 for heat exchange for the Yatir forest. This makes our PAI similar to a recent simulation study of Dias-Junior et al. (2015). In fact, as we already simulate a homogeneous canopy to show that the CCE appears more generically above vegetation canopies, we have decided to tailor our simulations following the examples of Patton et al. (2015) and Dias-Junior et al. (2015) in order to allow a better comparison of the LES data. The vertical distribution of plant area density (a) follows the pdf of a Beta distribution as described in Markkanen et al. (2003) and the parameters α and β controlling the vertical distribution of foliage are set as 3.0 and 2.0 respectively to simulate a PAD distribution similar to Dias-Junior et al. (2015). The parameters to drive the simulations for five different ~~(in)stability~~ instability classes namely near-neutral (NN), weakly unstable (WU), moderately unstable (MU), strongly unstable (SU) and free convection (FC) are similar to those of Patton et al. (2015) and are presented in table 2. Note that the canopy convective effect as a general phenomenon should not depend on water content in the soil-plant-atmosphere continuum and moreover, the PALM-LES does not take into account any physiological processes which normally happen with a larger time scale. Nevertheless, instead of simulating a specific dry water free environment, some moisture at the lower surface is provided and the boundary conditions for surface moisture content are taken similar to the simulations of Dias-Junior et al. (2015) as well. The initial conditions of the potential temperature (and moisture) profile as also taken similar to Dias-Junior et al. (2015). PALM’s canopy module allows sensible heat flux input at the canopy top only, and the sensible heat flux is attenuated exponentially due to the decay of the incoming energy by absorption

Table 2. Parameters to drive the simulations for five different ~~(in)stability~~-~~instability~~ classes namely near-neutral (NN), weakly unstable (WU), moderately unstable (MU), strongly unstable (SU) and free convection (FC) are similar to Patton et al. (2015). U_g and V_g denote geostrophic wind speeds, $\overline{w'T'_s}$ $\overline{w'T'_{top}}$ denotes ~~ground~~-~~canopy top~~ surface sensible heat flux, T_s denotes ground surface potential temperature and q_s denotes specific humidity at the ground surface.

Stability class	$(U_g, V_g) (ms^{-1})$	$\overline{w'T'_s} (Kms^{-1})$	$T_s (K)$	$q_s (g/g)$
NN -Near Neutral (NN)	20, 0	0.24 -0.18	307.7	0.02
WU -Weakly Unstable (WU)	10, 0	0.21 -0.18	307.7	0.02
MU -Moderately Unstable (MU)	5, 0	0.20 -0.18	307.7	0.02
SU -Strongly Unstable (SU)	2, 0	0.20 -0.18	307.7	0.02
FC -Free Convection (FC)	0, 0	0.23 -0.18	307.7	0.02

~~and reflection by the leaves. Thus the ground surface heat flux would be different from Patton et al. (2015).~~ Another important point to note is that instead of lowering the wind speeds while maintaining similar sensible heat fluxes, the different stability classes can also be achieved by maintaining the same wind speed and ramping up the surface sensible heat fluxes. However, this should not affect the generic feature of CCE as discussed at the end of section 2.1. ~~Further details and boundary conditions~~

5 ~~about the LES are discussed in appendix B.~~

~~It is worth highlighting again here that the large eddy simulations have been conducted with an explicit 3-D canopy. This means that the surface assumptions are not needed to develop a revised approximation approach for the surface-equivalence that accounts for the forest density effects. Only the outcomes of the LES are parameterized in a way that will allow resolving the canopy convective effect even in large-scale models.~~

10 4 Results and Discussions

4.1 Comparison with LES

The results of the LES simulations are presented in figure 1 as temporally and spatially averaged vertical profiles for all five stability classes, where the lightest cyan shade indicates near neutral and the most magenta shade indicates free convective conditions. Panel (a) shows the mean wind speed (U), panel (b) shows the standard deviation of longitudinal velocity fluctuations (σ_u) and panel (c) shows the friction velocity (which can be taken as a measure of turbulent intensity) (u_*) at every level for each simulation

$$u_* = (\overline{u'w'^2} + \overline{v'w'^2})^{1/4}. \quad (10)$$

In the second row, panel (d) shows profiles of temporally and spatially averaged potential temperature (T), panel (e) shows the kinematic sensible heat flux ($\overline{w'T'}$) and panel (f) shows the Prandtl number $P_{r0} = K_m/K_h$. The profiles (except P_{r0}) are shown in their dimensional form to clearly illustrate the differences between the different stability conditions. The simulation results closely follow the results presented in Patton et al. (2015) and Dias-Junior et al. (2015). It is interesting to observe that




Figure 1. Summary statistics of five LES simulations showing the variations between different stability classes in increasing order of instability- from near neutral to free convection color coded as indicated in the legend.

the magnitude of velocity, the velocity fluctuations and the turbulent intensity decreases gradually from the near neutral to free convective conditions, i.e., with increasing instability. The potential temperature also reduces with increasing instability at all heights. On the other hand, the sensible heat flux appears to increase with increasing stability, especially more above the forest ($z/h_c = 1$ indicates the canopy top). These results are physical consistent. The near neutral case is dominated by mechanical shear driven turbulence - given by the highest mean velocity. The free convection case is fully buoyancy driven and the motion is fully upwards-as evident by the near zero mean horizontal velocity. For the same reasons, the turbulent intensity and friction velocity follow the same pattern. The strongly unstable cases have highest heat fluxes, which is also physically consistent. Note that the canopy top sensible heat flux is similar to the imposed value of 0.18 Kms^{-1} that was used to drive the simulations.




Figure 2. Aerodynamic resistance to heat transfer exhibiting canopy convective effect. Panel (a): Difference between surface and air temperature ($T_s - T_a$); panel (b): stability parameter ζ ; panel (c): canopy aerodynamic resistance (r_H).

Figure 2 shows temporally and spatially averaged vertical profiles for the different stability conditions with the same color coding as figure 1. We investigate the vertical profile of r_H in order to assess the uncertainty that arises from varying the reference height for the air temperature under varying stability. The temperature of the canopy top is taken as the surface temperature (T_s) and thus results are shown from above the canopy top, i.e., $z/h_c = 1$. Panel (a) shows the difference of surface and air temperature ($T_s - T_a(z)$). Panel (b) shows the stability parameter ζ at every level computed as $\zeta = (z-d)/L$ as explained in section 2. Panel (c) plots canopy aerodynamic resistance to heat transfer (r_H) at every level computed from eq. 1. As evident from panel (c), the aerodynamic resistance reduces with increasing instability, confirming the hypothesis constructed earlier and thus clearly demonstrating the canopy convective effect (CCE). As noted by Zilitinkevich et al. (2008), with increasing instability, “convective updraughts developing at side walls of roughness elements extend upwards and provide extra resistances to the mean flow. Then the mean flow interacts with both solid obstacles and their virtual extensions (updraughts), which results in the increased roughness length”. This increased roughness can be recognized as the aerodynamic roughness. For the same physical roughness of the canopy, increase of instability increases this aerodynamic roughness and in turn, reduces r_H . The low aerodynamic resistance effectively allows larger eddies to form above the forest canopy which are more efficient to dissipate the sensible heat by promoting buoyancy. This description refers to a more general phenomenon as opposed to the ~~the~~ description




Figure 3. Variations of r_H with height across stability ranges and comparisons with different parameterization schemes as described in table 1.

by Rotenberg and Yakir (2011) which identifies the higher physical roughness of the canopy compared to the desert and is thus a more site specific description. Nevertheless, it is acknowledged that the more generic description presented here can be reconciled with the explanation from Rotenberg and Yakir (2011) by noting that increased physical roughness can also result in increased aerodynamic roughness. Also incidentally, [Rotenberg and Yakir \(2010\) RY10](#) reported a value of $r_H \approx 16$ for the

5 Yatir forest which is of similar order of magnitude as what is found in panel (c) of figure 2. One important point to note in figure 1 is the magnitude of the Prandtl number, which is almost fixed to about 0.335 above the canopy. This can be reconciled with the theoretical prediction of the variation of P_{r0} with stability by Li et al. (2015). For stability ranges $1 \leq -\zeta \leq 10$, P_{r0} is also estimated to be approximately 0.33, consistent with the stability ranges plotted in figure 2. The variation of P_{r0} with stability is discussed further in the appendix.

10 4.2 Testing different parameterizations

It is interesting to study if the different parameterizations capture the correct behavior of r_H at different heights across stability. To compute r_H variations, the LES generated profiles of mean velocity u , sensible heat flux, air temperature, Prandtl number (thus the diffusivities) are used where all of them have z variations. The friction velocity u_* and the roughness lengths are




Figure 4. Variations of r_H with height for different stability classes computed for each parameterization scheme as described in table 1.




Figure 5. Variations of r_H as given by the parameterization of Yang et al. (2001) with height across stability ranges and a wide range of z_{0m} . black '+' markers indicate the observed r_H from LES at any particular stability state.

fixed. Figure 3 plots the variation of r_H with height as obtained from the LES (black '+' markers), and the predicted r_H from different parameterizations for the different stability cases - near neutral (column 1), weakly unstable (column 2), mildly unstable (column 3) and strongly unstable (column 4). The top row compares the parameterizations by Thom (1975) (blue line), Yang et al. (2001) (red line), Choudhury et al. (1986) (black line) and Viney (1991) (pink line) which assume $z_{0m} \neq z_{0h}$. The bottom panel compares the parameterizations by Verma et al. (1976) (blue dashed line), Hatfield et al. (1983) (red dashed line), Mahrt and Ek (1984) (black dashed line) and Xie (1988) (pink dashed line). It should be noted that a single value of roughness $z_{0m} = 0.6 h_c$ has been chosen for all cases by trial and error to obtain a 'good' comparison in figure 3. As observed, none of the parameterizations can capture the correct height variations of r_H except the one by Yang et al. (2001) for more unstable cases. However, all parameterizations seems to do a decent job close to the canopy top. This clearly indicates that one single value for z_{0m} as suggested by these parameterizations is inadequate. To study if the different parameterization schemes can capture the canopy convective effect, r_H computed from each method is plotted for different heights for the different stability classes in figure 4. The title of each panel describes which parameterization is plotted and the color shades starting from cyan to purple indicates increasing instability. As evident, only the parameterization by Thom (1975) captures the canopy convective effect for weaker eases instabilities. The parameterization by Yang et al. (2001) also displays the signatures of CCE, however weakly.

The other formulations cannot capture the correct trend of CCE at all. Thus at this stage, it is clear that Yang et al. (2001) formulation, based on MOST and distinguishing between two different roughness lengths is the most promising candidate to parametrize r_H compared to the other formulations which apply some form of approximations or do not apply MOST.

4.3 Towards an improved parameterization for r_H

- 5 Until this stage, the momentum roughness length has been prescribed by trial and error and it warrants a more detailed investigation. To explore the effect of different roughness lengths, the parameterization by Yang et al. (2001) is computed across a wide range of z_{0m} and compared with the LES outputs for the different stability classes. As observed, an increase of z_{0m} with increasing instability captures the height variation better than a single roughness length for all stability classes further providing support for the notion put forward by Zilitinkevich et al. (2008). Hence the formulation by Yang et al. (2001) can be
- 10 modified to include the effects of stratification on several parameters. Zilitinkevich et al. (2008) suggested a stability dependent zero-plane displacement length as well as a stability dependent z_{0m} based on dimensional analysis, given by

$$d_s = \frac{d}{\left[1 + 0.56 \left(\frac{h_c}{-L}\right)^{1/3}\right]}, \quad (11)$$

and

$$z_{0ms} = z_{0m} \left[1 + 1.15 \left(\frac{h_c}{-L}\right)^{1/3}\right], \quad (12)$$

- 15 where d_s and z_{0ms} are the stability dependent zero-plane displacement length and roughness lengths for momentum respectively, d and z_{0m} being their neutral counterpart. $d = (2/3)h_c$ can be assumed as usual. The neutral z_{0m} can be assumed to be related to LAI as given by Shuttleworth and Gurney (1990). According to the relation used by Shuttleworth and Gurney (1990), for an LAI of 5, a $z_{0m} = 0.12h_c$ can be obtained (which is almost constant for a wide range of canopy drag coefficients and LAI). Moreover, if one uses the correct stability dependent Prandtl number $P_{r0}(\zeta)$ instead of setting it to unity, an improved
- 20 parameterization based on the Yang et al. (2001) can be written as

$$r_H = \frac{P_{r0}(\zeta)}{\kappa^2 u} \left[\ln \left(\frac{z - d_s}{z_{0ms}} \right) - \psi_m(\zeta, \zeta_{0ms}) \right] \left[\ln \left(\frac{z - d_s}{z_{0hs}} \right) - \psi_h(\zeta, \zeta_{0hs}) \right]. \quad (13)$$

- Note that z_{0ms} and z_{0hs} are still related by the same relation $\kappa B^{-1} = \ln(z_{0ms}/z_{0hs})$ with $\kappa B^{-1} = 2.0$ as discussed earlier and $\zeta = (z - d_s)/L$ and $\zeta_{0ms} = z_{0ms}/L$, $\zeta_{0hs} = z_{0hs}/L$. If a Prandtl number of unity is still assumed but the roughness lengths are assumed to be varying with stability as given by equation 13 with a neutral value of $z_{0m} = 0.2h_c$, the formulation by Yang
- 25 et al. (2001) is found to display the correct behavior of canopy convective effect with stability as observed in figure 6. Panel (a) shows the variation of r_H with height across stability according to the improved formulation as given by equation 13. Panel (b) shows similar variations of r_H computed from the LES repeated again for comparison. The profile for the near neutral case crosses over the more highly unstable cases at heights around $6h_c$, however, the general behavior of CCE is captured well. On the other hand, if the full complexity of equation 13 is used including a stability dependent Prandtl number (discussed in




Figure 6. (a) The variation of r_H with height across stability according to the improved formulation as given by equation 13; (b) similar variations of r_H computed from the LES repeated again for comparison.

appendix A), but using the canopy top surface value of the sensible heat flux for all computations involved, the variation of modeled r_H is shown on panel (a) of figure 7. r_H computed from the LES results using the surface value of the heat flux is shown in panel (b). This assumption of a constant sensible heat flux in the canopy sub layer or the atmospheric surface layer is a more realistic one than a monotonically reducing sensible heat flux with height as shown in panel (e) of figure 1. In fact, the surface layer is defined as a constant flux layer (Stull, 2012). The reducing flux profiles in LES are common features of large-eddy simulations since the top boundary of the LES domain is assumed stress free (Shaw and Schumann, 1992). Figure 7 correctly captures the order of magnitude of the r_H observed from the simulations, and also captures CCE correctly. However, it is acknowledged that the exact profiles of the observed r_H can not be captured. However, these different comparisons highlight the uncertainties involved in the parameterization of r_H .

10 5 Conclusion

The canopy aerodynamic resistance is a concept borrowed from the evapotranspiration literature where it represents the resistance between the idealized ‘big-leaf’ (a reduced order representation of the fully heterogeneous three dimensional




Figure 7. (a) The variation of r_H with height across stability according to the improved formulation as modeled by equation 13, but using the top-of-canopy surface flux throughout all heights;(b) similar variations of r_H computed from the LES using the top-of-canopy surface flux assumed to be constant in the surface layer.

canopy) and the atmosphere for heat or vapor transfer (Alves et al., 1998). ~~The Penman-Monteith equation to calculate evapotranspiration requires parameterization of the aerodynamic resistance which require information on roughness lengths for heat and momentum and stability (Penman, 1948; Allen et al., 1998; Cleverly et al., 2013). r_H parameterizations are also used in global climate models to describe the canopy-atmosphere interaction at the canopy surface layer (Walko et al., 2000).~~

- 5 In semi arid ecosystems, vegetation canopies maintain a relatively cool surface temperature in spite of the high sensible heat flux by reducing the canopy aerodynamic resistance to heat transfer (r_H) — a phenomenon named ‘canopy convective effect’ by Rotenberg and Yakir (2010). In the present work, a large-eddy simulation is used to examine this canopy convective effect and in the process, several existing parameterizations for r_H is examined. The objectives behind this exploration is twofold. The first one is to investigate if the existing parameterizations exhibit canopy convective effect and the second one is to identify the uncertainties associated with these different parameterizations since they are applied in different climate models often
- 10 under conditions of thermal stratification. As illustrated by the LES results, r_H above the canopy are found to reduce systematically as the strength of unstable stratification increases. This is deemed to be the core feature of canopy convective effect, since with increasing instability, more convective updraughts enhance the roughness over the canopy elements that the mean

flow encounters. The height variation of r_H is also found to have a highly nonlinear profile, thus any model prescribing a parameterization for r_H needs to employ considerable caution regarding the height it is prescribed. Existing parameterizations of r_H employ either Monin-Obukhov similarity theory (MOST) or Richardson number based empirical or semi-empirical formulations to account for thermal stratification. However, most of them are found to be unable to describe the correct trend of CCE. Among different formulations, the one by Yang et al. (2001) is found to be the most promising candidate. This parameterization employs MOST, and accounts for stability parameters associated with roughness lengths for momentum and heat transfer. It is found out that a stability dependent zero-plane displacement height as well as stability dependent roughness lengths for momentum and heat transfer can improve its performance. Moreover, if the surface layer or the canopy sublayer is assumed to have a constant sensible heat flux equal to the flux at the canopy top, and a stability dependent Prandtl number is used, the performance improves further. These ~~assumption also leads~~ assumptions also lead to a less nonlinear height variation. These explorations highlight the uncertainties associated with the parameterizations of r_H . One possible major source of uncertainty is the usage of Monin-Obukhov similarity theory in the canopy sublayer (CSL) (up to $3h_c$ to $6h_c$) since it is not expected to perform in the CSL (Kaimal and Finnigan, 1994). Nevertheless, MOST formulations are found to outperform other semi-empirical formulations using Richardson numbers. Thus future research work will involve studying these uncertainties of r_H parameterizations in regional and global climate models. The consequence of this CCE on local circulation, atmospheric moisture and tree physiology will also be investigated, extending the preliminary study of Eder et al. (2015). However, the fact that CCE is a more generic feature of canopy turbulence provides hope that also the afforestation of an area larger than the Yatir forest would be able to cope with a high-radiation load under water scarcity in semi-arid climates.

Acknowledgements. This research was supported by the German Research Foundation (DFG) as part of the project “Climate feedbacks and benefits of semi-arid forests (CliFF)” and the project “Capturing all relevant scales of biosphere-atmosphere exchange - the enigmatic energy balance closure problem”, which is funded by the Helmholtz-Association through the President’s Initiative and Networking Fund, and by KIT. The authors thank the PALM group at Leibniz University Hannover for their open-source PALM code and also thank Dan Yakir and Eyal Rotenberg at the Weizmann Institute of Science, Israel, for their support during the CliFF campaign. We ~~also~~ thank Prof. Thomas Foken, Bayreuth center of Ecology and environmental Research (BayCEER), University of Bayreuth, Germany, for his comments and suggestions. We also thank Prof. Gil Bohrer, (Ohio State University) and the other anonymous reviewer for their constructive suggestions to improve the manuscript during the discussion stage.

Appendix A: Appendix: Stability dependence of Prandtl number

The turbulent Prandtl number P_{r0} is defined as the ratio of the eddy diffusivities of momentum and heat (K_m/K_h). The variation of Prandtl number with stability ($P_{r0}(\zeta)$) was discussed in detail by Li et al. (2015) by using a spectral budget formulation and not repeated here. Only the predicted variation of Pr^{-1}/Pr_n^{-1} with stability ($\zeta = z/L$) is digitized and produced on figure A1, which was experimentally validated by Li et al. (2015). Pr_n^{-1} denotes the inverse of the neutral Prandtl




Figure A1. The variation of Pr^{-1}/Pr_n^{-1} with stability according to the spectral budget formulation of Li et al. (2015).

number which can assumed to be equal to 1. Note that for the stability ranges computed in the LES simulations in figure 2, this formulation predicts a $Pr_0 \approx 0.33$, which is also observed in the Pr_0 independently computed in figure 1.

Appendix B: [Some computational details of the of the LES](#)

B1 [surface heat flux formulation and boundary conditions](#)

5 [The ground surface heat flux for gridpoints with a canopy layer is given by:](#)

$$\overline{w'T'_s} = \overline{w'T'_{toc}} \times \exp\left(-\epsilon_c \int_0^{h_c} LAD(z) dz\right), \quad (\text{B1})$$

with $\epsilon_c = 0.6$ the extinction coefficient of light within the canopy. Within the canopy the plant-canopy heating rate is calculated as the vertical divergence of the canopy heat fluxes:

$$\overline{w'T'}_{toc} \frac{d}{dz} \exp \left(\epsilon_c \int_z^{h_c} LAD(z') dz' \right) \quad (\text{B2})$$

5 The bottom boundary condition for potential temperature is a Neumann condition, the boundary condition at the top of the domain is such that the initial temperature gradient is maintained at the top of the domain.

B2 Eddy diffusivity formulation

The computation for the eddy diffusivities in PALM follows the standard procedure for 1.5 order turbulence closure. Thus they are computed from the subgrid-scale turbulent kinetic energy, more precisely equations (13-14) from Maronga et al. (2015).

Eddy diffusivity for momentum:

$$10 \quad K_m = c_m l \sqrt{e} \quad (\text{B3})$$

Eddy diffusivity for heat:

$$K_h = \left(1 + \frac{2l}{\Delta} \right) K_m \quad (\text{B4})$$

15 With e the subgrid-scale turbulent kinetic energy (a prognostic variable), $c_m = 0.1$, and Δ the geometric mean of the grid spacings in x, y and z . Finally, l is the subgrid-scale mixing length depending on Δ , stability, and distance from the topography elements or ground surface.

Author contributions. T. Banerjee conceived the idea, conducted data analysis and wrote the paper. F. De Roo set up the Large Eddy Simulations. M. Mauder had written the proposal that funded this project, supervised the project and provided comments and suggestions.

Competing interests. The authors declare no conflict of interest.

References

- Allen, R. G., Pereira, L. S., Raes, D., and Smith, M.: Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56, FAO, Rome: Food and Agriculture Organization of the United Nations, 300, D05 109, 1998.
- Alves, I., Perrier, A., and Pereira, L.: Aerodynamic and surface resistances of complete cover crops: How good is the “big leaf”?, *Transact. ASAE*, 41, 345, 1998.
- 5 Baldocchi, D., Falge, E., Gu, L., Olson, R., Hollinger, D., Running, S., Anthoni, P., Bernhofer, C., Davis, K., Evans, R., et al.: FLUXNET: a new tool to study the temporal and spatial variability of ecosystem-scale carbon dioxide, water vapor, and energy flux densities, *Bull. Am. Met. Soc.*, 82, 2415–2434, 2001.
- Banerjee, T., Katul, G., Fontan, S., Poggi, D., and Kumar, M.: Mean flow near edges and within cavities situated inside dense canopies, *Bound.-Lay. Meteorol.*, 149, 19–41, 2013.
- 10 Chamberlain, A.: Transport of gases to and from surfaces with bluff and wave-like roughness elements, *Q. J. Roy. Meteor. Soc.*, 94, 318–332, 1968.
- Choudhury, B., Reginato, R., and Idso, S.: An analysis of infrared temperature observations over wheat and calculation of latent heat flux, *Agr. Forest Meteorol.*, 37, 75–88, 1986.
- 15 Cleverly, J., Chen, C., Boulain, N., Villalobos-Vega, R., Faux, R., Grant, N., Yu, Q., and Eamus, D.: Aerodynamic resistance and Penman-Monteith evapotranspiration over a seasonally two-layered canopy in semiarid central Australia, *J. Hydrometeorol.*, 14, 1562–1570, 2013.
- Deardorff, J. W.: Stratocumulus-capped mixed layers derived from a three-dimensional model, *Bound.-Lay. Meteorol.*, 18, 495–527, 1980.
- Dias-Junior, C. Q., Marques Filho, E. P., and Sá, L. D.: A large eddy simulation model applied to analyze the turbulent flow above Amazon forest, *J. Wind Eng. Ind. Aerodyn.*, 147, 143–153, 2015.
- 20 Dyer, A.: A review of flux-profile relationships, *Boundary-Layer Meteorology*, 7, 363–372, 1974.
- Dyer, A. and Hicks, B.: Flux-gradient relationships in the constant flux layer, *Quarterly Journal of the Royal Meteorological Society*, 96, 715–721, 1970.
- Eder, F., De Roo, F., Rotenberg, E., Yakir, D., Schmid, H. P., and Mauder, M.: Secondary circulations at a solitary forest surrounded by semi-arid shrubland and their impact on eddy-covariance measurements, *Agr. Forest Meteorol.*, 211, 115–127, 2015.
- 25 Foken, T.: 50 years of the Monin-Obukhov similarity theory, *Bound.-Lay. Meteorol.*, 119, 431–447, 2006.
- Foken, T., Dlugi, R., and Kramm, G.: On the determination of dry deposition and emission of gaseous compounds at the biosphere-atmosphere interface, *Meteorol. Z.*, 4, 91–118, 1995.
- Garratt, J. R.: Aerodynamic roughness and mean monthly surface stress over Australia, 29, Commonwealth Scientific and Industrial Research Organization, 1977.
- 30 Harman, I. N. and Finnigan, J. J.: A simple unified theory for flow in the canopy and roughness sublayer, *Boundary-layer meteorology*, 123, 339–363, 2007.
- Hatfield, J., Perrier, A., and Jackson, R.: Estimation of evapotranspiration at one time-of-day using remotely sensed surface temperatures, *Agr. Water. Manage.*, 7, 341–350, 1983.
- Kaimal, J. C. and Finnigan, J. J.: Atmospheric boundary layer flows: their structure and measurement, Oxford University Press, 1994.
- 35 Katul, G. G., Mahrt, L., Poggi, D., and Sanz, C.: One-and two-equation models for canopy turbulence, *Bound.-Lay. Meteorol.*, 113, 81–109, 2004.

- Li, D., Katul, G. G., and Zilitinkevich, S. S.: Revisiting the turbulent Prandtl number in an idealized atmospheric surface layer, *J. Atmos. Sci.*, 72, 2394–2410, 2015.
- Liu, S., Lu, L., Mao, D., and Jia, L.: Evaluating parameterizations of aerodynamic resistance to heat transfer using field measurements, *Hydrol. Earth Syst. Sci.*, 11, 769–783, 2007.
- 5 Mahrt, L. and Ek, M.: The influence of atmospheric stability on potential evaporation, *J. Clim. Appl. Meteorol.*, 23, 222–234, 1984.
- Markkanen, T., Rannik, Ü., Marcolla, B., Cescatti, A., and Vesala, T.: Footprints and fetches for fluxes over forest canopies with varying structure and density, *Bound.-Lay. Meteorol.*, 106, 437–459, 2003.
- Maronga, B., Gryschka, M., Heinze, R., Hoffmann, F., Kanani-Sühring, F., Keck, M., Ketelsen, K., Letzel, M., Sühring, M., and Raasch, S.: The Parallelized Large-Eddy Simulation Model (PALM) version 4.0 for atmospheric and oceanic flows: model formulation, recent
10 developments, and future perspectives, *Geosci. Model Dev.*, 8, 1539–1637, 2015.
- Maurer, K., Bohrer, G., Kenny, W., and Ivanov, V.: Large-eddy simulations of surface roughness parameter sensitivity to canopy-structure characteristics, *Biogeosciences*, 12, 2533–2548, 2015.
- Maurer, K. D., Hardiman, B. S., Vogel, C. S., and Bohrer, G.: Canopy-structure effects on surface roughness parameters: Observations in a Great Lakes mixed-deciduous forest, *Agricultural and forest meteorology*, 177, 24–34, 2013.
- 15 Moeng, C.-H. and Wyngaard, J. C.: Spectral analysis of large-eddy simulations of the convective boundary layer, *J. Atmos. Sci.*, 45, 3573–3587, 1988.
- Monteith, J.: *Principles of environmental physics*, Academic Press, London, England, 1973.
- Monteith, J. and Unsworth, M.: *Principles of Environmental Physics*, Academic Press, London, England, 2007.
- Owen, P. and Thomson, W.: Heat transfer across rough surfaces, *J. Fluid Mech.*, 15, 321–334, 1963.
- 20 Patton, E. G., Sullivan, P. P., Shaw, R. H., Finnigan, J. J., and Weil, J. C.: Atmospheric stability influences on coupled boundary-layer-canopy turbulence, *J. Atmos. Sci.*, 73, 1621–1648, 2015.
- Paulson, C. A.: The mathematical representation of wind speed and temperature profiles in the unstable atmospheric surface layer, *J. Appl. Meteorol.*, 9, 857–861, 1970.
- Penman, H. L.: Natural evaporation from open water, bare soil and grass, in: *Proc. Roy. Soc. A - Math. Phys.*, vol. 193, pp. 120–145, The
25 Royal Society, 1948.
- Raasch, S. and Schröter, M.: PALM—a large-eddy simulation model performing on massively parallel computers, *Meteorol. Z.*, 10, 363–372, 2001.
- Rotenberg, E. and Yakir, D.: Contribution of semi-arid forests to the climate system, *Science*, 327, 451–454, 2010.
- Rotenberg, E. and Yakir, D.: Distinct patterns of changes in surface energy budget associated with forestation in the semiarid region, *Glob. Change Bio.*, 17, 1536–1548, 2011.
- 30 Saiki, E. M., Moeng, C.-H., and Sullivan, P. P.: Large-eddy simulation of the stably stratified planetary boundary layer, *Bound.-Lay. Meteorol.*, 95, 1–30, 2000.
- Seginer, I.: Aerodynamic roughness of vegetated surfaces, *Bound.-Lay. Meteorol.*, 5, 383–393, 1974.
- Shaw, R. H. and Schumann, U.: Large-eddy simulation of turbulent flow above and within a forest, *Bound.-Lay. Meteorol.*, 61, 47–64, 1992.
- 35 Shuttleworth, W. J. and Gurney, R. J.: The theoretical relationship between foliage temperature and canopy resistance in sparse crops, *Q. J. Roy. Meteorol. Soc.*, 116, 497–519, 1990.
- Smagorinsky, J.: General circulation experiments with the primitive equations: I. the basic experiment, *Mon. Weather Rev.*, 91, 99–164, 1963.

- Stull, R. B.: An introduction to boundary layer meteorology, vol. 13, Springer Science & Business Media, 2012.
- Thom, A.: Momentum, mass and heat exchange of plant communities, vol. 1, Academic Press, London, 1975.
- Verma, S., Rosenberg, N., Blad, B., and Baradas, M.: Resistance-energy balance method for predicting evapotranspiration: Determination of boundary layer resistance and evaluation of error effects, *Agron. J.*, 68, 776–782, 1976.
- 5 Viney, N. R.: An empirical expression for aerodynamic resistance in the unstable boundary layer, *Bound.-Lay. Meteorol.*, 56, 381–393, 1991.
- Walko, R. L., Band, L. E., Baron, J., Kittel, T. G., Lammers, R., Lee, T. J., Ojima, D., Pielke Sr, R. A., Taylor, C., Tague, C., et al.: Coupled atmosphere-biophysics-hydrology models for environmental modeling, *J. Appl. Meteorol.*, 39, 931–944, 2000.
- Watanabe, T.: Large-eddy simulation of coherent turbulence structures associated with scalar ramps over plant canopies, *Bound.-Lay. Meteorol.*, 112, 307–341, 2004.
- 10 Webb, E.: On the correction of flux measurements for effects of heat and water vapour transfer, *Boundary-Layer Meteorology*, 23, 251–254, 1982.
- Xie, X.: An improved energy balance-aerodynamic resistance model used estimation of evapotranspiration on the wheat field, *Acta Meteorol. Sin.* (in Chinese), 46, 102–106, 1988.
- Yang, K., Tamai, N., and Koike, T.: Analytical solution of surface layer similarity equations, *J. Appl. Meteorol.*, 40, 1647–1653, 2001.
- 15 Zilitinkevich, S. S., Mammarella, I., Baklanov, A. A., and Joffre, S. M.: The effect of stratification on the aerodynamic roughness length and displacement height, *Bound.-Lay. Meteorol.*, 129, 179–190, 2008.