

The benefit of seamless forecasts for hydrological predictions over Europe

Fredrik Wetterhall¹ and Francesca Di Giuseppe¹

¹European Centre for Medium-range Weather Forecasts, Shinfield Park, Reading ,UK

Correspondence to: Fredrik wetterhall: fredrik.wetterhall@ecmwf.int

Abstract. Two different systems provide long range forecasts at ECMWF. On the sub-seasonal time scale, ECMWF issues an extended-range ensemble prediction system (ENS-ER) which runs a 46-day forecast integration issued twice weekly. On longer time scales the current seasonal forecasting system (SYS4) produces a 7-month outlook starting from the first of each month. SYS4 uses an older model version and has lower spatial and temporal resolution than ENS-ER, which is issued with the current operational ensemble forecasting system. Given the substantial differences between the ENS-ER and the SYS4 configurations and the difficulties of creating a seamless integration, applications that rely on weather forcing as input such as the European Flood Awareness System (EFAS) often follow the route of the creation of two separate systems for different forecast horizons. This study evaluates the benefit of a seamless integration of the two systems for hydrological applications and shows that the benefit of the new seamless system when compared to the seasonal forecast can be attributed to (1) the use of a more recent model version in the sub-seasonal range (first 46 days) and (2) the much more frequent updates of the meteorological forecast.

1 Introduction

ECMWF produces a range of forecasts, among them a 10 day deterministic high resolution forecast (HRES) and a lower resolution 15-day 51 member ensemble prediction system (ENS) that is extended to 46 days twice weekly (Mondays and Thursdays at 00UTC; Vitart et al. 2008). In this paper we refer to the extended ENS as ENS-ER. On longer time ranges ECMWF issues a seasonal ensemble forecast system (SYS4), operational since November 2011. SYS4 issues a 7-month prediction (extended to 13 months four times a year) once every month (Molteni et al., 2011). The ENS-ER forecast system benefits from frequent updates of the model physics and data assimilation system

(Vitart et al., 2008). ECMWF releases official model updates on average 2-3 times a year which typically include new improved schemes for physical processes, better use of observations and their assimilation and sometimes increase in model resolution. The seasonal forecast has a lower resolution, is an older model version than ENS-ER and is also updated much less frequently. This implies that the skill of the seasonal forecasting system is lower relative to ENS-ER in the overlapping first six weeks.

Applications that use numerical weather predictions as forcing, such as the operational European Flood Awareness System (EFAS; Thielen et al. 2009; Bartholmes et al. 2009; Smith et al. 2016), are often designed for a specific purpose. EFAS has since the start focused on early warning of floods in the medium-range forecast horizon, typically up to 15 days. Recently, a seasonal hydrological outlook forced by SYS4 was implemented operationally with a lead-time of 7 months (Arnal et al., 2017).

This extension to the monthly and seasonal scales is potentially very useful in order to; (i) produce products which extend the previous forecast horizon; (ii) benefit from hindcasts for pre- and post-processing to produce output of higher quality (e.g. model based return periods); and (iii) design completely new early warning frameworks complementing the existing ones. The extended lead time provided by running EFAS forced by weather prediction across different time scales could potentially provide added benefit in terms of very early planning, for example for agriculture, energy 40 Bazile et al. (2017) and transport sectors Meißner et al. (2017) as well as water resources management Sene et al. (2018). Such a forecast system would be a first step to close the identified gap between hydrological forecasts on the medium (up to 15 days) and seasonal range (White et al., 2017). These extended range systems may not be able to capture extremes of short-lived events like floods, but they are able to detect anomalous conditions on longer lead times, such as low flows 45 (Meißner et al., 2017) and droughts (Dutra et al., 2014).

The concept of seamless forecasts was first introduced by Palmer and Webster (1993). Palmer et al. (2008) formally expanded the idea showing how short-lived phenomena under certain conditions may persist and increase predictability at longer time scales. Since then the concept of a unified or seamless framework for weather and climate prediction has been vastly debated (Hurrell et al., 50 2009; Brunet et al., 2010). However as noticed by Hoskins (2013) in his seminal paper, while "the atmosphere knows no barriers in time-scales", often model implementation is segmented for practical reasons. Still, major efforts have been made to create unified systems. Indeed, the ENS-ER was the first attempt to create a seamless extension of the ECMWF medium-range forecast to the monthly scales(Vitart et al., 2008). Similarly, the UK Met Office has in the past twenty-five years worked to 55 create a unified model that could work across all time scales (Brown et al., 2012). Also the climate community has moved in the same direction. For example, the EC-Earth project shows that a bridge can be made between weather, seasonal forecasting and beyond (Hazeleger et al., 2010, 2012).

The latter projects went all the way to create new systems starting from existing components and were therefore costly and time demanding. In contrast, a practical and simpler approach could be 60 taken. The seamless idea could be translated into the simple concatenation of "the best" forecast at each lead-time. The clear advantage of this off-the shelf seamless prediction conversion is that it utilizes products that are already in place, thereby avoiding the complications of new developments while generating forecast products to meet different types of users (Pappenberger et al., 2013). There is however an underlying complexity in this simplification; the substantial difference in design be- 65 tween the various forecasting systems makes the concatenation technically difficult. As systems are designed for different users they often have non-matching temporal and spatial resolutions, different hindcast span and different ensemble sizes. One important consequence of this difference in design is that, for example, the much more frequent updates to the extended range compared to the seasonal system at ECMWF, implies that the bias characteristics of the two systems diverge over time, only 70 closing this gap when the seasonal system is updated (Di Giuseppe et al., 2013). Then model outputs either need to be bias-corrected to be useful forcing to drive sectoral models such as EFAS, or that final products should be provided in terms of anomalies calculated against the model climate, taken into consideration the bias of the seamless forecast system. In both cases the seamless system needs to account for the use of the hindcast dataset and the application of some bias correction algorithm.

75 In return, the advantage is in the gain in skill and the extension of the lead-time.

In this work the benefit of a seamless hydro-meteorological system was tested for a span of time ranges from 1 week to 6 months for stream flow forecasts over the European domain using the EFAS system. The aim was to test whether integrating medium-range forecasts with seasonal prediction contributes to enhance hydrological predictability on the seasonal scale. Specifically, the questions 80 addressed were: What is the gain in terms of hydrological forecasting of using a more recent model version in the first 46 days provided by the use of the ENS-ER? What is the skill gain provided by having more frequent forecast updates?

2 Methods

2.1 Hydrological model system

85 The hydrometeorological system used in this study was the European Flood Awareness System (EFAS Thielen et al. 2009; Bartholmes et al. 2009; Smith et al. 2016). EFAS is an operational early warning system covering most of the European domain and has been run operationally since October 2012 as part of the COPERNICUS Emergency Management Service (CEMS). The hydrological component of EFAS is the distributed rainfall-runoff model LISFLOOD (De Roo et al., 2000; Van 90 Der Knijff et al., 2010; Burek et al., 2013). LISFLOOD calculates the main hydrological processes on sub-daily and daily time-scales that generate runoff, such as soil and ground water interactions, for each grid cell. In the operational setup EFAS covers most of Europe on a 5x5 km equal-area

grid. The runoff is transformed through a routing scheme to estimate the river discharge at each grid cell along the river network. The routing scheme also takes into account water retention in lakes and reservoirs. This study will concentrate on the forecast of river discharge at the outlets of the sub basins of the river network that were used for calibration of the current EFAS system (Smith et al., 2016; Zajac and Bianchi., 2013). The total number of outlets used were 679, and they represent river basins of all sizes and characteristics across the EFAS domain.

In its operational implementation the latest calibration (referred to as tuning in the NWP nomenclature) of LISFLOOD used an observational dataset of meteorological forcing data (precipitation and temperature) and observed discharge covering the model domain over the period 1990-2013 (Smith et al., 2016; Zajac and Bianchi., 2013). The meteorological dataset comprises more than 5000 synoptic stations that have been interpolated to a 5x5 km Lambert azimuthal equal-area projection (Ntegeka et al., 2013). The high resolution gridded observation of precipitation and temperature were used for the calibration of LISFLOOD. The observational meteorological dataset was also used to generate a reference modeled climatology of discharge (hereafter called water balance, WB) which is used as; (i) initial conditions for the operational forecast and hindcasts and (ii) reference model run to assess the performance of the forecasts. Using the WB run as proxy observation simplifies the interpretation of the skill scores as it avoids the complication of having to assess the bias against observed discharge. The purpose of this paper is rather to assess the skill of the two forecasts used for forecasts rather than the total skill of the forecasting system.

2.2 Seamless integration of meteorological forcing data

Every Monday and Thursday ECMWF issues an extended-range ensemble forecast (ENS-ER) by continuing the integration time beyond day 15 up to day 46, with a lower-resolution model (Figure 1, Table 1). Each ENS-ER integration comes with an 11-member hindcast set produced for the same dates over the previous 20 years. This hindcast set provides identical integrations as the current operational forecast with the difference that ERA-Interim reanalysis (ERA1; Dee et al. (2011)) and ERAI land reanalysis (Balsamo et al., 2015) are used to provide the initial conditions, whereas the operational ensemble forecast uses the operational analysis. The hindcast data together with observations can be used in many applications, for example to calibrate the forecast in an operational setting (Di Giuseppe et al., 2013).

The operational seasonal forecast (SYS4) issues a new forecast at the beginning of each month with a lead-time up to 7 months, four times a year extended to 13 months (Figure 2). SYS4 has a hindcast consisting of 30 years started at each month and consisting of 15 members. The new seamless forecasting system (hereafter called SEAM) was created by concatenating each ENS-ER ensemble member with a randomly selected SYS4 ensemble member at day 46, which is the last day of the ENS-ER (Figure 2). SEAM benefits from the frequent updates of the ENS-ER and has the seven months horizon of the seasonal system.

Since the two systems have different resolutions (table 1) the horizontal resolution was homogenized to the 5x5 km equal-area grid through a mass-conserving interpolation for precipitation and a bilinear for temperature before it was used as input to the hydrological model in EFAS. The time step was reduced to daily by averaging (accumulating for precipitation and evapotranspiration) the three hourly outputs of the ENS-ER and the six hourly outputs of SYS-4. Since the ENS-ER has a reduced hindcast (20 years) and number of members (11), SEAM has the same number of members and hindcast period. Note that in real-time mode, a full 51-member SEAM is possible. The technical details of the forecast and the hindcast used in this experiment are presented in table 1. For simplicity SYS4 and SEAM will from now on refer to the full hydro-meteorological integrations for the remainder of this paper.

2.3 Experimental set-up

This study focuses on the performance of SYS4 and SEAM over the hindcasts of the operational forecast. The hindcasts starting from 2015-05-14 (the first available date with 11-member hindcast for ENS-ER) to 2016-06-02 were used as input to the full EFAS modeling chain. This provided 13 monthly starting dates for SYS4 and 111 biweekly starting dates for SEAM with a corresponding hindcast set covering all seasons over the previous 20-year period, each with 15 and 11 ensemble members respectively (Fig. 1). The output was averaged to weekly means before the skill score analysis. Since the starting dates of the SEAM and SYS4 were not always in sync (the starting date of the SYS4 integrations are only sometimes on a Monday or Wednesday), it is impossible to do a completely like-for like comparison since the validation periods would be slightly different. However, this error will be random and given the sample size (260 and 2220) it was not considered to have a big impact on the results.

SEAM was validated against the runs with SYS4 to assess the added value of the merged forecast. Further, both model systems were compared against a climatological benchmark simulation (hereafter called CLIM). CLIM was constructed by forcing the LISFLOOD with 15 randomly selected time series of observed meteorological forcing from the period 1990-2014, excluding the modeled year. CLIM has the advantage of having the same initial conditions as the SYS4 and SEAM hindcasts, but has no expected predictive skill beyond the horizon of the initial conditions. The advantage of CLIM is that in theory it has near perfect reliability with regards to the WB runs since it is produced with the same unbiased forcing data. It should therefore score better or equal as the hindcasts as predictor on time ranges beyond their respective limits of predictability.

2.4 Score metrics

The performance of the two forecast systems was compared against modeled discharge using observations at the 679 sub basin outlets using deterministic and probabilistic scores. We refer to this run as observations in the paper. The scores used were the continuous ranked probability score (CRPS;

Hersbach 2000), mean relative error (MRE) and forecast reliability. CRPS is a common tool to validate probabilistic forecasts and can be seen as generalization of the mean absolute error to the probabilistic realm of ensemble forecasts. It is defined as:

$$CRPS = \frac{1}{N} \sum_{t=1}^N \int_{-\infty}^{+\infty} [F_t(x(n)) - H_t(x(n) - x_0)^2] dx \quad (1)$$

where $x(n)$ is the forecast at time step t of N number of forecasts and x_0 is the observed value. The CRPS is the continuous extension of RPS where $F(x)$ is the cumulative distribution function (CDF) $F(x) = p(X - x)$ and $H(x - x_0)$ is the Heaviside function, which has the value 0 when $x - x_0 < 0$ and 1 otherwise.

The CRPS compares the cumulative probability distribution of the discharge forecasted by the ensemble forecast system to an observation. It is sensitive to the mean forecast biases as well as the spread of the ensemble. Since the SEAM has 11 members and SYS4 and CLIM has 15 members in the hindcast, the CRPS are not directly comparable. Ferro et al. (2008) showed that for two ensemble distributions with different ensemble sizes, M and m , the unbiased estimate for $CRPS_M$ based on CRPS calculated from the ensemble size m is:

$$CRPS_M = CRPS_m - \frac{M - m}{2Mmn} \sum_{t=1}^n \Delta_t \quad (2)$$

where

$$180 \quad \Delta_t = \frac{1}{m(m-1)} \sum_{i \neq j} |X_{t,i} - X_{t,j}| \quad (3)$$

is Gini's mean difference of ensemble members $[X_{t,1}, \dots, X_{t,m}]$ at time t . From the CRPS a skill score (CRPSS) can be derived by comparing CRPS of the verified forecast against a reference forecast.

$$SS_{CRPS} = 1 - \frac{CRPS_{fc}}{CRPS_{rf}} \quad (4)$$

185 Mean relative error was measured as the forecast error against observations normalised against observations. The reliability was assessed through a reliability diagram, where the forecast probability of exceeding the percentiles of its climatology is compared against the observed frequencies Weisheimer and Palmer (2014). All scores were calculated for SYS4, SEAM and CLIM over the hindcast period.

190 **3 Results and discussion**

3.1 Overall forecast skill

The forecast skill gain provided by SEAM with respect to SYS4 is mostly concentrated to the first six weeks (Figure 3,a) when the forcing data are from the ENS-ER. The difference in CRPSS is 0.6 at week one, which then decreases to 0.2 by week six. All river points show a gain in skill up 195 until week three, then some points show a benefit of using the SYS4 instead of SEAM. However, in some catchments there is skill up further than eight weeks. The overall better performance of SEAM with respect to SYS4 is partly because of the use of a more recent model version and partly because of the more frequent update of the atmospheric and hydrological initial conditions. It is possible to disentangle the relative contributions between these two factors by only considering a 200 reduced number of starting dates for the SEAM forecast; i.e dates that are the closest to the SYS4 starting dates (figure 3,b). This reduced statistic provides a measure of the expected contribution of *only* employing a newer model cycle in the first weeks while both simulations benefits from the same hydrological initialization. In this case the skill gain in CPRS reduces to between 0 and 0.4 (median 0.2) against SYS4 for the first week, reducing to neutral around week four. Therefore the 205 most relevant gain comes from the more frequent initializations of the hydrological model.

To put these increments into context we also look at the improvement in skill of the two system SYS4 and SEAM against the CLIM benchmark forecast (Figure 3c-d). The gain from having an improved initial conditions in SEAM is similar in comparison with CLIM (Figure 3c) as with SYS4 (Figure 3a) in the first week, but the skill deteriorates quicker and the median CRPSS is negative after 210 5 weeks. Without the increase in skill due to the advantage in the better initial conditions, SEAM still shows a gain against the CLIM forecast with a CRPSS of 0.4 for the first week, although the spread is quite large (Figure 3d). Also SYS4 shows an increase of skill against the CLIM forecast. Both forecasts are less skillful than CLIM for most river points after week four. It can also be noted that SEAM has a higher spread than SYS4 on longer lead times even though they are forced 215 with the same data from day 47 and onwards. An explanation can be that the ensembles from the two meteorological forecasts are not matched in terms of their relative attributes with regards to their ensemble mean. If two extreme driving forecasts from the two meteorological forecasts are combined it can lead to members that are further away from the ensemble mean than when only one driving forecast is used.

220 **3.2 Geographical variation of forecast skill**

The geographical distribution of skill gain provided by the SEAM and SYS4 prediction reveals a coherent picture with good scores against the CLIM run over most of Europe (Figure 4 a-b). The gain in the figure is expressed as a difference in the number of weeks into the forecast needed for the CRPSS to drop below zero (i.e. there is no skill in the forecast in comparison with CLIM), which

225 gives an indication of the expected time gain in terms of information provided by the forecast against
the reference forecast. Both SYS4 and SEAM are better than CLIM, and SEAM has higher skill
than SYS4 for most of Europe. There is a small negative affect over the Alps, southeastern Europe
and northern Finland (Figure 4d). The performance of the operational EFAS in these regions is
generally poor, which is caused by the difficulty of having good observations of precipitation in high
230 altitude stations and the atmospheric models difficulty in resolving steep orography (Alfieri et al.,
2014).“The snow accumulation and snowmelt are further divided into three elevation zones within
a grip in LISFLOOD to better account for orographic effects in mountainous regions. However, this
increase in sub grid resolution is not likely to be high enough to capture the snow variability during
the snow accumulation and snowmelt in mountainous regions. Further, precipitation forecasts have
235 documented biases in steep orography ((Haiden et al., 2014)).

Another interesting aspect to showcase is the relevance of more frequent model version updates
is the overall improvement on river discharge for all stations in proximity to the western coasts. This
can be attributed to recent developments of the precipitation forecasts, for example a new diagnostic
closure introduced in the convection scheme (Bechtold et al., 2014) and a new parameterization of
240 precipitation formation (Haiden et al., 2014).

3.3 Bias and reliability

The relatively sharp decline in CRPSS can to some extent be explained by the negative bias (too
wet forecast) for both SEAM and SYS4 forecast (Figure 5). SEAM has lower bias than SYS4, also
when the analysis is confined to the first few weeks (Figure 5b). The slightly better bias in SEAM
245 disappears quickly after the merge (week 7). The bias of the forecast is not spatially consistent, it is
generally larger west- and mid-Europe (Figure 6). The figure shows the bias for SEAM (a-c) but the
pattern is similar for SYS4. SEAM has generally a smaller bias than SYS4 (Figure 6d). SYS4 has
lower bias south of the alps, where it is also performs better than SEAM.

Reliability of a forecast is terms of its usefulness for decision making. A reliable forecast can be
250 trusted to predict the correct probability of certain events, regardless of the accuracy. An unreliable
forecast is in practice of no use and can lead to poor decisions Weisheimer and Palmer (2014). Both
forecast systems are over-confident (Figure 7, which is similar to a previous study of 2 m temperature
and precipitation over Europe with SYS4 Weisheimer and Palmer (2014). The skill of the forecasts
from any of the system could be potentially higher by performing a bias correction either of the
255 atmospheric input and/or of the discharge. However in this paper we concentrate on the difference
in skills provided by the various configurations and no bias correction has been applied.

3.4 Added value of the seamless forecast

Even though the increase in the overall skill provided by the SEAM in comparison with SYS4 is
noticeable, the justification for its use in an operational context also depends on the actionable time

260 gain in a response situation. More frequent forecast updates could potentially be useful in decision making. As an example we analyze the predicted stream flow for the Rhine river at a station just upstreams Cologne, Germany, during the European heat wave in the summer of 2003. It was an exceptional meteorological event which combined significant precipitation deficits with record-setting high temperatures (García-Herrera et al., 2010). At its peak in August, extremely low discharge levels of rivers were reported in large parts of Europe. Economic losses were huge in many primary economic sectors including transportation (Ciais et al., 2005). For several months inland navigation was heavily impaired and the major European transport routes in the Danube and Rhine basins ceased completely (Jonkeren et al., 2008).

265
270 Despite the fact that 2003 conditions were extreme from the meteorological point of view, the upcoming deficit in precipitation and the high temperatures were well predicted by the ECMWF seasonal systems operational at that time (System-3; Weisheimer et al. 2011). The good predictability of the event is confirmed by the low discharge prediction provided by SYS4 at the Rhein upstreams of Cologne (figure 8). More than 30 % of the ensemble members forecast extreme low-flow conditions. In fact the observed discharge confirms that the river flow on two separate occasions, 275 event one on August 17-27 and event two September 18-28 2003, went below the 3% percentile of its climatological value for the season (figure 8). While most of SYS4 ensemble members mark the extreme condition three to four weeks ahead, there is no information of the recovery period observed between event one and two in the forecast starting the first of August. SYS4 predicts a swift recovery back to normal conditions on the forecast issued 1 September. A more detailed picture of 280 this intermediate recovery is instead conveyed by the seamless system. Thanks to the more frequent updates, the temporary increase in river flow is correctly picked-up giving a potential advantage of two to three weeks for planning actions. SYS4 does indicate the second low flow with a longer lead time than SEAM. However, SYS4 misses the timing of the event.

285 Even if this was a good forecast for SYS4, the information it provides is more informative (anomaly condition) than "actionable" (White et al., 2017). In the above example, a decision maker would have to make a decision based on a forecast that was issued 2.5 weeks earlier, which would inherently make the decision more uncertain if you only had the seasonal forecast. With the seamless system available a decision maker would gain the same early indication of a hazardous event and also have the benefit of frequent updates. In this particular case, the SEAM forecast for the first 290 event was more unstable for some ensemble members, but in general the event was well captured (Figure 8). The SEAM could also correctly capture the recovery with higher water levels between the extreme low flow events. The onset of the second low period was correctly modeled by the SEAM system, whereas the timing of the low flow was missed by SYS4. It should be said that using other less extreme thresholds (<90 and <95 percentiles) even further strengthened the case for using the 295 SEAM.

4 Conclusions

This study compared a set of hydrological hindcast experiments over the European domain with two meteorological forcings; ECMWF's seasonal forecasting system (SYS4) and a merged system of ECMWF extended range forecast and seasonal forecast system (SEAM). The latter showed a better overall skill and lower bias over most areas in Europe with lead times up to seven weeks. This increase in skill could be attributed to better initial conditions of the hydrological and meteorological model as well as a better atmospheric model version in SEAM. In some areas, particularly in the Alps and northern Finland, the seasonal forecast outperformed the merged forecast. However, in these areas the predictability the hydrological model is generally poor which makes these results quite uncertain. Given that the skill in the sub-seasonal range over Europe is in the range of the extended-range ensemble forecast would motivate to use the ENS-ER instead of SYS-4 for hydrometeorological predictions.

Still, there is an added benefit of using a seamless forecast over the extended range due to the extension of forecast horizon for the early detection of upcoming anomalous conditions. Indeed, as an example this study also highlighted the potential for the use of a sub-seasonal to seasonal forecast in the case of an extreme low-flow situation in the River Rhine. The higher frequency and skill of SEAM has the advantage of being a more "actionable" forecast than seasonal forecasts, given that a decision maker would be able to make use of the extra information. Care should be taken when using the forecasts in decision making since the reliability over Europe is "marginally useful" Weisheimer and Palmer (2014). It is therefore important to assess the reliability and skill of SEAM at the location it is to be implemented over the season of interest.

Future work with the seamless forecasting system is to further explore the limits of predictability, reliability and bias to assess the strengths and limitations of the current setup. The assumption that the forecasts can be randomly concatenated would also need to be tested against a system where the forecasts are matched according to their respective climatology. Bias correction of the forecasts might be a necessity, and the advantage of the extended-range and seasonal forecasts from ECMWF is that the availability of hindcasts which are enables just that.

Acknowledgements. This paper was financed through the Framework service contract for operating the EFAS computational centre in support of to the Copernicus Emergency Management Service (EMS)/Early Warning Systems (EWS) 198702. The authors would like to acknowledge Blazej Krzeminski for setting up the computational framework and Florian Pappenberger for the discussions regarding the seamless forecast system. We would also like to thank Kean Foster, Bastian Klein and Mike Hardeker for useful comments on the discussion paper.

References

330 Alfieri, L., Pappenberger, F., Wetterhall, F., Haiden, T., Richardson, D., and Salamon, P.: Evaluation of ensemble streamflow predictions in Europe, *Journal of Hydrology*, doi:<http://dx.doi.org/10.1016/j.jhydrol.2014.06.035>, <http://www.sciencedirect.com/science/article/pii/S0022169414004958>, 2014.

335 Arnal, L., Cloke, H. L., Stephens, E., Wetterhall, F., Prudhomme, C., Neumann, J., Krzeminski, B., and Pappenberger, F.: Skilful seasonal forecasts of streamflow over Europe?, *Hydrology and Earth System Sciences Discussions*, 2017, 1–27, doi:10.5194/hess-2017-610, <https://www.hydrol-earth-syst-sci-discuss.net/hess-2017-610/>, 2017.

340 Balsamo, G., Albergel, C., Beljaars, A., Boussetta, S., Brun, E., Cloke, H., Dee, D., Dutra, E., Muñoz Sabater, J., Pappenberger, F., de Rosnay, P., Stockdale, T., and Vitart, F.: ERA-Interim/Land: a global land surface reanalysis data set, *Hydrology and Earth System Sciences*, 19, 389–407, doi:10.5194/hess-19-389-2015, <http://www.hydrol-earth-syst-sci.net/19/389/2015/>, 2015.

345 Bartholmes, J. C., Thielen, J., Ramos, M. H., and Gentilini, S.: The european flood alert system EFAS – Part 2: Statistical skill assessment of probabilistic and deterministic operational forecasts, *Hydrol. Earth Syst. Sci.*, 13, 141–153, doi:10.5194/hess-13-141-2009, <http://www.hydrol-earth-syst-sci.net/13/141/2009/>, 2009.

350 Bazile, R., Boucher, M.-A., Perreault, L., and Leconte, R.: Verification of ECMWF System 4 for seasonal hydrological forecasting in a northern climate, *Hydrology and Earth System Sciences*, 21, 5747–5762, doi:10.5194/hess-21-5747-2017, <https://www.hydrol-earth-syst-sci.net/21/5747/2017/>, 2017.

355 Bechtold, P., Semane, N., Lopez, P., Chaboureau, J.-P., Beljaars, A., and Bormann, N.: Representing equilibrium and nonequilibrium convection in large-scale models, *Journal of the Atmospheric Sciences*, 71, 734–753, 2014.

Brown, A., Milton, S., Cullen, M., Golding, B., Mitchell, J., and Shelly, A.: Unified modeling and prediction of weather and climate: A 25-year journey, *Bulletin of the American Meteorological Society*, 93, 1865–1877, 2012.

360 Brunet, G., Shapiro, M., Hoskins, B., Moncrieff, M., Dole, R., Kiladis, G. N., Kirtman, B., Lorenc, A., Mills, B., Morss, R., et al.: Collaboration of the weather and climate communities to advance subseasonal-to-seasonal prediction, *Bulletin of the American Meteorological Society*, 91, 1397–1406, 2010.

Burek, P., Van Der Knijff, J. M., and De Roo, A.: LISFLOOD - Distributed Water Balance and Flood Simulation Model - Revised User Manual 2013, EUR - Scientific and Technical Research Reports 978-92-79-33191-6 (print); 978-92-79-33190-9, doi:10.2788/24982, 2013.

365 Ciais, P., Reichstein, M., Viovy, N., Granier, A., et al.: Europe-wide reduction in primary productivity caused by the heat and drought in 2003, *Nature*, 437, 529, 2005.

De Roo, A. P. J., Wesseling, C. G., and Van Deursen, W. P. A.: Physically based river basin modelling within a GIS: the LISFLOOD model, *Hydrological Processes*, 14, 1981–1992, doi:10.1002/1099-1085(20000815/30)14:11/12<1981::AID-HYP49>3.0.CO;2-F, [http://dx.doi.org/10.1002/1099-1085\(20000815/30\)14:11/12<1981::AID-HYP49>3.0.CO](http://dx.doi.org/10.1002/1099-1085(20000815/30)14:11/12<1981::AID-HYP49>3.0.CO), 2000.

370 Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L.,

Källberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J. J., Park, B. K.,
 370 Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J. N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Quarterly Journal of the Royal Meteorological Society*, 137, 553–597, doi:10.1002/qj.828, <http://dx.doi.org/10.1002/qj.828>, 2011.

Di Giuseppe, F., Molteni, F., and Tompkins, A. M.: A rainfall calibration methodology for impacts modelling based on spatial mapping, *Quarterly Journal of the Royal Meteorological Society*, 139, 1389–1401, 2013.

375 Dutra, E., Pozzi, W., Wetterhall, F., Di Giuseppe, F., Magnusson, L., Naumann, G., Barbosa, P., Vogt, J., and Pappenberger, F.: Global meteorological drought – Part 2: Seasonal forecasts, *Hydrol. Earth Syst. Sci.*, 18, 2669–2678, doi:10.5194/hess-18-2669-2014, <http://www.hydrol-earth-syst-sci.net/18/2669/2014/>, 2014.

Ferro, C. A. T., Richardson, D. S., and Weigel, A. P.: On the effect of ensemble size on the discrete and continuous ranked probability scores, *Meteorological Applications*, 15, 19–24, doi:10.1002/met.45, <http://dx.doi.org/10.1002/met.45>, 2008.

380 García-Herrera, R., Díaz, J., Trigo, R., Luterbacher, J., and Fischer, E.: A review of the European summer heat wave of 2003, *Critical Reviews in Environmental Science and Technology*, 40, 267–306, 2010.

Haiden, T., Magnusson, L., Tsonevsky, I., Wetterhall, F., Alfieri, L., Pappenberger, F., de Rosnay, P., Muñoz-Sabater, J., Balsamo, G., Albergel, C., Forbes, R., Hewson, T., Malardel, S., and Richardson, D.: ECMWF forecast performance during the June 2013 flood in Central Europe, Report, European Centre for Medium-Range Weather Forecasts, 2014.

385 Hazeleger, W., Severijns, C., Semmler, T., Ștefănescu, S., Yang, S., Wang, X., Wyser, K., Dutra, E., Baldasano, J. M., Bintanja, R., et al.: EC-Earth: a seamless earth-system prediction approach in action, *Bulletin of the American Meteorological Society*, 91, 1357–1363, 2010.

390 Hazeleger, W., Wang, X., Severijns, C., Ștefănescu, S., Bintanja, R., Sterl, A., Wyser, K., Semmler, T., Yang, S., Van den Hurk, B., et al.: EC-Earth V2. 2: description and validation of a new seamless earth system prediction model, *Climate dynamics*, 39, 2611–2629, 2012.

Hersbach, H.: Decomposition of the Continuous Ranked Probability Score for Ensemble Prediction Systems, [http://dx.doi.org/10.1175/1520-0434\(2000\)0152.0.CO;2](http://dx.doi.org/10.1175/1520-0434(2000)0152.0.CO;2), doi:10.1175/1520-0434(2000)0152.0.CO;2, [http://journals.ametsoc.org/doi/abs/10.1175/1520-0434\(2000\)015%3C0559%3ADOTCRP%3E2.0.CO%3B2](http://journals.ametsoc.org/doi/abs/10.1175/1520-0434(2000)015%3C0559%3ADOTCRP%3E2.0.CO%3B2), 2000.

395 Hoskins, B.: The potential for skill across the range of the seamless weather-climate prediction problem: a stimulus for our science, *Quarterly Journal of the Royal Meteorological Society*, 139, 573–584, 2013.

Hurrell, J., Meehl, G. A., Bader, D., Delworth, T. L., Kirtman, B., and Wielicki, B.: A unified modeling approach to climate system prediction, *Bulletin of the American Meteorological Society*, 90, 1819–1832, 2009.

400 Jonkeren, O., van Ommeren, J., and Rietveld, P.: Effects of low water levels on the river Rhine on the inland waterway transport sector, in: *Economics and Management of Climate Change*, pp. 53–64, Springer, 2008.

Meißner, D., Klein, B., and Ionita, M.: Development of a monthly to seasonal forecast framework tailored to inland waterway transport in Central Europe, *Hydrology and Earth System Sciences Discussions*, 2017, 1–31, doi:10.5194/hess-2017-293, <http://www.hydrol-earth-syst-sci-discuss.net/hess-2017-293/>, 2017.

405 Molteni, F., Stockdale, T., Balmaseda, M., Balsamo, G., Buizza, R., Ferranti, L., Magnusson, L., Mogensen, K., Palmer, T., and Vitart, F.: The new ECMWF seasonal forecast system (System 4), Report, ECMWF, 2011.

Ntegeka, V., Salamon, P., Gomes, G., Sint, H., Lorini, V., Thielen del Pozo, J., and Zambrano, H.: EFAS-Meteo: A European daily high-resolution gridded meteorological data set for 1990 - 2011, Report, 410 doi:10.2788/51262, 2013.

Palmer, T. and Webster, P.: Towards a unified approach to climate and weather prediction, in: Proceedings of 1st Demetra Conference on Climate Change, 1993.

Palmer, T. N., Doblas-Reyes, F. J., Weisheimer, A., and Rodwell, M. J.: Toward seamless prediction: Calibration of Climate Change Projections Using Seasonal Forecasts, *Bull. Amer. Meteor. Soc.*, 89, 459–470, 2008.

415 Pappenberger, F., Wetterhall, F., Dutra, E., Di Giuseppe, F., Bogner, K., Alfieri, L., and Cloke, H. L.: Seamless forecasting of extreme events on a global scale, pp. 3–10, missing, 2013.

Sene, K., Tych, W., and Beven, K.: Exploratory studies into seasonal flow forecasting potential for large lakes, *Hydrology and Earth System Sciences*, 22, 127–141, doi:10.5194/hess-22-127-2018, <https://www.hydrol-earth-syst-sci.net/22/127/2018/>, 2018.

420 Smith, P., Pappenberger, F., Wetterhall, F., Thielen, J., Krzeminski, B., Salamon, P., Muraro, D., Kalas, M., and Baugh, C.: On the operational implementation of the European Flood Awareness System (EFAS), Report 778, European Centre for Medium-Range Weather Forecasting, <http://www.ecmwf.int/en/elibrary/16337-operational-implementation-european-flood-awareness-system-efas>, 2016.

Thielen, J., Bartholmes, J., Ramos, M. H., and de Roo, A.: The European Flood Alert System - Part 1: 425 Concept and development, *Hydrol. Earth Syst. Sci.*, 13, 125–140, doi:10.5194/hess-13-125-2009, <http://www.hydrol-earth-syst-sci.net/13/125/2009/>, 2009.

Van Der Knijff, J. M., Younis, J., and De Roo, A. P. J.: LISFLOOD: a GIS-based distributed model for river basin scale water balance and flood simulation, *International Journal of Geographical Information Science*, 24, 189–212, doi:10.1080/13658810802549154, <http://dx.doi.org/10.1080/13658810802549154>, 2010.

430 Vitart, F., Buizza, R., Alonso Balmaseda, M., Balsamo, G., Bidlot, J. R., Bonet, A., Fuentes, M., Hofstadler, A., Molteni, F., and Palmer, T. N.: The new VarEPS-monthly forecasting system: A first step towards seamless prediction, *Q. J. R. Meteorol. Soc.*, 134, 1789–1799, 2008.

Weisheimer, A. and Palmer, T. N.: On the reliability of seasonal climate forecasts, *Journal of The Royal Society Interface*, 11, doi:10.1098/rsif.2013.1162, <http://rsif.royalsocietypublishing.org/content/11/96/20131162>, 435 2014.

Weisheimer, A., Doblas-Reyes, F. J., Jung, T., and Palmer, T.: On the predictability of the extreme summer 2003 over Europe, *Geophysical Research Letters*, 38, 2011.

White, C. J., Carlsen, H., Robertson, A. W., Klein, R. J., Lazo, J. K., Kumar, A., Vitart, F., Coughlan de Perez, E., Ray, A. J., Murray, V., Bharwani, S., MacLeod, D., James, R., Fleming, L., Morse, A. P., 440 Eggen, B., Graham, R., Kjellström, E., Becker, E., Pegion, K. V., Holbrook, N. J., McEvoy, D., Depledge, M., Perkins-Kirkpatrick, S., Brown, T. J., Street, R., Jones, L., Remenyi, T. A., Hodgson-Johnston, I., Buontempo, C., Lamb, R., Meinke, H., Arheimer, B., and Zebiak, S. E.: Potential applications of subseasonal-to-seasonal (S2S) predictions, *Meteorological Applications*, pp. n/a–n/a, doi:10.1002/met.1654, <http://dx.doi.org/10.1002/met.1654>, 2017.

445 Zajac, Z., Z.-B. M. S. P. B. P. G. A. and Bianchi, A.: Calibration of the lisflood hydrological model for europe - calibration round 2013., 2013.

Table 1. technical details of the forecast and the hindcast used in this paper.

System	T Res	Spatial Res	Horizon	Ens size	Issue frequency	Hindcast set	Hindcast Ens size
ENS-ER	3h/6h	18/36 km ¹	46 days	51	Twice weekly	20 years	11 members
SYS4	6h	80 km	7/13 months	51	Monthly	30 years	15/51 members
SEAM	6h	5 km	6 months	51	Twice weekly	20 years	11 members




Figure 1. Schematic overview of the operational ECMWF ensemble forecast for the extended range and its associated hindcast. The hindcasts consists of a reduced ensemble forecast (11 members) with the same starting date of year as the current forecast, but run for the previous 20 years.




Figure 2. Schematic overview of the seasonal, extended-range forecast and merged systems. The Extended forecast is issued every Monday and Thursday and extends up until 46 days, the seasonal forecasts is issued on the first of each month and extends up until 7 months (13 months in February, May, August and November). The merged forecasts concatenates the latest extended forecast with the latest seasonal forecast.




Figure 3. Continuous ranked probability skill score (CRPSS) for a) merged forecast against seasonal forecast for all start dates; b) as in a) but only for the first merged forecast of each month; c) merged forecast against climatology for all lead times in blue and d) as in c) but for the first merged forecast in the month. The shaded blue area denotes the 10-90 percentile of the CRPSS and the blue line the median. The black solid (dotted) lines in figure c and d denote the mean and 10-90th percentile of the CRPSS of the seasonal against the climatological forecast.




Figure 4. The number of weeks before the CRPSS goes below zero using only the first forecast of the month for a) SEAM against CLIM; b) SYS4 against CLIM c) SEAM against SYS4; and d) difference between SEAM against CLIM and SYS4 against CLIM expressed in weeks. The dimension of the circles is proportional to the number of days while the color scale refers to progressive weeks.




Figure 5. Mean relative error over all outlet points as a functionality of lead time in weeks for a) all starting dates of the forecasts and b) for the starting dates close to the beginning of the months. Negative values denote that the forecast is too wet in comparison with the CLIM run. The SEAM (SYS4) forecast is in blue (black) where the solid line denotes the median and the filled area (area between dotted lines) denote the 10th to 90th percentile.




Figure 6. Mean relative error for each of the outlet points for the SEAM forecast over the outlet points for a) week 2, b) week 4 and c) week 6. Red indicates where the forecast is too wet, and blue where it is too dry. Figure d) shows the difference in absolute error between SEAM and SYS, where blue (red) denotes points where SEAM has a smaller (larger) MAE than SYS4.




Figure 7. Reliability diagram for SEAM (blue) and SYS4 (black) for week 5 for all outlet points. The solid lines indicate the median reliability and the dotted lines the 25th and 75th percentiles.




Figure 8. Percentage of ensemble members predicting low flow anomaly (< 97%) on the Rhine river north of Cologne for summer 2003. The two starting dates in August and September from SYS4 are compared to the 17 starting dates of the seamless forecasting system. In two separate events the discharge was recorded below the 97 % percentile, event 1 on 17-27 of August and event 2 on 18-28 of September 2003.