

Dear Prof. De Michele,

I hereby resubmit the manuscript hess-2017-572, with first submission on 21 Sep 2017:

Seasonal Drought Prediction for Semiarid Northeast Brazil: Verification of Six Hydro-Meteorological Forecast Products

We introduced some major revisions addressing the requests of the referees:

- #1 The methods section was expanded
- #2 Brier skill score was added to the verification (Figs. 4, 8 and 11 in the revised manuscript)
- #3 We further explained the reasons for using the ECHAM and ECMWF models
- #4 Focus of the manuscript moved from longer to shorter time scales of the meteorological drought indices (1, 3 and 6 months)
- #5 A new paragraph explains the choice of drought thresholds
- #6 Methods section was expanded regarding SWSI including its formal expression
- #7 Methods section was expanded to explain the truncation of index values close to zero when deriving HDIs
- #8 Conclusions were reformulated and no longer compare ROCSS and RMSE

Another major revision, not requested by the referees, was removing the verification of the forecasted reservoir volume variation. Instead, we added that same forecast (months January to June) to the measured December reservoir volumes. This way we obtained a forecasted reservoir volume, comparable to the direct regression of the reservoir volume of MDIs. The methodology is explained in the reviewed manuscript.

Other revisions:

- # improved color legend of figures 3 and 5 (revised manuscript)
- # moved Table 2 (first version of the manuscript) to the appendix

Minor revisions to the text are marked in the attached document and listed in pages 27 to 31.

Best regards,
Jose Miguel Delgado

Seasonal Drought Prediction for Semiarid Northeast Brazil: Verification of Six Hydro-Meteorological Forecast Products

José Miguel Delgado¹, Sebastian Voss¹, Gerd Bürger¹, Klaus Vormoor¹, Aline Murawski², José Marcelo Rodrigues Pereira³, Eduardo Martins³, Francisco Vasconcelos Júnior³, and Till Francke¹

¹Institute of Earth and Environmental Sciences, University of Potsdam, Germany

²German Research Centre of Geosciences GFZ Potsdam, Germany

³Research Institute for Meteorology and Water Resources - FUNCEME, Fortaleza, Brazil

Correspondence to: José Miguel Delgado (martinsd@uni-potsdam.de)

Abstract. A set of seasonal drought forecast models was assessed and verified for the Jaguaribe River in semiarid northeast Brazil. Meteorological seasonal forecasts were provided by the operational forecasting system used at FUNCEME (Ceará's research foundation for meteorology) and by the European Centre for Medium-Range Weather Forecasts (ECMWF). Three downscaling approaches (*empirical quantile mapping, extended downscaling and weather pattern classification*)^{JMD} were tested and combined with the models in hindcast mode for the period 1981 to 2014. The forecast issue time was January and the forecast period was January to June. Hydrological drought indices were obtained by fitting a *multivariate linear regressiongeneralized linear model*^{JMD1} to observations. In short, it was possible to obtain forecasts for *a) monthly precipitation, b) meteorological drought indices, and c) hydrological drought indices.*

The skill of the forecasting systems was evaluated with regard to root mean square error (RMSE), *the Brier skill score (BSS)*^{JMD} and the relative operating characteristic *skill score (ROCSS)(ROC-skill-score)*^{JMD}. *The tested forecasting products showed similar performance in the analyzed metrics.*^{JMD} Forecasts of monthly precipitation had little or no skill considering RMSE and mostly no skill with BSS^{JMD}. *Still, the forecast of extreme events of low monthly precipitation showed skill for the rainy season (ROC-skill-score of 0.24 to 0.33)*^{JMD}. A similar picture was seen when forecasting meteorological drought indices: low skill regarding RMSE and BSS^{JMD} and significant skill when *discriminating hit rate and false alarm rate given by the ROCSS (forecasting drought events of e.g. SPEI₀₁ showed a ROCSS of around 0.5)*^{JMD}. *foreeasting drought events of e.g. SPEI₀₁ (ROC-skill-score of 0.53 to 0.61)*^{JMD}. *Similar results were obtained for low regional reservoir storage forecasts.*^{JMD} Regarding *the temporal variation of the forecast skill of the meteorological indecesthe skill in the forecasted months*^{JMD}, it was greatest^{JMD} for April, when compared to *February and March (the remaining months of the rainy season)*^{JMD}, while the *skill of reservoir volume forecasts decreased with lead time*^{JMD}.

This work showed that a multi-model ensemble can forecast drought events of time scales relevant to water managers in northeast Brazil with skill. But no or little skill could be found in the forecasts *of the whole range*^{JMD} of monthly precipitation or drought indices *of lower scales, like SPI₀₁(e.g. forecasting average years)*^{JMD}. Both this work and those here revisited showed that major steps forward are needed in forecasting the rainy season in northeast Brazil.

¹JMD: stricter definition

1 Introduction

Northeastern Brazil has historically been the epicenter of major drought events. Fioreze et al. (2012) identified 100 severe droughts since the 16th century in this region, while Marengo et al. (2016) identified 68 major events for the same period. Within this region, the state of Ceará has been in the frontline of the fight against this natural hazard. This has been both due to the impacts suffered in the past and to the measures taken to improve its resilience.

Droughts in Ceará reflect a meteorological anomaly over the tropical Atlantic Ocean. Dry years are generally related to a positive sea surface temperature (SST) anomaly on the tropical North Atlantic, associated with a negative anomaly on the tropical South Atlantic and over the equator. This forces a northward shift of the intertropical convergence zone, taking the rainbelt to northern latitudes. The causes for this anomaly are linked to the occurrence of the El Niño Southern Oscillation and to the North Atlantic Oscillation (Hastenrath, 2012).

Past famines and mass migrations triggered large investments in infrastructure in recent decades. These investments brought hundreds of strategic reservoirs and thousands of small dams to a semi-arid landscape, which are being managed according to a transparent water allocation process (Formiga-Johnsson and Kemper, 2005). In order to support water allocation and management, the state runs a seasonal drought forecasting system and issues annual quantitative and qualitative forecasts of the magnitude of the rainy season. These predictions can support decisions ranging from agricultural management (choice of crop, planning of seeding time) to water distribution and reservoir operation.

Currently, the forecasting system in Ceará is based on the general circulation model ECHAM4.6 (Roeckner et al., 1992). It runs from January to August on persisted SSTs (observed SST anomalies $SSTs^{JMD}$ which are assumed invariant), covering each year's rainy season (February to April). The forecasts produced by this model are generally downscaled with the NCEP regional spectral model (Juang et al., 1997), in order to resolve the spatial variability of Ceará. Verification and the current forecast can be retrieved under <http://www3.funcceme.br/previsao-climatica/>. For downscaled forecasts check <http://seca-vista.geo.uni-potsdam.de/JMD>.

In this study we intend to evaluate and extend this prediction system by employing

1. an additional underlying GCM,
2. a statistical approach based on the classification of weather patterns,
3. empirical-statistical downscaling methods to increase the spatial resolution and temporal fidelity of the predictions, and
4. drought indices as powerful integrative descriptors for the description of drought severity.

By these means, we aim to address the following questions:

What skill do the seasonal *meteorological drought* forecasts have?

While the term *meteorological drought* focuses on the atmospheric forcing causing water shortage, its effective implications for society are more specifically accounted for by the term *hydrological drought* (Araújo and Bronstert, 2016). Since the aim of the prediction system is to support water management, we sharpen the previous question in this regard:




Figure 1. Flowchart explaining the methodology used for predicting meteorological data, meteorological drought indices (MDI) and hydrological drought indices (HDI).

Can we forecast *hydrological* droughts in Ceará based on these seasonal meteorological forecasts?

2 Methods

2.1 General approach

This work employed a cascade of models and algorithms ranging from two general circulation models (one atmospheric and one coupled) at the top to hydrological indices at the bottom (Fig. 1). Each step involved different types of target variables being forecasted: The *meteorological forecasts* (Fig. 1, top) refer to meteorological variables ("meteo data") from GCM-forecasts and the subsequent downscaling and bias correction to match the spatial and temporal resolution. The *meteorological indices* (same figure, centre), refer to the indices that were used to describe the magnitude of the forecasted meteorological drought. Finally, the *hydrological indices* (same figure, bottom) were calculated based on meteorological indices in an attempt to infer the magnitude of a hydrological drought characterized by meteorological and hydrological properties. To allow for the comparison with observations, we use results of GCM hindcast, i.e. a model that has been run with data only known until the specified time in the past. As these are supposed to represent and technically resemble true forecasts, they are referred to as "forecasts" henceforward. All results and computations after the statistical downscaling have a monthly time step. Similarly, all results and computations here presented were aggregated to selected subbasins (Fig. 2).




Figure 2. Left panel shows the location of the Jaguaribe River basin in South America. Right panel shows the Jaguaribe River together with its main tributaries, division into subcatchments used in this work, and meteorological and rainfall observations stations.

2.2 Study area

The spatial domain chosen for this analysis is the Jaguaribe river basin. Due to the river's regional importance, a lot has been written about its hydrology and development (see e.g. Araújo and Bronstert, 2016; Araújo, 1990). The Jaguaribe is the most important river in Ceará. Its catchment has an area of 70000km² and is home to about 2.7 million people (IPECE, 2017). Annual precipitation amounts to 755 mm, of which about 90% fall in the months January to June. **The rainfall season comprising the months January to May is often considered key in securing water reserves for the whole year. June contributes with considerably less rainfall.**^{JMD²}

Average potential evapotranspiration is estimated to 2100mm. Due to its dominant geology composed of a crystalline complex, aquifers in the region are unproductive. Runoff is practically the only source of drinking water for people and animals as well as irrigation. To that end, most of the water is stored in thousands of reservoirs of all scales across the watershed.

The main tributaries are the Banabuiú river in central Ceará and the Salgado river in southern Ceará. We aggregated the results of this research into five subcatchments: the aforementioned tributaries Banabuiú and Salgado, the upper (upstream of Orós Reservoir), middle (upstream of Castanhão Reservoir) and lower (downstream of Castanhão Reservoir) Jaguaribe. An overview of the state and location of these catchments and tributaries is given in Fig. 2.

2.3 Seasonal Forecast Models ("GCM output")

To address the first research question we employed different combinations of dynamical and statistical models and a weather pattern classification methodology to produce meteorological drought indices. The dynamical seasonal forecast models were

²JMD: quadra chuvosa

provided by FUNCEME and ECMWF in the form of hindcasts for the period 1981 to 2014. Details like resolution, reference and short description are given in Table 1.

The ECMWF operational seasonal forecasting system S4 has 51 ensemble members and six months lead time. ~~It is a fully coupled atmosphere-ocean model~~~~It is fully coupled to an ocean circulation model~~^{JMD}. The system has been systematically verified (Vitart, 2013; Molteni et al., 2011; Richardson et al., 2012). The hindcast version of the system has the same specifications of the operational model but only 15 ensemble members. ~~It is available for academic purposes and is here employed as a benchmark for the verification of the regional forecasting system in operation in Ceará.~~^{JMD3}

The seasonal forecasting system ~~implemented at FUNCEME (Ceará's hydro-meteorological agency)~~~~used by Ceará's meteorological agency~~^{JMD} ~~is are~~^{JMD} based on the general circulation model ECHAM4.6. Details on this model can be found in Table 1. The operational and hindcast version have 20 ensemble members and are run on initial sea surface temperature (SST) anomalies persisted during the forecasting period (8 months). ~~The initial state represents a typical (but random) realization of late December as derived from AMIP-type runs (Gates et al., 1999)~~~~The forecast model is initialized in January with the conditions of the atmosphere modeled by an AMIP-type run for December 31st.~~^{JMD} The AMIP-type^{JMD} run starts in 1961 and is forced by monthly observed SSTs (NOAA Optimum Interpolation SST V2). ~~Therefore, potential forecast skill is solely based on oceanic memory.~~~~This means that only modeled and no observed atmospheric conditions are used to initialize the forecast model.~~^{JMD} The forecasting system of FUNCEME is in operational use and seasonal forecasts are released monthly.^{JMD4}

2.4 Downscaling of GCM output

In order to predict precipitation over particular locations it is necessary to downscale the GCM forecasts. Three statistical downscaling approaches were employed: expanded downscaling (XDS), empirical quantile mapping (EQM) and weather pattern classification (WP, see Table 1 for details and references). To differentiate between two fundamentally different downscaling approaches, weather pattern classification will not be referred to as downscaling approach/method throughout the text.

The downscaling approaches used here yielded a full set of meteorological variables distributed across the catchment at points where observations were available (daily mean temperature, relative humidity, wind speed and daily total precipitation and radiation). The forecasting products obtained from the combinations of GCM and downscaling will be named after their components: *XDS:ECHAM*, *XDS:ECMWF*, *EQM:ECHAM*, *EQM:ECMWF*, *WP:ECHAM*, and *WP:ECMWF*.

Weather patterns were classified using the SANDRA methodology described in Philipp et al. (2016). The selection of the *optimal* classification was done visually in respect to the explained variation of the observed meteorological drought indices. The classification itself was independent of the MDIs, so that no artificial skill was to be expected from forecasting the stations. Only MDI scales of 1, 12 and 36 months were calculated.

³JMD: why ecwfw?

⁴JMD: why echam?

Table 1. Output variables of each prediction model used in this paper

Model/Method	Short description	Reference	Spatial Scale
FUNCEME Seasonal Forecast System	A 20 member ECHAM4.6 ensemble. Atmospheric circulation model, initial SSTs persisted for 6 months. Initial conditions of the atmosphere modeled by AMIP-type run (starting in 1961). AMIP run is forced by monthly observed SSTs (NOAA Optimum Interpolation SST V2).	Roeckner et al. (1992)	approx. 2.8 degree longitude/latitude
ECMWF Seasonal Forecast System	A multi-model 15 member ensemble including ocean circulation. Initial conditions coming from ERA Interim.	e.g. Stockdale et al. (1998)	approx. 0.7 degree latitude/longitude
expanded downscaling	Simulates local events consistent with prevailing atmospheric circulation while preserving local covariability	Bürger (1996)	network of monitoring stations
empirical quantile mapping	Improves systematic biases throughout the statistical distribution by mapping the empirical cumulative distributions of the observed and modelled variable	e.g. Wetterhall et al. (2012)	network of monitoring stations
weather pattern classification	Including pre-selection of variables, variable combinations and spatial domain.	e.g. Murawski et al. (2016); Philipp et al. (2007)	network of monitoring stations

2.5 Drought Quantification using Drought Indices

Meteorological droughts were quantified in magnitude and temporal scale using meteorological drought indices (MDI). After careful appraisal regarding data demand and current conventions, the following indices were selected: SPI_{01} , $SPI_{03}SPI_{T2}^{JMD5}$, $SPI_{06}SPI_{T36}^{JMD6}$ (Svoboda et al., 2012; McKee et al., 1993), $SPEI_{01}$, $SPEI_{03}SPEI_{T2}^{JMD7}$ and $SPEI_{06}SPEI_{T36}^{JMD8}$ (Vicente-Serrano et al., 2009). The numbers in the index name (e.g. SPI_{01}) refer to the temporal scale in months for which the index was computed.

⁵JMD: changes scales

⁶JMD: changes scales

⁷JMD: changes scales

⁸JMD: changes scales

The forecast is generated at the beginning of ~~always takes place in~~^{JMD} January for the period from January until June. Indices obtained by downscaling forecasts with temporal scale greater than the lead time of the forecast will include values from the observation set. SPI_6 , for example, will contain five month of measured precipitation in the January forecast. In June, the same index will be calculated exclusively with forecasted precipitation. The skill of a SPI_6 forecast for some months is therefore expected to be greater than the skill of a SPI_1 forecast beforehand. ~~SPI_{12} , for example, will contain at least six months of measured precipitation and a maximum of six months forecasted precipitation. The skill of a SPI_{12} forecast is therefore expected to be greater than the skill of a SPI_{01} forecast beforehand.~~^{JMD9} This feature does not apply to WP classification.

Time scales greater than 6 months are of no value for the verification of the forecasting system in terms of meteorology, as rainfall in the preceding dry season is usually negligible. However, the hydrology of Ceará is characterized by long-term memory introduced by a vast network of reservoirs. Additionally, drought events in this region are known to be long and creeping phenomena that must be quantified on large temporal scales. MDIs with long temporal scales will therefore have to be considered when designing the hydrological drought index (HDI) forecast model in the next section. To that end, we will employ shorter time scales for MDI verification, but keep longer time scales (greater than 6 months) in the regression of hydrological drought indices (HDIs), since they provide a better fit for the forecast model. ~~Although time scales greater than 6 months are in strict terms of no value for the verification of the forecasting system, they put the system into perspective. Droughts are long, creeping phenomena that must be quantified on large temporal scales. A forecasting system should at least be able to improve the ability to predict well the onset of a drought given the previous months rainfall deficit.~~^{JMD10}

Regarding hydrological droughts, various hydrological drought indices (HDI) were reviewed and two were considered suitable for this work. All other indices ~~reviewed~~^{JMD} either *a*) require consumptive data for water use, which is impractical for the given settings or *b*) focus on streamflow, which misses the most important features (ephemeral rivers, role of reservoirs) of the hydrological system of Ceará and many other semi-arid regions. The only index chosen from the literature was the surface water supply index (SWSI) as formulated in Doesken et al. (1991) with a weight of 0.5 for precipitation within the reservoir catchment and 0.5 for reservoir volume;^{JMD}

$$SWSI = \frac{0.5P(rs) + 0.5P(pr) - 50}{12} \text{JMD11} \quad (1)$$

where $P(x)$ is the non-exceedance probability of x based on available historical records of x , rs is mean monthly reservoir storage in the respective catchment and pr is the monthly precipitation averaged for the respective catchment.^{JMD12} The second index, V , was defined as the regional reservoir volume at the end of each month relative to the total regional reservoir storage capacity. ~~the regional reservoir volume divided by the maximum regional reservoir volume; and the monthly variation of the regional reservoir volume.~~^{JMD13}

⁹JMD: replacing use of SPI12

¹⁰JMD: clarifying use of long scales in HDI

¹²JMD: introducing swsi

¹³JMD: introducing V

In terms of event prediction, the event considered for the meteorological drought indices in use in this work is “dry spells of moderate to extreme magnitude”, translated by values lower than or equal to -1 in the SPI/SPEI scale. For precipitation a threshold based on the 30th percentile of the series of observed monthly precipitation was used. The threshold for defining HDI drought events was based on the 30th percentile of the series of observed monthly HDI. The reason for using the 30th percentile was the classification used by the regional agencies to separate between a “dry”, a “wet” (above the 70th percentile) and a “normal” year. ~~Drought events were here considered to be the periods that fall below the 30th percentile of each index.~~^{JMD14}

2.6 Regression of Hydrological Drought indices

Forecasts of hydrological drought indices were obtained by searching and fitting a ~~multivariate regression model~~^{JMD} ~~generalized linear model~~^{JMD} to observations of hydrological drought indices and reservoir volume changes. ^{JMD}As candidate predictors, meteorological drought indices of all temporal scales were used, ~~using meteorological drought indices as predictors~~^{JMD15}.

For predicting SWSI the multivariate linear regression was fit directly to the hydrological index. For the regional reservoir volume, V , two different approaches were followed. With the first approach $M1$, the multivariate linear regression was fit directly to the values of V , analogous to SWSI. With $M2$, the second approach, the multivariate linear regression was first fit to the monthly changes of V . Then the predicted value of V was calculated by adding its predicted monthly changes to the most recent measured value in December. The regional reservoir volume predicted by the two regression models $M1$ and $M2$ are denoted V_{M1} and V_{M2} respectively.^{JMD16}

Model parsimony was enforced by predictor selection comprising a heuristic search for the best Akaike information criterion (AIC) under the constraint of checking the predictors for multi-collinearity. ~~Predictor selections comprised a heuristic search for the best Akaike information criterion. Model parsimony was enforced by the constraint of checking the predictors for multi-collinearity.~~^{JMD} To eliminate multi-collinearity between predictors, correlated predictors were replaced by their ratios.^{JMD}

Possible forms of multilinear regression include predictors as denominators of fractions. This implies that these predictors must not take the value zero, in order to exclude the division by zero. To enforce this condition, the MDIs in question were removed from the time-series when approaching zero, in particular values in $] - 0.1, 0.1[$.^{JMD17}

2.7 Forecast Verification

At each level of Fig. 1, a verification of the forecast was performed. Three metrics were employed: the root mean square error (RMSE), the relative operating characteristic skill score (ROCSS) and the Brier skill score (BSS) (Wilks, 2005). Root mean square error is a scalar accuracy measure applied to the realisations of the ensemble forecast. The Brier score is also a scalar accuracy measure, though for verification of probabilistic forecasts of predefined events. The relative operating characteristic is a

¹⁴JMD: explaining thresholds

¹⁵JMD: expanded in new sentence

¹⁶JMD: explains prediction

¹⁷JMD: describing truncation of indices close to zero

discrimination-based verification metric for forecasts of defined events. For more information on these metrics, we recommend chapter 8 of Wilks (2005) ^{JMD18}

5 ~~RMSE~~Root-mean-square-error (RMSE) ^{JMD} was computed for each member, ensemble mean and climatology, i.e. long-term mean annual cycle. Climatology was considered the reference forecast. The mean square error was computed for monthly values in the forecast period (1981-2014, January-June) and averaged over the entire period. The square root of this measure is the RMSE. It shows the capability of the model to correctly forecast monthly ~~values~~precipitation ^{JMD}, but it does not quantify the skill to predict particular events of water scarcity. January to June precipitation represents over 90% of the annual precipitation in the Jaguaribe basin.

10 Another important metric employed was the Brier skill score. The Brier score can be seen as the sum of three terms *reliability*, *resolution* and *uncertainty*. The term *reliability* measures the differences between forecast probabilities and relative frequencies of the observed event. Thus, low values of this score correspond to high reliability. *Resolution* measures the ability of the forecast to discern periods in which observed frequencies depart from average. Finally the term *uncertainty* quantifies the variability of the observations: when the event being forecasted almost never, or almost always happens, the uncertainty of the forecast is small. The Brier score is here understood as in Wilks (2005) as: ^{JMD19}

15
$$BS = \frac{1}{n} \sum_{k=1}^n (y_k - o_k)^2$$
 ^{JMD20} (2)

where k denotes the numbering of the n forecast-event pairs (y_k and o_k respectively). The Brier skill score (BSS) is computed with respect to the reference forecast (BS_{ref}), here the climatological relative frequencies of the event: ^{JMD21}

$$BSS = 1 - \frac{BS}{BS_{ref}}$$
 ^{JMD22} (3)

20 BSS can take any value lower than or equal to one and a forecast is said to have skill if its Brier skill score is greater than zero. ^{JMD23}

The last metric employed was the relative operating characteristic (~~ROC~~) ^{JMD} skill score (ROCSS) ^{JMD}. The relative operating characteristic describes the ability to discriminate between true positives and false positives when forecasting a given event. It is normally calculated for a set of forecast probability bins ~~with varying forecasting probabilities~~ ^{JMD}, therefore having great importance for decision makers ^{JMD}. ROC skill score was calculated as

25
$$ROCSS = 2 \cdot AUC - 1$$
 (4)

as in Wilks (2005), where AUC is the area under the ROC curve. The ROCSS can have values between -1 and 1 , where anything below zero means *no skill*. A ROCSS of 0 corresponds to the skill of a reference random forecast.

¹⁸JMD: intro to brier score

¹⁹JMD: explaining brier score

²¹JMD: explaining brier score

²³JMD: explaining brier score




Figure 3. Root mean square error of the forecast of monthly precipitation. On the left panels, box plots show the spread of the RMSE of each model. The asterisks (*) show the RMSE of the ensemble mean. The RMSE of using climatology as a forecasting product is given by the grey dashed line. The four panels on the right show RMSE for each individual station for each model. Note that in general the ensemble mean ranks better than the best of the ensemble members.

3 Results and Discussion

3.1 Forecasting precipitation

The RMSE of the precipitation forecast are presented in Fig. 3. ECMWF ranks better than ECHAM, while EQM:ECMWF results in the lowest RMSEs and XDS:ECHAM in the greatest. Still, the best results in terms of RMSE are comparable to the climatology, meaning that there is limited skill in forecasting monthly precipitation. The spatial distribution of RMSE of the ensemble mean in April shows a concentration of high RMSEs in the lower Jaguaribe catchment for EQM and in the Salgado catchment for XDS.

The ensemble mean of the forecast, shown by the asterisks in Fig. 3 as well as in other figures below, always displayed a lower RMSE than *any of the ensemble members*. This happens because the ensemble mean “smooths out unpredictable detail and presents the more predictable elements of the forecast” ((WMO) World Meteorological Organization, 2012). Despite its usefulness, the ensemble mean is not entirely appropriate for predicting drought events. Ensemble means do not provide any information on the probability of an extreme event.




Figure 4. BSS of the forecast of a^{JMD} monthly low precipitation event. On the left side the BSS is shown for each model/downscaling combination and for the forecasting months averaged over the respective subcatchments. BSS values below zero were assigned a “no skill” category in order to improve readability. The grey line is the BSS of the multi-model ensemble. The four panels on the right show BSS averaged over all forecasting months for each station. Note that in most cases the forecast of monthly precipitation has no skill.

Other than RMSE, which does not provide any information on the skill of extreme^{JMD} event forecasts, the BSSROCSS^{JMD} is explicitly suited for that purpose and is, as^{JMD} shown in Fig. 4. One remarkable observation is to be made regarding the BSS: skill is mostly absent when forecasting drought events based on precipitation and its thresholds. The only exception is the forecast for April, where the multi-model ensemble shows limited skill in the three regions considered. This results will be
 5 discussed further in face of the greater skill shown when forecasting drought events based on MDIs.^{JMD24}

Still, all forecasting systems here presented show skill in discriminating events against false alarm forecasts. This is expressed by the ROC-curve shown in Fig. 5^{JMD25} based on the hit rate and false alarm rate of occurrence of a predefined event^{JMD}. The variation of the ROCSS over time can be attributed to lead time (skill decreasing with increasing lead time) and to low or no precipitation in the months before and after the rainy season. Months of typically low precipitation showed poor ROCSS
 10 (Fig. 5: January, May, June). When comparing downscaling techniques and GCMs, EQM mostly outperformed XDS, while the skill was less affected by using different GCMs.

²⁴JMD: discussion p bss

²⁵JMD: discussion p rocss




Figure 5. ROCSS of the forecast of a^{JMD} monthly low precipitation event. On the left side the ROCSS is shown for each model/downscaling combination and for the forecasting months averaged over the respective subcatchments. The four panels on the right show ROCSS averaged over all forecasting months for each station.

To put our results into context, we could find three reports with a statement of verification concerning precipitation forecast in Ceará. Castro et al. (2013) presents a RMSE of between 120 and 130 mm for the *Sertão Interior de Inhamuns*, using an empirical model with forecasts issued in January for the period February to June. Moura and Hastenrath (2004), with a forecast issued in end of February for the period of March to June, i.e. with shorter lead-time than our work, shows a RMSE of 50 to 70 mm (Hastenrath and Greischar (1993) obtained similar results).

3.2 Forecasting meteorological drought indices

A time series of seasonal MDI forecasts was plotted to illustrate the forecast spread given by model EQM:ECMWF (Fig. 6). The improvement provided by the ensemble mean, when compared to each member, is clearly visible. Also visible are several observed events of moderate to severe drought (below the dashed grey line). The ensemble mean is able to forecast at least a few of these events.

A measure of the general agreement (for all kinds of conditions, dry, average or wet) between forecasted and observed MDI is given by the RMSE in Fig. 7. The relationship between forecast probability and relative frequency of a drought event (i.e. the




Figure 6. Time series of the seasonal forecast of SPI_{01} in the Castanhão subcatchment given by ECMWF:EQM. Only periods from January to June are shown. T_{τ}^{JMD} the threshold “moderate drought event” is given by the grey dashed line.




Figure 7. box plots of the root mean square error of forecasted meteorological drought index. Asterisk (*) shows the RMSE of the ensemble mean and box plots show the spread of the individual members. Note that in general the ensemble mean ranks better than the best of the ensemble members.

BSS) is provided in Fig. 8, whereas the balance between hit rate and false alarm rate for the same event can be seen in the form of ROCSS in Fig. 9 below.^{JMD26}

The RMSE of MDI forecasts is shown in Fig. 7. With the exception of the predictions produced by the WP approach for SPI_{01} , the general ranking of the approaches is quite consistent among the three subbasins. As with precipitation, the

²⁶JMD: intro to plots




Figure 8. Brier Skill Score of forecasted meteorological drought event based on an event of index lower than -1. The grey line shows the result of the multi-model ensemble mean.

RMSE of SPI_{01} and $SPEI_{01}$ generally does not differ from that of the climatology and is greatest for ECHAM and EQM. EQM:ECMWF and XDS:ECMWF show consistently lowest RMSE and XDS:ECMWF performs better than the climatology. Interestingly, ECMWF consistently outperforms ECHAM on all scales.^{JMD}

For longer time scales, i.e. to the right of SPI_{12} in Fig. 7, the WP approach shows notably poorer performance than the other methods. A reason could be the way MDIs of longer time scales are computed: when using GCM downscaling approaches, observed data has to be used for computing indices of longer time scales. Still, here all models except WP classification show an improvement regarding climatology.^{JMD27}

RMSE reflects the prediction skill for the whole range of the indices, including wet spells and dry spells/droughts. When aiming primarily at forecasting drought events, this verification may be misleading. Nevertheless, this metric shows which models are most appropriate for this domain and confirms the plausibility of the forecasting system also for wet years.

As for the Brier skill score (BSS), Fig. 8 shows this indicator of skill for time scales of 1, 3 and 6 months in three regions of the Jaguaribe river. For the one month time scale, it is noteworthy that the first three months of the forecast display the lowest skill. In particular the March forecast shows no skill in most models, March being a key contributing month in the rainy season. The second half of the rainy season, April/May/June, has generally more skill. The same BSS minimum can be observed on the

²⁷JMD: not relevant any longer




Figure 9. ROCSS of forecasted meteorological drought event based on an event of index lower than -1. The grey line shows the result of the multi-model ensemble mean.

SPI_{03} and SPI_{06} panels, but this time with slightly greater value than for SPI_{01} , since these indices entail some measured data. Another interesting observation is that, contrary to RMSE, here no product can be considered a clear winner.^{JMD28}

For the ECHAM model a possible explanation for lower skill on the onset of the rainy season may lie on its initial conditions. Since the initial conditions for each model run are provided by the output of an AMIP-type run (Gates et al., 1999), they may depart considerably from actual atmospheric conditions. According to this hypothesis, the model would come closer to atmospheric conditions only through the SST forcing, which could explain a certain lag in the forecasting skill. Still, this explanation can only account for ECHAM and not the ECMWF model, which is fully coupled and whose initial conditions are derived from ERA Interim.^{JMD29}

The ROCSS for the different months of the forecasting period shows a slightly different picture than the RMSE and BSS^{JMD} previously presented. Fig. 9 shows ROCSS for time scales of 1, 312^{JMD30} and 636^{JMD31} months in three regions of the Jaguaribe river. For a time scale of one month,^{JMD} There is no clear pattern concerning the relationship between lead time and skill for

²⁸JMD: discussion mdi bss

²⁹JMD: discussion mdi bss

³⁰JMD: changed MDI scales

³¹JMD: changed MDI scales

anyone^{JMD} of the forecasting models. As in previous plots, the forecasting skill for different MDIs tend to display a minimum in March.^{JMD} MDIs with time scales of 12 and 36 months show a clear decrease in skill with increasing lead time.^{JMD32}

Contrary to the results for RMSE, EQM:^{JMD}ECHAM shows^{JMD} comparably good ROCSS and BSS^{JMD} in forecasting MDI drought events of all scales $SPI_{01}/SPEI_{01}$ drought events for January to May^{JMD} in all three regions. Still, the comparably low skill of the March forecast is problematic, March being the month of greatest precipitation in most of the catchment. WP:ECHAM features the best BSSs for $SPI_{01}/SPEI_{01}$ in April and May, whereas EQM:ECHAM features generally the highest ROCSS in April for the same scale. April features a maximum ROCSS for EQM:ECHAM within the main months of the rainy season.^{JMD}

The good results of the $SPI_{36}/SPEI_{36}$ forecasts are not surprising, given that by definition they are computed with monthly precipitation from the previous 36 months, where most of the contributing months are coming from past observations and not within the forecast period. More interesting are the results for $SPI_{12}/SPEI_{12}$, in particular for June. Here, six wet months contributing to the index are forecasted and the remaining six dry months (July to December of the previous year) come from observations. Since the variability of precipitation in July to December is very low (mostly close to zero), this index measures well the skill of the forecasting system concerning long term droughts. Here, a greater differentiation is visible, in particular EQM:ECHAM and EQM:ECMWF standing out. WP:ECHAM also compares relatively well to other models.^{JMD33}

It is worth to look at the BSS of $SPI_6/SPEI_6$, even if they partly encompass measured values. BSS in June in particular is a valuable indicator of the ability of the models in forecasting the whole rainy season. Here, most products display some skill in forecasting a drought event. XDS:ECMWF is the only displaying no skill for all three regions in $SPEI_6$. Generally the skills are higher with $SPEI_6$ than with SPI_6 . Regarding $SPEI_6$, EQM:ECHAM and EQM:ECMWF display skill in all three regions. In the important region of Castanhão, where the largest reservoir and most infrastructure is located, EQM:ECHAM and XDS:ECHAM perform best in forecasting $SPEI_6$ for June, although with a low value of BSS.^{JMD34}

The multi-model ensemble skill, shown by the grey line is generally close to the upper envelope formed by that of the individual models. For $SPEI_{01}$ in the months January to May (rainy season) the ROCSSskill^{JMD} of the multi-model ensemble is always positive and oscillates around 0.5. An interesting result is the improvement in skill when SPI_{01} is replaced by $SPEI_{01}$. The gray line, which shows the ROCSS/BSS^{JMD} for the multi-model ensemble, has an increase in skill at all scales and regions during the rainy season^{JMD}.

In both RMSE and ROCSS, WP classification ranks comparably well for time scales of one month ($SPI_{01}/SPEI_{01}$), but poorly for longer time scales. There are two methodological reasons for that. First, the variables for each WP classification are selected from a pool of candidate variables according to their skill in predicting SPEI or SPI. The selected variables are therefore different among different time scales of SPI/SPEI. In other words, the SPI/SPEI with longer time scales are not predicted simply as an accumulation of previous SPI/SPEI states, but as a fundamentally distinct WP classification. The second

³²JMD: changed MDI scales

³³JMD: changes MDI scales

³⁴JMD: discussion on MDI6

reason is that the EQM/XDS downscaled forecasts of SPI/SPEI of longer time scales (12 and 36 months) are based in part on observed values, since they include data belonging to the period before the forecast issue time.^{JMD35}

A similar forecast assessment has been reported by e.g. Dutra et al. (2013). Events were defined by a SPI_{03} lower than -1 , with a lead time of 3 months. ROCSS obtained were in the order of 0.6 for the Blue Nile basin^{JMD}, which is comparable with the results presented in this paper, but much lower for other rivers^{JMD} basins^{JMD} e.g. Zambezi.

3.3 Forecasting hydrological drought indices

The multivariate regression model~~generalized linear model~~^{JMD} equations obtained and their respective R^2 are shown in the appendix, Table A1. Long scale MDIs (like SPI_{12} or SPI_{36}) prevail as predictors of reservoir volume, whereas short scale MDIs like SPI_1 are mostly present as predictors of reservoir volume change~~HDI~~^{JMD}. This reflects the time scale of reservoir storage variations. At a given moment in time, the reservoir storage results from several months of inflow. Similarly, the effect of a month of high inflow in the reservoir storage level is likely to be only residual,~~which is given by the comparably low factors associated to SPI_{01} and $SPEI_{01}$ in the Jaguaribe region Table A1.~~^{JMD}

The forecast of the three HDIs shows notable differences between downscaling techniques EQM/XDS and the WP classification (Fig. 10). WP classification has lower RMSE than EQM/XDS when predicting SWSI or V_{M1} . For V_{M2} , the difference between WP and EQM/XDS is much smaller. The ensemble spread of WP classification shrinks considerably from V_{M1} to V_{M2} . All methods show a decrease in RMSE from V_{M1} to V_{M2} .^{JMD} EQM/XDS have more ROC skill and lower RMSE. Notably, WP classification has lower RMSE when forecasting reservoir volume change, which reflects the results found in Sect. 3.2, i.e. greater skill in forecasting SPI_{01} and $SPEI_{01}$ than longer time scales^{JMD}.

Again, WP classification considers by design only a range of discrete MDIs, which can affect RMSE. MDIs were limited to nine values, of which -0.75 , 0 and 0.75 are the closest to zero. The continuous values of MDI derived by the other products downscaling^{JMD} are problematic, because the multilinear regression also considers division by the meteorological drought index. When the MDI are close to zero, outliers arise and skew the RMSE. These datapoints were therefore removed from the verification metrics.^{JMD}

Regarding the prediction of HDI drought events, Fig. 11 clearly points out that prediction performs best when targeting reservoir volume with model M2 (adding predicted monthly value to the December observed regional reservoir volume). Here, all products show reasonable performance for most regions, but a decreasing skill with increasing lead time. Another important observation is that WPs do not display skill in forecasting HDI events as shown in Fig. 11.^{JMD36}

Contrary to the MDIs, the BSS of the HDIs do not feature a minimum in March. A slight tendency of lower skills towards the end of the rainy season is observable in V_{M1} forecasts. V_{M2} shows comparably good results for all GCM/downscaling combinations in predicting HDI events, confirming the results in Fig. 10.^{JMD37}

³⁵JMD: SPI12 and SPI36 not shown in this version

³⁶JMD: discussion of hdi bss

³⁷JMD: discussion of hdi bss




Figure 10. Root mean square error of the forecast of SWSI, V predicted with $M1$ and V predicted with $M2$ (based on month-to-month variation). The forecast period is January to June. Three regions are presented: Lower Jaguaribe, Orós and Castanhão. The horizontal grey dashed line shows the RMSE of the climatology.

Forecast quality is only weakly affected by different GCMs or downscaling techniques. SWSI and reservoir volume can be forecasted with less error using the statistical downscaling approaches EQM and XDS. The WP approach apparently ranks better when forecasting HDI with high variation in short timescales as the monthly variation of the regional reservoir volume. This especially applies to Orós and lower Jaguaribe catchments. In the Castanhão catchment the RMSE is similar for all approaches.^{JMD}

This latter issue affecting the RMSE of the hydrological drought index is not felt by the ROCSS, since it considers only the distinction of an event between dry and not dry.^{JMD} The ROCSS shows small differences between GCMs or downscaling methods (Fig. 12). V_{M2} features the highest ROCSS of the different indexes used and very little variability among downscaling approaches and GCMs employed. As with BSS, the ROCSS of V_{M2} decreases with increasing lead time. The results of SWSI and V_{M1} are very similar, with SWSI showing higher variability among downscaling approaches and GCMs. V_{M2} could be predicted by WP classification with high ROCSS, whereas V_{M1} and SWSI show no skill.^{JMD38}

It was possible to predict any of the indexes with skill in most modelling approaches and catchments. Still, V_{M2} was predicted with the greatest BSS and ROCSS, even though it showed worse R^2 when fitting the regression model on which the prediction

³⁸JMD: discuss V M2




Figure 11. BSS of the forecast of drought events as defined by SWSI, V predicted with $M1$ and V predicted with $M2$ (based on month-to-month variation). An event is defined as an index of magnitude lower than the 30th percentile of the observations. The forecast period is January to June. Three regions are presented: Lower Jaguaribe, Orós and Castanhão. The horizontal grey dashed line shows the RMSE of the climatology.

is based (Table A1). This result hints at better HDI predictability when the predictant is a change in reservoir volume than the reservoir volume itself. One reason for the improved predictability of V_{M2} is surely the importance of persistence in reservoir storage dynamics. By adding the predicted change to the measured reservoir volume we are providing valuable measured information to the forecast model that SWSI and V_{M1} do not have.^{JMD39}

- 5 ~~Dry conditions can be most skillfully forecasted for the SWSI and the reservoir volume. The skill is mostly constant during the season and differs between the regions mainly for the reservoir volume with^{JMD}~~

We could not find reports on streamflow/reservoir forecasting systems for the region of Ceará stating BSS,^{JMD} ROCSS, RMSE or other verification measure. Still, for other semi-arid regions of the world, similar skill values could be found in the literature. Trambauer et al. (2015) forecasted events of standardized runoff index of 6 months lower than -0.5 with variable
 10 lead times. Their best catchment point to a ROCSS of 0.7 with a lead time of 5 months. Seibert et al. (2017) forecasted events with a standardized streamflow index (Vicente-Serrano et al., 2012) below -0.5, reporting ROCSS of 0 at the outlet of a large river (the Limpopo in southern Africa) to close to 1 in its headwater catchments.




Figure 12. ROCSS of the forecast of drought events as defined by SWSI, V predicted with $M1$ and V predicted with $M2$ (based on month-to-month variation). An event is defined as an index of magnitude lower than the 30th percentile of the observations. The forecast period is January to June. Three regions are presented: Lower Jaguaribe, Orós and Castanhão. The horizontal grey dashed line shows the RMSE of the climatology.

Table 2. BSS of January-May multi-model ensemble forecast. The ensemble includes ECHAM and the ECMWF seasonal forecast model, as well as the EQM and XDS downscaling techniques. The BSS are averaged over each region. Columns show different indices used for the forecast: P is seasonal precipitation, SPI_{01} and $SPEI_{01}$ are standardized precipitation indices with scale 1 month, and *Reservoir Volume* stands for regional reservoir volume in percentage of regional storage capacity. The BSS refers to meteorological and hydrological drought events described in ROCSS of January-May multi-model ensemble forecast. The ensemble includes ECHAM and the ECMWF seasonal forecast model, as well as the EQM and XDS downscaling techniques. The ROCSS are averaged over each region. Columns show different indices used for the forecast: P is seasonal precipitation, SPI_{01} and $SPEI_{01}$ are standardized precipitation indices with scale 1 month, and *Reservoir Volume* stands for regional reservoir volume in percentage of regional storage capacity. The ROC refers to an event described in ^{JMD40} Sect. 2

region	BROC ^{JMD} SS						
	P	SPI_{01}	$SPEI_{01}$	$SPEI_{06}$	$SPEI_{12}$ ^{JMD}	<i>Reservoir Volume (M1)</i>	<i>Reservoir Volume (M2)</i>
Orós	no skill	no skill	0.11	0.38		0.43	0.66
Castanhão	no skill	0.08	0.11	0.32		0.23	0.52
Lower Jaguaribe	0.06	no skill	0.16	0.37		0.12	0.71

3.4 Multimodel ensemble forecast

Finally, we present the skill score of the multi-model ensemble forecast in Table 2. Each type of index considered (precipitation, meteorological drought index and hydrological drought index) is presented. Results of the WP classification were excluded from the multi-model ensemble, because they ~~did not cover all the indeces addressed in this worke~~ contributed very negatively to the overall skill^{JMD41}.

The BSS of ^{JMD} forecasts of low precipitation events (given in column P), as well as that of ^{JMD} the forecasts of drought defined by the SPI_{01} show either very low or none or very low^{JMD} skill. Forecasts of $SPEI$ and reservoir storage show greatest skill^{JMD} generally display greater skill than the forecasts of SPI ^{JMD}. This points to a possible bias in the forecasting that is compensated by introducing temperature in the equation of $SPEI$. Still it is possible to forecast events of low relative reservoir storage with similar skill as for $SPEI_{01}$ ^{JMD}

The best skill obtained by the multi-model ensemble was forecasting drought events related to reservoir storage in the Lower Jaguaribe region^{JMD}. The good skill of the reservoir storage forecast is likely related to the long memory of the reservoir system. The forecasted precipitation will affect the reservoir only marginally, since most of its storage is accumulated throughout several years. Most importantly, BSS increases when V_{M2} is used instead of V_{M1} , i.e. when reservoir volume is forecasted by adding forecasted reservoir volume change to measured December reservoir volume. In fact, the multilinear regression used for predicting reservoir storage employs as predictors $SPEI_{12}$ as well as other MDIs of shorter time scales^{JMD}. In this sense, the reservoir storage forecast has less skill than would be expected, knowing that the skill of $SPEI_{12}$ was very high.^{JMD}

Since the $SPEI_{12}$ forecasts are calculated based on a combination of past observed data and the forecasted months, it is clear that the reservoir storage forecast is not a good indicator for streamflow. Reservoir inflow depends more on short time scales and contrary to reservoir storage is not affected by interannual memory. Note the comparably lower skill of the reservoir storage variation (between 0.11 and 0.15), a variable which is expected to closely reflect reservoir inflow.^{JMD}

Table 2 reveals an interesting pattern in this work: additional information to the forecast model tends to increase forecast skill. $SPEI_1$ is based on temperature and precipitation data and was forecasted with greater skill than SPI_1 , which is only based on precipitation. Similarly, $SPEI_6$, which combines forecasted and measured precipitation and temperature from months prior to the forecasting period have more skill than $SPEI_1$ forecasts. The greatest BSS is given by V_{M2} , a HDI that requires measured initial reservoir volumes as well as a combination of several MDIs. This last point stresses the importance of assimilating prior hydrological conditions into the forecast products.^{JMD42}

Improvements in the skill of hydrological drought forecasting can still be obtained. Both reservoir storage's and reservoir inflow's forecast skill showed considerable differences to the skills of the corresponding meteorological forecasts. The multilinear regression employed for estimating hydrological drought is likely the reason for such a drop in forecast skill. A way of circumventing this multilinear regression would be to add a process-based hydrological model to the forecasting system.^{JMD}

³⁹JMD: why is V_{M2} so good?

⁴¹JMD: only WP for scales 1, 12 and 36

⁴²JMD: increasing skill with complexity

4 Conclusions

The plausibility and skill of a set of drought forecasting models was presented. Different types of drought events were considered: a rainfall anomaly during the rainy season, standardized precipitation indices below a given threshold and anomalies in regional reservoir storage. The forecast products considered were combinations of two models, ECHAM and the ECMWF seasonal forecast, two downscaling techniques, XDS and EQM, and a weather pattern classification approach.

Each model provided an ensemble of predictions, so deterministic and probabilistic measures of skill could be used. Results showed that models with little to no skill by a deterministic measure (RMSE) showed skill under a probabilistic skill score. This underlines the importance of having at least one deterministic and one probabilistic measure of skill.^{JMD43} The deterministic measure also^{JMD} allowed to see the significant improvement introduced by the ensemble mean: the ensemble mean had in most cases lower a greater^{JMD44} root mean square error than the climatology. The RMSE of the ensemble mean however was comparable to the climatology and in some cases greater lower^{JMD45}. Still, no approach model^{JMD46} had an^{JMD} RMSE that significantly departed from the RMSE of the climatology.

It was possible to provide a probabilistic forecast based on a multi-model ensemble.^{JMD} A multi-model ensemble forecast (The probabilistic forecast^{JMD} was obtained by binding all members of all models into one product. The skill of this forecast is given in Figs. 4, 8, 11, and Table 25, 9, 12, and Table ??^{JMD}. Multimodel ensembles^{JMD} These^{JMD} can be considered to be our best guess of a probabilistic drought forecast, since they are consistently among the best forecast skills provided by the individual models. Individual members surpassed the multi-model ensemble skill only occasionally, for particular combinations of regions, months and indices.

The skill of the hydrological drought forecast, namely the relative reservoir storage V_{M2} was 0.66, 0.52 and 0.71 for the regions of Orós, Castanhão and lower Jaguaribe, respectively was comparable to the skill of the $SPEI_{OT}$ forecast, around 0.6 for the regions of Castanhão and Orós and 0.5 for lower Jaguaribe^{JMD}. The skill obtained for the hydrological drought forecast is likely inflated by the long memory of the reservoir system and the use of observed reservoir volume to define the conditions prior to each forecast^{JMD} and by the dependence on meteorological drought indices of long time scales^{JMD47}. Still, the R^2 of the regression that provides the reservoir variation underlying V_{M2} was lower than that of V_{M1} , indicating that a regression might be a poor prediction of reservoir inflow. The hydrological drought index that most resembles streamflow, i.e. reservoir storage variation, was forecasted with much lower skill^{JMD48}. Improvements are expected by coupling a process-based hydrological model to the seasonal forecasting system.

This work showed that a multi-model ensemble can forecast drought events of time scales relevant to water managers in northeast Brazil with skill. But none or little skill could be found in the forecasts of the whole range^{JMD} of monthly precipitation

⁴³JMD: this is wrong, as the reviewer correctly said

⁴⁴JMD: in fact the opposite is true, I suppose this was a lapse

⁴⁵JMD: same as previous remark, probably a lapse

⁴⁶JMD: improves clarity

⁴⁷JMD: not true with M2

⁴⁸JMD: refer to regression

Table A1. Regression used for predicting regional reservoir volume and regional reservoir volume change using a set of MDIs as predictors. Regional reservoir volume was taken at the end of each month relative to the total regional reservoir storage capacity. Regional reservoir volume change refers to the difference between the given and the previous month.

Region	Predictant	Formula	R^2
Orós	Reservoir Volume	$59.0 - 22.9SPEI_{36} + 6.67SPEI_{12} + 45.4SPEI_{36} + 6.00SPEI_{12}SPEI_1 - 5.30SPEI_{36}SPEI_{36}$	0.64
Orós	Reservoir Volume Change	$0.416 + 2.43SPEI_1SPEI_1 + 2.23SPEI_1SPEI_{12} - 0.173SPEI_{36} \frac{SPEI_{36}}{SPEI_{12}} + 4.20SPEI_1 \frac{SPEI_{12}}{SPEI_{12}} - 0.00334 \frac{SPEI_{36}}{SPEI_{12}} \frac{SPEI_{36}}{SPEI_{12}}$	0.36
Castanhão	Reservoir Volume	$55.4 + 12.5SPEI_{36} + 12.4 \frac{SPEI_{36}}{SPEI_{36}} - 3.12SPEI_{12} \frac{SPEI_{12}}{SPEI_{36}} - 3.19 \frac{SPEI_{12}}{SPEI_{36}} \frac{SPEI_{12}}{SPEI_{12}} + 10.19 \frac{SPEI_{36}}{SPEI_{36}} SPEI_{12}$	0.41
Castanhão	Reservoir Volume Change	$2.95 + 3.47SPEI_1 - 1.15 \frac{SPEI_1}{SPEI_1} - 1.27 \frac{SPEI_1}{SPEI_1} SPEI_{36} - 0.791SPEI_1 \frac{SPEI_{36}}{SPEI_{12}} + 1.41SPEI_{12} \frac{SPEI_{36}}{SPEI_{36}}$	0.21
Lower Jaguaribe	Reservoir Volume	$33.4 + 16.3SPEI_{36} + 16.6SPEI_{12} + 5.65SPEI_{36}SPEI_{12} - 13.5SPEI_{12} \frac{SPEI_{12}}{SPEI_{12}} + 0.877SPEI_{12} \frac{SPEI_{12}}{SPEI_{36}}$	0.60
Lower Jaguaribe	Reservoir Volume Change	$0.689 + 2.22SPEI_1 + 0.0353 \frac{SPEI_{36}}{SPEI_{12}} + 2.12SPEI_1SPEI_{12} + 1.08SPEI_{12} \frac{SPEI_{12}}{SPEI_{12}} + 0.286SPEI_1 \frac{SPEI_{12}}{SPEI_{36}}$	0.38

or drought indices of smaller temporal scales, like SPI_{01} (e.g. forecasting average years)^{JMD}. Both this work and others here revisited showed that major steps forward are needed in forecasting the rainy season in northeast Brazil.

Appendix A: Appendix A

Author contributions. Conceived and designed the experiments: TF. Performed the experiments: SV, MR, GB, KV, AM, JD, FVJ, EM.

5 Analyzed the data: JD, SV, TF. Wrote the paper: JD, TF, SV.

Competing interests. There are no competing interests present.

Acknowledgements. This work was funded by the Federal Ministry of Education and Research of Germany under grant number *01DN14013*. The first author was also supported by the German Research Foundation under grant number *BR1731/18-1*. One of the hindcast datasets was kindly provided by the European Centre for Medium-Range Weather Forecasts. We would like to acknowledge the support of the Open Access Publication Fund of the University of Potsdam.

References

- Araújo, J. A. d. A.: Barragens no Nordeste do Brasil, DNOCS, Fortaleza, José Amáury de Aragão Araújo edn., 1990.
- Araújo, J. C. d. and Bronstert, A.: A method to assess hydrological drought in semi-arid environments and its application to the Jaguaribe River basin, Brazil, *Water International*, 41, 213–230, doi:10.1080/02508060.2015.1113077, <http://dx.doi.org/10.1080/02508060.2015.1113077>, 2016.
- 5 Bürger, G.: Expanded downscaling for generating local weather scenarios, *Climate Research*, 7, 111–128, <http://www.int-res.com/abstracts/cr/v07/n2/p111-128/>, 1996.
- Castro, T. d., Souza, F. B., J. M., Alves, Pontes, R. S. T., Reis, D., L. n., L., and Daher, S.: Neo-fuzzy neuron model for seasonal rainfall forecast: A case study of Ceara's eight homogenous regions, *Journal of Intelligent & Fuzzy Systems*, 25, 389–394, doi:10.3233/IFS-2012-10645, <http://content.iospress.com/articles/journal-of-intelligent-and-fuzzy-systems/ifs645>, 2013.
- 10 Doesken, N., McKee, T., and Kleist, J.: Development of a Surface Water Supply Index for the Western United States, Tech. Rep. Climatology Report 91-3, Colorado Climate Center, Department of Atmospheric Science, Colorado State University, http://climate.colostate.edu/pdfs/climo_rpt_91-3.pdf, 1991.
- Dutra, E., Di Giuseppe, F., Wetterhall, F., and Pappenberger, F.: Seasonal forecasts of droughts in African basins using the Standardized Precipitation Index, *Hydrol. Earth Syst. Sci.*, 17, 2359–2373, doi:10.5194/hess-17-2359-2013, <https://www.hydrol-earth-syst-sci.net/17/2359/2013/>, 2013.
- 15 Fioreze, A. P., Bubel, A. P. M., Callou, A. P., Mendonça, B. C. d. S., Nunes, C. M., Pinto, C. G., Viana, C. F. G., Junior, D. S. R., Martins, E. S. P. R., Rodrigues, F. S. F., Filho, F. d. A. d. S., Teixeira, F. J. C., Viana, F. L., Nascentes, J. C. d. M., Filho, J. G. C. G., Júnior, J. A. d. L., Campos, J. N. B., Carvalho, J. O. d., Gonçalves, J. Y. d. B., Burte, J., Silva, L. M. C. d., Azevedo, L. G. T. d., Bursztyn, M., Cerqueira, M. R. S., Coimbra, Thomas, P., Nobre, P., Vieira, R. F., Alves, R. F. F., Chacon, S. S., and Paulino, W. D.: A questão da água no Nordeste, Ministério da Ciência e Tecnologia (MCT), <http://livroaberto.ibict.br/handle/1/669>, 2012.
- Formiga-Johnsson, R. M. and Kemper, K.: Institutional and Policy Analysis of River Basin Management: The Jaguaribe River Basin, Ceara, Brazil, SSRN Scholarly Paper ID 757424, Social Science Research Network, Rochester, NY, <https://papers.ssrn.com/abstract=757424>, 2005.
- 25 Gates, W. L., Boyle, J. S., Covey, C., Dease, C. G., Doutriaux, C. M., Drach, R. S., Fiorino, M., Gleckler, P. J., Hnilo, J. J., Marlais, S. M., Phillips, T. J., Potter, G. L., Santer, B. D., Sperber, K. R., Taylor, K. E., and Williams, D. N.: An Overview of the Results of the Atmospheric Model Intercomparison Project (AMIP I), *Bulletin of the American Meteorological Society*, 80, 29–56, doi:10.1175/1520-0477(1999)080<0029:AOOTRO>2.0.CO;2, <https://journals.ametsoc.org/doi/abs/10.1175/1520-0477%281999%29080%3C0029%3AAOOTRO%3E2.0.CO%3B2>, 1999.
- 30 Hastenrath, S.: Exploring the climate problems of Brazil's Nordeste: a review, *Climatic Change*, 112, 243–251, doi:10.1007/s10584-011-0227-1, <https://link.springer.com/article/10.1007/s10584-011-0227-1>, 2012.
- Hastenrath, S. and Greischar, L.: Further Work on the Prediction of Northeast Brazil Rainfall Anomalies, *Journal of Climate*, 6, 743–758, doi:10.1175/1520-0442(1993)006<0743:FWOTPO>2.0.CO;2, <http://journals.ametsoc.org/doi/abs/10.1175/1520-0442%281993%29006%3C0743%3AFWOTPO%3E2.0.CO%3B2>, 1993.
- 35 IPECE: Anuário Estatístico do Ceará, <http://www.ipece.ce.gov.br/index.php/anuario-estatistico-do-ceara>, 2017.

- Juang, H.-M. H., Hong, S.-Y., and Kanamitsu, M.: The NCEP Regional Spectral Model: An Update, *Bulletin of the American Meteorological Society*, 78, 2125–2143, doi:10.1175/1520-0477(1997)078<2125:TNRMSMA>2.0.CO;2, <http://journals.ametsoc.org/doi/abs/10.1175/1520-0477%281997%29078%3C2125%3ATNRMSMA%3E2.0.CO%3B2>, 1997.
- Marengo, J. A., Torres, R. R., and Alves, L. M.: Drought in Northeast Brazil—past, present, and future, *Theoretical and Applied Climatology*, pp. 1–12, doi:10.1007/s00704-016-1840-8, <https://link.springer.com/article/10.1007/s00704-016-1840-8>, 2016.
- McKee, T. B., Doesken, N. J., Kleist, J., and others: The relationship of drought frequency and duration to time scales, Anaheim, California, http://clima1.cptec.inpe.br/~rclima1/pdf/paper_spi.pdf, 1993.
- Molteni, F., Stockdale, T., Balmaseda, M., Balsamo, G., Buizza, R., Ferranti, L., Magnusson, L., Mogensen, K., Palmer, T., and Vitart, F.: The new ECMWF seasonal forecast system (System 4), *European Centre for Medium-Range Weather Forecasts*, https://www.researchgate.net/profile/Frederic_Vitart/publication/230688826_The_new_ECMWF_seasonal_forecast_system_System_4/links/0912f51028aa651240000000.pdf, 2011.
- Moura, A. D. and Hastenrath, S.: Climate Prediction for Brazil's Nordeste: Performance of Empirical and Numerical Modeling Methods, *Journal of Climate*, 17, 2667–2672, doi:10.1175/1520-0442(2004)017<2667:CPFBNP>2.0.CO;2, <http://journals.ametsoc.org/doi/full/10.1175/1520-0442%282004%29017%3C2667%3ACPFBNP%3E2.0.CO%3B2>, 2004.
- Murawski, A., Bürger, G., Vorogushyn, S., and Merz, B.: Can local climate variability be explained by weather patterns? A multi-station evaluation for the Rhine basin, *Hydrol. Earth Syst. Sci.*, 20, 4283–4306, doi:10.5194/hess-20-4283-2016, <https://www.hydrol-earth-syst-sci.net/20/4283/2016/>, 2016.
- Philipp, A., Della-Marta, P. M., Jacobeit, J., Fereday, D. R., Jones, P. D., Moberg, A., and Wanner, H.: Long-Term Variability of Daily North Atlantic–European Pressure Patterns since 1850 Classified by Simulated Annealing Clustering, *Journal of Climate*, 20, 4065–4095, doi:10.1175/JCLI4175.1, <http://journals.ametsoc.org/doi/abs/10.1175/JCLI4175.1>, 2007.
- Philipp, A., Beck, C., Huth, R., and Jacobeit, J.: Development and comparison of circulation type classifications using the COST 733 dataset and software, *International Journal of Climatology*, 36, 2673–2691, doi:10.1002/joc.3920, <http://onlinelibrary.wiley.com/doi/10.1002/joc.3920/abstract>, 2016.
- Richardson, D. S., Bidlot, J., Ferranti, L., Ghelli, A., Haiden, T., Hewson, T., Janousek, M., Prates, F., and Vitart, F.: Verification statistics and evaluations of ECMWF forecasts in 2011–2012, *European Centre for Medium-Range Weather Forecasts*, 2012.
- Roeckner, E., Arpe, K., Bengtsson, L., Brinkop, S., D{u}menil, L., Esch, M., Kirk, E., Lunkeit, F., Ponater, M., Rockel, B., Sausen, R., Schlese, U., Schubert, S., and Windelband, M.: Simulation of the present-day climate with the {ECHAM-3} model: impact of model physics and resolution, 1992.
- Seibert, M., Merz, B., and Apel, H.: Seasonal forecasting of hydrological drought in the Limpopo Basin: a comparison of statistical methods, *Hydrol. Earth Syst. Sci.*, 21, 1611–1629, doi:10.5194/hess-21-1611-2017, <https://www.hydrol-earth-syst-sci.net/21/1611/2017/>, 2017.
- Stockdale, T. N., Anderson, D. L. T., Alves, J. O. S., and Balmaseda, M. A.: Global seasonal rainfall forecasts using a coupled ocean–atmosphere model, *Nature*, 392, 370–373, doi:10.1038/32861, <http://www.nature.com/nature/journal/v392/n6674/abs/392370a0.html>, 1998.
- Svoboda, M., Hayes, M., Wood, D. A., and World Meteorological Organization, W.: Standardized Precipitation Index User Guide, WMO, WMO, http://library.wmo.int/pmb_ged/wmo_1090_en.pdf, 2012.
- Trambauer, P., Werner, M., Winsemius, H. C., Maskey, S., Dutra, E., and Uhlenbrook, S.: Hydrological drought forecasting and skill assessment for the Limpopo River basin, southern Africa, *Hydrology and Earth System Sciences*, 19, 1695–1711, doi:10.5194/hess-19-1695-2015, wOS:000353877000007, 2015.

Vicente-Serrano, S. M., Beguería, S., and López-Moreno, J. I.: A Multiscalar Drought Index Sensitive to Global Warming: The Standardized Precipitation Evapotranspiration Index, *Journal of Climate*, 23, 1696–1718, doi:10.1175/2009JCLI2909.1, <http://journals.ametsoc.org/doi/abs/10.1175/2009JCLI2909.1>, 2009.

Vicente-Serrano, S. M., López-Moreno, J. I., Beguería, S., Lorenzo-Lacruz, J., Azorin-Molina, C., and Morán-Tejeda, E.: Accurate Computation of a Streamflow Drought Index, *Journal of Hydrologic Engineering*, 17, 318–332, doi:10.1061/(ASCE)HE.1943-5584.0000433, <http://ascelibrary.org/doi/abs/10.1061/%28ASCE%29HE.1943-5584.0000433>, 2012.

Vitart, F.: Evolution of ECMWF sub-seasonal forecast skill scores over the past 10 years, ECMWF Research Department Technical Memorandum, p. 28, 2013.

Wetterhall, F., Pappenberger, F., He, Y., Freer, J., and Cloke, H. L.: Conditioning model output statistics of regional climate model precipitation on circulation patterns, *Nonlin. Processes Geophys.*, 19, 623–633, doi:10.5194/npg-19-623-2012, <https://www.nonlin-processes-geophys.net/19/623/2012/>, 2012.

Wilks, D. S.: *Statistical Methods in the Atmospheric Sciences*, Academic Press, google-Books-ID: _b4R9j2Iy7EC, 2005.

(WMO) World Meteorological Organization: *Guidelines on Ensemble Prediction Systems and Forecasting*, WMO, WMO, http://library.wmo.int/pmb_ged/wmo_1091_en.pdf, 2012.

15 **Changes**

Author: anonymous

Added 0
 Deleted 0
 Replaced 0

20 **Author: JMD (JM Delgado)**

Added 53
 Deleted 33
 Replaced 56

25 **List of changes**

	Added (JMD): (empirical quantile mapping, exte ...	1
	Replaced (JMD): multivariate linear regression	1
	Added (JMD): , the Brier skill score (BSS)	1
	Replaced (JMD): skill score (ROCSS)	1
30	Added (JMD): The tested forecasting products sh ...	1
	Added (JMD): and mostly no skill with BSS	1
	Deleted (JMD): . Still, the forecast of extreme eve ...	1

	Added (JMD): and BSS	1
	Replaced (JMD): discriminating hit rate and false al...	1
	Deleted (JMD): Similar results were obtained for 1...	1
	Replaced (JMD): the temporal variation of the forec...	1
5	Replaced (JMD): st	1
	Deleted (JMD): February and March (1
	Deleted (JMD):)	1
	Added (JMD): ,while the skill of reservoir volum...	1
	Deleted (JMD): of the whole range	1
10	Replaced (JMD): of lower scales, like SPI_{01}	1
	Replaced (JMD): SST anomalies	2
	Added (JMD): For downscaled forecasts check...	2
	Added (JMD): The rainfall season comprising the...	4
	Replaced (JMD): It is a fully coupled atmosphere-...	5
15	Added (JMD): It is available for academic purpos...	5
	Replaced (JMD): implemented at FUNCEME (Cear...	5
	Replaced (JMD): is	5
	Replaced (JMD): The initial state represents a typic...	5
	Deleted (JMD): -type	5
20	Replaced (JMD): Therefore, potential forecast skill...	5
	Added (JMD): The forecasting system of FUNCE...	5
	Replaced (JMD): SPI_{03}	6
	Replaced (JMD): SPI_{06}	6
	Replaced (JMD): $SPEI_{03}$	6
25	Replaced (JMD): $SPEI_{06}$	6
	Replaced (JMD): is generated at the beginning of	7
	Replaced (JMD): SPI_6 , for example, will contain fi...	7
	Replaced (JMD): Time scales greater than 6 months...	7
	Deleted (JMD): reviewed	7
30	Replaced (JMD): :	7
	Added (JMD):	7
	Added (JMD): where $P(x)$ is the non-exceedance...	7
	Replaced (JMD): The second index, V , was defined...	7
	Replaced (JMD): In terms of event prediction, the e...	8
35	Replaced (JMD): multivariate regression model	8

	Added (JMD): and reservoir volume changes.	8
	Replaced (JMD): As candidate predictors, meteorol...	8
	Added (JMD): For predicting SWSI the multivari...	8
	Replaced (JMD): Model parsimony was enforced by...	8
5	Added (JMD): To eliminate multi-collinearity bet...	8
	Added (JMD): Possible forms of multilinear regr...	8
	Added (JMD): At each level of Fig. 1, a verificati...	9
	Replaced (JMD): RMSE	9
	Replaced (JMD): values	9
10	Added (JMD): Another important metric employ...	9
	Added (JMD):	9
	Added (JMD): where k denotes the numbering of...	9
	Added (JMD):	9
	Added (JMD): BSS can take any value lower than...	9
15	Deleted (JMD): (ROC)	9
	Added (JMD): (ROCSS)	9
	Replaced (JMD): . It is normally calculated for a set...	9
	Added (JMD): , therefore having great importanc...	9
	Added (JMD): a	11
20	Deleted (JMD): extreme	11
	Replaced (JMD): BSS	11
	Replaced (JMD): and is	11
	Added (JMD): One remarkable observation is to...	11
	Added (JMD): Still, all forecasting systems here...	11
25	Deleted (JMD): based on the hit rate and false alar...	11
	Added (JMD): a	12
	Replaced (JMD): T	13
	Added (JMD): A measure of the general agreeme...	13
	Added (JMD): Interestingly, ECMWF consistentl...	14
30	Deleted (JMD): For longer time scales, i.e. to the...	14
	Added (JMD): As for the Brier skill score (BSS),...	15
	Added (JMD): For the ECHAM model a possible...	15
	Added (JMD): and BSS	15
	Replaced (JMD): 3	15
35	Replaced (JMD): 6	15

	Deleted (JMD): For a time scale of one month,	15
	Replaced (JMD): any	16
	Added (JMD): As in previous plots, the forecasti	16
	Deleted (JMD): MDIs with time scales of 12 and	16
5	Deleted (JMD): EQM:	16
	Added (JMD): s	16
	Added (JMD): and BSS	16
	Replaced (JMD): MDI drought events of all scales	16
	Replaced (JMD): WP:ECHAM features the best BS	16
10	Deleted (JMD): The good results of the	16
	Added (JMD): It is worth to look at the BSS of	16
	Replaced (JMD): ROCSS	16
	Added (JMD): /BSS	16
	Replaced (JMD): at all scales and regions	16
15	Deleted (JMD): In both RMSE and ROCSS, WP c	17
	Added (JMD): basin	17
	Deleted (JMD): s	17
	Added (JMD): basins	17
	Replaced (JMD): multivariate regression model	17
20	Replaced (JMD): reservoir volume, whereas short s	17
	Deleted (JMD): , which is given by the comparably	17
	Added (JMD): WP classification has lower RMS	17
	Deleted (JMD): EQM/XDS have more ROC skill a	17
	Replaced (JMD): the other products	17
25	Added (JMD): These datapoints were therefore re	17
	Added (JMD): Regarding the prediction of HDI d	17
	Added (JMD): Contrary to the MDIs, the BSS of	17
	Deleted (JMD): Forecast quality is only weakly aff	18
	Deleted (JMD): This latter issue affecting the RM	18
30	Added (JMD): V_{M2} features the highest ROCSS	18
	Added (JMD): It was possible to predict any of th	19
	Deleted (JMD): Dry conditions can be most skillfu	19
	Added (JMD): BSS,	19
	Replaced (JMD): BSS of January-May multi-model	20
35	Replaced (JMD): B	20

	Replaced (JMD): $SPEI_{06}$	20
	Replaced (JMD): did not cover all the indices addre...	21
	Added (JMD): BSS of	21
	Added (JMD): that of	21
5	Replaced (JMD): very low or no	21
	Deleted (JMD): and reservoir storage show greater...	21
	Added (JMD): generally display greater skill than...	21
	Deleted (JMD): Still it is possible to forecast event...	21
	Added (JMD): The best skill obtained by the...	21
10	Replaced (JMD): Most importantly, BSS increases...	21
	Deleted (JMD): In this sense, the reservoir storage...	21
	Deleted (JMD): Since the $SPEI_{12}$ forecasts are c...	21
	Added (JMD): Table 2 reveals an interesting patte...	21
	Deleted (JMD): Improvements in the skill of hydr...	21
15	Deleted (JMD): Results showed that models with 1...	22
	Deleted (JMD): also	22
	Replaced (JMD): lower	22
	Replaced (JMD): greater	22
	Replaced (JMD): approach	22
20	Deleted (JMD): n	22
	Deleted (JMD): It was possible to provide a proba...	22
	Replaced (JMD): A multi-model ensemble forecast	22
	Replaced (JMD): 4, 8, 11, and Table 2	22
	Replaced (JMD): Multimodel ensembles	22
25	Replaced (JMD): V_{M2} was 0.66, 0.52 and 0.71 for...	22
	Added (JMD): and the use of observed reservoir...	22
	Deleted (JMD): and by the dependence on meteor...	22
	Replaced (JMD): Still, the R^2 of the regression that...	22
	Deleted (JMD): of the whole range	22
30	Replaced (JMD): of smaller temporal scales, like...	23