

## AUTHOR'S RESPONSE

We would like to thank the reviewers for the thoughtful and interesting comments, which we address in detail in the revised version of our manuscript. Below you find the complete list of the reviewer's comments, our replies and the respective changes made, as well as a marked-up version of the revised manuscript for your consideration.

## REPLIES AND CHANGES IN MANUSCRIPT

### REFEREE COMMENT #1

Comment:

*The paper is interesting and well written.*

Reply:

We highly appreciate your positive evaluation!

### **(1.1) "Compound triggering"**

Comment (1.1):

*However, being based on modelling results (that, as also the authors acknowledge, is an oversimplification of reality) the assessments presented in the results and discussion section are somehow speculative.*

*I believe that less emphasis (i.e. by not mentioning it in the title, for instance) should be given to the so-called "compound triggering concept" that, in my perspective, is quite obvious and possibly over-rated. As matter of fact, Authors have honestly demonstrated (and clearly synthetized in Fig. 7) that in the majority of the debris flows cases they have considered there is a "dominant" trigger (which in most cases is, as usual, precipitation). Thus, despite their modelling effort, I have the feeling that still it is impossible to demonstrate/quantify, without having field monitoring data, the extent to which the other factors where co-influent at the time of triggering.*

Reply (1.1):

In general we agree with this observation and we will put less emphasis on the term „compound trigger concept“, also removing it from the title. Indeed, for the majority of debris flows precipitation has been identified as the "dominant" trigger. Yet, our results (see especially figure 6) clearly suggest how an analysis based on precipitation only is not sufficient for regions as the inner Pitztal and that complementary hydrological information helps to better understand the debris flow initiation process.

We also agree that it is not entirely possible to quantify the extent to which the factors were co-influent (which is the reason why we used three indicative classes of relevance (high, moderate, low) rather than providing numerical values). We would nevertheless here also like to put this

comment into some other perspective. Of course a model (actually any model) describing environmental systems is subject to various sources of uncertainty, which were also quantified in our analysis. The actual problem here, from our point of view, is not the model per se as it captures the main features and dynamics of the hydrological response relatively well. Rather, we think that much of the inconsistencies are driven by epistemic errors in the available precipitation and temperature data, linking to the ever-recurring problem that debris flows are highly localized and thus need highly localized information on these variables –which are typically not available. The same problem will therefore be present in any type of debris flow initiation study, no matter if a model is used or not.

#### Changes in manuscript (1.1):

We removed the terms “compound” and “complementary” triggers and changed the main emphasis towards temporally varying roles of different triggers throughout the manuscript.

#### **(1.2) Probabilistic concept**

##### Comment (1.2):

*In general Figure 4 - together with fig. 5 (and other similar graphs and plots provided in supplementary material) are the “key” to estimate how significant are the Authors findings. However, there is little or no description and discussion in the paper about the NON-EVENT days. It is actually quite clear already from Fig. 4, that the days with debris flows are not that much different (in terms of the analyzed parameters) from many other days in the series. So, please, integrate the discussion.*

##### Reply (1.2):

We are not entirely sure to fully understand the reviewer’s comment here. Does the comment mean that (a) it is not clear how we include the non-event days in our analysis or that (b) we should discuss why no debris flows have been observed on days with system variables similar to those on debris flow days?

If (a), we have realized in discussions that the probabilistic concept (cf. Berti et al., 2012) used in our paper may not be obvious at first. The key figure is figure 5, where we compare the system variables on the event days (i.e. date of event no. 1, no. 2, etc.) with the marginal distribution of the variables (May 15<sup>th</sup> to October 15<sup>th</sup>, 1953-2012). We will try to clarify this in the revised paper.

Thereof, non-event days are always included in our analysis, as these days (together with the event days) were used to calculate each system variable’s marginal distribution, which is plotted on the upper x-axis of the plot. For all events where our interpretation is marked with a ++-confidence, the conditional probability is significantly increased, i.e. the relevant system variable at debris flow occurrence is substantially different from non-debris flow triggering days. For example, for the events identified as primarily triggered by snowmelt (and interpreted with ++-confidence), 16% of the debris flow events (4 out of 25) were observed when snowmelt was more than ca. 7 mm/d, while the marginal probability for such snowmelt to occur would be only 1%. In Bayesian terms this means that the posterior (conditional) probability of a debris flow to occur if the snowmelt exceeds the threshold value would be 16 times higher than the prior probability (in absolute numbers: 4.5% vs. 0.28%;  $p(\text{Fisher's exact test}) = 0.000095$ ).

Of course, the posterior probabilities for the events interpreted with less confidence, are not as clear. For rainfall, this is due to the temporal (and also spatial) averaging, as we have outlined in the paper (page 11, lines 6-16; page 15, line 32 – page 16, line 6). Here our interpretations were also

based on the absence of other system variables that were notably increased (i.e. neither high snowmelt, nor high antecedent soil moisture), which indirectly again considers the non-event days, as the assessment of “no high snowmelt” or “no high antecedent soil moisture” is based on the respective marginal distributions. Conversely, a non-clear attribution of a dominant trigger points, besides potential effects the influence of epistemic errors, towards compound triggering, which we indicated in table 2 by listing triggers by their relevance and visualized in figure 6.

Please note, that due to the limited sample size and the focus of the paper not being on providing probabilities for debris flow occurrence (and thus a blueprint for a prediction model), but to analyse the event’s triggering conditions, we did not explicitly provide the posterior probabilities (as demonstrated above) in the paper for our detailed analysis (fig. 5 resp. table 2).

If (b), we on purpose did not address this issue since (as stated above) providing posterior probabilities is not the key focus of our paper. Of course on the majority of days no debris flows occurred although the system variables have been similar to those on debris flow days. This can largely be attributed to non-hydrological factors such as sediment availability. Actually, a Bayesian approach (fig. 6) explicitly considers the fact that not all potentially triggering events do lead to debris flow initiation (cf. Berti et al., 2012, page 16-17).

#### Changes in manuscript (1.2):

We clarified our approach by adding two passages to section 3.2 (p.9, l.4-7 and p.10, l.4-10).

Also, we added the term “marginal distribution” and a reference to section 3.2 to figure 5 (now figure 6).

### **(1.3) Observed vs. measured runoff**

#### Comment (1.3):

*Moreover, Fig. 5, plot “f” clearly indicated substantial difference between the modelled and recorded runoff on 3 out of 6 debris flow events during which observed runoff was available. I believe that, also this fact, deserves some comments/discussion.*

#### Reply (1.3):

While we did address this issue shortly on page 16, lines 9-11, we agree that this fact deserves a more detailed discussion, which we will include in the revised version of the paper.

#### Changes in manuscript (1.3):

We added a section discussing the difference between modelled and recorded runoff to the discussion, page 17, lines 15-20.

### **(1.4) Exceedance probabilities time period**

#### Comment (1.4):

*I also somehow question the fact that (as mentioned in page 8, lines 24 to 28) the exceedance probability of precipitation was analyzed over the limited period May-October. This choice should be more clearly explained/justified. Also: (i) it is not clear if this probability is based only on the 15 may-15 oct period of years with debris flows or – rather – of any year in the series. (ii) Is May 15 as lower*

*limit correct ??, as the plots in fig 4 and supplementary material, seem to start in march 15. Please check.*

Reply (1.4):

The May 15<sup>th</sup> to October 15<sup>th</sup> period represents the typical debris flow season in an Alpine environment (e.g. Stoffel et al., 2011), i.e. the period in which debris flows have been reported for the study region. As we base our analysis on comparing the system variables (precipitation, snowmelt, etc.) of days with debris flow occurrence with the marginal distribution of these variables, an analysis only comprising the debris flow season to generate the data for the general distributions was found more applicable. This was stated in the original paper (page 8, lines 24 to 27) as: “Due to the generally very low occurrence probability of 25 debris flow events (i.e. 25 events over 60 years), which potentially may in the following lead to instable and overly discontinuous statistical models, we limited the definition of exceedance probabilities (and all other probabilities estimated hereafter) to the period of the year in which all debris flow events occurred [...]”. We will further clarify this.

(i) The probability is based on all days of the May 15 – Oct 15 period in all years, i.e. 1953-2012, as stated in the caption of figure 5. We will add this to the text.

Changes in manuscript (1.4):

To address these two issues we changed the sentence quoted above to: “[...] we limited the definition of exceedance probabilities (and all other probabilities estimated hereafter) to the period of the year in which all debris flow events occurred (**“debris flow season”**), i.e. from May 15<sup>th</sup> to October 15<sup>th</sup>, **1953-2012.**”

Reply (1.4) continued:

(ii) Yes, May 15<sup>th</sup> is correct. We provide the time series plots (fig. 4) from March 15<sup>th</sup> to show a more “complete” picture, i.e. start and amount of rainfall and snowmelt, as this does subsequently considerably influence the antecedent soil moisture of the corresponding year. (We did not include late fall and winter, though, as the information of how much snow fell during these seasons is already implicitly expressed in the snowmelt data and would only have decreased graph readability). However, we did not realize that this indeed leads to some confusion. We will thereof add a note to the caption of figure 4 to clarify this issue.

Changes in manuscript (1.4) continued:

We added the following sentence to the caption of figure 4: “Please note that the plots display the period March 15th to October 15th to depict the start and amount of rainfall and snowmelt, however, the analysis (Figs. 6 and 7) is based on the period May 15th to October 15th.”

## **(1.5) Picture of study area**

Comment (1.5):

*At least one Figure (picture) showing the physiographic setting of the study area should be added.*

Reply (1.5):

This is a great suggestion! We agree and will include a picture in the revised paper.

Changes in manuscript (1.5):

We added figure 2 showing a picture of the inner Pitztal. (Subsequently, Figs. 2-7 have been renumbered to Fig. 3-8).

**(1.6) Abbreviations Figure 3**

Comment (1.6):

*In caption of Figure 3, please include descriptions of Abbreviations (now, the reader is posted to Table1 and sect.2.2, thus making it difficult to follow in case – during editorial setup - these elements are placed in different pages )*

Reply (1.6):

We agree and, as suggested by reviewer No. 2, will include all abbreviations used in this figure resp. section 3.1 in Table 1.

Changes in manuscript (1.6):

We added the table as Table 1a, in which we list all model storages and fluxes. Subsequently, in figure 3 (now figure 4), we refer to Table 1 only, in which every abbreviation used in the figure is listed.

**(1.7) Figure 6**

Comment (1.7):

*Figure 6 should, in my opinion, be eliminated, as it does not really add much real information, as the concept of combined probability is quite easily understandable even without such scheme.*

Reply (1.7):

We would strongly prefer to keep this figure in, as it provides an intuitive visualization of the potential (simultaneous) influences of the different hydrological conditions at the occurrence of debris flows in the region. As it does not include any interpretation, Fig. 6 (now Fig. 7), is actually very close to the data and summarizes the outcomes of our modelling efforts.

Changes in manuscript (1.7):

None.

## REFeree COMMENT #2

### Comment:

*The paper presents results of the analysis of potential factors responsible for debris flow initiation. The study explores possibilities of the use of semi-distributed conceptual rainfall-runoff modelling results to identify possible critical values of triggering factors which could indicate or lead to occurrence of debris flows. The authors use measured (e.g. rainfall data) and modelled factors (e.g. snowmelt, different underground storages) and try to identify their potential role as debris flow triggering factors in view of corresponding exceedance probabilities.*

*I find the manuscript in line with aims and scope of HESS. Generally, the paper is well structured. However, there are some issues that need to be solved in order to improve the presentation and discussion of the results.*

### Reply:

We appreciate the reviewer's generally positive evaluation and tried to address his/her concerns as completely as possible in the replies below and by adjusting the relevant parts of the manuscript.

### **General comments:**

#### **(2.1) Geological setting**

### Comment (2.1):

*While the main topic of the paper are triggering factors related to hydrological conditions, the authors should give some stress (in the Introduction section) also to other possible factors especially related to geological or hydrogeological conditions. These are only briefly mentioned on Page 3 (line 10). Namely, the geological setting strongly pre-define the possible effects of all the hydrological conditions discussed in the paper.*

### Reply (2.1):

We agree and will add this to the Introduction and Discussion sections. Indeed, it is an inherent strength of our method that – while the geological condition does influence the hydrological condition – the geological condition is not needed as explicit a-priori input. Rather, the role of geology is implicitly encapsulated in the posterior parameter distributions of the calibrated hydrological model. While geology is clearly a highly relevant control on debris flow occurrence, its effect does only become relevant once regions in distinct geological settings are compared. In the study area, the geology is rather homogenous – it is thus reasonable to assume that all parts of that region can be expected to trigger debris flows under similar conditions. However, we fully agree that if the results of this study want to be generalized to different areas, the differences in geology need to be factored in for a meaningful understanding of the underlying processes.

### Changes in manuscript (2.1):

We added a paragraph in the introduction section, explicitly acknowledging the importance of geology and related factors (p.3, l.10-15).

## **(2.2) Model structure and abbreviations figure 3**

### Comment (2.2):

*The Study area and Data presentation (Section 2) as well as the Model structure and model calibration and validation process (Section 3.1) is concise and informative. Additional information on past successful applications of the proposed hydrological model structure for any other purposes (besides the analysis of debris flow triggering) would be helpful. There seems to be some discrepancies in the abbreviations used for the model parameters in section and the ones listed in Table 1 (e.g. metlf, M, Mgalcier etc). If I understand correctly, only free calibration parameters are listed in Table 1. All the model parameters mentioned in section 3.1 and in Figure 3 should be listed together in one place (Table) in order to enable reader easier understanding of the model structure. Otherwise, it is extremely difficult to follow the explanation of the role of different parameters that could be potentially considered as important in view of debris flow triggering analysis (Section 3.2).*

### Reply (2.2):

Modular and flexible modelling strategies, as has the chosen model structure that is functionally equivalent to the wide-spread HBV model and similar models (e.g. Seibert 1999; Seibert and Beven, 2009; Fenicia et al., 2014; Berghuijs et al., 2014; Birkel et al., 2015; Nijzink et al., 2016) and which has been rigorously implemented and tested for the study area of this manuscript.

### Changes in manuscript (2.2):

We added references and some more explanation on p.6, l.22-30.

### Reply (2.2) continued:

We agree that the different abbreviations should be listed in one place. As suggested, we will include all abbreviations of figure 3 resp. section 3.1 in Table 1. As for the discrepancies, we will add Mglacier to figure 3, in which it has been erroneously missing.

### Changes in manuscript (2.2) continued:

We added the table as Table 1a, in which we list all model storages and fluxes. Subsequently, in figure 3 (now figure 4), we refer to Table 1 only, in which every abbreviation used in the figure is listed.

Likewise, we added Mglacier to figure 3 (4). In addition, we added SI and Qobs (which we used in our analysis) to figure 3 (4) and Table 1a for improved clarity (see also comment 2.10).

## **(2.3) Section 4.3, classification of events**

### Comment (2.3):

*The discussion on the relevance of potential triggering factors in Section 4.3 is relatively lengthy and extremely difficult to follow. It seems that most of the discussion relies on the authors pre-knowledge about the particular characteristics of the debris flow events and, unfortunately, many of the statements on authors speculations. I believe author should put more effort in extracting the most relevant information from the data analysis instead of commenting particular events in view of available measured and modelled data. On possible solution could be classification of the events based on some predefined criteria, one of them could be e.g. seasonality, as this could lead to*

*possible easier identification of the relevance of discussed triggering factors during particular debris flow events (e.g. convective storms occur mostly in the late spring, summer or early autumn; snowmelt occurs in spring). Sections (4.3.1-4.3.2) discussing the role of high-intensity rainfall events and snowmelt could in my view directly fit into some predefined classification criteria (e.g. seasonality). The influence of seasonality is indicated in several parts of the manuscript but should be more clearly pointed out. Data shown in Figure 7 and discussion in section 4.3.4 could be very useful for developing further discussion in that direction.*

Reply (2.3):

While we agree that some aspects of our interpretations (e.g. triggering by snowmelt with or without additional rainfall?) remain ambiguous, we think that in the vast majority of cases the results as extracted from the hydrological model did allow a conclusive reading. We clearly provided *indicative* levels of confidence for the interpretation with the “direct support by daily data” column in Table 2, which suggests that for 20 out of 25 debris flow event days, our classification displays quite strong direct support by data. In the case of the interpretation of high-intensity short-duration rainfall as debris flow trigger, we also had information on the 10/15 min. precipitation intensities for several events, which provided additional supporting evidence for our interpretations of high-intensity rainfall as dominant debris flow trigger solely based on the daily precipitation data (described in manuscript on page 11-12). To avoid the notion that the interpretation provided is speculative, we think it is necessary to keep the admittedly quite long description, as only from this description our reasoning for direct data support becomes evident.

Concerning the structure, we agree that a classification based on seasonality would be an equally valid option. However, we on purpose decided on the structure at hand based on the dominant trigger (as derived from the results) since this appeared as the most logical structure for us. Of course, convective storms most commonly occur during late spring / summer / early autumn, which is the typical debris flow season. However, in such a high alpine environment, snowmelt does also occur in summer oder autumn, same as long-lasting rainfall. In these cases we believe this should be listed at the same place where snowmelt in spring / long-lasting rainfall in autumn is discussed since the snowmelt / rainfall, rather than the season, would be the most important characteristic. In any case, as the reviewer points out, the dominant trigger and seasonality are closely connected in our study area. As suggested, we will elaborate on this in more detail and more clarity in the revised manuscript.

Changes in manuscript (2.3):

We added an explicit discussion of the effects of seasonality in snow melt/high-intensity precipitation (p.13, l.9-11; p.14, l.21-24; p.16, l.8-11) and changed section title 4.3.4 to “**Seasonally varying importance of the different trigger contributions**” to better reflect the seasonally varying roles.

## **(2.4) “Compound triggering”**

Comment (2.4):

*Although the proposed approach of using semi-distributed hydrological model in combination with relatively scarce data is interesting, my overall concern is, that the complementary effect of different triggering factors is not clearly demonstrated (the so called “compound triggering concept”). Namely, in many parts of the manuscript, authors clearly state that for many debris flow events, only single*



*triggering factor was recognized as the prevailing one. It seems the complementary effect of different triggering factors has much smaller role as the authors try to present.*

Reply (2.4):

We agree that the “compound triggering concept” has not been adequately demonstrated in the paper. As also suggested by the first reviewer, we will significantly reduce the emphasis on this concept, while simultaneously better mapping out the complementary effect when applicable. For this reason, we also will skip the phrase “compound trigger concept” in the title.

Changes in manuscript (2.4):

We removed the terms “compound” and “complementary” triggers and changed the main emphasis towards temporally varying roles of different triggers throughout the manuscript.

**Specific comments:**

**(2.5) Term “Hydrological disposition”**

Comment (2.5):

*Page 2, Line 5: What is meant by “hydrological disposition”?*

Reply (2.5):

The term “hydrological disposition” was adopted from Kienholz (1995) and refers to the antecedent hydrological conditions of a watershed or hillslope. This was also mentioned in the original paper (page 2, lines 11-12) as: “Yet, little is known about the influence of other factors such as snowmelt or the antecedent soil moisture, which may increase a catchment's susceptibility for debris flow initiation (“the disposition concept”; Kienholz, 1995).” We did not realize that this term is not commonly known (probably mostly by the natural hazard community) and will therefore clarify this in the revised manuscript.

Changes in manuscript (2.5):

We added a short clarification on page 2, lines 3-4, reading “hydrologic disposition (**i.e. the general wetness state**)”.

Also, we changed the sentence quoted above, so that it now reads: “Yet, little is known about the influence of other factors such as snowmelt or the antecedent soil moisture, which may increase a catchment's susceptibility for debris flow initiation by reducing the additional water input needed to trigger a debris flow (“the disposition concept”; Kienholz, 1995). “

**(2.6) Hourly rainfall thresholds**

Comment (2.6):

*Page 11, lines 12: I believe it is not useful to discuss possible hourly threshold rainfall intensities derived from daily rainfall data.*

#### Reply (2.6):

Our explanation here was insufficient and thus seems to have sparked a bit of confusion. The hourly rainfall intensities are not “derived”, which would imply some degree of uncertainty around them. Rather, they are the physically lowest possible limit to hourly rainfall intensities. In other words, e.g. during a day with observed rainfall of 24 mm/d there must be at least one hour during which rainfall intensities are equal or exceed 1 mm/hr. We believe that the discussion with perspective to hourly rainfall intensities adds some value here, as debris flows are mostly linked to spatially and temporally highly localized events. Demonstrating and illustrating that the observed daily intensities also map to sufficiently high intensities (without further assumptions or uncertainties involved) to exceed previously estimated hourly thresholds therefore helps in our opinion to provide a more robust justification of our interpretation.

#### Changes in manuscript (2.6):

We added a clarification on p.9, l.20-22.

### **(2.7) Precipitation, seasonality in triggering**

#### Comment (2.7):

*Page 13, line18: Related to general comments above, could precipitation be generally considered as a factor of low relevance for debris flow triggering during some seasons or maybe months?*

#### Reply (2.7):

Yes, for the study region we definitely think that the data and the results presented in the manuscript strongly suggest that there are situations, when precipitation alone is maybe a necessary, but probably not sufficient factor for debris flow initiation. As discussed in the previous comment concerning seasonality, we will clarify this in the revised version.

#### Changes in manuscript (2.7):

See replies to comment 2.3.

### **(2.8) Convective cells**

#### Comment (2.8):

*Page 13, lines 19-20: Do authors have any data that would support the speculations about the occurrence of convective cells?*

#### Reply (2.8):

The lines in question discuss the occurrence of event no. 2 and 20, which we interpreted as triggered by snowmelt. As we write, however, “the absence of observed precipitation and – in case of No. 2 – only moderate maximum temperature, suggests that precipitation is likely to be of low relevance [...], although the occurrence of small convective shower cells cannot be fully dismissed.” We disagree that this is a speculative statement as we merely do not – and, given the possibility of epistemic observational errors, – cannot exclude the possibility of the occurrence of convective (i.e. small scale) shower cells, as the occurrence of un- / underrecorded convective rainfall is a common debris flow

trigger (section 3.2.1; see also the study mentioned of Borga et al. (2014) as pointed out on page 16, lines 4-6). This uncertainty is also reflected in the “direct support by daily data” column, where we marked no. 20 with “strong support” (as considerable snowmelt was recorded), but no. 2 only with “moderate support” (as snowmelt may not have been sufficient for debris flow initiation).

Changes in manuscript (2.8):

None.

**(2.9) “Compound triggering” example**

Comment (2.9):

*Page 14, line 14: In my view, the complementary nature of triggering factors is not so evident or significant as the authors try to present. Could they clearly demonstrate (e.g. for a particular debris flow event) possible evidences of the “complementary” effect?*

Reply (2.9):

See reply to general comment on “compound triggering concept”. We agree and will adapt this in the revised manuscript.

Changes in manuscript (2.9):

See reply to general comment on “compound triggering concept” (comment 2.4).

**(2.10) Miscancelous**

Comments (2.10):

*Page 1, lines 24-27: The last sentence of the abstract is extremely long, contain too much information and is consequently unclear. I suggest to rewrite and shorten the sentence.*

*Page 3, line 17: “Meteorological conditions” instead of “meteorological forcing”?*

*Page 8, line 13: Related to my general comment on presentation of model parameters. What is parameter SI? As far as a can see, here it is mentioned for the first time and its explanation is give in line 9 (Page 9).*

*Page 9, line 12: ... on days when a specific variable ... (add when).*

*Figure 5: The meaning of red vertical lines should be explained in the figure’s caption.*

Reply (2.10):

We agree and will rewrite / include information as suggested.

Changes in manuscript (2.10):

As suggested we replaced “forcing” with “conditions” on page 3, now line 24, and added “when” on page 9, now line 34.

We rewrote the last sentence of the abstract (page 1, now lines 24-25) as: „This highlights in particular the relevance of snowmelt contributions and the switch between mechanisms during early- to mid-summer in snow dominated systems.”

Parameter  $SI$  is the total liquid water availability, i.e. the antecedent soil moisture  $S_u$  plus the water input (rain, melt) of the respective day. We added  $SI$  to figure 3 (now figure 4) as well as Table 1a for improved clarity.

We added “The days where a debris flow event has been documented are marked with red vertical lines” to the caption of figure 4 (now figure 5).

## REFeree COMMENT #3

### Comment:

*This paper aimed to clarify dominant triggers and their interactions that can initiate debris flow in a snow dominated mountainous area. The authors rigorously investigated results generated by their semi-distributed hydrological model as well as observed hydrometeorological variables and deduced several concepts on mechanism of debris flow initiation. Their reasoning on the concepts is interesting and deserves thoughtful consideration. I have several comments that are hopefully helpful for their further advancement.*

### Reply:

We appreciate the reviewer's generally positive evaluation and tried to address his/her concerns as completely as possible in the replies below and by adjusting the relevant parts of the manuscript.

### **Major comments:**

#### **(3.1) Spatial differentiation**

##### Comment (3.1):

*Their aim was to identify triggering factors for debris flow initiation, and they classified past debris flow events into several groups of which trigger is different each other. Their attempt looks successful within the framework used in this study. However, as they realized and discussed in 4.3.4, their framework is based on the semi-distributed model, and thus it falls short of capability in identifying the differences between locations, while debris flow depends on the hydrological, meteorological and geographical conditions at the specific location of their initiation. For example, in 4.3.3, they discussed a difference between the events occurred in lower elevations and in higher elevations, and found the reason in the difference of soil moisture conditions in relation to the difference of surface soil layer. It may be true, but is a matter of speculation. This example clearly shows a limitation of the framework used in this study. A spatial explicit modeling in combination with a semi-distributed model may be helpful to give a solution to this question.*

##### Reply (3.1):

We fully agree that our approach lacks spatial differentiation. However, the root cause for this lack of spatial differentiation is not the method (i.e. the model) used. Rather, it is the (un-)available data that limits a meaningful spatial differentiation.

The most crucial meteorological input, namely precipitation, is typically (and also here) not available on a spatially distributed basis, let alone for the actual source area of a specific debris flow. Remotely sensed precipitation will be very valuable to somewhat alleviate that problem in the future, but currently available remotely sensed time series (going back 15 years or so at best) are at this point insufficient given the very low debris flow occurrence frequencies. Given the highly localized nature of precipitation, especially in summer, calibrating a spatially distributed model on the basis of the available time series of – in our case – three measuring stations, all located at the valley bottom, would not generate reliable additional information. Furthermore, the calibration of a more distributed model would be more problematic and – in the case of fully distributed physically-based models – would encounter many other sources of uncertainties (e.g. model/parameter equifinality,

scale of available field observations of physical parameters vs. scale of the modelling application/grid size, the suitability of the model equations for the scale of the applications (e.g. the Darcy-Richards formulation assumes equilibrium over the grid cell, which is only a valid assumption for scales  $< \sim 1\text{m}$  as recently demonstrated by Or et al., 2015), etc.). These limitations have been acknowledged for a quite some time but no real progress to close the gap between simplicity and complexity has yet been made (e.g. Dooge, 1986; Beven, 1989, 2006; Jakeman and Hornberger, 1993; Sivapalan, 2005; McDonnell et al., 2007; Zehe et al., 2007, 2014; Clark et al., 2011, 2017; Hrachowitz and Clark, 2017). Further, the locations of the debris flows' initiation points are not known (the locations indicated in Figure 1 show the "center of deposition"). Thus even if spatially distributed input and model output was available, this could not be readily linked to the debris flows. In summary, while we agree with the reviewer that the applied model type is not perfect, by using a spatially explicit, fully-coupled model we would only trade-in the limitations mentioned by the reviewer against the (equally relevant) limitations listed above.

In spite of the uncertainties involved we believe that the results of our study allow us to gain more insight into debris flow trigger mechanisms, even though spatially distributed information is not available. As a side note, the very same problem does occur for every analysis that deals with debris flows – no matter if purely statistical or with help of a hydrological model (any model!). In statistical analyses, such as the intensity-duration thresholds, the implicit assumption is that the rainfall observed at some point is representative of the rainfall at the source area of a debris flow. This assumption is in many cases likely to be violated. We will add more discussion on this point in the revised version of the manuscript, better highlighting the advantages and limitations of the chosen modelling strategy.

As for our remarks on the difference of soil moisture conditions at lower vs. higher elevations (section 4.3.3, page 14, lines 4-11), we agree that there is some uncertainty around our reasoning. However, we somewhat disagree with the term "speculative" as this suggests that the statements were made without any supporting evidence. Clearly, soil moisture build-up can be subject to spatial differences. However, the large scale pattern in soil moisture dynamics (not necessarily the absolute values, though) are very likely to be similar within a region as the soil acts as a low-pass filter, attenuating a considerable part of the high (temporal and spatial) frequency fluctuations of incoming precipitation (e.g. Oudin et al., 2004; Fenicia et al., 2008; Euser et al., 2015). The gradual soil moisture build-up in feedback with drainage and evaporation in the wet seasons is therefore a metric for the general wetness state of a system. In an extreme, illustrative example, it can be considered highly unlikely that when one location in the system is at the highest soil moisture content of the year, that another location is at the lowest soil moisture content.

#### Changes in manuscript (3.1):

We added a paragraph in the Discussion concerning spatial differentiation (page 17, lines 23-33).

#### **Minor comments:**

#### **(3.2) Model layout**

##### Comment (3.2):

1. *I did not get the whole picture of the semi-distributed model used in this study. It seems to divide a target area into several zones based on elevations as is shown in Figure 3. If so, the model should*

*have different parameter values for each of different zones, while the parameter values seem to be applied uniformly over the entire study area as is summarized in Table 1. Clear and straightforward description on the model structure is strongly recommended.*

Reply (3.2):

As indicated in Figure 3, the snow routine (fluxes P, PI, Ps, M, storage S<sub>snow</sub>) is distributed into 100 m elevation zones, while the remaining processes are modelled on a lumped scale. Please note that a “distribution” does not necessarily mean a distribution of model parameters, but can equally refer to the distribution of moisture accounting (i.e. different values for input variables; e.g. Ajami et al., 2004; Fenicia et al., 2008; Euser et al., 2015), which was done here by elevation-adjusting the observed temperatures with an environmental lapse rate, which allows different snow accumulation and melt dynamics at different elevations. We will outline the model structure more clearly in the revised text.

Elevation-stratification of the snow processes and then lumping the remaining processes is the standard layout of a semi-distributed conceptual hydrological model. When necessary, the lumped part would include parallel components, modelling different hydrologically similar parts of the study area, for example differentiating between plateau, hillslope and wetland (Savenije et al., 2010) or other hydrological response units. In our case, we tested different levels of spatial distribution due to different hydrological response units, including for example a parallel wetland component. This did neither improve model performance, nor notably influence the runoff behavior. Thus we decided to go for the most parsimonious feasible model architecture, resulting in a model that consists of an elevation-stratified snow routine and a hillslope component.

Changes in manuscript (3.2):

We adapted the sentence describing the transition from the snow routine to the remaining water routing to: “Rain (i.e. liquid precipitation P<sub>l</sub>) and meltwater M (**areally-weighted sum from all elevation zones**) directly enter the unsaturated root zone (S<sub>u</sub>), [...]” and added a section on the choice of our modelling architecture to the following paragraph (page 7, lines 14-17).

### **(3.3) Exceedance probability definition**

Comment (3.3):

*2. Lines 22-23 on Page 8: The authors introduce "exceedance probability (Pe)" and several classes based on Pe. I got confused about the classes, in which  $1 \geq Pe > 0.5$  is defined as "high" and  $Pe \leq 0.01$  as "very low". In my understanding,  $Pe \leq 0.01$  is "very high" because Pe is exceedance probability (not NON-exceedance probability). In Result and Discussion part, they use "high" for  $Pe \leq 0.1$  (see Line 32 on Page 11), which is not consistent with their definition.*

Reply (3.3):

Lines 22-23 on page 8 refer to the exceedance probabilities, which would be highest if  $Pe=1$ , and lowest if  $Pe=0$ . However,  $Pe=1$  corresponds to the lowest value ever measured/modelled in the study period (e.g. snowmelt  $M=0$ ), which would suggest a low contribution of the system variable (in this example, snowmelt) to the debris flow triggering, while  $Pe=0$  corresponds to the highest value and would suggest a high contribution. This is stated in the original manuscript on page 9, lines 12-15 as: “On days when a specific variable reached values that correspond with a high exceedance probability

(see above), the relative contribution of this variable to trigger debris flows was classified as having low relevance, while on days with moderate, low or very low exceedance probabilities, the relative contribution of this variable to trigger debris flows were correspondingly classified as having moderate, high and very high relevance.” We will clarify this in the revised version of the manuscript.

Changes in manuscript (3.3):

We rewrote the sentence to: “On days a specific variable reached values that correspond with a high exceedance probability ( $1 \geq P_e > 0.5$ ; see above), the relative contribution of this variable to trigger debris flows was classified as having low relevance, while on days with moderate ( $0.5 \geq P_e > 0.1$ ), low ( $0.1 \geq P_e > 0.01$ ) or very low ( $P_e \leq 0.01$ ) exceedance probabilities, the relative contribution of this variable to trigger debris flows were correspondingly classified as having moderate, high and very high relevance.”



## REFERENCES

- Ajami, N. K., Gupta, H., Wagener, T., and Sorooshian, S.: Calibration of a semi-distributed hydrologic model for streamflow estimation along a river system, *J. Hydrol.*, 298, 112-135, 2004.
- Berghuijs, W. R., Sivapalan, M., Woods, R. A., and Savenije, H. H.: Patterns of similarity of seasonal water balances: A window into streamflow variability over a range of time scales, *Water Resour. Res.*, 50(7), 5638-5661, doi:10.1002/2014WR015692, 2014.
- Berti, M., Martina, M. L. V., Franceschini, S., Pignone, S., Simoni, A., and Pizzio, M.: Probabilistic rainfall thresholds for landslide occurrence using a Bayesian approach, *J. Geophys. Res.*, 117, F04006, doi:10.1029/2012JF002367, 2012.
- Beven, K.: Changing ideas in hydrology – the case of physically based models, *J. Hydrol.*, 105, 157-172, 1989.
- Beven, K.: Searching for the Holy Grail of scientific hydrology:  $Q_t = (S, R, dt)A$  as closure, *Hydrol. Earth Syst. Sci.*, 10, 609-618, doi:10.5194/hess-10-609-2006, 2006.
- Birkel, C., Soulsby, C., and Tetzlaff, D.: Conceptual modelling to assess how the interplay of hydrological connectivity, catchment storage and tracer dynamics controls nonstationary water age estimates, *Hydrol. Process.*, 29(13), 2956-2969, doi:10.1002/hyp.10414, 2015.
- Borga, M., Stoffel, M., Marchi, L., Marra, F., and Jakob, M.: Hydrogeomorphic response to extreme rainfall in headwater systems: flash floods and debris flows, *J. Hydrol.*, 518, 194-205, doi:10.1016/j.jhydrol.2014.05.022, 2014.
- Clark, M. P., Slater, A. G., Rupp, D. E., Woods, R. A., Vrugt, J. A., Gupta, H. V., Wagener, T., and Hay, L. E.: Framework for Understanding Structural Errors (FUSE): A modular framework to diagnose differences between hydrological models, *Water Resour. Res.*, 44, W00B02, doi:10.1029/2007WR006735, 2008.
- Clark, M. P., Kavetski, D., and Fenicia, F.: Pursuing the method of multiple working hypotheses for hydrological modeling, *Water Resour. Res.*, 47, W09301, doi:10.1029/2010WR009827, 2011.
- Clark, M. P., Bierkens, M. F. P., Samaniego, L., Woods, R. A., Uijlenhoet, R., Bennett, K. E., Pauwels, V. R. N., Cai, X., Wood, A. W., and Peters-Lidard, C. D.: The evolution of process-based hydrologic models: historical challenges and the collective quest for physical realism, *Hydrol. Earth Syst. Sci.*, 21, 3427-3440, doi:10.5194/hess-21-3427-2017, 2017.
- Dooge, J. C.: Looking for hydrologic laws, *Water Resour. Res.*, 22, 46-58, doi:10.1029/WR022i09Sp0046S, 1986.
- Euser, T., Hrachowitz, M., Winsemius, H. C., and Savenije, H. H. G.: The effect of forcing and landscape distribution on performance and consistency of model structures: Distribution of forcing and model structures, *Hydrol. Process.*, 29(17), 3727-3743. doi:10.1002/hyp.10445, 2015.
- Fenicia, F., Savenije, H. H., Matgen, P., and Pfister, L.: Understanding catchment behavior through stepwise model concept improvement, *Water Resour. Res.*, 44, W01402, doi:10.1029/2006WR005563, 2008.
- Fenicia, F., Kavetski, D., Savenije, H. H. G., Clark, M. P., Schoups, G., Pfister, L., and Freer, J.: Catchment properties, function, and conceptual model representation: is there a correspondence?, *Hydrol. Process.*, 28(4), 2451-2467, doi:10.1002/hyp.9726, 2014.

Fenicia, F., Kavetski, D., Savenije, H. H. G., and Pfister, L.: From spatially variable streamflow to distributed hydrological models: Analysis of key modeling decisions, *Water Resour. Res.*, 52(2), 954-989, doi:10.1002/2015WR017398, 2016.

Gharari, S., Hrachowitz, M., Fenicia, F., Gao, H., and Savenije, H. H.: Using expert knowledge to increase realism in environmental system models can dramatically reduce the need for calibration, *Hydrol. Earth Syst. Sci.*, 18, 4839-4859, doi:10.5194/hess-18-4839-2014, 2014.

Hrachowitz, M. and Clark, M. P.: HESS Opinions: The complementary merits of competing modelling philosophies in hydrology, *Hydrol. Earth Syst. Sci.*, 21, 3953-3973, doi:10.5194/hess-21-3953-2017, 2017.

Hrachowitz, M., Fovet, O., Ruiz, L., Euser, T., Gharari, S., Nijzink, R., Freer, J., Savenije, H. H., and Gascuel-Oudou, C.: Process consistency in models: The importance of system signatures, expert knowledge, and process complexity, *Water Resour. Res.*, 50, 7445-7469, doi:10.1002/2014WR015484, 2014.

Jakeman, A. J. and Hornberger, G. M.: How much complexity is warranted in a rainfall-runoff model?, *Water Resour. Res.*, 29, 2637-2649, 1993.

Kienholz, H.: Gefahrenbeurteilung und -bewertung – auf dem Weg zu einem Gesamtkonzept, *Schweizerische Zeitschrift für Forstwesen*, 146, 701-725, 1995.

Leavesley, G. H., Markstrom, S. L., Brewer, M. S., and Viger, R. J.: The modular modeling system (MMS) – The physical process modeling component of a database-centered decision support system for water and power management, in: Chow, W., Brocksen, R. W., and Wisniewski, J.: *Clean Water: Factors that Influence Its Availability, Quality and Its Use*, 303-311, Springer Netherlands, 1996.

McDonnell, J. J., Sivapalan, M., Vaché, K., Dunn, S., Grant, G., Haggerty, R., Hinz, C., Hooper, R., Kirchner, J., Roderick, M. L., Selker, J., and Weiler, M.: Moving beyond heterogeneity and process complexity: A new vision for watershed hydrology, *Water Resour. Res.*, 43, W07301, doi:10.1029/2006WR005467, 2007.

Nijzink, R.C., Samaniego, L., Mai, J., Kumar, R., Thober, S., Zink, M., Schäfer, D., Savenije, H. H. G., and Hrachowitz, M.: The importance of topography-controlled sub-grid process heterogeneity and semi-quantitative prior constraints in distributed hydrological models, *Hydrol. Earth Syst. Sci.*, 20, 1151-1176, doi:10.5194/hess-20-1151-2016, 2016.

Or, D., Lehmann, P., and Assouline, S.: Natural length scales define the range of applicability of the Richards equation for capillary flows, *Water Resour. Res.*, 51, 7130-7144, doi:10.1002/2015WR017034, 2015.

Oudin, L., Andréassian, V., Perrin, C., and Anctil, F.: Locating the sources of low-pass behavior within rainfall-runoff models, *Water Resour. Res.*, 40, W11101, doi:10.1029/2004WR003291, 2004.

Savenije, H. H. G.: HESS Opinions "Topography driven conceptual modelling (FLEX-Topo)", *Hydrol. Earth Syst. Sci.*, 14, 2681-2692, doi: 10.5194/hess-14-2681-2010, 2010.

Seibert, J.: Regionalisation of parameters for a conceptual rainfall-runoff model, *Agricultural and forest meteorology*, 98, 279-293, 1999.

Seibert, J. and Beven, K. J.: Gauging the ungauged basin: how many discharge measurements are needed?, *Hydrol. Earth Syst. Sci.*, 13(6), 883-892, doi:10.5194/hess-13-883-2009, 2009.

Sivapalan, M.: Pattern, process and function: Elements of a new unified theory of hydrologic at the catchment scale, in: Encyclopedia of Hydrological Sciences, edited by: Anderson, M. G., John Wiley & Sons Australia Ltd, UK, vol. 1, 193-220, 2005.

Stoffel, M., Bollschweiler, M., and Beniston, M.: Rainfall characteristics for periglacial debris flows in the Swiss Alps: past incidences–potential future evolutions, *Climatic Change*, 105, 263-280, doi:10.1007/s10584-011-0036-6, 2011.

Wagner, T., Boyle, D. P., Lees, M. J., Wheeler, H. S., Gupta, H. V., and Sorooshian, S.: A framework for development and application of hydrological models, *Hydrol. Earth Syst. Sci. Discussions*, 5(1), 13-26, doi: 10.5194/hess-5-13-2001, 2001.

Zehe, E., Elsenbeer, H., Lindenmaier, F., Schulz, K., and Blöschl, G.: Patterns of predictability in hydrological threshold systems, *Water Resour. Res.*, 43, W07434, doi:10.1029/2006WR005589, 2007.

Zehe, E., Ehret, U., Pfister, L., Blume, T., Schröder, B., Westhoff, M., Jackisch, C., Schymanski, S. J., Weiler, M., Schulz, K., Allroggen, N., Tronicke, J., van Schaik, L., Dietrich, P., Scherer, U., Eccard, J., Wulfmeyer, V., and Kleidon, A.: HESS Opinions: From response units to functional units: a thermodynamic reinterpretation of the HRU concept to link spatial organization and functioning of intermediate scale catchments, *Hydrol. Earth Syst. Sci.*, 18, 4635-4655, doi:10.5194/hess-18-4635-2014, 2014.

# The temporally varying roles of rainfall, snowmelt and soil moisture for debris flow initiation in a snow dominated system: ~~the compound trigger concept~~

Karin Mostbauer<sup>1</sup>, Roland Kaitna<sup>1</sup>, David Prenner<sup>1</sup>, and Markus Hrachowitz<sup>2</sup>

<sup>1</sup> Institute of Mountain Risk Engineering, University of Natural Resources and Life Sciences, Vienna, Austria

<sup>2</sup> Water Resources Section, Faculty of Civil Engineering and Geosciences, Delft University of Technology, Netherlands

*Correspondence to:* Karin Mostbauer (karin.mostbauer@students.boku.ac.at)

**Abstract.** Debris flows represent ~~a severe hazard~~ frequent hazards in mountain regions. Though significant effort has been made to predict such events, the trigger conditions as well as the hydrologic disposition of a watershed at the time of debris flow occurrence are not well understood. Traditional intensity-duration threshold techniques to establish trigger conditions generally do not account for distinct influences of rainfall, snowmelt, and antecedent moisture. To improve our knowledge on the connection between debris flow initiation and the hydrologic system ~~and to overcome the above limitations~~ at a regional scale, this study explores the use of a semi-distributed conceptual rainfall-runoff model, linking different system variables such as soil moisture, snowmelt, or runoff with documented debris flow events in the inner Pitztal watershed, western Austria. The model was run on a daily basis between 1953 and 2012. Analyzing a range of modelled system state and flux variables at days on which debris flows occurred, three distinct dominant trigger mechanisms could be clearly identified. While the results suggest that for 68% (17 out of 25) of the observed debris flow events during the study period high-intensity rainfall was the dominant trigger, snowmelt was identified as dominant trigger for 24% (6 out of 25) of the observed debris flow events. In addition, 8% (2 out of 25) of the debris flow events could be attributed to the combined effects of low-intensity, long-lasting rainfall and transient storage of this water, causing elevated antecedent soil moisture conditions. The results also suggest a relatively clear temporal separation between the distinct trigger mechanisms, with high-intensity rainfall as trigger being limited to mid- and late summer. The dominant trigger in late spring/early summer is snowmelt. Based on the discrimination between different modelled system states and fluxes and more specifically, their temporally varying importance relative to each other, ~~rather than their absolute values~~, this exploratory study demonstrates that already the use of a relatively simple hydrological model can prove useful to gain some more insight into the importance of distinct debris flow trigger mechanisms ~~in a compound trigger concept, highlighting~~. This highlights in particular the relevance of snowmelt contributions and the switch between mechanisms during early- to mid-summer in snow dominated systems.

## 1 Introduction

Debris flows are rapidly flowing mixtures of sediment and water transiting steep channels (Hungr et al., 2014) and often represent a severe hazard in mountain regions. In alpine regions the mechanism of debris flow initiation typically ranges from distinct slope failures transforming into a flow like movement to intensive sediment bulking due to channel erosion (e.g. Rickenmann and Zimmermann, 1993; Prancevic et al., 2014). Hereafter we refer to debris flow as channel-based mass flows that can be either triggered from landsliding or channel erosion. In contrast to the effect of a region's geomorphological and geological disposition ~~of~~ debris flows ~~to occur~~ (e.g. Nandi and Shakoor, 2008; von Ruetten et al., 2011) and in spite of significant efforts in the past (e.g. Guzzetti et al., 2008), neither the effect of hydrologic disposition (i.e. the general wetness state) of a specific region at the time of ~~mass movement, and in particular~~ debris flow, initiation nor the actual triggering hydro-meteorological conditions are well understood. Reliable regional predictions of debris flow events so far therefore remain essentially elusive.

There is a widespread consensus that high-intensity, short-duration rainfall is the primary trigger of debris flows in Alpine environments (e.g. Berti et al., 1999; Marchi et al., 2002; McArdell et al., 2007; McCoy et al., 2012; Kean et al., 2013), while longer duration precipitation is of minor, but not negligible importance (e.g. Moser and Hohensinn, 1983; Stoffel et al., 2011). Yet, little is known about the influence of other factors such as snowmelt or the antecedent soil moisture, which may increase a catchment's susceptibility for debris flow initiation by reducing the additional water input needed to trigger a debris flow ("the disposition concept"; Kienholz, 1995).

While antecedent wetness, quantified as pre-storm rainfall, has been widely observed as an important factor for triggering debris flows (e.g. Napolitano et al., 2016), there is little agreement on the specific water volumes and/or time periods required for the build-up of debris flow-relevant antecedent soil moisture (Wieczorek and Glade, 2005). Similarly, there is no consensus on the level of soil moisture, i.e. the water volume stored in near-surface layers of the unsaturated substrate, required to trigger debris flows under different rainfall conditions (Johnson and Sitar, 1990; Montgomery et al., 2009). Essentially omitting the temporally variable yet cumulative influences of evaporation, transpiration and drainage on the soil wetness state, these concepts of antecedent wetness should be treated with caution and may hold only limited information. Interestingly, Aleotti (2004) and Berti et al. (2012) found no significant influence of antecedent rainfall, as a proxy for soil moisture, on the triggering of landslides and debris flows in different regions in Italy. This is somewhat surprising, as ~~we would expect~~ slope failures are to be expected to occur more readily ~~after building up of~~ under situations with elevated pore fluid pressures (Iverson, 2000). Such somewhat contrasting interpretations probably arose from slightly different definitions of antecedent rainfall, which mask what is effectively the role of soil moisture (see discussion in Berti et al., 2012). In the specific cases where the triggering rainfall was restricted to the rainfall on the event day (e.g. Glade et al., 2000), the role of antecedent rainfall was interpreted to be higher than in cases where the definition of events was widened to longer durations (e.g. Berti et al., 2012). However, other research has identified catchments where the antecedent wetness does not have

substantial impact on the triggering of different types of mass movements, including landslides and debris flows (Deganutti et al., 2000; Coe et al., 2008; Ciavolella et al., 2016; Chitu et al., 2017).

Similarly, snowmelt, often combined with rainfall ("rain-on-snow"), is recognized as a common triggering factor of debris flows (Church and Miles, 1987) and shallow landslides (which may subsequently transform into debris flows) (Bil et al., 2015). In spite of this general understanding, there is little systematic effort to quantify its influence and its role may often be under-estimated (Decaulne et al., 2005).

Detailed, direct observations of these two (e.g. Johnson and Sitar, 1990; Coe et al., 2008; Montgomery et al., 2009) and other potentially relevant ~~variables~~system components, such as canopy interception (e.g. Sidle and Ziegler, 2017), are typically not available at sufficient spatial and temporal resolutions. This is ~~especially~~in particular true ~~in~~for debris flow-prone, mountainous environments, and if measurements are available, they are mostly limited to point observations in small, experimental catchments over relatively short time periods, including, if any, only a few debris flow events. Notwithstanding these limitations, estimates of spatial distributions of soil water storage from relatively low-resolution observations or at least relative differences in its spatial occurrence are often used for the identification of locations more susceptible to mass movements, including shallow landslides, and less often, debris flows, than others in regional hazard assessments (cf. Bogaard and Greco, 2016).

Besides liquid water input and subsurface water storage a region's susceptibility to debris flows is also strongly influenced by its landscape and the past evolution thereof (Takahashi, 1981; Rickenmann and Zimmermann, 1993; Reichenbach et al., 2014; Sidle and Ziegler, 2017). More specifically, the type of underlying bedrock and its resistance to weathering are, together with the associated soil formation/erosion processes (i.e. sediment availability), vegetation cover (i.e. reduction of effective rainfall intensities and "reinforcement" of soil) in constant feedback with the resulting topography (i.e. gradient), another first order control on debris flows.

Since the pioneering work of Montgomery and Dietrich (1994), considerable progress has been made in understanding and describing the interplay between the above hydrological and geomorphological/geological susceptibility of hillslopes and small catchments to mass movements based on elegant, spatially explicit, high resolution mechanistic model frameworks (e.g. Dhakal and Sidle, 2004; Simoni et al., 2008; Lehmann and Or, 2012; Mancarella et al., 2012; von Ruetten et al., 2013; Anagnostopoulos et al., 2015). Despite their outstanding value for developing our understanding of the detailed processes and feedbacks involved in the initiation of mass movement events as well as for local predictions of such (mainly shallow landslides) at the study sites, these models have at the present and for the foreseeable future limited value for larger scale applications (cf. Hrachowitz and Clark, 2017). In order for being meaningful descriptions of reality, they need to rely on detailed descriptions of the spatial and temporal natural heterogeneity of both the meteorological ~~forcing~~conditions and the subsurface. For example, Fan et al. (2016) demonstrated the spatial variations in soil properties, without changing other boundary conditions, lead to considerable variations in landslide occurrence characteristics. While ever-improving remote sensing products continue to alleviate the problems of the availability of suitable meteorological data, a meaningful and detailed characterization of the multi-scale subsurface heterogeneity is out of reach for the vast majority of regions

worldwide. Without this information, though, such models cannot be adequately calibrated (i.e. equifinality; Beven, 2006a) nor rigorously tested (i.e. the boundary flux problem; Beven, 2006a), making them problematic to use as debris flow prediction tools at the spatial scales and extent of relevance for operational early-warning systems.

In contrast, efforts to provide meaningful and feasible debris flow prediction tools are largely limited to statistical model frameworks with little explicit consideration of the physical processes involved (e.g. Baum and Godt, 2010; Papa et al., 2013; Berenguer et al., 2015). The vast majority of these applications rely exclusively on the well-established concept of intensity-duration thresholds (e.g. Aleotti 2004, Guzzetti et al., 2007, 2008 and references therein), or apply other probabilistic assessments of rainfall characteristics (Berti et al., 2012; Braun and Kaitna, 2016; Turkington et al., 2016; van den Heuvel et al., 2016). Either approaches work under the implicit conjecture that rainfall is the only hydrological factor controlling debris flow initiation. While this is likely to hold in rainfall dominated, warm, humid climates (e.g. Köppen-Geiger climate classes Af, Am, Cfa, Csb), it may carry substantial uncertainty in cooler, snow or rain-on-snow dominated climates, often characterized by lower precipitation intensities (e.g. Dfa, Dfb, Dsa, Dsb), as both, relatively high-intensity snowmelt in spring to mid-summer and gradual soil moisture build-up through the warm season by persistent, lower-intensity rainfall and snowmelt, can add significant additional liquid water volumes to the subsurface of the system. This very likely leads to much less sharply defined rainfall intensity thresholds for debris flow initiation, as also to some degree reflected in the concept of variable hydrological disposition (Kienholz, 1995).

To circumvent the problem of data scarcity in mechanistic models to a certain degree while at the same time bringing some more process knowledge into the traditional intensity-duration thresholds and antecedent rainfall model approaches, we here analyse the value of describing debris flow initiation as a function of several contributing and potentially complementary hydrological and meteorological variables. To do so, we here explore the potential of zooming out to the macro-scale (cf. Savenije and Hrachowitz, 2017), using a well-constrained, semi-distributed conceptual rainfall-runoff model to analyse and quantify these individual variables and their potentially temporally varying importance as complementary additional contributions for the initiation of debris flows, i.e. a compound trigger concept. Briefly, such a model generates time series of different system state and flux variables, such as soil moisture or snow melt. As these variables explicitly reflect the combined and temporally integrated influences of different interacting individual processes, this approach allows a more complete and detailed picture of the processes involved. For example, as recently emphasized by Bogaard and Greco (2016), using the modelled soil moisture to replace the general concept of antecedent wetness has the advantage of both, explicitly *accounting for* and *integrating* the temporally varying effects of precipitation, soil and interception evaporation, plant transpiration and drainage on the level of water storage in different components of the system (e.g. unsaturated root zone, groundwater). Such a continuous model must not be confounded with previous approaches such as the "antecedent soil water status model" (Crozier, 1999; Glade, 2000), which was designed for porous soils in a maritime climate and only takes an antecedent period of up to 10 days into account.

In this exploratory, proof-of-concept paper we test for a catchment in the Austrian Alps (Köppen-Geiger class Dfb) the hypotheses that time series of system state and flux variables generated with a semi-distributed model, used together with

observed meteorological variables, can contain enough information (1) to discriminate between distinct, ~~complementary contributions~~ contributing factors to debris flow trigger mechanisms, ~~i.e. the compound trigger concept~~, and (2) to identify intra-annual shifts in the relative importance of these distinct mechanisms to understand at which time in the year traditional rainfall intensity-duration thresholds (e.g. Guzzetti et al., 2008) may exhibit reduced predictive power.

## 2 Study area and data

### 2.1 Study area

The Pitztal, situated in south-western Austrian province of Tyrol, is a side valley of the river Inn. The longitudinal inner Pitztal (Figs. 1 [and](#) 2) features a narrow valley bottom with steep hillslopes. The study area (approximately encompassing the inner Pitztal) is about 20 km long in its north-east extension with an average width of 6.5 km, covering an area of 133 km<sup>2</sup>. Only 25% of the study area are forested, while 35% are covered by pasture or natural grassland and the remaining 40% are sparsely vegetated, bare rocks or glaciers (glaciers 2.5%). Elevation ranges from 1093 m a.s.l. at the flow gauge *Ritzenried* up to 3340 m a.s.l. at the mountain ridge. The Pitztal is part of the Ötztal-Stubai-crystalline and mainly consists of para- and orthogneiss rocks mostly overlain by sandy Podzols.

Mean annual precipitation in the inner Pitztal is about 1330 mm a<sup>-1</sup>, of which – on average – 42% fall as snow. The inner-alpine dry valley ranks among the driest regions of the Austrian Alps as it is located in the rain shadow of the Northern Limestone Alps and the main Alpine ridge. Mean yearly runoff totals ca. 930 mm a<sup>-1</sup> (runoff coefficient: 0.7), displaying a nivo-glacial regime with the highest flows in June (river regime definition after Mader et al., 1996).

### 2.2 Data

Available hydro-meteorological data included daily time series of precipitation (P), mean temperature ( $T_{\text{mean}}$ ) and potential evapotranspiration ( $E_p$ ) for the period 1952-2012 ~~and~~ as model input, while daily stream flow data (Q) for the period 1986-2013 were available for model calibration and validation (Fig. [23](#)). The data were provided by national hydrological and meteorological services (HD Tirol, ZAMG) and a hydropower plant operator (TIWAG). Supplementing the daily precipitation sums, 15-min precipitation totals were available for stations *St. Leonhard im Pitztal-Neurur (Tiwig)* and *Taschachbach* from 1987 and 10-min totals for station *St. Leonhard im Pitztal-Neurur (Zamg)* from 2007 [onwards](#). These high-frequency data were in the following used as supporting information to interpret dominant debris flow triggers. The catchment outline and elevation zones for the semi-distributed model were obtained from a digital elevation model with 10 m resolution (Data.gv.at).

The daily precipitation input was calculated as the weighted mean of the stations *Jerzens-Ritzenried*, *St. Leonhard im Pitztal* and *Plangeröß* and – as all stations are located at the valley bottoms – was adjusted for elevation (Valéry et al., 2010; Beven, 2012), using high-resolution gridded vertical precipitation gradients provided by Mergili and Kerschner (2015) for the study area. The temperature data were, likewise, elevation corrected using an environmental lapse rate determined in relation to the



nearby climate station *Innsbruck Flugplatz* (cf. Auer et al., 2007). For the estimation of the potential evapotranspiration, the Hargreaves (1985) equation was applied.

We restricted the hydrological modelling to the relevant study area, specifically adapting the hydrological model to the geomorphologically homogeneous inner Pitztal. We thereby avoided the need to model the extensively glaciated valley head and the outer Pitztal, where no significant debris flow activity was recorded. To do so, daily discharge data from the stations *Pitz- & Taschachbach*, located at the upstream boundary of the study area, were used as additional inflow to the model (Fig. 1). In contrast, daily discharge data from the flow gauge *Ritzenried* at the catchment outlet was used for model calibration and validation. At the stations *Pitz- & Taschachbach* flow is measured in an artificial structure, providing very reliable data. The discharge data from the downstream gauge at *Ritzenried* was plausibility-checked against additional data from station *St. Leonhard im Pitztal*.

In addition, daily snow depth measurements for the whole study period 1953-2012 were available from stations *Jerzens-Ritzenried*, *St. Leonhard im Pitztal* and *Plangeroß*. Annual glacier extent data were obtained from the Austrian Glacier Inventory (Lambrecht and Kuhn, 2007), while annual glacier melt time series from three glaciers in the adjacent Ötztal catchment were accessible for the whole study period (*Hintereisferner*, *Kesselwandferner*), resp. from 1965 (*Vernagtferner*) from the WGMS.

Within the study period, 1953-2012, 81 debris flow events in the inner Pitztal have been documented by the Austrian Federal Ministry of Agriculture, Forestry, Environment and Water Management (BMLFUW) (Hübl et al., 2008). For 43 debris flows (Fig. 1) occurring on 25 individual event days (hereafter referred to as “events”) the date of occurrence was known (Fig. 23) and could thus be used for the detailed analysis of the trigger conditions in this study. For the statistical assessment of debris flow occurrence, however, the full set of 81 debris flow events, i.e. also including those for which only the year or month of occurrence was known, was taken into account.

### 3 Methods

#### 3.1 The hydrological model

To estimate otherwise unavailable hydrological state and flux variables at the time of debris flow occurrences, we implemented a semi-distributed conceptual rainfall-runoff model on a daily basis.

##### 3.1.1 Model structure

Adopting a flexible modeling strategy (Clark et al., 2011; Fenicia et al., 2011; ~~Hrachowitz et al., 2014~~), which has proven highly valuable for many studies worldwide in the past (e.g. Leavesley et al., 1996; Wagener et al., 2001; Clark et al., 2008; Fenicia et al., 2014, 2016; Gharari et al., 2014; Hrachowitz et al., 2014) we customized and extensively tested a range of functionally different model structures and parameterizations (not shown). The most suitable of these tested model structures, which was subsequently used for the study catchment (Fig. 34), has 9 free calibration parameters (Table 1b) and

resembles the wide-spread HBV-type of models, which were previously successfully applied over a wide range of environmental conditions (e.g. Seibert 1999; Seibert and Beven, 2009; Fenicia et al., 2014; Berghuijs et al., 2014; Birkel et al., 2015; Hrachowitz et al., 2015; Nijzink et al., 2016b). All model equations are provided in Table S1 in the Supplementary Material.

Briefly, the model was implemented with a semi-distributed snow routine, stratified into 100\_m elevation zones. In the absence of more detailed data, the volume of water falling as snow (i.e. solid precipitation  $P_s$ ) and eventually stored in the snow pack ( $S_{\text{snow}}$ ) was based on a simple temperature threshold method (e.g. Gao et al., 2017). Due to their minor importance in the snowmelt dominated study catchment (Böhm et al., 2007) and in spite of their potentially distinct accumulation and ablation dynamics, glaciers were included in the snow module by allowing continued release of meltwater ( $M_{\text{glacier}}$ ) after the depletion of the transient annual snow pack at elevations with observed perennial glaciers.

Rain (i.e. liquid precipitation  $P_l$ ) and meltwater  $M$  (areally-weighted sum from all elevation zones) directly enter the unsaturated root zone ( $S_u$ ), where a runoff coefficient ( $C_r$ ) controls the proportion of incoming water directly released as preferential percolation ( $Q_{\text{up}}$ ) to the slow responding groundwater storage ( $S_s$ ) or as influx ( $Q_{\text{uf}}$ ) to a fast responding model component ( $S_f$ ) and the proportion transiently stored as soil moisture in  $S_u$ . Water can then leave  $S_u$  either through an evaporative flux ( $E_a$ ), comprising plant transpiration and evaporation, or through percolation ( $Q_{\text{us}}$ ) that eventually recharges the groundwater storage  $S_s$ . Stream flow is then generated from the combined outflow of  $S_f$  and  $S_s$ , both implemented as linear reservoirs with storage coefficients  $K_f$  and  $K_s$ , respectively.

The model at hand thus consists of a semi-distributed, elevation-stratified, snow routine and a lumped hillslope component. While we tested different levels of spatial distribution due to different hydrological response units, including for example a parallel wetland component, we decided to go for the most parsimonious feasible model architecture, since more complex models did neither improve model performance, nor notably influence the runoff behavior. As flow velocities are very ~~short~~ high, due to the elevated elevation gradients, and flow distances are relatively short, channel routing was considered negligible on the timescale of the implementation. Similarly, interception was neglected due to the limited amount of forested areas.

### **3.1.2 Model calibration and validation**

Model calibration, based on Monte-Carlo sampling with  $10^6$  realizations from uniform prior parameter distributions (Table 1), was performed for 1987-2007. For a robust model that can reproduce several aspects of the hydrological response simultaneously, thereby ensuring consistency of the internal processes (e.g. Gupta et al., 2008; Euser et al., 2013; Hrachowitz and Clark, 2017), a multi-objective calibration approach was applied. This was done by combining three objective functions, i.e. the Nash-Sutcliffe efficiencies (Nash and Sutcliffe, 1970) of flow ( $E_{\text{NS},Q}$ ) and the logarithm of flow ( $E_{\text{NS},\log(Q)}$ ) as well as the volume error of flow ( $V_{\text{E},Q}$ ; Criss and Winson, 2008) into the Euclidean Distance  $D_E$  to the "perfect" model as overall objective function (e.g. Schoups et al., 2005; Hrachowitz et al., 2014; Fovet et al., 2015; Nijzink et al., 2016a):

$$D_E = \sqrt{(1 - E_{NS,Q})^2 + (1 - E_{NS,\log(Q)})^2 + (1 - V_{E,Q})^2} \quad (\text{Eq. 1})$$

In the absence of more detailed information all three objective functions in  $D_E$  were given equal weights. Note that in contrast to the three individual objective criteria,  $D_E=0$  indicates a perfect fit.

The best performing 0.1% of parameter sets in terms of  $D_E$ , roughly corresponding to a performance threshold of 0.75 for each of the three individual performance metrics (see results section), were retained as behavioural solutions. These solutions were subsequently used to construct ensemble solutions and thus envelopes for the modelled variables, reflecting their respective sensitivities to parameter uncertainty.

The period 2007-2012 was thereafter used for post-calibration model testing and evaluation (“validation”; Fig. 23), based on the set of retained solutions and their performance metrics  $D_E$  for that period. In addition, for a post-calibration plausibility check and evaluation of the snow routine at low elevations, we compared the timing of the presence of an observed snow pack (snow present yes/no) at the three climate stations with the modelled timing of the presence of snow storage at corresponding elevations in the model. Note that in the absence of time series of snow density, no more detailed evaluation could be done. For higher elevations we correlated the modelled annual glacier melt dynamics with the annual glacier melt time series from the three glaciers in the adjacent Ötztal valley.

### 3.2 Debris flow initiation analysis

To identify potentially different triggers for debris flow initiation, we then explored a range of hydro-meteorological system variables at days  $t$  when debris flows occurred. These included observed variables, such as daily precipitation  $P(t)$  [ $\text{mm d}^{-1}$ ], daily runoff  $Q_{\text{obs}}(t)$  [ $\text{mm d}^{-1}$ ] and daily maximum temperature  $T_{\text{max}}$  [ $^{\circ}\text{C}$ ], as well as modelled state and flux variables such as unsaturated soil moisture  $S_u(t)$  [ $\text{mm}$ ] to account for antecedent moisture ~~input~~, daily snowmelt  $M(t)$  [ $\text{mm d}^{-1}$ ], daily runoff  $Q_{\text{mod}}(t)$  [ $\text{mm d}^{-1}$ ] and the total liquid water present at the near-surface, calculated as  $S_l(t)=S_u(t)+P_l(t)\Delta t+M(t)\Delta t$  [ $\text{mm}$ ], which is to be interpreted as an upper bound of near-surface storage as it does not consider drainage and evaporation at that time step.

For the observed system variables  $P$  (1953-2012) and  $Q_{\text{obs}}$  (1986-2012), analysis was based on the actual values recorded at the respective observation points for the day of occurring debris flows. Specifically, this involved use of  $Q_{\text{obs}}$  for each debris flow event measured at the gauge *Ritzenried*. For precipitation, the individual raw values recorded at the three weather stations *Jerzens-Ritzenried*, *St. Leonhard im Pitztal* and *Plangeroß* were used for initial analysis to account for and illustrate the spatial variation in precipitation within the catchment. The subsequent estimation of debris flow probabilities (see below) was then based on the elevation-corrected, weighted areal mean precipitation. For temperature, the aeriially weighted (according to elevations zones) temperature distributions as estimated from applying environmental lapse rates (see section 2.2) were used.

The analysis of the modelled system variables was based on the behavioural parameter sets, which were used to generate distributions of values for each variable at the days of debris flow events occurring. The material presented hereafter is

limited to  $M$ ,  $S_u$ ,  $S_l$ , and  $Q_{\text{mod}}$ . All other tested variables (not shown), such as groundwater storage, recharge, preferential flow or evaporative fluxes did not exhibit distinguishable patterns with respect to debris flow events; some of which may be attributed to poorly identifiable parameters and the resulting elevated uncertainty in these variables, i.e. the variation of the modelled variables generated with the suite of behavioural parameter sets was so high that for the same debris flow event this variable could take on either, a low or a high value, depending on which parameter set is considered (for examples see Supplementary Material Fig. S1a-b). Note that the state variables  $S_u$  and  $S_l$  were normalized and the analysis thus based on their respective relative water content. This allowed more insights as the model parameter representing the absolute storage capacity of  $S_u$ , i.e.  $S_{u,\text{max}}$ , varied within some range, which in turn is likely to mask relevant pattern (cf. Fig. S1c-d).

To be able to assess the variables' magnitude at debris flow initiation, we compared the magnitude of each system variable with the marginal distributions (i.e. distributions generated with the time series of all days, namely event days and non-event days; see also below) of the respective variables, allocating an "exceedance probability" to each value, rather than looking at the absolute numbers. Due to the generally very low occurrence probability of debris flow events and gaps in the data records (i.e. 25 well documented events over 60 years), which potentially may in the following lead to instable and overly discontinuous statistical models, we limited the definition of exceedance probabilities (and all other probabilities estimated hereafter) to the period of the year in which all debris flow events occurred ("debris flow season"), i.e. from May 15<sup>th</sup> to October 15<sup>th</sup>, 1953-2012. In other words, all probabilities reported hereafter are conditional on that period.

To facilitate a more objective and quantifiable comparison of the system variables, classes of exceedance probabilities were defined for the individual variables, with exceedance probabilities  $1 \geq P_e > 0.5$  hereafter loosely referred to as high,  $0.5 \geq P_e > 0.1$  as moderate,  $0.1 \geq P_e > 0.01$  as low, and  $P_e \leq 0.01$  as very low, i.e. corresponding to extreme events and for precipitation to a lower bound of heavy precipitation events (cf. Schimpf, 1970). ~~Due to the generally very low occurrence probability of debris flow events (i.e. 25 events over 60 years), which potentially may in the following lead to instable and overly discontinuous statistical models, we limited the definition of exceedance probabilities (and all other probabilities estimated hereafter) to the period of the year in which all debris flow events occurred, i.e. from May 15<sup>th</sup> to October 15<sup>th</sup>. In other words, all probabilities reported hereafter are conditional on that period.~~

These classes of exceedance probabilities were subsequently used to systematically analyse if patterns of different dominant trigger mechanisms emerge from the observed and modelled data, i.e. daily precipitation  $P$  as a proxy of short duration, high intensity moisture input to the system, snowmelt  $M$  and  $S_u$  as a metric of longer duration, lower intensity moisture input to the system, under different hydrological conditions. Due to the unavailability of historical sub-daily precipitation totals before 1987, the daily precipitation  $P$  was here used for the overall analysis as a proxy for precipitation intensities. Here the  $P_e < 0.01$ , equivalent to  $P = 45 \text{ mm d}^{-1}$ , implies a ~~lower~~ lowest physically possible limit for precipitation intensity of approximately  $1.9 \text{ mm h}^{-1}$ , (i.e. during at least one hour of that day a precipitation intensity of  $1.9 \text{ mm h}^{-1}$  was reached or even exceeded) which is consistent with the intensity thresholds for 24 h rainfall that were observed to trigger shallow landslides and debris flows in mountain areas as reported by Guzzetti et al. (2008). The high-resolution precipitation data (available from 1987 onwards; see sect. 2.2) allowed, at least to some degree, a plausibility check of the identification of

observed high-intensity rainfalls based on daily rainfall records during that time period. Please note, however, that exact exceedance probabilities for high-resolution precipitation data could not be determined due to the limited time frame of high-resolution data availability. Thus we provide conservative estimates of minimum exceedance probabilities.

Using the exceedance probabilities of the three system variables daily precipitation  $P$ , daily snowmelt  $M$  and relative soil moisture  $S_u$  at the days when debris flows occurred, then allowed together with a qualitative consideration of the total liquid water availability  $S_l$ , daily runoff  $Q_{\text{mod}}$  (and  $Q_{\text{obs}}$ ) and daily maximum temperature  $T_{\text{max}}$  (as an indicator for the likelihood of a local convective rainfall event), a relative assessment of which variable contributed most to trigger an event -and how the relative influences of the three individual variables varied over time, depending on the prevailing meteorological conditions. On days when a specific variable reached values that correspond with a high exceedance probability ( $1 > P_e > 0.5$ ; see above), the relative contribution of this variable to trigger debris flows was classified as having low relevance, while on days with moderate ( $0.5 > P_e > 0.1$ ), low ( $0.1 > P_e > 0.01$ ) or very low ( $P_e < 0.01$ ) exceedance probabilities, the relative contribution of this variable to trigger debris flows were correspondingly classified as having moderate, high and very high relevance.

By comparing the values reached at debris flow initiation with the marginal distribution of the variables we applied a probabilistic concept (cf. Berti et al., 2012), which does not only consider the days where debris flows were reported, but also the non-event days. This, in turn, allowed an assessment of whether the respective variables were significantly increased, and thus likely to be (partially) responsible for the debris flow triggering. Please note that we on purpose do not provide any explicit posterior probabilities for debris flows in this paper, due to the limited sample size and the focus of the paper not being on providing probabilities of debris flow occurrence (and thus a blueprint for a prediction model), but to analyse the event's triggering conditions.

## 4 Results and Discussion

### 4.1 Hydrological model

The retained behavioural parameter sets (see posterior parameter distributions in Table 1) generated model outputs that reproduced the features of the hydrological response in a generally plausible way, as can be seen in Fig. 45 for some selected years and in Supplementary Material Fig. S2 for the remaining years of the study period. This is on the one hand reflected in the rather elevated performance metrics for stream flow. The models' best fit overall objective function reached  $D_E=0.25$  for the twenty year calibration period, with  $E_{NS,Q}=0.85$ ,  $E_{NS,\log(Q)}=0.93$ , and  $V_{E,Q}=0.81$ . The model similarly produced adequate performance levels for the validation period with  $D_E=0.26$  (5/95<sup>th</sup> percentiles  $0.25 \leq D_E \leq 0.31$ ),  $E_{NS,Q}=0.86$  ( $0.82 \leq E_{NS,Q} \leq 0.87$ ),  $E_{NS,\log(Q)}=0.93$  ( $0.91 \leq E_{NS,\log(Q)} \leq 0.93$ ) and  $V_{E,Q}=0.79$  ( $0.76 \leq V_{E,Q} \leq 0.80$ ). On the other hand, post-calibration evaluation (cf. Hrachowitz et al., 2014) also indicated that the overall pattern in snow and glacier dynamics, which the model was not trained for, were adequately captured. Comparing the information whether snow has been present (yes/no) at the three climate stations *Jerzens-Ritzenried*, *St. Leonhard im Pitztal* and *Plangeroß* with the model's results at corresponding elevations shows that the (non-)presence of snow corresponds reasonably well, with correlation coefficients reaching  $r=0.77$ ,

0.87 and 0.88, respectively (with  $p < 0.001$  throughout), for the best model fit. Likewise, the observed glacier melt dynamics correlated well with the modelled snowmelt dynamics at higher elevations with best fit model's correlation coefficients  $r = 0.85$ ,  $0.81$  and  $0.91$  ( $p < 0.001$  throughout) for the *Hintereisferner*, the *Kesselwandferner* and the *Vernagtferner*, respectively.

#### 4.2 System variables at debris flow initiation

In the following the values of hydro-meteorological variables at the days of debris flow occurrences were extracted from the observed and modelled time series. On 3 out of the 25 days with debris flows (No. 7, 11, 19), the observed precipitation at all three rain gauges exceeded  $P = 45 \text{ mm d}^{-1}$ , corresponding to a precipitation exceedance probability  $P_e = 0.01$  over the study period (Fig. 5a6a). This threshold was exceeded for at least one gauge on 2 further event days (No. 21, 24). In addition, precipitation recorded at all three gauges reached exceedance probabilities  $0.01 < P_e \leq 0.1$  ( $\sim 17 \text{ mm d}^{-1}$ ) for 3 event days (No. 1, 16, 22) and for at least one gauge on 4 days (No. 3, 12, 23, 25). On 9 more event days precipitation with  $0.1 < P_e \leq 0.5$  was recorded for at least one gauge, while on 4 days (No. 2, 8, 9, 20) no precipitation was observed at any gauge.

High modelled snowmelt rates with  $P_e \leq 0.01$  for almost all behavioural solutions, corresponding to  $M = 15 \text{ mm d}^{-1}$ , occurred on 4 event days (No. 8, 9, 10, 17; Fig. 5e6c), while snowmelt plotted between  $0.01 < P_e \leq 0.1$  for one event (No. 20). All remaining events, except for No. 25, for which no snowmelt was generated by the model, occurred on days with at least some degree of snowmelt.

Similarly, the mean modelled antecedent soil moisture  $S_u$  (Fig. 5d6d) from behavioural parameter sets was exceptionally high on 4 event days (No. 8, 9, 10, 13), i.e. at each event day at least 75% of the behavioural solutions indicate  $P_e \leq 0.01$ , and at least moderately elevated on at least 7 additional days (No. 6, 7, 11, 12, 18, 19, 20). For completeness and as support for the following analysis, the maximum daily temperature ( $T_{\max}$ ) distribution over all elevation zones in the catchment (Fig. 5b6b), the near-surface total liquid water storage  $S_l$  (Fig. 5e6e), the observed and modelled runoff  $Q_{\text{obs}}$  and  $Q_{\text{mod}}$  (Fig. 5f6f), respectively, are also shown. While  $S_l$ ,  $Q_{\text{obs}}$  and  $Q_{\text{mod}}$  broadly reflect the combined pattern of  $P$ ,  $M$  and  $S_u$ , the pattern of  $T_{\max}$  suggests that almost 50% of the events (11 out of 25) occurred on days with high or very high temperatures (i.e.  $P_e < 0.1$ ).

#### 4.3 Dominant debris flow triggers

The above presented exceedance probabilities of several system variables at days of debris flow occurrence allowed to estimate the changing relative relevance of  $P$ ,  $M$  and  $S_u$ , respectively, for triggering the observed debris flows on the 25 event days and to classify the debris flows according to the variable that is the most relevant (i.e. “dominant”) contributor for triggering debris flows on the individual event days (Table 2).

##### 4.3.1 The role of high-intensity precipitation

On the three event days with precipitation totals observed at all three stations  $P > 45 \text{ mm d}^{-1}$  and thus  $P_e \leq 0.01$  (No. 7, 11, 19), being a lower limit of traditional rainfall intensity-duration thresholds for debris flow initiation (see above; Guzzetti et al.,

2008), this heavy (cf. Schimpf, 1970), although not necessarily high-intensity and short-duration convective rainfall, is very likely to have a very high relevance as contributor to initiate the debris flows (Table 2). The values of  $S_u$  for these events, with exceedance probabilities  $P_e \leq 0.25$ , suggest some moderately relevant additional contributions from previous water input that left the soil at above-average moisture conditions. Although present at these event days, snowmelt is likely to have low relevance ( $P_e \geq 0.40$ ) as a contributor to these debris flow events. Interestingly, while temperatures have been moderate ( $0.1 < P_e \leq 0.5$ ) for No. 11 and 19, they have been rather low for event No. 7 (Figs. 4a, 5b, 5a, 6b). Thus, for this event, the precipitation only fell as rain at lower elevations ( $< 2000$  m a.s.l.) and the debris flows are therefore likely to have been initiated at lower elevations, which is in accordance with the [associated observation of these debris flow's initiation points](#) flows, located at the lowest section of the inner Pitztal (Fig. 1).

For the events No. 21 and 24, heavy precipitation was likely to have a very high relevance as contributor to trigger debris flows, as well (Table 2). This is in spite of the catchment average observed precipitation on these days being less extreme with  $0.01 < P_e \leq 0.1$ . Rather, as shown in Fig. 5a6a, both debris flows occurred close to the rain gauge with the respective highest precipitation recorded on that day, i.e. station *Plangeroß* for No. 21 and *St. Leonhard im Pitztal* for No. 24 (Fig. 1), both of which reached  $P_e \leq 0.01$ . Together with the high temperatures (Fig. 5b6b), this suggests that the precipitation on these days very likely occurred as highly localized and temporally concentrated convective rainstorms (“thunderstorms”), which potentially exhibited precipitation intensities far above the  $\sim 1.9$  mm h<sup>-1</sup> threshold (as derived as lower limit from the observed

45 mm d<sup>-1</sup> if precipitation is uniformly distributed over one day) for debris flow initiation in mountain areas (Guzzetti et al., 2008), at these two stations. In fact, the available high-resolution precipitation data shows that exceptionally high maximum intensities (6.3 mm 15 min<sup>-1</sup> and 10.8 mm 10 min<sup>-1</sup>; corresponding to exceedance probabilities of  $P_e < 0.0001$ ) occurred on these two days. Snowmelt had some moderate additional contribution to event No. 24, while its relevance was low for No. 21 (Fig. 5e6c). Similarly, the largely below-average  $S_u$  indicates a low relevance of antecedent soil moisture for these two events (Fig. 5d6d). A similar reasoning applies to events No. 3 and 12, albeit somewhat less unambiguous (Table 2). For both events, catchment averaged observed precipitation fell within exceedance probabilities  $0.01 < P_e \leq 0.1$ , and thus below the empirical trigger threshold. However, also in this case, the rain stations recording the highest daily precipitation totals were largely the ones closest to the observed debris flows, i.e. *Plangeroß* for No. 3 and *Jerzens-Ritzenried* for No. 12 (Fig. 1). Although the precipitation recorded at these stations for the two event days did not reach the  $P_e \leq 0.01$  threshold (Fig. 5a6a), the high to very high temperatures on these days plausibly suggest the presence of convective precipitation cells and thus of temporally and spatially concentrated and thus high-intensity rainfall. In contrast, while the temperatures for events No. 1, 16, 22 and 23 were only somewhat above average, the precipitation recorded at gauges close to the respective events (Fig. 1) was mostly closer to the threshold  $P_e = 0.01$  than for the above discussed events No. 3 and 12 (Fig. 5a6a), implying that already a moderate temporal concentration of these values to precipitation durations  $\leq 12$  h (and thus not necessarily convective) on the respective event days would result in precipitation intensities exceeding the threshold for debris flow initiation. Again, for No. 22 and 23 the high-resolution precipitation intensity data shows that clear intensity peaks have

occurred (Table 2). Conversely, only rather moderate precipitation ( $0.1 < P_e \leq 0.5$ ), for both the catchment average and the gauge with the respective highest recorded values, was observed for events No. 4, 5 and 14, albeit most of them with the highest values for the gauges closest to the debris flows. The high temperatures ( $P_e \leq 0.1$ ) indicate that localized and temporally highly concentrated precipitation from convective events and above the necessary trigger thresholds is not unlikely for these days. Similarly and although the precipitation data do not give any direct evidence, the merely moderate snowmelt and antecedent soil moisture together with maximum temperatures nearly reaching the  $P_e < 0.1$  threshold for events No. 15 and 18 suggest that highly localized (and thus potentially not adequately recorded) and/or temporally concentrated precipitation may have generated sufficient local precipitation intensities to trigger these debris flows, as well. Lastly, relatively elevated precipitation values ( $0.01 < P_e \leq 0.1$ ) were observed for event No. 25, therefore suggesting triggering by precipitation, even though temperatures have been – atypically – very low ( $P_e = 0.99$ , corresponding to maximum temperatures of  $-5^\circ\text{C}$  to  $+7^\circ\text{C}$  (Fig. 5b6b)). This interpretation is supported by the available high-resolution precipitation data ( $P_e < 0.01$ ). Please note that the output from the hydrological model suggests that all of the precipitation has fallen as snow (and would therefore not be likely to trigger any debris flow at all); however this is due to the mean temperature amounting to  $-3.8^\circ\text{C}$  and an inherent limitation of using a daily averaged temperature input. The above points suggest, together with the generally low antecedent moisture storage  $S_u$  from preceding and potentially more persistent rain and snowmelt (Fig. 5d6d), that very intense, relatively short-duration precipitation was likely a highly relevant contributor to the events No. 1, 3, 4, 5, 12, 14, 15, 16, 18, 22, 23, and 25 although the level to which this assessment is fully warranted by the available data varies between the events. In addition, debris flow initiation was supported by ~~moderate~~ contributions of snowmelt (No. 1, 3, 12, 14, 15, 16, 18; Fig. 5e6c) for several events. However, as most of the above events occurred during summer (i.e. July and August) after the snow melt peaks, which typically occur much earlier in the season (i.e. May and June; see Fig. 5 and Supplementary Material Fig. S2) and thus when only relatively little snow was left, the snow melt contributions to these events remained quite moderate.

#### 4.3.2 The role of snowmelt

The events No. 8, 9 and 10 occurred on days when the modelled snowmelt reached exceedance probabilities of  $P_e \leq 0.01$  (Fig. 5e6c, Table 2) and only very little to no additional precipitation has been recorded. In spite of these exceedance probabilities, the total median melt volumes of about 18-23 mm  $\text{d}^{-1}$  on these days, equivalent to melt intensities of 0.75-0.96 mm  $\text{h}^{-1}$  for uniform 24 h melt, fall short of the debris flow initiation threshold for precipitation intensities of  $\sim 1.9$  mm  $\text{h}^{-1}$ . However, and importantly, it is very likely that the required intensity threshold was exceeded locally. The reasons are that on the one hand most of the melt water on the event days was generated at high elevations ( $> 2000$  m), leading to locally considerably elevated melt rates and thus intensities at these higher elevations (up to 38 mm  $\text{d}^{-1}$  for No. 8 and 10 and up to 46 mm  $\text{d}^{-1}$  for No. 9), which are the source area of debris flows. On the other hand, melt is unlikely to occur uniformly over a 24 hour period. This causes further temporal concentrations of melt water generation, and thus higher peak melt intensities, within individual days which will roughly reflect daily temperature variations, yet in an attenuated, temporally lagged



manner due to the thermal capacity of the snow pack. Based on the above reasoning, the snowmelt contribution is thus likely to have a very high relevance for the initiation of debris flows on these event days (Table 2). In addition, antecedent soil moisture was also at very high levels, i.e.  $P_e \leq 0.01$  (Fig. 5d6d). This continuous build-up of antecedent soil moisture by persistent snowmelt and some moderate rainwater input over the preceding days (Fig. 4a5a), resulting in catchment-wide almost fully saturated conditions, is thus also likely to provide highly relevant contributions to trigger the debris flow events No. 8, 9 and 10. Indeed, total liquid water availability and also modelled runoff have been at least as high ( $P_e \leq 0.003$ ) as those of events No. 7, 11, and 19, which have been identified as triggered by heavy precipitation with a high confidence (sect. 3.2.1, Table 2). In contrast, the precipitation totals observed on the three days exceed  $P_e > 0.1$ , with no precipitation recorded at all for No. 8 and 9. Although, localized, high intensity precipitation missed by the precipitation gauges cannot be ruled out for these event days, given the already high melt rates of up to  $46 \text{ mm d}^{-1}$  and the fact that for No. 8, 9 and 10 most gauges did not observe any precipitation, rainfall is thus considered to have not more than a moderate additional contribution to the initiation of these debris flows.

For No. 17, an extremely **high/low** snowmelt exceedance probability of  $P_e = 0.0001$  was estimated, resulting from the highest snowmelt rate that was modelled within the study period 1953-2012. Yet a maximum local melt intensity of “only”  $38 \text{ mm d}^{-1}$  has been calculated which equals those of event No. 8 and 10, due to the snowmelt occurring over a wider range of elevations ( $> 1700 \text{ m a.s.l.}$ ) on that day. As at all three climate stations, moderate ( $0.1 < P_e \leq 0.5$ ) precipitation was recorded, rainfall will have played a more prominent role than for events No. 8, 9 and 10, making this event a classical rain-on-snow triggered event (cf. Church and Miles, 1987).

Mirroring the reasoning for events No. 8, 9 and 10, the snowmelt exceedance probabilities of  $0.01 < P_e \leq 0.1$  for event No. 20 and  $0.1 < P_e \leq 0.5$  for No. 2 suggest at least high and moderate snowmelt contributions, respectively, for triggering the associated debris flow. Interestingly, for both events, the snowmelt has been restricted to a smaller elevation band ( $> 2400 \text{ m a.s.l.}$ ) than for the other events described above, thus rendering higher **local** melt intensities. Indeed, for No. 20 maximum melt intensities of ca.  $39 \text{ mm d}^{-1}$ , equalling those of events No. 8, 10 and 17 were modelled and for No. 2, maximum melt intensities of up to  $16 \text{ mm d}^{-1}$ , which – given a **catchment** mean snowmelt of only  $4 \text{ mm d}^{-1}$  – are also quite noteworthy. Similarly, the absence of observed precipitation and – in case of No. 2 – only moderate maximum temperature, suggests that precipitation is likely to be of low relevance for the initiation of debris flow events No. 2 and 20, although the occurrence of small convective shower cells cannot be fully dismissed. Note, however, that the direct evidence provided by data in particular for No. 2 is less strong than for events No. 8, 9, 10 and 17, leaving the assessment of the relative relevance of the individual contributors less robust.

To sum up, events No. 2, 8, 9, 10, 17, and 20 have been associated with snowmelt as the primary trigger, while the assumed additional influence of rainfall (i.e. “rain-on-snow”) and antecedent soil moisture varies between the events. [Additional supporting evidence for the above reasoning is that the general timing of the above events coincides well with the snow melt season. Snow melt typically peaks during May and June in the study region \(Fig. 5, Supplementary Material Fig. S2\), while high-intensity, convective rainfall is mostly only observed later in the season \(i.e. July and August\).](#)

### 4.3.3 The role of antecedent soil moisture

For event No. 13, the gradual build-up of soil moisture  $S_u$  by considerable precipitation in the days before as well as by persistent, low-intensity snowmelt in the weeks before the event to nearly fully saturated levels (Supplementary Material Fig. S2f), resulted in a soil moisture level with exceedance probability of  $P_e \leq 0.01$  (Fig. ~~Sd~~ 6d, Table 2). This suggests that soil moisture had likely a very high relevance to trigger this event. Precipitation and snowmelt rates corresponding to  $0.1 < P_e \leq 0.5$  provided additional moderate contributions to initiate event No. 13.

A similar pattern can be found for event No. 6, albeit with a lower relative contribution from soil moisture, whose contribution to trigger the event was moderately relevant ( $P_e = 0.24$ ), as were the contributions of precipitation ( $P_e = 0.19$ ) and snowmelt ( $P_e = 0.40$ ).

Interestingly, both events, No. 6 and 13, occurred in the lowest part of the study area, where relatively large parts are vegetated (Fig. 1), while most of the events associated with high-intensity precipitation (No. 1, 3, 4, 5, 12, 14, 15, 16, 18, 21, 22, 23, 24) took place at higher elevations. For these events, the antecedent soil moisture estimates have been mostly below average, which not only backs the interpretation of high-intensity precipitation as dominant trigger (as discussed in sect. 4.3.1), but may also indicate that the antecedent soil moisture is in general of minor significance at higher elevations, as in its headwaters the catchment is dominated by lower-permeability surfaces (bare rock, sparsely vegetated areas) and shallow soils that only provide limited storage capacities (cf. Berti and Simoni, 2005; Coe et al., 2008; Gregoretti and Fontana, 2008).

### 4.3.4 ~~Complementary~~ Seasonally varying importance of the different trigger contributions

The above analysis illustrated quite clearly that water inputs originating from different individual “sources” can significantly contribute to generate ~~compound~~ trigger conditions in the study area. The data further suggest that the relative relevance of each these variables contributing to the actual trigger conditions does vary over time. Even more, there is some evidence that among the three tested variables, high-intensity and potentially short-duration precipitation  $P$  may be not the consistently most relevant (or “dominant”) contributing factor for all events. Rather, it is not unlikely that also high-intensity snowmelt  $M$  and similarly, although with some lower degree of confidence, persistent, lower intensity water input, building up antecedent soil moisture content  $S_u$  and eventually causing saturated conditions, can generate the most relevant contributions to reach trigger conditions. More specifically, high-intensity precipitation was likely to be the dominant contributor to ~~the compound~~ trigger debris flows on 17 out of 25 event days (68%). This corroborates previous studies that this type of precipitation is the prevalent trigger in such environments (e.g. Berti et al., 1999; Marchi et al. 2002; Berti and Simoni, 2005; Coe et al., 2008; Gregoretti and Fontana, 2008; Braun and Kaitna 2016; Ciavolella et al., 2016). In addition, however, high-intensity snowmelt was likely the dominant contributor on 6 days, corresponding to 24% of the observed events and antecedent soil moisture on 2 event days (8%), highlighting their critical ~~complementary~~ individual contributions to debris flow initiation.

A somewhat different, more quantitative perspective is given by Fig. 67, showing the joint conditional posterior probabilities of a debris flow event  $E$  occurring, given the exceedance probability of each individual variable  $P$ ,  $M$  and  $S_u$ , i.e.  $p(E | P, M, S_u)$ . Note that  $p(E | P, M, S_u)$  is shown in classes of exceedance probabilities with an increment of 0.25 to allow a meaningful visualization of the clustering effects. High probabilities of debris flow events predominantly cluster at low exceedance probabilities of precipitation or in other words, on days with high precipitation totals which were exceeded only in 25% of all days in the study period (i.e. the right-most slice in Fig. 67). Under such conditions, additional contributions from snowmelt or antecedent soil moisture are not necessarily required to trigger debris flows (e.g. Aleotti 2004, Berti et al., 2012), which is also reflected in the elevated  $p(E | P, M, S_u)$  for low  $M$  and  $S_u$  in that class of precipitation exceedance probability. However, elevated event probabilities can also occur when little to no precipitation is observed, i.e. at exceedance probabilities of  $P > 25\%$ , which is roughly equivalent to  $P < 6 \text{ mm d}^{-1}$ , but when instead higher melt rates and/or, albeit to a lesser extent, antecedent moisture levels are likely to be present, as suggested by the model results. Although both, the relative proportions of the different dominant triggers as well as actual values of  $p(E | P, M, S_u)$  as shown in Fig. 67, may be subject to some change over time due to the relatively low absolute number of events with respect to the 60 year study period, the general pattern strongly underline the ~~complementary nature~~ varying roles of the three variables under consideration as individual and potentially dominant contributors to debris flow trigger conditions in the study region.

Most debris flow events in the study area occur between mid- and late summer (Fig. 78), when spring precipitation and persistent snowmelt have developed above-average soil moisture levels and when the frequency of high-intensity, convective rain storms increases (Fig. 45, Supplementary Material Fig. S2). Further analysis also revealed a relatively clear pattern in the seasonally changing relative relevance of the three considered variables as contributors to debris flow trigger conditions. In general, three distinct seasonal debris flow trigger regimes emerge from the analysis, which to a high degree reflect both the seasonal cycle in the hydro-meteorological conditions and in debris flow occurrence, from snow melt to convective rainfall dominated debris flow triggers. While late spring and early summer events are mostly associated with snowmelt in combination with elevated soil moisture, and only very minor contributions of high-intensity precipitation, the latter is, for the above reasons, the dominant trigger ~~later in the year~~ summer and early autumn. While the former may be trivial given that significant snowmelt is less common from July onwards, it is interesting to observe that high-intensity precipitation may be, though also sometimes occurring in spring and early summer, less relevant for triggering debris flows in that -time of the year. This pattern mostly arises from a combination of two factors, namely that in spring considerable proportions of precipitation observed at lower elevations (1) still fall as snow, in particular at higher elevations and (2) are, if falling as rain, intercepted by, transiently stored in and/or potentially refrozen in the snow pack, in particular if the snow pack has not yet reached isothermal conditions at  $0^\circ\text{C}$  throughout the region of interest. Although a mature snow pack later in the melt season may reverse the latter into a positive feedback, i.e. actually reinforcing intensive precipitation in rain-on-snow events (e.g. Harr, 1981; Conway and Raymond, 1993; Cohen et al., 2015), both factors above can, in principle, also cause an attenuation of the observed precipitation intensity as water will be released from the snow pack with some time lags and potentially over longer time, i.e. at lower rates than the observed ones. The immediate implications are then that thresholds

for debris flow initiation estimated from traditional rainfall (but also precipitation) intensity-duration approaches may be suitable for some regions as for example demonstrated by Berti et al. (2012), who showed that antecedent soil moisture is of limited importance in their study region, but will be unreliable for certain hydrological conditions, in particular in snow dominated regions (cf. Decaulne et al., 2005), and thus insufficient for meaningful predictions of debris flows. As a step forward, it may therefore be beneficial to move towards understanding the problem in a more comprehensive and thus multivariate way, expressing and combining the varying relative relevance of complementary different water “sources” in terms of *total liquid water availability*  $S_1$  (see Fig. ~~5e6e~~ as example) in the source zone of debris flows, ~~i.e. a compound trigger approach~~ as recently also emphasized by Bogaard and Greco (2017).

#### 4.3.4 Discussion

We would like to reiterate here that, as in any hydrological study at scales larger than the hillslope scale, the issue of epistemic errors in data (Beven, 2012; Beven et al., 2017a,b), arising from the typically insufficient spatial but also temporal resolutions of the available observations (mostly precipitation) can introduce considerable uncertainty in the interpretation of a specific hydrological system (e.g. Valéry et al., 2010; Nikolopoulos et al., 2014; Marra et al., 2017) and which is further exacerbated by complex, mountainous terrain (e.g. Hrachowitz and Weiler, 2011). This is in particular relevant for debris flows as they depend on the hydrological conditions at the specific location of their initiation, which is frequently of very limited spatial extent. Borga et al. (2014), for instance, reported the occurrence of several debris flows that were triggered by highly localized, high-intensity rainfall  $> 100 \text{ mm h}^{-1}$ , which remained completely unrecorded by rain gauges at 5-10 km distance.

We also explicitly acknowledge additional uncertainties arising from the use of a simple, semi-distributed model to represent the hydrological system of the study area. Such models are clearly oversimplification of the detailed processes controlling the storage and release of water. Together with the effect of the above discussed data errors, this explains, why the model cannot fully reproduce some of the features in the observed hydrograph (e.g. Figs. ~~4b, c and 5f~~ 5b, c and 6f), in spite of its adequate overall performance. Indeed, out of the 6 debris flow days, where both modelled and measured runoff values were available, the modelled runoff was in 3 cases not corresponding particularly well with the measured runoff (Fig. 6f), although in those cases where the runoff was underestimated by the model (No. 20), this was most likely due to un- or underrecorded precipitation. While this ambiguity equally affects the estimates for total soil moisture, the modelled snowmelt and the antecedent soil moisture, respectively, can be assumed to be more correct, as these variables are integrations over time in which case erroneous precipitation measurements are likely to be compensated and thus of less consequence (e.g. Hrachowitz and Weiler, 2011). In addition, the spatial integration of local processes is likely to result in a misrepresentation of hydrological conditions for the locations of debris flow initiation.

However, even though the model is rather simple with limited spatial differentiation, we would like to point out that our approach is not due to an ill-advised oversimplification. Rather, it is the (un-)available data that limits a meaningful spatial differentiation. The most crucial meteorological input, namely precipitation, is very often (and also here) not available on a

spatially sufficiently distributed basis (see above), let alone for the actual source area of a specific debris flow. Furthermore, the calibration of a more distributed model would be more problematic and – in the case of fully distributed physically-based models – would encounter many other sources of uncertainties (e.g. model/parameter equifinality, scale of available field observations of physical parameters vs. scale of the modelling application/grid size, the suitability of the model equations for the scale of the applications, etc.). These issues have been acknowledged for a quite some time but no real progress to close the gap between simplicity and complexity has yet been made (e.g. Dooge, 1986; Beven, 1989, 2006b; Jakeman and Hornberger, 1993; Sivapalan, 2005; McDonnell et al., 2007; Zehe et al., 2007, 2014; Clark et al., 2011, 2017; Hrachowitz and Clark, 2017).

More specifically and notwithstanding these limitations, the catchment-wide considerable melt rates  $M$ , together with the generally elevated soil moisture  $S_u$  during snowmelt dominated events generated by the model suggest that, in spite of the potential presence of un- or under-recorded precipitation, these two sources contribute considerable volumes of water to the required trigger threshold. Additional precipitation may then further contribute, but this does neither imply that these contributions were actually necessary nor, and even less so, that they were dominant for triggering these events. Moreover, although modelled melt rates and soil moisture levels may not be fully representative for the location of the debris flow initiation, they provide most likely conservative estimates, as their real values are likely to be higher at the location and moment of debris flow initiation due to spatial and temporal concentration effects. Furthermore, soil water storage, besides being largely controlled by low-intensity, larger-scale water input, also acts as a low-pass filter. As such it attenuates spatio-temporal variability in precipitation to some degree and is thus more homogeneous than the precipitation itself (e.g. Oudin et al., 2004; Euser et al., 2015).

## **Conclusions**

The results of this study suggest that the available, relatively scarce data and the semi-distributed model together contained sufficient information to facilitate an analysis that allowed the identification of general, large scale pattern and thus the distinction of three different relevant, ~~complementary~~ “sources” of water, i.e. high-intensity precipitation, snowmelt and antecedent soil moisture, that contribute with varying relative importance to the initiation of debris flows in the study region. This highlights the value of a more holistic perspective, ~~i.e. a compound trigger concept~~, for developing a better understanding for debris flow formation and, may provide a first step towards more reliable debris flow predictions, in particular for snow dominated regions.

### *Code availability*

The model code used can be made available by the first author upon request.

### *Data availability*

Hydrological data may be requested from HD Tirol ([www.tirol.gv.at](http://www.tirol.gv.at)), TIWAG ([www.tiwag.at](http://www.tiwag.at)) and ZAMG ([www.zamg.ac.at](http://www.zamg.ac.at)). Data on debris flow events must be directly requested from the Austrian Federal Ministry of Agriculture, Forestry, Environment and Water Management (BMLFUW). The rainfall multipliers are available from Martin Mergili. The digital elevation model, land cover and glacier data can be freely downloaded (links see reference entries for Data.gv.at, CORINE, Austrian Glacier Inventory and WGMS).

### *Author contributions*

KM, RK and MH designed the study, KM and DP carried out the analysis, and MH, KM and RK wrote the paper.

### *Competing interests*

No competing interests to declare.

### *Acknowledgements*

We thank HD Tirol, TIWAG and ZAMG for supplying the climate datasets and Martin Mergili for readily sharing his rainfall lapse rate data. This project receives financial support from the Austrian Climate and Energy Fund and is carried out within the framework of the ‘ACRP’ Programme. .

## References

- Aleotti, P.: A warning system for rainfall-induced shallow failures, *Eng. Geol.*, 73(3), 247-265, doi:10.1016/j.enggeo.2004.01.007, 2004.
- Anagnostopoulos, G. G., Faticchi, S., and Burlando, P.: An advanced process-based distributed model for the investigation of rainfall-induced landslides: The effect of process representation and boundary conditions, *Water Resour. Res.*, 51, 7501-7523, doi:10.1002/2015WR016909, 2015.
- Auer, I., Böhm, R., Jurkovic, A., Lipa, W., Orlik, A., Potzmann, R., ... and Jones, P.: HISTALP – historical instrumental climatological surface time series of the Greater Alpine Region, *Int. J. Climatol.*, 27(1), 17-46, doi:10.1002/joc.1377, 2007.
- Austrian Glacier Inventory: Institute of Atmospheric and Cryospheric Sciences, University of Innsbruck, <http://acinn.uibk.ac.at/research/ice-and-climate/projects/agi>, 1969, 1997, downloaded: 2 February 2016.
- Baum, R. L. and Godt, J. W.: Early warning of rainfall-induced shallow landslides and debris flows in the USA, *Landslides*, 7(3), 259-272, doi:10.1007/s10346-009-0177-0, 2010.
- Berenguer, M., Sempere-Torres, D., and Hürlimann, M.: Debris-flow forecasting at regional scale by combining susceptibility mapping and radar rainfall, *Nat. Hazards Earth Syst. Sci.*, 15(3), 587-602, doi:10.5194/nhess-15-587-2015, 2015.
- [Berghuijs, W. R., Sivapalan, M., Woods, R. A., and Savenije, H. H.: Patterns of similarity of seasonal water balances: A window into streamflow variability over a range of time scales, \*Water Resour. Res.\*, 50\(7\), 5638-5661, doi:10.1002/2014WR015692, 2014.](#)
- Berti, M. and Simoni, A.: Experimental evidences and numerical modelling of debris flow initiated by channel runoff, *Landslides*, 2(3), 171-182, doi:10.1007/s10346-005-0062-4, 2005.
- Berti, M., Genevois, R., Simoni, A., and Tecca, P. R.: Field observations of a debris flow event in the Dolomites, *Geomorphology* 29(3-4), 265-274, doi:10.1016/S0169-555X(99)00018-5, 1999.
- Berti, M., Martina, M. L. V., Franceschini, S., Pignone, S., Simoni, A., and Pizziolo, M.: Probabilistic rainfall thresholds for landslide occurrence using a Bayesian approach, *J. Geophys. Res.*, 117, F04006, doi:10.1029/2012JF002367, 2012.
- Beven, K.: [Changing ideas in hydrology – the case of physically based models, \*J. Hydrol.\*, 105, 157-172, 1989.](#)
- [Beven, K. J.: A manifesto for the equifinality thesis, \*J. Hydrol.\*, 320\(1-2\), 18-36, doi:10.1016/j.jhydrol.2005.07.007, 2006a.](#)
- [Beven, K.: Searching for the Holy Grail of scientific hydrology:  \$Q\_t=\(S, R, dt\)A\$  as closure, \*Hydrol. Earth Syst. Sci.\*, 10, 609-618, doi:10.5194/hess-10-609-2006, 2006b.](#)
- Beven, K. J.: *Rainfall-Runoff Modelling: The Primer*, 2nd edition, John Wiley & Sons Inc., Chichester, UK, ISBN:978-0-470-71459-1, 2012.
- Beven, K. J., Almeida, S., Aspinall, W. P., Bates, P. D., Blazkova, S., Borgomeo, E., Goda, K., Hall, J. W., Phillips, J. C., Simpson, M., Smith, P. J., Stephenson, D. B., Wagener, T., Watson, M., and Wilkins, K. L.: Epistemic uncertainties and natural hazard risk assessment. 1. A review of different natural hazard areas, *Nat. Hazards Earth Syst. Sci. Discuss.*, doi:10.5194/nhess-2017-250, in review, 2017a.
- Beven, K. J., Aspinall, W. P., Bates, P. D., Borgomeo, E., Goda, K., Hall, J. W., Page, T., Phillips, J. C., Simpson, M., Smith, P. J., Wagener, T., and Watson, M.: Epistemic uncertainties and natural hazard risk assessment. 2. What should constitute good practice?, *Nat. Hazards Earth Syst. Sci. Discuss.*, doi:10.5194/nhess-2017-251, in review, 2017b.
- Bíl, M., Andrášik, R., Zahradníček, P., Kubeček, J., Sedoník, J., and Štěpánek, P.: Total water content thresholds for shallow landslides, Outer Western Carpathians, *Landslides*, 13, 337-347, doi:10.1007/s10346-015-0570-9, 2015.

- [Birkel, C., Soulsby, C., and Tetzlaff, D.: Conceptual modelling to assess how the interplay of hydrological connectivity, catchment storage and tracer dynamics controls nonstationary water age estimates, \*Hydrol. Process.\*, 29\(13\), 2956-2969, doi:10.1002/hyp.10414, 2015.](#)
- Bogaard, T. A. and Greco, R.: Landslide hydrology: from hydrology to pore pressure, *WIREs Water*, 3, 439-459, doi:10.1002/wat2.1126, 2016.
- Bogaard, T. A. and Greco, R.: Invited perspectives. A hydrological look to precipitation intensity duration thresholds for landslide initiation: proposing hydro-meteorological thresholds, *Nat. Hazards Earth Syst. Sci. Discuss.*, doi:10.5194/nhess-2017-241, in review, 2017.
- Böhm, R., Schöner, W., Auer, I., Hynek, B., Kroisleitner, C., Weyss, G., and Jurkovic, A.: Gletscher und Abflussverhalten, Bericht zu Zielvereinbarung 2008/31, Zentralanstalt für Meteorologie und Geodynamik, 2007.
- Borga, M., Stoffel, M., Marchi, L., Marra, F., and Jakob, M.: Hydrogeomorphic response to extreme rainfall in headwater systems: flash floods and debris flows, *J. Hydrol.*, 518, 194-205, doi:10.1016/j.jhydrol.2014.05.022, 2014.
- Braun, M. and Kaitna, R.: Analysis of meteorological trigger conditions for debris flows on a daily time scale., in: Makarov, S. A., Atutova, J. V. and Shekhovtsov, A. I. (Eds.): Debris flows: risks, forecast, protection: Materials of IV International Conference (Russia, Irkutsk – Arshan village (The Republic of Buriatia), Irkutsk: Publishing House of Sochava Institute of Geography SB RAS; ISBN:978-5-94797-273-3, 2016.
- BMLFUW: Austrian Federal Ministry of Agriculture, Forestry, Environment and Water Management.
- Chitu, Z., Bogaard, T. A., Busuioc, A., Burcea, S., Sandric, I., and Adler, M. J.: Identifying hydrological pre-conditions and rainfall triggers of slope failures at catchment scale for 2014 storm events in the Ialomita Subcarpathians, Romania, *Landslides*, 14, 419-434, doi:10.1007/s10346-016-0740-4, 2017.
- Church, M. and Miles, M. J.: Meteorological antecedents to debris flow in southwestern British Columbia: some case studies, in: Costa, J. E. and Wiczorek, G. F. (Eds.): Debris flows/avalanches: Process, Recognition and Mitigation, *Reviews in Engineering Geology*, 7, 63-80, Geological Society of America, Boulder, Colorado, doi:10.1130/REG7-p63, 1987.
- Ciavolella, M., Bogaard, T. A., Gargano, R., and Greco, R.: Is there predictive power in hydrological catchment information for regional landslides hazard assessment?, *Proced. Earth Plan. Sc.*, 16, 195-203, doi:10.1016/j.proeps.2016.10.021, 2016.
- [Clark, M. P., Slater, A. G., Rupp, D. E., Woods, R. A., Vrugt, J. A., Gupta, H. V., Wagener, T., and Hay, L. E.: Framework for Understanding Structural Errors \(FUSE\): A modular framework to diagnose differences between hydrological models, \*Water Resour. Res.\*, 44, W00B02, doi:10.1029/2007WR006735, 2008.](#)
- Clark, M. P., Kavetski, D., and Fenicia, F.: Pursuing the method of multiple working hypotheses for hydrological modeling, *Water Resour. Res.*, 47, W09301, doi:10.1029/2010WR009827, 2011.
- [Clark, M. P., Bierkens, M. F. P., Samaniego, L., Woods, R. A., Uijlenhoet, R., Bennett, K. E., Pauwels, V. R. N., Cai, X., Wood, A. W., and Peters-Lidard, C. D.: The evolution of process-based hydrologic models: historical challenges and the collective quest for physical realism, \*Hydrol. Earth Syst. Sci.\*, 21, 3427-3440, doi:10.5194/hess-21-3427-2017, 2017.](#)
- Coe, J. A., Kinner, D. A., and Godt, J. W.: Initiation conditions for debris flows generated by runoff at Chalk Cliffs, central Colorado, *Geomorphology* 96, 270-297, doi:10.1016/j.geomorph.2007.03.017, 2008.
- Cohen, J., Ye, H., and Jones, J.: Trends and variability in rain-on-snow events, *Geophys. Res. Lett.*, 42, 7115-7122, doi:10.1002/2015GL065320, 2015.
- Conway, H. and Raymond, C. F.: Snow stability during rain, *J. Glaciol.*, 39(133), 1993.



- CORINE Land cover: European Environment Agency, European Union, [www.eea.europa.eu/data-and-maps/data/corine-land-cover-1990-raster-3](http://www.eea.europa.eu/data-and-maps/data/corine-land-cover-1990-raster-3), .../corine-land-cover-2000-raster-2, .../corine-land-cover-2006-raster-3, downloaded: 2 February 2016.
- Criss, R. E. and Winston, W. E.: Do Nash values have value? Discussion and alternate proposals, *Hydrol. Process.*, 22, 2723-2725, doi:10.1002/hyp.7072, 2008.
- Crozier, M. J.: Prediction of rainfall-triggered landslides: a test of the antecedent water status model, *Earth Surf. Proc. Land.*, 24, 825-833, doi:10.1002/(SICI)1096-9837(199908)24:9<825::AID-ESP14>3.0.CO;2-M, 1999.
- Data.gov.at: Digital Elevation Model (DEM) “b5de6975-417b-4320-afdb-eb2a9e2a1dbf”, [www.data.gov.at/katalog/dataset/b5de6975-417b-4320-afdb-eb2a9e2a1dbf](http://www.data.gov.at/katalog/dataset/b5de6975-417b-4320-afdb-eb2a9e2a1dbf), downloaded: 2 February 2016.
- Decaulne, A., Sæmundsson, Þ., and Pétursson, O.: Debris flow triggered by rapid snowmelt: a case study in the Gleidárhjalli area, northwestern Iceland, *Geografiska Annaler*, 87A(4), 487-500, doi:10.1111/j.0435-3676.2005.00273.x, 2005.
- Deganutti, A. M., Marchi, L., and Arattano, M.: Rainfall and debris-flow occurrence in the Moscardo basin (Italian Alps), in: Wieczorek, G. F., and Naeser, N. D. (Eds.): *Debris-flow Hazards Mitigation: Mechanics, Prediction and Assessment*, Proceedings of the second international conference on debris flow hazards mitigation, Taipei, Taiwan, August 16-18, 62-72, 2000.
- Dhakal, A. S. and Sidle, R. C.: Distributed simulations of landslides for different rainfall conditions, *Hydrol. Process.*, 18, 757-776, doi:10.1002/hyp.1365, 2004.
- [Dooge, J. C.: Looking for hydrologic laws, \*Water Resour. Res.\*, 22, 46-58, doi:10.1029/WR022i09Sp0046S, 1986.](#)
- Euser, T., Winsemius, H. C., Hrachowitz, M., Fenicia, F., Uhlenbrook, S., and Savenije, H. H. G.: A framework to assess the realism of model structures using hydrological signatures, *Hydrol. Earth Syst. Sci.*, 17, 1893-1912, doi:10.5194/hess-17-1893-2013, 2013.
- Euser, T., Hrachowitz, M., Winsemius, H. C., and Savenije, H. H. G.: The effect of forcing and landscape distribution on performance and consistency of model structures: Distribution of forcing and model structures, *Hydrol. Process.*, 29(17), 3727-3743, doi:10.1002/hyp.10445, 2015.
- Fan, L., Lehmann, P., and Or, D.: Effects of soil spatial variability at the hillslope and catchment scales on characteristics of rainfall-induced landslides, *Water Resour. Res.*, 52(3), 1781-1799, doi:10.1002/2015WR017758, 2016.
- Fenicia, F., Kavetski, D., and Savenije, H. H. G.: Elements of a flexible approach for conceptual hydrological modeling: 1. Motivation and theoretical development, *Water Resour. Res.*, 47, W11510, doi:10.1029/2010WR010174, 2011.
- [Fenicia, F., Kavetski, D., Savenije, H. H. G., Clark, M. P., Schoups, G., Pfister, L., and Freer, J.: Catchment properties, function, and conceptual model representation: is there a correspondence?, \*Hydrol. Process.\*, 28\(4\), 2451-2467, doi:10.1002/hyp.9726, 2014.](#)
- [Fenicia, F., Kavetski, D., Savenije, H. H. G., and Pfister, L.: From spatially variable streamflow to distributed hydrological models: Analysis of key modeling decisions, \*Water Resour. Res.\*, 52\(2\), 954-989, doi:10.1002/2015WR017398, 2016.](#)
- Fovet, O., Ruiz, L., Hrachowitz, M., Faucheux, M., and Gascuel-Oudou, C.: Hydrological hysteresis and its value for assessing process consistency in catchment conceptual models, *Hydrol. Earth Syst. Sci.*, 19, 105–123, doi:10.5194/hess-19-105-2015, 2015.
- Gao, H., Ding, Y., Zhao, Q., Hrachowitz, M., and Savenije, H. H. G.: The importance of aspect for modelling the hydrological response in a glacier catchment in Central Asia, *Hydrol. Process.*, 31, 2842-2859, doi:10.1002/hyp.11224, 2017.
- [Gharari, S., Hrachowitz, M., Fenicia, F., Gao, H., and Savenije, H. H.: Using expert knowledge to increase realism in environmental system models can dramatically reduce the need for calibration, \*Hydrol. Earth Syst. Sci.\*, 18, 4839-4859, doi:10.5194/hess-18-4839-2014, 2014.](#)
- Glade, T.: Modelling landslide-triggering rainfalls in different regions of New Zealand – the soil water status model, *Zeitschrift für Geomorphologie*, 122, 63-84, 2000.

- Glade, T., Crozier, M., and Smith, P.: Applying Probability Determination to Refine Landslide-triggering Rainfall Thresholds Using an “Empirical Antecedent Daily Rainfall Model”, *Pure Appl. Geophys.*, 157, 1059-1079, doi:10.1007/s000240050017, 2000.
- Gregoretti, C. and Fontana, G. D.: The triggering of debris flow due to channel-bed failure in some alpine headwater basins of the Dolomites: analyses of critical runoff, *Hydrol. Process.*, 22(13), 2248-2263, doi:10.1002/hyp.6821, 2008.
- Gupta, H. V., Wagener, T., and Liu, Y.: Reconciling theory with observations: Elements of a diagnostic approach to model evaluation, *Hydrol. Process.*, 22, 3802-3813, doi:10.1002/hyp.6989, 2008.
- Guzzetti, F., Peruccacci, S., Rossi, M., and Stark, C. P.: Rainfall thresholds for the initiation of landslides in central and southern Europe, *Meteorol. Atmos. Phys.*, 98(3-4), 239-267, doi:10.1007/s00703-007-0262-7, 2007.
- Guzzetti, F., Peruccacci, S., Rossi, M., and Stark, C. P.: The rainfall intensity–duration control of shallow landslides and debris flows: an update, *Landslides*, 5, 3-17, doi:10.1007/s10346-007-0112-1, 2008.
- Hargreaves, G. H. and Samani, Z. A.: Reference crop evapotranspiration from temperature, *Appl. Eng. Agric.*, 1(2), 96-99, doi:10.13031/2013.26773, 1985.
- Harr, R. D.: Some characteristics and consequences of snowmelt during rainfall in Western Oregon, *J. Hydrol.*, 53, 277-304, 1981.
- HD Tirol: Hydrographischer Dienst Tirol (Hydrographic Service Tyrol), Sachgebiet Hydrographie und Hydrologie, Amt der Tiroler Landesregierung, data received: 25 August 2015.
- Hrachowitz, M. and Clark, M. P.: HESS Opinions: The complementary merits of competing modelling philosophies in hydrology, *Hydrol. Earth Syst. Sci.*, 21, 3953-3973, doi:10.5194/hess-21-3953-2017, 2017.
- Hrachowitz, M. and Weiler, M.: Uncertainty of Precipitation Estimates Caused by Sparse Gauging Networks in a Small, Mountainous Watershed, *J. Hydrol. Eng.*, 16(5), 460-471, doi:10.1061/(ASCE)HE.1943-5584.0000331, 2011.
- Hrachowitz, M., Fovet, O., Ruiz, L., Euser, T., Gharari, S., Nijzink, R., Freer, J., Savenije, H.H.G., and Gascuel-Oudou, C.: Process consistency in models: The importance of system signatures, expert knowledge, and process complexity, *Water Resour. Res.*, 50, 7445-7469, doi:10.1002/2014WR015484, 2014.
- [Hrachowitz, M., Fovet, O., Ruiz, L., and Savenije, H. H.: Transit time distributions, legacy contamination and variability in biogeochemical 1/f \$\alpha\$  scaling: how are hydrological response dynamics linked to water quality at the catchment scale?. \*Hydrol. Process.\*, 29\(25\), 5241-5256, doi: 10.1002/hyp.10546, 2015.](#)
- Hübl, J., Totschnig, R., Sitter, F., Mayer, B., and Schneider, A.: Historische Ereignisse - Band 2: Auswertung von Wildbach Schadereignissen in Westösterreich auf Grundlage der Wildbachaufnahmeblätter, IAN Report 111, Band 2, University of Natural Resources and Life Sciences, Vienna, 2008.
- Hungr, O., Leroueil, S., and Picarelli, L.: The Varnes classification of landslide types: an update, *Landslides* 11(2), 167-194, doi:10.1007/s10346-013-0436-y, 2014.
- Iverson, R. M.: Landslide triggering by rain infiltration. *Water Resour. Res.* 36(7), 1897-1910, doi:10.1007/3-540-27129-5\_1, 2000.
- [Jakeman, A. J. and Hornberger, G. M.: How much complexity is warranted in a rainfall-runoff model?, \*Water Resour. Res.\*, 29, 2637-2649, 1993.](#)
- Johnson, K. and Sitar, N.: Hydrologic conditions leading to debris-flow initiation, *Can. Geotech. J.*, 27(6), 789-801, doi:10.1139/t90-092, 1990.
- Kean, J. W., McCoy, S. W., Tucker, G. E., Staley, D. M., and Coe, J. A.: Runoff-generated debris flows: Observations and modeling of surge initiation, magnitude, and frequency, *J. Geophys. Res. - Earth*, 118(4), 2190-2207, doi:10.1002/jgrf.20148, 2013.

- Kienholz, H.: Gefahrenbeurteilung und -bewertung – auf dem Weg zu einem Gesamtkonzept, Schweizerische Zeitschrift für Forstwesen, 146, 701-725, 1995.
- Lambrecht, A. and Kuhn, M.: Glacier changes in the Austrian Alps during the last three decades, derived from the new Austrian glacier inventory, *Ann. Glaciol.*, 46, 177-184, 2007.
- [Leavesley, G. H., Markstrom, S. L., Brewer, M. S., and Viger, R. J.: The modular modeling system \(MMS\) – The physical process modeling component of a database-centered decision support system for water and power management, in: Chow, W., Brocksen, R. W., and Wisniewski, J.: Clean Water: Factors that Influence Its Availability, Quality and Its Use, 303-311, Springer Netherlands, 1996.](#)
- Lehmann, P., and Or, D.: Hydromechanical triggering of landslides: From progressive local failures to mass release, *Water Resour. Res.*, 48(3), W03535, doi:10.1029/2011WR010947, 2012.
- Mader, H., Steidl, T., and Wimmer, R.: Abflussregime Österreichischer Fließgewässer: Beitrag zu einer bundesweiten Fließgewässertypologie, Umweltbundesamt, Monographien Vol. 82, Wien, ISBN:3-85457-336-7, 1996.
- Mancarella, D., Doglioni, A., and Simeone, V.: On capillary barrier effects and debris slide triggering in unsaturated layered covers, *Eng. Geol.*, 147-148, 14-27, doi:10.1016/j.enggeo.2012.07.003, 2012.
- Marchi, L., Arattano, M., and Deganutti, A. M.: Ten years of debris-flow monitoring in the Moscardo Torrent (Italian Alps), *Geomorphology*, 46(1), 1–17, doi:10.1016/S0169-555X(01)00162-3, 2002.
- Marra, F., Destro, E., Nikolopoulos, E. I., Zoccatelli, D., Dominique, J., Creutin, F. G., Guzzetti, F., and Borga, M.: Impact of rainfall spatial aggregation on the identification of debris flow occurrence thresholds. *Hydrol. Earth Syst. Sci.*, doi:10.5194/hess-21-4525-2017, 2017.
- McArdell, B. W., Bartelt, P., and Kowalski, J.: Field observations of basal forces and fluid pore pressure in a debris flow, *Geophys. Res. Lett.*, 34(7), L07406, doi:10.1029/2006GL029183, 2007.
- McCoy, S. W., Kean, J. W., Coe, J. A., Tucker, G. E., Staley, D. M., and Wasklewicz, T. A.: Sediment entrainment by debris flows: In situ measurements from the headwaters of a steep catchment, *J. Geophys. Res. - Earth*, 117, F03016, doi:10.1029/2011JF002278, 2012.
- [McDonnell, J. J., Sivapalan, M., Vaché, K., Dunn, S., Grant, G., Haggerty, R., Hinz, C., Hooper, R., Kirchner, J., Roderick, M. L., Selker, J., and Weiler, M.: Moving beyond heterogeneity and process complexity: A new vision for watershed hydrology, \*Water Resour. Res.\*, 43, W07301, doi:10.1029/2006WR005467, 2007.](#)
- Mergili, M. and Kerschner, H.: Gridded precipitation mapping in mountainous terrain combining GRASS and R, *Norsk Geografisk Tidsskrift - Norwegian Journal of Geography*, 69(1), 2-17, doi:10.1080/00291951.2014.992807, 2015.
- Montgomery, D. R. and Dietrich, W. E.: A physically based model for the topographic control on shallow landsliding, *Water Resour. Res.*, 30(4), 1153-1171, doi:10.1029/93WR02979, 1994.
- Montgomery, D. R., Schmidt, K. M., Dietrich, W. E., and McKean, J.: Instrumental record of debris flow initiation during natural rainfall: Implications for modeling slope stability, *J. Geophys. Res. - Earth*, 114, F01031, doi:10.1029/2008JF001078, 2009.
- Moser, M. and Hohensinn, F.: Geotechnical aspects of soil slips in Alpine regions. *Eng. Geol.* 19, 185-211, doi:10.1016/0013-7952(83)90003-0, 1983.
- Nandi, A. and Shakoor, A.: Application of logistic regression model for slope instability prediction in Cuyahoga River Watershed, Ohio, USA, *Georisk*, 2(1), doi:10.1080/17499510701842221, 2008.

- Napolitano, E., Fusco, F., Baum, R. L., Godt, J. W., and De Vita, P.: Effect of antecedent-hydrological conditions on rainfall triggering of debris flows in ash-fall pyroclastic mantled slopes of Campania (southern Italy), *Landslides*, 13(5), 967-983, doi:10.1007/s10346-015-0647-5, 2016.
- Nash, J. E. and Sutcliffe, J. V.: River flow forecasting through conceptual models: Part I, a discussion of principles, *J. Hydrol.*, 10(3), 282-290, doi:10.1016/0022-1694(70)90255-6, 1970.
- Nijzink, R.C., Samaniego, L., Mai, J., Kumar, R., Thober, S., Zink, M., Schäfer, D., Savenije, H. H. G., and Hrachowitz, M.: The importance of topography-controlled sub-grid process heterogeneity and semi-quantitative prior constraints in distributed hydrological models, *Hydrol. Earth Syst. Sci.*, 20, 1151-1176, doi:10.5194/hess-20-1151-2016, 2016a.
- [Nijzink, R., Hutton, C., Pechlivanidis, I., Capell, R., Arheimer, B., Freer, J., Han, D., Wagener, T., McGuire, K., Savenije, H. and Hrachowitz, M.: The evolution of root-zone moisture capacities after deforestation: a step towards hydrological predictions under change?, \*Hydrol. Earth Syst. Sci.\*, 20\(12\), 4775-4779, doi:10.5194/hess-20-4775-2016, 2016b.](#)
- Nikolopoulos, E. I., Crema, S., Marchi, L., Marra, F., Guzzetti, F., and Borga, M.: Impact of uncertainty in rainfall estimation on the identification of rainfall thresholds for debris flow occurrence, *Geomorphology*, 221, 286-297, doi:10.1016/j.geomorph.2014.06.015, 2014.
- Oudin, L., Andréassian, V., Perrin, C., and Anctil, F.: Locating the sources of low-pass behavior within rainfall-runoff models, *Water Resour. Res.*, 40, W11101, doi:10.1029/2004WR003291, 2004.
- Papa, M. N., Medina, V., Ciervo, F., and Bateman, A.: Derivation of critical rainfall thresholds for shallow landslides as a tool for debris flow early warning systems. *Hydrol. Earth Syst. Sci.*, 17(10), 4095-4107, doi:10.5194/hess-17-4095-2013, 2013.
- Prancevic, J. P., Lamb, M. P., and Fuller, B. M.: Incipient sediment motion across the river to debris-flow transition, *Geology*, 42(3), 191-194, doi:10.1130/G34927.1, 2014.
- [Reichenbach, P., Mondini, A. C., and Rossi, M.: The influence of land use change on landslide susceptibility zonation: the Briga catchment test site \(Messina, Italy\), \*Environmental management\*, 54\(6\), 1372-1384, 2014.](#)
- Rickenmann, D. and Zimmermann, M.: The 1987 debris flows in Switzerland: documentation and analysis, *Geomorphology*, 8(2), 175-189, 1993.
- Savenije, H.H.G. and Hrachowitz, M.: HESS Opinions "Catchments as meta-organisms - a new blueprint for hydrological modelling", *Hydrol. Earth Syst. Sci.*, 21(2), 1107-1116, doi:10.5194/hess-21-1107-2017, 2017.
- Schimpf, H.: Untersuchung über das Auftreten beachtlicher Niederschläge in Österreich, *Österreichische Wasserwirtschaft*, 22(5/6), 121-127, 1970.
- Schoups, G., Hopmans, J. W., Young, C. A., Vrugt, J. A. and Wallender, W. W.: Multi-criteria optimization of a regional spatially-distributed subsurface water flow model, *J. Hydrol.*, 311, 20-48, doi:10.1016/j.jhydrol.2005.01.001, 2005.
- [Seibert, J.: Regionalisation of parameters for a conceptual rainfall-runoff model, \*Agricultural and forest meteorology\*, 98, 279-293, 1999.](#)
- [Seibert, J. and Beven, K. J.: Gauging the ungauged basin: how many discharge measurements are needed?, \*Hydrol. Earth Syst. Sci.\*, 13\(6\), 883-892, doi:10.5194/hess-13-883-2009, 2009.](#)
- Sidle, R. C. and Ziegler, A. D.: The canopy interception-landslide initiation conundrum: evidence from a secondary tropical rainforest in northern Thailand, *Hydrol. Earth Syst. Sci.*, 21, 651-667, doi:10.5194/hess-21-651-2017, 2017.
- Simoni, S., Zanotti, F., Bertoldi, G., and Rigon, R.: Modelling the probability of occurrence of shallow landslides and channelized debris flows using GEOTop-FS. *Hydrol. Process.*, 22, 532-545, doi:10.1002/hyp.6886, 2008.

- [Sivapalan, M.: Pattern, process and function: Elements of a new unified theory of hydrologic at the catchment scale, in: Encyclopedia of Hydrological Sciences, edited by: Anderson, M. G., John Wiley & Sons Australia Ltd, UK, vol. 1, 193-220, 2005.](#)
- Stoffel, M., Bollschweiler, M., and Beniston, M.: Rainfall characteristics for periglacial debris flows in the Swiss Alps: past incidences– potential future evolutions, *Climatic Change*, 105, 263-280, doi:10.1007/s10584-011-0036-6, 2011.
- [Takahashi, T.: Estimation of potential debris flows and their hazardous zones: Soft countermeasures for a disaster, \*Natural disaster science\*, 3\(1\), 57-89, 1981.](#)
- TIRIS: Tiroler Rauminformationssystem, State of Tyrol, [www.tirol.gv.at/data/datenkatalog](http://www.tirol.gv.at/data/datenkatalog), downloaded: 29 July 2015.
- TIWAG: Tiroler Wasserkraft AG (Tyrolean Hydropower Corporation), data received: 11 September 2015.
- Turkington, T., Remaître, A., Ettema, J., Hussin, H., and Westen, C.: Assessing debris flow activity in a changing climate, *Climatic Change*, 1, 1-13, doi:10.1007/s10584-016-1657-6, 2016.
- Valéry, A., Andréssian, V., and Perrin, C.: Regionalization of precipitation and air temperature over high-altitude catchments – learning from outliers, *Hydrolog. Sci. J.*, 55(6), 928-940, doi:10.1080/02626667.2010.504676, 2010.
- van den Heuvel, F., Goyette, S., Rahman, K., and Stoffel, M.: Circulation patterns related to debris-flow triggering in the Zermatt valley in current and future climates, *Geomorphology*, 272, 127-136, doi:10.1016/j.geomorph.2015.12.010, 2016.
- von Ruetze, J., Papritz, A., Lehmann, P., Rickli, C., and Or, D.: Spatial statistical modeling of shallow landslides – Validating predictions for different landslide inventories and rainfall events, *Geomorphology*, 133(1-2), 11-22, doi:10.1016/j.geomorph.2011.06.010, 2011.
- von Ruetze, J., Lehmann, P., and Or, D.: Rainfall-triggered shallow landslides at catchment scale: Threshold mechanics-based modeling for abruptness and localization, *Water Resour. Res.*, 49, 6266-6285, doi:10.1002/wrcr.20418, 2013.
- [Wagner, T., Boyle, D. P., Lees, M. J., Wheeler, H. S., Gupta, H. V., and Sorooshian, S.: A framework for development and application of hydrological models, \*Hydrol. Earth Syst. Sci.\*, 5\(1\), 13-26, doi: 10.5194/hess-5-13-2001, 2001.](#)
- WGMS: Fluctuations of Glaciers Database, World Glacier Monitoring Service, Zurich, Switzerland, doi:10.5904/wgms-fog-2017-06, downloaded: 12 August 2017.
- Wieczorek, G. F. and Glade, T.: Climatic factors influencing occurrence of debris flows, in: Jakob, M., and Hungr, O. (Eds.): *Debris-flow Hazards and Related Phenomena*, Springer Berlin Heidelberg, 325-362, doi:10.1007/3-540-27129-5\_14, 2005.
- ZAMG: Zentralanstalt für Meteorologie und Geodynamik (Central Institute for Meteorology and Geodynamics), data received: 6 October 2015.
- [Zehe, E., Elsenbeer, H., Lindenmaier, F., Schulz, K., and Blöschl, G.: Patterns of predictability in hydrological threshold systems, \*Water Resour. Res.\*, 43, W07434, doi:10.1029/2006WR005589, 2007.](#)
- [Zehe, E., Ehret, U., Pfister, L., Blume, T., Schröder, B., Westhoff, M., Jackisch, C., Schymanski, S. J., Weiler, M., Schulz, K., Allroggen, N., Tronicke, J., van Schaik, L., Dietrich, P., Scherer, U., Eccard, J., Wulfmeyer, V., and Kleidon, A.: HESS Opinions: From response units to functional units: a thermodynamic reinterpretation of the HRU concept to link spatial organization and functioning of intermediate scale catchments, \*Hydrol. Earth Syst. Sci.\*, 18, 4635-4655, doi:10.5194/hess-18-4635-2014, 2014.](#)

## Figure captions

### Figure 1.

Study area with locations of observed debris flows (center of [mass deposition](#)), location of stream gauges and weather stations (debris flows: BMLFUW; gauging stations: TIWAG; weather stations: HD Tirol, TIWAG, ZAMG; land cover data: CORINE Land cover; glacier data: Austrian Glacier Inventory; rivers & lakes: TIRIS).

### Figure 2.

Photograph of the inner Pitztal, located next to Plangeroß.

10 changes in figure: added one photograph of the inner Pitztal

### Figure 23.

Data availability, modelled study period and number of days with known debris flow occurrence. Only those debris flow events are plotted of which the exact date of occurrence was known i.e. which were used for this study.

15

### Figure 34.

Structure of the semi-distributed (stratification into 100 m elevation zones) hydrological model. Black symbols indicate fluxes and states, black underlined symbols indicate model input, and grey symbols indicate model parameters (abbreviations see Table 1 ~~and sect. 2.2, 3.1.1, 3.2~~).

20 changes in figure: added Mglacier, Sl, Oobs

### Figure 45.

Observed daily stream flow  $Q_{\text{obs}}$  (black solid line), daily mean temperature  $T_{\text{mean}}$  at mean elevation (grey solid line) and maximum temperature  $T_{\text{max}}$  at mean elevation (black solid line) as well as, based on observed precipitation data, modelled daily rainfall  $P_1$  (dark blue downward columns for 5<sup>th</sup> percentile, incl. grey downward columns for 95<sup>th</sup> percentile), daily snowfall  $P_s$  (light blue downward columns for 5<sup>th</sup> percentile, incl. grey downward columns for 95<sup>th</sup> percentile) and daily snowmelt  $M$  (dark blue upward columns for 5<sup>th</sup> percentile, incl. grey upward columns for 95<sup>th</sup> percentile), modelled stream flow (dark blue line for the median and the grey shaded area for the 5/95<sup>th</sup> percentiles of all behavioral model solutions) and modelled relative soil moisture (solid blue line for the median and the grey shaded area for the 5/95<sup>th</sup> percentiles) for the three selected years (a) 1965, (b) 1989 and (c) 2011 (all remaining years with debris flow occurrence are provided in Supplementary Material Fig. S1). The days where a debris flow event has been documented are marked with red vertical lines. Please note that the plots display the period March 15<sup>th</sup> to October 15<sup>th</sup> to depict the start and amount of rainfall and

30

[snowmelt, however, the analysis \(Figs. 6 and 7\) is based on the period May 15<sup>th</sup> to October 15<sup>th</sup>.](#)

**Figure 56.**

Plots of relevant system variables: (a) Precipitation P elevation adjusted for mean catchment elevation, (b) Maximum temperature  $T_{\max}$  for all catchment elevations (blue bars) and mean elevation (white dots), (c-f) Modelled snowmelt M, antecedent soil moisture  $S_u$ , total liquid water availability  $S_l$ , and runoff  $Q_{\text{mod}}$ , (and where available  $Q_{\text{obs}}$ ). Boxplots comprise all behavioral models. For event numbering see Table 2.  $P_e$  is the observed/modelled probability of exceedance ([i.e. marginal distribution, see section 3.2](#)) for a specific variable considering all days between May 15<sup>th</sup> and October 15<sup>th</sup> within the study period 1953-2012.

10

**Figure 67.**

Individual exceedance probabilities  $P_e$  of precipitation (P; x-axis), snowmelt (M; y-axis) and relative antecedent soil moisture ( $S_u$ ; z-axis) as well as the corresponding joint conditional posterior probabilities of an event occurring given specific values (expressed as classes of exceedance probabilities) of precipitation, snowmelt and antecedent soil moisture,  $p(E | P, M, S_u)$ .

15

**Figure 78.**

Debris flow events by month of occurrence and likely dominant trigger; shades indicate the relative strength (the darker the stronger) of the dominant trigger in terms of (1) its relative relevance compared to the other contributing variables and (2) the extent to which it is directly supported by data (see also Table 2).

20

## Tables

**Table 1.** Model storages, fluxes and parameters (model structure see fig. 4).

**Table 1a.** Model storages and fluxes.

<u>Abbreviation</u>	<u>Unit</u>	<u>Description</u>	<u>Abbreviation</u>	<u>Unit</u>	<u>Description</u>
<u>Storages</u>			<u>Fluxes (cont.)</u>		
$S_{\text{snow}}$	mm	snow storage	$M$	mm/d	snowmelt
$S_{\text{glacier}}$	mm	glacier storage	$M_{\text{glacier}}$	mm/d	glacier melt
$S_u$	mm	unsaturated storage, “antecedent soil moisture”	$E_p$	mm/d	potential evapotranspiration
$S_l$	mm	total liquid water availability $= S_u + PI + M (+M_{\text{glacier}})$	$E_a$	mm/d	actual evapotranspiration
$S_f$	mm	fast responding model component	$Q_{\text{uf}}$	mm/d	influx to fast responding model component
$S_s$	mm	slow responding groundwater storage	$Q_{\text{up}}$	mm/d	preferential percolation
<u>Fluxes</u>			$Q_{\text{us}}$	mm/d	percolation
$P$	mm/d	precipitation	$Q_f$	mm/d	fast runoff
$T_{\text{mean}}$	°C	mean daily temperature	$Q_s$	mm/d	slow runoff
$P_s$	mm/d	solid precipitation, i.e. snow	$Q_{\text{mod}}$	mm/d	modelled total runoff
$P_l$	mm/d	liquid precipitation, i.e. rain	$Q_{\text{obs}}$	mm/d	observed total runoff

**Table 1b.** Model calibration parameters with their uniform prior parameter distributions and the median as well as the 5/95<sup>th</sup> percentiles of the posterior parameter distributions of the set of behavioral solutions.

Abbreviation	Unit	Description	Uniform prior parameter distribution		Posterior parameter distribution percentiles		
			lower	upper	5 <sup>th</sup>	50 <sup>th</sup>	95 <sup>th</sup>
$T_{\text{temp}}$	°C	threshold temperature	0.5	1.5	0.8	1.3	1.5



$melt_f$	$mm\ ^\circ C^{-1}\ d^{-1}$	melt factor	2.5	5	2.7	3.6	4.6
$L_p$	–	transpiration coefficient	0.3	1	0.6	0.8	1.0
$S_{u,max}$	mm	unsaturated storage capacity	40	300	218	276	297
$\beta$	–	shape parameter	0.1	1	0.3	0.6	1.0
$P_{max}$	$mm\ d^{-1}$	percolation capacity	0.1	4	1.1	1.7	2.5
$D$	–	partitioning coefficient	0	1	0.1	0.7	1.0
$K_f$	$d^{-1}$	storage coefficient	0.05	3	0.1	0.3	2.4
$K_s$	$d^{-1}$	storage coefficient	0.001	0.3	0.05	0.09	0.14

---

**Table 2.** The 25 recorded debris flow events in the inner Pitztal that occurred at known dates since 1953. For each event the exceedance probabilities  $P_e$  associated to the observed variables daily precipitation  $P$ , daily maximum temperature  $T_{\max}$  and daily mean stream flow  $Q_{\text{obs}}$  as well as to the modelled variables daily snow melt  $M$ , daily antecedent moisture content  $S_u$ , daily total near-surface water availability  $S_f$  and the daily stream flow  $Q_{\text{mod}}$  at the day of the respective events are given. Bold and underlined values indicate a very high relevance ( $P_e \leq 0.01$ ) and bold values a high relevance ( $0.01 < P_e \leq 0.1$ ) of each individual variable for a given event; normal values indicate moderate relevance ( $0.1 < P_e \leq 0.5$ ) and italic values indicate a low relevance ( $P_e > 0.5$ ). The columns indicating the relevance of contributing variables show the likely level of importance of the three variables that directly affect debris flow initiation ( $P$ ,  $M$ ,  $S_u$ ), after consideration of supporting evidence from variables, such as  $T_{\max}$ , that do not directly affect the triggering of debris flows. As additional plausibility check of our interpretation, information on high-resolution precipitation data is provided (column  $P_{\max,10/15\text{min}}$ ) when available. The direct support by data column indicates to which extent the classification of the contributing variables into very high/high, moderate and low is directly supported by daily data (++: excellent support, +: strong support, ~: moderate support) and thus provides an indicative quality check of how likely this interpretation reflects the real conditions during debris flow initiation.

[ TABLE SEE NEXT PAGE ]

Event No.	Date	Contributing variable											Direct support by daily data	Dominant contributing variable	
		Observed variables				Modelled variables				Relevance					
		P	P <sub>max,10/15min.</sub>	T	Q <sub>obs</sub>	M	S <sub>u</sub>	S <sub>l</sub>	Q <sub>mod</sub>	Very high/high	Moderate	Low			
7	a-b	10/06/1965	<b>0.0002</b>	-	0.98	-	0.85	0.24	<b>0.03</b>	<b>0.04</b>	P	S <sub>u</sub>	M		
11	-	08/08/1966	<b>0.006</b>	-	0.42	-	0.42	0.13	<b>0.004</b>	<b>0.04</b>	P	S <sub>u</sub> , M	-	++	
19	-	06/08/1985	<b>0.001</b>	-	0.28	-	0.83	0.13	<b>0.0005</b>	<b>0.001</b>	P	S <sub>u</sub>	M		
21	-	22/08/1989	<b>0.08</b>	<b>&lt; 0.0001</b> <sup>(21)</sup>	<b>0.06</b>	0.67	0.58	0.52	0.26	0.36	P	-	S <sub>u</sub> , M		
24	-	11/07/2010	<b>0.02</b>	<b>&lt; 0.0001</b> <sup>(24)</sup>	<b>0.01</b>	0.27	0.37	0.85	0.30	0.34	P	M	S <sub>u</sub>		
3	a-d	22/07/1963	0.13	-	<b>0.01</b>	-	0.37	0.59	0.39	0.34	P	M	S <sub>u</sub>		
12	a-b	14/08/1966	<b>0.07</b>	-	<b>0.07</b>	-	0.29	0.14	<b>0.04</b>	0.16	P	S <sub>u</sub> , M	-		
1	-	14/07/1958	<b>0.04</b>	-	0.18	-	0.31	0.57	0.15	0.20	P	M	S <sub>u</sub>		
16	a-e	28/07/1971	<b>0.05</b>	-	0.14	-	0.39	0.84	0.49	0.43	P	M	S <sub>u</sub>	+	<b>P</b>
22	a-d	04/08/1998	<b>0.03</b>	<b>&lt; 0.0001</b> <sup>(22)</sup>	0.36	0.14	0.68	0.57	0.15	0.17	P	-	S <sub>u</sub> , M		
23	-	17/07/2003	<b>0.09</b>	<b>&lt; 0.01</b> <sup>(23)</sup>	0.33	0.60	0.69	0.89	0.68	0.58	P	-	S <sub>u</sub> , M		
4	-	14/07/1964	0.27	-	<b>0.08</b>	-	0.54	0.74	0.69	0.61	P	-	S <sub>u</sub> , M		
5	a-b	19/07/1964	0.24	-	<b>0.02</b>	-	0.55	0.87	0.81	0.72	P	-	S <sub>u</sub> , M		
14	a-b	23/07/1969	0.23	-	<b>0.06</b>	-	0.35	0.89	0.82	0.78	P	M	S <sub>u</sub>		
15	-	26/07/1969	0.38	-	0.11	-	0.34	0.89	0.87	0.76	P	M	S <sub>u</sub>		
18	-	20/07/1982	0.28	-	0.10	-	0.45	0.46	0.40	0.44	P	S <sub>u</sub> , M	-	~	
25	-	09/10/2011	<b>0.08</b>	<b>&lt; 0.01</b> <sup>(25)</sup>	0.99	0.94	0.85	0.71	0.75	0.81	P	-	S <sub>u</sub> , M		
8	a-d	24/06/1965	1.00	-	<b>0.08</b>	-	<b>0.001</b>	<b>0.009</b>	<b>0.003</b>	<b>0.003</b>	M, S <sub>u</sub>	-	P		
9	-	25/06/1965	1.00	-	<b>0.004</b>	-	<b>0.0002</b>	<b>0.008</b>	<b>0.002</b>	<b>0.002</b>	M, S <sub>u</sub>	-	P		
10	-	26/06/1965	0.43	-	<b>0.06</b>	-	<b>0.002</b>	<b>0.007</b>	<b>0.003</b>	<b>0.001</b>	M, S <sub>u</sub>	P	-	++	
17	-	20/05/1979	0.16	-	0.14	-	<b>0.0001</b>	0.97	0.70	0.29	M	P	S <sub>u</sub>		<b>M</b>
20	-	30/06/1987	1.00	1.00	<b>0.01</b>	<b>0.004</b>	<b>0.02</b>	0.22	0.13	0.12	M	S <sub>u</sub>	P	+	
2	-	13/07/1962	1.00	-	0.39	-	0.20	0.64	0.64	0.51	-	M	P, S <sub>u</sub>	~	
13	a-b	21/08/1966	0.24	-	0.92	-	0.36	<b>0.002</b>	<b>0.004</b>	<b>0.007</b>	S <sub>u</sub>	P, M	-	+	
6	-	09/06/1965	0.19	-	0.90	-	0.40	0.24	0.18	0.15	-	S <sub>u</sub> , P, M	-	~	<b>S<sub>u</sub></b>

<sup>(21)</sup> Taschachbach: 6.3 mm 15 min<sup>-1</sup> (; St. Leonhard im Pitztal-Neurur (Tiwag): 0.7 mm 15 min<sup>-1</sup>)

<sup>(24)</sup> St. Leonhard im Pitztal-Neurur (Zamg): 10.8 mm 10 min<sup>-1</sup> (; St. Leonhard im Pitztal-Neurur (Tiwag): 5.2 mm 15 min<sup>-1</sup>; Taschachbach: 2.1 mm 15 min<sup>-1</sup>)

<sup>(22)</sup> Taschachbach: 6.4 mm 15 min<sup>-1</sup>. (; St. Leonhard im Pitztal-Neurur (Tiwag): 0 mm)

<sup>(23)</sup> St. Leonhard im Pitztal-Neurur (Tiwag): 0.9 mm 15 min<sup>-1</sup> (; Taschachbach: 0.4 mm 15 min<sup>-1</sup>)

<sup>(25)</sup> St. Leonhard im Pitztal-Neurur (Zamg): 0.9 mm 10 min<sup>-1</sup>, (; St. Leonhard im Pitztal-Neurur (Tiwag): 0.5 mm 15 min<sup>-1</sup>; Taschachbach: 0.5 mm 15 min<sup>-1</sup>)