

Saturated Hydraulic Conductivity and Textural Heterogeneity of Soils

Carlos García-Gutiérrez¹, Yakov Pachepsky², and Miguel Ángel Martín¹

¹Universidad Politécnica de Madrid, Madrid, Spain.

²USDA-ARS Environmental Microbial and Food Safety Laboratory.

Correspondence to: Carlos García-Gutiérrez (carlos.garciagutierrez@upm.es)

Abstract. Saturated hydraulic conductivity (K_{sat}) is an important soil parameter that highly depends on soil's particle size distribution (PSD). The nature of this dependency is explored in this work in two ways, (1) by using the Information Entropy as a heterogeneity parameter of the PSD and (2) using descriptions of PSD in forms of textural triplets, different than the usual description in terms of the triplet of sand, silt and clay contents. The power of this parameter, as a descriptor of $\ln K_{sat}$, was tested on a database larger than 19,000 soils. Bootstrap analysis yielded coefficients of determination of up to 0.977 for $\ln K_{sat}$ using a triplet that combines very coarse, coarse, medium and fine sand as coarse particles, very fine sand and silt as intermediate particles, and clay as fines. The power of the correlation was analysed for different textural classes and different triplets using a bootstrap approach. Also, it is noteworthy that soils with finer textures had worse correlations, as their hydraulic properties are not solely dependant on soil PSD.

10 This heterogeneity parameter can lead to new descriptions of soil PSD, other than usual clay, silt and sand, that can describe better different soil physical properties, that are texture dependant.

1 Introduction

Saturated hydraulic conductivity (K_{sat}) is the measure of soil's ability to conduct water under saturation conditions (Klute and Dirksen, 1986). It is an essential parameter of soil hydrology. Soil K_{sat} affects many aspects of soil functioning and soil ecological services, like infiltration, runoff, groundwater recharge and nutrients transport. Knowing values of soil K_{sat} appears to be essential in designing management actions and practices, such as irrigation scheduling, drainage, flood protection, and erosion control.

The dependence of K_{sat} on soil texture has been well documented (Hillel, 1980). Different parameterizations of particle size distributions (PSDs) were suggested to relate K_{sat} and soil texture. It was proposed to use d_{10} , d_{20} , and d_{50} particle diameters (Chapuis, 2004; Odong, 2007) or slope and intercept of the particle size distribution curve (Arya and Paris, 1980; Alyamani and Sen, 1993). Also various functions were fitted to PSDs, and the fitting parameters were related to K_{sat} . For example, Chapuis et al. (2015) proposed to use two lognormal distributions to fit the detailed particle size distribution and to use the lognormal distribution parameters to predict the K_{sat} .

A common way to parameterize the PSD for K_{sat} estimation purposes is using the textural triplet that provides the percentage of coarse particles (sand), intermediate particles (silt), and fine particles (clay). K_{sat} values are estimated using the contents of one or two triplet fractions or just the textural class (Rawls et al., 1998). Representing PSD by textural triplets is the common way to estimate a large number of soil parameters (Pachepsky and Rawls, 2004). The coarse, intermediate, and fine fractions need not to be sand, silt and clay. Martín et al. (2017) showed that different definitions of the triplet, e.g. coarse sand, sand, and medium sand as coarse, fine sand, very fine sand as intermediate, and silt and clay as fine triplet fractions, provide much better inputs for bulk density estimation compared with the standard textural triplet. These different parametrizations of soil texture might put the focus on different soil physical properties, depending on the different particle sizes represented in the triplet.

The heterogeneity of particle size distributions appears to be an important factor affecting hydraulic parameters of soils, including the saturated hydraulic conductivity. Values of K_{sat} depend on both distribution of sizes of soil particles, i.e. soil texture, and the spatial arrangement of these particles, i.e. soil structure. Soil structure can be to some extent controlled by soil texture, since packing of particles is affected by the particle size distributions (e.g., Gupta and Larson, 1979; Assouline and Rouault, 1997; Horn et al., 1994; Jorda et al., 2015). It was recently proposed to use the information entropy as the parameter of the PSD heterogeneity for predicting soil water retention (Martín et al., 2005) and soil bulk density (Martín et al., 2017). Previously, information entropy was used, together with other predictor variables to estimate K_{sat} , using multivariate analysis (Boadu, 2000).

The objective of this work was to test the hypothesis that combining two recent developments -the description of the PSD by different textural triplets, that may represent different soil physical properties dependent on the particle sizes present in the triplet, and the information entropy, as a PSD heterogeneity parameter, that depends on the triplet used- may linearly correlate with $\ln K_{sat}$ and may be seen as a step forward to study the effect of heterogeneity widely recognized in the majority of works that studied the particle size - hydraulic conductivity relationships. By describing the PSD in terms of different triplets, the input information would possibly have different physical interpretations. We wanted to link the heterogeneity of this physical information to the hydraulic behaviour of the soil. Therefore, we explored the possible relationships between $\ln K_{sat}$ values and an entropy metric of soil texture heterogeneity using different size limits of coarse intermediate and fine fractions, using the large USKSAT database on laboratory measured K_{sat} containing more than 19,000 samples. The triplets with highest correlations will be understood as the physical sizes that influence the most in the packing of particles yielding the particular hydraulic behaviour. While PTFs are a useful tool to predict difficult-to-measure soil properties, they sometimes exhibit highly non-linear relationships which are difficult to interpret. While the objective of this paper was the exploration of the physical relation of the new tools and the saturated hydraulic conductivity, the future development of PTFs for prediction purposes is promising avenue for expanding this research. We note that research in this work is a descriptive one. It does not include explanation of what we have observed. However, any explanatory research with mechanisms, models, etc. was historically preceded with the descriptive research.

2 Materials and Methods

2.1 Database description and textural triplet selection

For this study we used USKSAT database in which detailed information can be found in Pachepsky and Park (2015). This database consists on soils from different locations of the USA and contains soils from 45 different sources. We selected only those sources which (a) had data on both K_{sat} and on the seven textural fractions, and (b) presented measurements of K_{sat} made in laboratory with the constant head method. From those, we subset those soils whose sum of mass in the seven textural fractions, i.e. (1) very coarse sand, (2) coarse sand, (3) medium sand, (4) fine sand, (5) very fine sand, (6) silt and (7) clay ranged from 98 to 102%. The final number of soils considered was 19,121. By USDA textural classes the total number of soils are: 12,068 sands, 1,780 loamy sands, 2,123 sandy loams, 104 loams, 135 silt loams, 36 silts, 2,004 sandy clay loams, 78 clay loams, 41 silt clay loams, 345 sandy clays, 0 silty clays and 407 clays. All the samples in the database used are undisturbed soil samples.

We used all possible triplets formed from seven textural fractions. Triplets consisted of coarse, intermediate, and fine fractions. The symbols for triplet showed how the fractions were grouped. For example the “coarse” fraction for the triplet ‘3-2-2’ included very coarse sand, coarse sand and medium sand, the “intermediate” fraction included fine sand and very fine sand, and “fine” included silt and clay; triplet ‘5-1-1’ was the standard one where “coarse” included all five sand fractions, “intermediate” included silt, and “fine” included clay. The amount of possible triplets with 7 textural fractions was 15.

2.1.1 Heterogeneity metric calculation

The Entropy based parametrization of textures introduced in Martín et al. (2001) has as central concept in the Information Entropy (Shannon, 1948). Assuming the texture interval divided into k textural size ranges and that the respective textural fraction contents are p_1, p_2, \dots, p_k , $1 \leq i \leq k$, with $\sum_{i=1}^k p_i = 1$, the Shannon Information Entropy (IE) (Shannon, 1948) is defined by

$$IE = - \sum_{i=1}^k p_i \log_2 p_i \quad (1)$$

where $p_i \log_2 p_i = 0$ if $p_i = 0$. The IE is a widely accepted measure of the heterogeneity of distributions (Khinchin, 1957). In case of three fractions, the minimum value of IE is zero when only one fraction is present, and the maximum value is 1.57 when three fractions are present in equal amounts (see Fig. 1).

For each soil in this study, we grouped the 7 available textural fractions in the 15 possible triplet combinations and calculated the respective triplet’s IE using formula (1). Fig 2 shows ternary graphs of IE calculated for all the soils available in this study but using two different triplets as input. It is clear that, by changing the triplet, the calculated IE values vary differently along the same textural triangle. IE is a measure of heterogeneity, but the triplet used is the substrate for this measure: $(IE, triplet)$, i.e., $(IE, '5-1-1')$.

As we want to compare the linearity, i.e. the proportionality between the heterogeneity of the particular physical sizes chosen and the hydraulic behaviour, we used coefficient of determination, R^2 , as a comparison statistic. As this statistic is highly sensitive to the number of points in the regression, we followed the binning method of Martín et al. (2017) to research the relationship between $\ln K_{sat}$ and soil heterogeneity. Specifically, the range of values of IE was divided into ten bins, the average value of $\ln K_{sat}$ was plotted against the average IE for the bin, i.e. the bin midpoint. This way, the number of points in each relationship was always the same. We want to state that this way, the particular value of R^2 is irrelevant, but it is to be only used as a comparison tool among these regressions.

Linear regressions ‘bin midpoint vs. average bin $\ln K_{sat}$ ’ were computed. Besides the coefficient of determination value for comparison purposes, the goodness-of-fit of these regressions was tested using the Root-mean-square error, RMSE

$$RMSE = \sqrt{\frac{\sum_{i=1}^n (\hat{y}_t - y_t)^2}{n}}$$

where \hat{y}_t are the predicted and y_t are the measured values of $\ln K_{sat}$, and n is the number of soils.

In order to make some inference on these parameters we employed the bootstrap method, which has been used in a very similar context by Schaap and Leij (2000). The bootstrap method is a tool for assessing statistical accuracy. It assumes that one can obtain multiple samples from a single data set, by randomly drawing data with replacement from the original sample. Thus, one can perform the same statistical analysis multiple times in different data sets, obtaining slightly different regression models, thus resulting in an uncertainty in each of the parameters of the model. All of the samples used have the same size as the original sample they were drawn from, so they are generated by random sampling with replacement. We used 1,000 bootstrap data sets, resulting in 1,000 linear regression models. In particular we obtained not just one R^2 and one $RMSE$ value for each IE vs. triplet regression, but one thousand of them, that were summarized into a mean and a standard deviation values. More information on this method can be found in (Efron and Tibshirani, 1993; T. Hastie, 2003).

We took 1,000 samples with size equal the total amount of soils, with repetition, and calculated, for each sample, the coefficient of determination, R^2 and the Root-mean-square error, RMSE. Finally, the mean and standard deviation from these two values, for the 1,000 samples were calculated.

These regressions were obtained for each of 15 triplets and for those of USDA textural classes that were represented in the selected database by more than 50 samples, i.e. all of them except silty clay loams and silts.

2.2 IE variation in the textural triangle

Ternary graphs were used to visually correlate the IE values calculated with the $\ln K_{sat}$ values of the soils in the study. Also, a less visual, but more quantifiable approach, to find out how much of $\ln K_{sat}$ could be explained through IE variation was to find out what ranges of IE are available for soils in different textural classes and compare them to the range of $\ln K_{sat}$ values of soils inside those same textural classes. Also, in order to compare the new tool (IE triplet), we compared these ranges to the ranges computed for (IE , ‘5-1-1’), i.e. to the values of the IE computed with the usual description of soil texture. We wanted to find out if, by changing the triplet, we would obtain a wider range of variation in IE for a given range of $\ln K_{sat}$. This

way we compared if the new descriptions of texture, in form of different triplets, might be suitable for explaining soil physical properties, in particular $\ln K_{sat}$.

For each textural class, we did a sensitivity analysis by calculating the ratio of the range of $\ln K_{sat}$ values inside the textural class versus the range of $\ln K_{sat}$ values of all the soils in the study. The same was done for IE for each triplet.

5 3 Results and Discussion

3.1 The Dataset Overview

Fig. 3 presents the 19,121 soils used in this study in the USDA textural triangle and in the modified '3-2-2' triangle. The density of points reflects the dominance of coarse textural soils in the database. When the triplet is changed, the distribution of points across the triangle changes. By setting the textural fractions to be the '3-2-2' triplet, the distribution of points/soils in the new textural triangle spreads. While there is still a high concentration of soils in the stripe of bigger than 85% of the Coarse fraction, where Coarse 3 includes very Coarse Sand, Coarse sand a Medium Sand, now those soils spread fully from 10 to 100% of the Intermediate-2 fraction, where Intermediate-2 contains Fine and Very Fine Sand. On the USDA textural triangle, most of the soils are clustered in the subtriangle limited by the lines "more than 70% sand" and "less than 20% silt". This new textural triangle allows for a finer look into the sand fraction, revealing the distribution of soils within the USDA sandy textural classes. This finer look might prove itself useful to study physical properties of these soils that are mainly related to the type and amount of sand in them.

Table 1 shows the K_{sat} statistics for the soils in the study. A total of 19,420 soils were used in this study, from which 299 (1.53%) had to be rejected due to missing values. The textural class sand comprises the 63.1% of all the soils, followed by sandy loam (11.1%) and sandy clay loam (10.48%). Five textural classes were poorly represented with percentages less than 1% of the total soils. The K_{sat} values varied between 0.0005 and 841 cm/h being 22.57 the mean value.

3.2 Regression in binned data: IE as a predictor of K_{sat} and $\ln K_{sat}$

Linear regressions for $\ln K_{sat}$ were done to find out the predictive power of the proposed parameter, (IE , triplet), with the 15 possible different triplets that could be archived by grouping the available textural data. Table 2 shows the computed R^2 and $RMSE$ values for the linear regressions using 10 interval bins.

The best triplet in terms of highest mean R^2 value was '4-2-1', with a mean of 0.977 and a standard deviation of 0.002, but the lowest mean $RMSE$ ($\ln(cm/h)$) value (mean=0.207, std = 0.030) was attained with the '1-2-4' triplet. Figure 4 shows a ternary representation representation of the $\ln K_{sat}$ values of the soils of the study on the textural triangle compared to a ternary representation of the IE values of the same soils computed using the '4-2-1' triplet. There is a high visual similarity between these two images, with high $\ln K_{sat}$ value zones, near the lower corners (sandy and silty soils) that correspond to low (IE , '4-2-1') values. The $\ln K_{sat}$ values tend to decrease towards the centre of the triangle. On the other hand, the (IE , '4-2-1') values tend to increase around this point.

The standard triplet ('5-1-1') yielded, for the $\ln K_{sat}$ regression, the R^2 value with this triplet had a mean of 0.960 and a standard deviation of 0.005; the RMSE mean value was 0.339 with a standard deviation of 0.021. The average of the R^2 mean values of the regressions with all possible triplets for $\ln K_{sat}$ was 0.727.

3.3 Predictive power of IE among the USDA textural classes

5 In this section we show how IE works differently among textural classes: using different triplets we can find that the textural classes are predicted differently; what works for some, for others is counterproductive.

Table 3 shows the best triplet, chosen in terms of highest mean R^2 value of all the possible regressions, for each textural class that had $N > 50$. In the table are shown the mean and standard deviation for R^2 , of the 1,000 bootstrap samples for $\ln K_{sat}$ linear regressions. The best R^2 values were obtained for the regression of the sand textural class against the (IE , '5-1-1'), i.e., the IE computed with the standard '5-1-1' clay-silt-sand USDA triplet. The mean value was equal to $R^2=0.987$ for all the regressions. A possible explanation for this triplet being the best among all the other possible triplets, is that sandy soils are the ones that contain percentages of the sand fraction higher than 70%, so their distribution is highly heterogeneous. Minor fractions are now silt and clay, and the information about this two fractions could be very important for the hydraulic properties of the soil, thus the (IE , '5-1-1') triplet yielded the best regression result. One might think that, having such a high concentration of sand particles, is now silt and clay the fractions that made the difference in the packing properties, thus in the saturated hydraulic conductivity values. The high value of R^2 indicates that the relation is very strong in this case.

Almost all sandy textural classes had the highest regression coefficients. Table 3 suggested grouping the textural classes into two superclasses: $SC1$, comprising the textures sandy, sandy clay loam, sandy loam and loamy sand; and $SC2$, with sandy clay, clay, clay loam, loam and silty loam. Soils in $SC1$ are mostly sandy soils, with the exception of the sandy clay textural class which is within the $SC2$ soils which are mostly clayey and loamy soils. The lowest mean R^2 value for the $\log K_{sat}$ regressions in the $SC1$ superclass was 0.742 and the highest one for the $SC2$ class was 0.604. Total number of soils in $SC1$ was 17,975 (94.06% of total soils in the database). $SC2$ contained 1,069 soils (5.59% of total). Tables 4 and ?? show the R^2 and RMSE values for all regressions for the soils in $SC1$ and $SC2$.

For the $SC1$ we observed that the best regression ($R^2=0.986$, $RMSE=0.184$) against $\ln K_{sat}$ was reached with (IE , '4-1-2'). This triplet creates a division among the sand fractions, grouping together very coarse, coarse, medium and fines, and leaving alone the very fines sand. Finally, the fines fractions contains only the silt and clays. Comparing this to the sandy textural class results, where the best triplet was '5-1-1', we observed that now more information from the sandy fraction was required to infer hydraulic properties. The area that the $SC1$ soils cover in the textural triangle and the hydraulic property variation of these soils can be related with a heterogeneity metric associated to triplets that distinguish well among the predominant fraction in that area of the triangle, i.e., sand.

For the $SC2$, best triplet was '1-1-5', with $R^2=0.623$. Regression results were worse than for $SC1$, but this might be just provoked by the nature of $SC2$ itself: these are soils with less sand, thus higher content in clays and aggregating particles. The packing -and consequently the K_{sat} - of these soils is not just mainly affected by the PSD, but also by aggregation, which cannot be accounted for in the IE value, regardless of the triplet used.

Furthermore, the best triplet, '1-1-5', also pointed in this direction: the fines fraction contains medium sand, fine sand, very fine sand, silt and sand particles, while the intermediate fraction contains only the coarse sand, leaving the coarse fraction with the very coarse sand, thus giving more importance to the possibly aggregating particles than a triplet like, '1-4-2' which had R^2 values equal to and 0.033.

5 3.4 Triplets and Scaling Break

In the regressions made with all the soils, it was noteworthy the behaviour of the (IE , '3-1-3'). The average value of all triplets was 0.727, but (IE , '3-1-3') gave an exceptionally low R^2 value of 0.087, being the next lowest (IE , '2-2-3') with a mean R^2 value of 0.235.

The '3-2-3' triplet groups fine sand with silt and clay, and coarse and very coarse sand with medium sand. Kravchenko and Zhang (1998); Wu et al. (1993); Tyler and Wheatcraft (1992) reported the break in scaling where the powerlaw scaling of soil texture occurred in the size range of fine sand. The Particle size distribution scales in a different way in two different regions of the size intervals, and that the change of scaling is produced around the fine sands. The triplet '3-1-3' separates these two regions, maybe bringing forth this scaling break effect. Fig 5. shows how the relationship between $\ln K_{sat}$ and (IE , '3-1-3') could be nonlinear, maybe due to the absence of global selfsimilarity showed in the scaling break.

On the other hand, it is also noteworthy that regressions against (IE , '3-1-3') were actually quite good ($R^2=0.939$) in the SC1, while in the SC2 they were moderate ($R^2=0.045$).

When all the soils are considered together, then (IE , '3-1-3') might fail, due to the scaling break, but when we restrict the study to a certain part of the textural triangle, that effect might diminish to a point where this triplet is even useful to predict some textural derived properties, or maybe the scaling break effect is also restricted to some textural classes and should be further investigated.

As results show, IE is not a powerful $\ln K_{sat}$ predictor by itself, but combined with an input triplet. By changing the triplet, we may focus on certain physical aspects of the soils, but it is also important to keep in mind that this might not work statistically for random groupings of soils that belong to different textures.

3.5 IE variation as a spatial function in the textural triangle

Table 6 shows, for each textural class, the ratio of the percentage of (IE , '5-1-1') against the percentage of $\ln K_{sat}$ range. The same ratio was also calculated using IE for the triplet that gave the best R^2 value in the linear regression against $\ln K_{sat}$. These values can be thought of as how much range of (IE , triplet) can be used to explain a certain variation of $\ln K_{sat}$ inside each textural class, i.e. as how much parametrizing power is available by the IE . In all the textural classes the parametrizing power of the alternative triplet was higher than the one by using the usual clay-silt-sand triplet. For the sand textural class, the triplet which gave the best R^2 regression was '5-1-1' thus the results are the same; the average value of the parametrizing power for the usual triplet was 0.50, while when we change the triplet we obtained 0.79. This shows how, by considering different triplets, combined with IE , a better description/parametrization of $\ln K_{sat}$ can be reached.

3.6 Final Comments

Textural heterogeneity is a crucial factor affecting soil K_{sat} , but it acts along many other ecological factors, as animal activity, root exudates, soil aggregation, etc. In this work we showed that a proper representation of textural heterogeneity, by IE , allows one to (1) demonstrate its effect on $\ln K_{sat}$ by binning samples based on the textural heterogeneity and (2) to statistically
5 parametrize this effect for some textures.

This work has limitations, in particular, the limited available texture data of only seven fractions in the database. The boundaries between coarse, intermediate, and fine fractions can be moved with data from continuous measurements of texture in the fine sand-silt-clay range of particle sized. This may bring the improvements in mean bin $\ln K_{sat}$ estimates for non-sandy soils that could not be achieved in this work.

10 Although globally the IE computed from different triplets show a potential to reflect the effect of soil texture on the $\ln K_{sat}$ values, the different relationship between the IE and the $\ln K_{sat}$ depending on the triplet used might have different possible explanations. While the $IE/\ln K_{sat}$ relationship is found satisfactory in some textural classes, results seem to indicate that the IE parameter cannot reflect with the same efficiency the $\ln K_{sat}$ values in other classes predominating fine particles, in which other processes as aggregation or weathering can not be elucidated by the single textural data input.

15 Overall, the heterogeneity parameter, IE , combined with the different triplet information, appears to be a strong candidate as an input for the development of new pedotransfer functions (PTFs) to predict $\ln K_{sat}$ and probably other soil physical parameters which are strongly dependant on soil particle size distribution.

4 Conclusions

The PSD coarse, intermediate, and fine fractions in soil textural triplets can be redefined from standard 'sand-silt-clay' to other
20 fraction size ranges. The textural heterogeneity parameters obtained for some of the new triplets correlate with soil saturated hydraulic conductivity averaged by ranges of the heterogeneity parameters. This approach allows one to quantify the effect of the textural heterogeneity of saturated hydraulic conductivity of soils. Given that size boundaries of sand, silt, and clay fractions have not originally been established for the purposes of prediction of soil hydraulic conductivity, it may be beneficial to look for other size based subdivisions of particle size distributions which, when used along with other soil properties such
25 as bulk density and organic matter content, may provide better predictions of the saturated hydraulic conductivity.

Acknowledgements. This research work was funded by Spain's Plan Nacional de Investigación Científica, Desarrollo e Innovación Tecnológica (I+D+I), under ref. AGL2015-69697-P.

References

- Alyamani, M. and Sen, Z.: Determination of Hydraulic Conductivity from Complete Grain-Size Distribution Curves, *Groundwater*, 1993.
- Arya, L. and Paris, J.: A physicoempirical model to predict soil moisture characteristics from particle-size distribution and bulk density data, *Soil Sci. Soc. Am. J.*, 1980.
- 5 Assouline, S. and Rouault, Y.: Modeling the relationships between particle and pore size distributions in multicomponent sphere packs: Application to the water retention curve, *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 1997.
- Boadu, F.: Hydraulic Conductivity of Soils from Grain-Size Distribution: New Models, *Journal of Geotechnical and Geoenvironmental Engineering*, 2000.
- Chapuis, R.: Predicting the saturated hydraulic conductivity of sand and gravel using effective diameter and void ratio, *Canadian geotechnical*
- 10 *journal*, 2004.
- Chapuis, R., Weber, S., and Duhaime, F.: Permeability test results with packed spheres and non-plastic soils, *Geotechnical Testing Journal*, 2015.
- Efron, B. and Tibshirani, R.: Improved prediction of unsaturated hydraulic conductivity with the Mualem-van Genuchten model, *Monographs on statistics and applied probability*, Champan and Hall, New York, 1993.
- 15 Gupta, S. and Larson, W.: A Model for predicting Packing density of soils using particle-size distribution, *Soil Sci. Soc. Am. J.*, 1979.
- Hillel, D.: *Fundamentals of Soil Physics*, Academic Press, Inc., <http://gen.lib.rus.ec/book/index.php?md5=497FB6D1C553C06632ADEF0EB958394>, 1980.
- Horn, R., Taubner, H., Wuttke, M., and Baumgartl, T.: Soil physical properties related to soil structure, *Soil Tillage Research*, 30, 187–216, 1994.
- 20 Jorda, H., Bechtold, M., Jarvis, N., and Koestel, J.: Using boosted regression trees to explore key factors controlling saturated and near-saturated hydraulic conductivity, *European Journal of Soil Science*, 2015.
- Khinchin, A.: *Mathematical Foundation of Information Theory*, Dover Publications, 1957.
- Klute, A. and Dirksen, C.: *Hydraulic conductivity and diffusivity: laboratory methods*, 1986.
- Kravchenko, A. and Zhang, R.: Estimating the soil water retention from particle-size distributions: a fractal approach, *Soil Science*, 1998.
- 25 Martín, M., Pachepsky, Y., Rey, J., and Taguas, F.: Balanced entropy index for soil water retention estimation, *Soil Sci.*, 2005.
- Martín, M., Pachepsky, Y., García-Gutiérrez, C., and Reyes, M.: On soil textural classifications and soil texture-based estimations, *Solid Earth Discuss.*, <https://doi.org/https://doi.org/10.5194/se-2017-84>, 2017.
- Martín, M., Rey, J., and Taguas, F.: An entropy-based parametrization of soil textures via fractal modelling of particle-size soil distribution, *Proc. R. Soc. Lond.*, 2001.
- 30 Martín, M., Reyes, M., and Taguas, F.: Estimating soil bulk density with information metrics of soil texture, *Geoderma*, doi:10.1016/j.geoderma.2016.09.008, 2017.
- Odong, J.: Evaluation of empirical formulae for determination of hydraulic conductivity based on grain-size analysis, *Journal of American Science*, 2007.
- Pachepsky, Y. and Park, Y.: Saturated Hydraulic Conductivity of US Soils Grouped According to Textural Class and Bulk Density, *Soil Sci. Soc. Am. J.*, <https://doi.org/doi:10.2136/sssaj2015.02.0067>, 2015.
- 35 Pachepsky, Y. and Rawls, W., eds.: *Development of Pedotransfer Functions in Soil Hydrology*, vol. 30, Elsevier Science, 2004.

- Rawls, W. J., Gimenez, D., and Grossman, R.: Use of soil texture, bulk density, and slope of the water retention curve to predict saturated hydraulic conductivity, Transactions of the ASAE, 1998.
- Schaap, M. and Leij, F.: Improved prediction of unsaturated hydraulic conductivity with the Mualem-van Genuchten model, Soil Sci. Soc. Am. J., 2000.
- 5 Shannon, C.: A mathematical theory of communication, I. Bell Syst. Technol. J., 1948.
- T. Hastie, R. Tibshirani, J. H. F.: The Elements of Statistical Learning, Springer, 2003.
- Tyler, S. and Wheatcraft, S.: Fractal Scaling of Soil Particle-Size Distributions: Analysis and Limitations, Soil Sci. Soc. Am. J., <https://doi.org/http://dx.doi.org/10.2136/sssaj1992.03615995005600020005x>, 1992.
- 10 Wu, Q., Borkovec, M., and Sticher, H.: On particle-size distributions in soils, Soil Sci. Soc. Amer. J., <https://doi.org/doi:10.2136/sssaj1993.03615995005700040001x>, 1993.

IE





Figure 1. *IE* numerical approximation ternary representation: *IE* is computed for a sample of 5051 evenly distributed soils in the USDA textural triangle using the clay, silt and sand fractions as input triplet. This distribution of *IE* is repeated for any textural triangle, when the fractions used for its calculations are the ones at the axes of the triangle. The lowest values for the *IE* are near the vertex of the triangle, i.e. where one fraction dominates above the others. Biggest values are located towards the centre of the triangle, where the distribution fractions are more balanced.

(IE, '1-1-5')



(IE, '5-1-1')




Figure 2. Ternary representations for *IE* calculated for the soils of the study but using different triplets. The usual clay, silt and sand triplet ('5-1-1') was used at the left and the grouping seven textural fractions into '1-1-5' was used as input for the right.




Figure 3. Representation in the USDA textural triangle of the 19,193 soils used in this study. (a) standard sand-silt-clay, i.e. ‘5-1-1’ triplet. (b) the ‘3-2-2’ triplet.




Figure 4. Ternary representations for $\ln K_{sat}$ and $(IE, '4-2-1')$ represented in the USDA textural triangle.




Figure 5. $\ln K_{sat}$ values against IE calculated with the '3-1-3' triplet in 10 interval binnings.

Table 1. Statistical description of K_{sat} (cm/h) values by classes. Soils have also been grouped into two super classes, SC1 and SC2, which can be interpreted as the *sandy class* and not sandy class, respectively. Legend: N , the number of soils in each class; sd, the standard deviation; skew, the skewness number and se, the standard error.

	N	min	1stQ	Median	Mean	3rdQ	max	sd	skew	kurtosis	se
Sandy clay	345	0.00	0.09	0.41	2.72	1.29	60.60	8.29	4.78	24.12	0.45
Sandy clay loam	2.004	0.00	0.12	0.50	3.23	1.67	405.00	17.11	14.04	244.07	0.38
Sandy loam	2.123	0.00	0.28	1.10	4.92	3.67	504.00	18.26	15.60	348.22	0.40
Loamy sand	1.780	0.01	1.37	5.00	9.84	13.80	189.00	13.35	3.86	29.54	0.32
Sand	12.068	0.01	11.80	23.95	32.97	43.40	841.00	32.83	4.01	51.12	0.30
Clay	407	0.00	0.04	0.16	4.07	0.73	421.00	25.49	13.12	196.18	1.26
Clay loam	78	0.01	0.04	0.22	1.26	0.71	38.20	4.56	7.27	57.93	0.52
Silty clay loam	41	0.00	0.08	0.34	18.02	1.67	159.00	43.36	2.60	5.69	6.77
Loam	104	0.01	0.17	0.72	5.77	2.89	52.60	11.26	2.43	5.42	1.10
Silty loam	135	0.00	0.17	0.69	5.20	4.42	53.90	9.65	2.90	9.40	0.83
Silt	36	0.27	1.27	5.21	19.16	22.54	213.00	40.62	3.88	16.30	6.77
SC1	17975	0.00	2.85	14.50	24.05	32.90	841.00	31.28	4.18	51.00	0.23
SC2	1069	0.00	0.07	0.31	3.74	1.36	421.00	17.21	16.54	360.54	0.53
All	19.121	0.00	1.92	13.10	22.89	31.60	841.00	31.06	4.25	51.47	0.22

Table 2. Computed mean and standard deviation (std) for R^2 and RMSE ($\ln cm/h$) values using the bootstrap method for the binned lineal regression of $\ln K_{sat}$ against all possible $(IE, triplet)$.

Triplet	R^2		RMSE	
	mean	std	mean	std
'1-1-5'	0.872	0.048	0.230	0.040
'1-2-4'	0.884	0.029	0.207	0.030
'1-3-3'	0.885	0.042	0.434	0.061
'1-4-2'	0.637	0.084	0.837	0.083
'1-5-1'	0.735	0.051	0.745	0.063
'2-1-4'	0.870	0.017	0.227	0.015
'2-2-3'	0.235	0.064	0.879	0.068
'2-3-2'	0.744	0.019	0.595	0.020
'2-4-1'	0.760	0.023	0.611	0.026
'3-1-3'	0.087	0.031	0.766	0.072
'3-2-2'	0.519	0.075	0.582	0.050
'3-3-1'	0.765	0.009	0.558	0.014
'4-1-2'	0.975	0.004	0.245	0.018
'4-2-1'	0.977	0.002	0.263	0.011
'5-1-1'	0.960	0.005	0.339	0.021
average	0.727		0.501	

Table 3. Summary of triplets for $\ln K_{sat}$ with highest R^2 mean values for regressions using 10 interval binnings. Both the mean value and the standard deviation of R^2 are shown.

Text class	n soils	triplet	mean R2	std R2
Sandy clay	345	'1-2-4'	0.386	0.194
Sandy clay loam	2004	'2-3-2'	0.879	0.054
Sandy loam	2123	'1-1-5'	0.917	0.046
Loamy sand	1780	'2-2-3'	0.742	0.073
Sand	12068	'5-1-1'	0.987	0.005
Clay	407	'1-2-4'	0.604	0.149
Clay loam	78	'4-1-2'	0.276	0.081
Loam	104	'3-1-3'	0.235	0.185
Silty loam	135	'5-1-1'	0.412	0.207

Table 4. R^2 and RMSE ($\ln cm/h$) values for linear regressions of IE vs $\ln K_{sat}$ using the 15 different triplets for the SC1 selection.

triplet	R^2		RMSE	
	mean	std	mean	std
'1-1-5'	0.833	0.056	0.240	0.039
'1-2-4'	0.794	0.046	0.213	0.035
'1-3-3'	0.915	0.031	0.402	0.059
'1-4-2'	0.646	0.088	0.786	0.081
'1-5-1'	0.651	0.055	0.789	0.046
'2-1-4'	0.748	0.028	0.252	0.015
'2-2-3'	0.807	0.015	0.485	0.017
'2-3-2'	0.769	0.019	0.548	0.020
'2-4-1'	0.739	0.028	0.568	0.026
'3-1-3'	0.939	0.007	0.191	0.011
'3-2-2'	0.799	0.009	0.429	0.011
'3-3-1'	0.720	0.009	0.592	0.013
'4-1-2'	0.986	0.002	0.184	0.013
'4-2-1'	0.977	0.002	0.255	0.011
'5-1-1'	0.927	0.011	0.426	0.030
max	0.986		0.184	min
average	0.817		0.424	

Table 5. R^2 and RMSE ($\ln \text{cm}/h$) values for linear regressions of IE vs $\ln K_{sat}$ using the 15 different triplets for the SC2 class.

triplet	R^2		RMSE	
	mean	std	mean	std
'1-1-5'	0.623	0.092	0.634	0.103
'1-2-4'	0.476	0.094	0.556	0.105
'1-3-3'	0.105	0.074	0.770	0.121
'1-4-2'	0.033	0.043	0.822	0.125
'1-5-1'	0.268	0.173	0.633	0.129
'2-1-4'	0.462	0.116	0.478	0.082
'2-2-3'	0.085	0.072	0.728	0.117
'2-3-2'	0.025	0.032	0.741	0.120
'2-4-1'	0.156	0.091	0.760	0.087
'3-1-3'	0.045	0.051	0.480	0.094
'3-2-2'	0.142	0.112	0.591	0.125
'3-3-1'	0.285	0.083	0.717	0.108
'4-1-2'	0.331	0.188	0.465	0.135
'4-2-1'	0.108	0.124	0.503	0.115
'5-1-1'	0.078	0.082	0.570	0.145
max	0.623		0.465	min
avg	0.215		0.630	

Table 6. Comparison of parametrizing power of (IE , '5-1-1') against IE calculated with other triplets. The ranges of variation of IE calculated with the different triplets are compared to the ranges of variation of K_{sat} for the textural classes. The triplets are chosen to be the ones that gav the highest R^2 values at the linear regressions for K_{sat} .

Textural class	%range '5-1-1' / % range $\ln K_{sat}$	% range best triplet / % range $\ln K_{sat}$
silty loam	0.818	1.157
sandy loam	0.561	0.670
sandy clay loam	0.581	0.718
sandy clay	0.385	0.444
sand	0.528	0.528
loamy sand	0.426	0.875
loam	0.274	0.953
clay loam	0.124	1.011
clay	0.816	0.797