

1 Enrique Morán Tejada, on behalf of the co-authors

2 Dear Editor, we are pleased to submit a revised version of the manuscript entitled “Recent
3 evolution and associated hydrological dynamics of a vanishing Tropical Andean glacier”. In this
4 document we respond to every comment done by the reviewers, emphasizing the changes done
5 in the manuscript, or the reasons why some suggested changes (the least) were not implemented.
6 We have undertaken a thorough revision, including changes in figures, with new analyses, large
7 portions of the text, new paragraphs in the introduction and discussion sections as well as all the
8 line-specific changes suggested by the reviewers.

9 In light of the recommendations by the two reviewers, suggesting an English revision of the
10 manuscript (albeit the original manuscript had been already revised by a native English speaker),
11 the manuscript has been carefully checked by a professional English editor

12 We believe that the revised version of the manuscript, thanks to all the changes suggested by the
13 reviewers, is more solid and compelling and hope that the editor takes this revised version for
14 publication.

15

16 **Referee #1**

17 *This paper presented glacier dynamics of a Colombian glacier, which is close to ex-tinction. In*
18 *general, the results are well presented and the paper, of course, fits well into a series of articles*
19 *published recently (e.g. Rabatel et al. 2017). I do recommend publication of this paper after*
20 *some minor revisions, such as adding a few references. I believe that proof reading by a native*
21 *English speaker would be an advantage.*

22 *It is notable that the authors tried to give some attention to volcanic activities; more or less*
23 *similar to that occurring in Mexico (even though not exactly belong to “tropical” glaciers).*

24 *A 57% reduction in the last three decades is comparable to the Peruvian and Bolivian glaciers*
25 *in the eastern cordilleras. However, as the authors mentioned, it is highly dependent on size and*
26 *altitude (and many more).*

27 *References:*

28 *Granados, H.D., Miranda, P.J., Nunez, G.C., Alzate, B.P., Mothes, P., Roa, H.M., Correa,*
29 *B.E.C., Ramos, J.C., 2015. Hazards at ice-clad volcanoes: phenomena, processes, and example*
30 *from Mexico, Colombia, Ecuador, and Chile. In: Haeberli, W., Whiteman, C., Shroder, J.F.*
31 *(Eds.), Chapter 17, Snow and Ice-related Hazards. Risks and Disasters, pp. 607-646*

32 *Morris, J.N., Poole, A.J., Klein, A.G., 2006. Retreat of tropical glaciers in Colombia and*
33 *Venezuela from 1984 to 2004 as measured from ASTER and Landsat images. In: Proceedings of*
34 *the 63rd Eastern Snow Conference, Newark, Delaware, USA, 181-191.*

35 *Rekowsky, I.C., 2016. Variações de área das geleiras da Colômbia e da Venezuela entre 1985 e*
36 *2015, com dados de sensoriamento. MSc Dissertation. Federal University of Rio Grande do Sul*
37 *(UFRGS), Brazil.*

38 Authors: We are grateful by the words of the reviewer. We have included the suggested
39 references, which were not on our original list

40 **Referee #2**

41 The authors greatly appreciate the exhaustive and constructive revision done by Referee #2,
42 who have deeply understood the strengths and shortcomings of our research. In the next
43 paragraphs we will try to respond to every comment done by the referee, explaining in detail the
44 changes done in the manuscript.

45

46 Reviewer:

47 *Despite an appropriately titled section (1.2), the paper lacks a clearly listed and justified*
48 *hypothesis or set of objectives. A reference to “one of” the main objectives is made later in*
49 *description of statistics, but these should be clearer up front in this section. Instead, the premise*
50 *of this study is framed by first summarizing the list of five specific characteristics that Hock et*
51 *al. outlined in 2005, and showing which are not relevant for inner tropics. Hock’s list provides*
52 *a description of likely sources of time variable signals under warming conditions. This does*
53 *provide some sort of guide; the original list of characteristics by Hock et al. was not claimed to*
54 *be exhaustive, as implied by the authors here, but serve as useful heuristic. What are the actual*
55 *hypotheses here?*

56

57 Authors:

58 In order to make the hypothesis and objectives clearer, we have changed the paragraph with
59 objectives descriptions, by the next paragraph (English has been revised):

60 **.... It is expected that changes will be observed in the hydrological dynamics of vanishing**
61 **glaciers, independently of climate drivers. Such hydrological changes may serve as indicators**
62 **of glacier shrinkage, complementing others such as mass balance or areal observations. The**
63 **objective of this work is to provide a comprehensive analysis of the hydrological dynamics of**
64 **a glacierized basin, with the glacier in its last stages prior to extinction. Considering the**
65 **abovementioned characteristics of the hydrology of retreating glaciers, the specific aim is to**
66 **explore changes on time of streamflow dynamics, focusing on the daily cycle, and to discern**
67 **whether such changes are driven by climate or are a result of the diminishing glacierized area**
68 **within the basin.**

69

70

71 Reviewer:

72 *Likewise, a more convincing case should be made about the motivations to examine the details*
73 *of this small glacier close to extinction, especially since the analyses are restricted to the upper*
74 *catchment. What makes these near-extinct ice bodies meritorious objects of analyses with only*
75 *short time series of observations? Moreover, how do the hydrological dynamics impact*
76 *streamflow further downstream? Is there any novel or more generalizable method developed*
77 *here that can find broader application? What is it about the PCA on diurnal timeseries that can*
78 *transcend climatic gradients and be applied to near-extinction glaciers throughout the Andes,*
79 *as is claimed (L161)?*

80 Authors:

81 We are thankful to the reviewer for pointing out these questions. We are positive that there are
82 enough reasons that make this case study scientifically interesting, it is our fault to not have
83 made them clear in the original manuscript. In the next lines we will try to respond to such
84 questions:

85 We are studying this small isolated glacier because is one of the very few monitored glaciers
86 that are in tropical mountains. The fact that the study is restricted to the upper catchment,
87 although it can limit the scientific relevance, is important for the objectives of the work. This is
88 because it allows isolating the hydrological signal of the glacier from the impact of any other
89 environmental factor that may take place in the downstream larger catchment which would
90 make it more difficult to discern the origin of the observed hydrological changes.

91 There is a lot literature about the diurnal cycles of streamflows, but this is the first study in a
92 tropical periglacial environment. Our approach allows to isolate these cycles to just the behavior
93 of the glacier, prevent it from being masked by other diurnal cycles, such that the plants
94 transpiration, for instance. The use of the PCA and the computed hydrological indices, has
95 allowed to neatly identify the timings of flows under the influence of glacier melt or glacier
96 melt + rainfall. The approach used is perfectly applicable to other studies at glacier outlets, so
97 we believe that the results are comparable to other tropical glaciers.

98 Even though the time series is short, in terms of number of years analyzed, the sub-daily
99 resolution of data allows characterizing the daily cycle for 1614 days, in which 3 daily patterns
100 (3 Principal Components) repeat throughout the years. In this sense, we don't think that more
101 years of data would give any other different pattern. The limitation of the short period is that it
102 doesn't allow investigating the long-term changes related to climate trends or patterns of multi-
103 year variability, however the climatology exposed in section 4.1. shows a neat relation between
104 mass balance and temperature/ENSO

105 As we acknowledge in the introduction, the fact that we are in a very humid area, and given
106 the small size of the glacier, the hydrological impact on streamflow further downstream is very
107 limited (in terms of water resources). We don't think this fact diminishes the value of our
108 research, because our approach can be taken as an example in tropical areas with a marked
109 dry season (Peru or Bolivia), where glacier retreat can have a serious impact on communities
110 and ecosystems living downstream. There can be other impacts on streamflow downstream:
111 macroinvertebrates that depend on glacier melt; release of heavy metals or other products
112 from volcanic activity accumulated in the ice during centuries (López-Moreno et al., 2017).
113 However, the consideration of the latter is way far from the objectives of our research.
114

115 All these previous considerations have been included at different parts of the revised manuscript
116 (objectives section and discussion section)

117 The next paragraph has been included in the "hypothesis and objectives" section: (English has
118 been revised)

119 **The case study is a small glacier (see description in Section 2) in the Central Colombian**
120 **Andes and the catchment that drains the water at the snout of the glacier. It is one of the**
121 **very few monitored glaciers in the tropical Andes (Mölg et al., 2017; Rabatel et al., 2017)**
122 **and represents an ideal case, where the hydrological signal of the glacier can be studied in**
123 **isolation from any environmental factors that may occur in the downstream areas. For**
124 **this reason, the approach used (see Section 3.3) can be applied to similar environments,**
125 **and the obtained results can be representative of expected hydrological dynamics in other**
126 **glacierized areas in the Andes, with glaciers close to extinction.**

127

128 The next paragraph has been included at the end of the “discussion and conclusions” section:
129 (English has been revised)

130 **The added value of studying the hydrology related to this small-sized and near-extinct glacier**
131 **is that the changes observed in the hydrology of the catchment could be directly attributed to**
132 **the dynamics of the glacier and the climate that occurs at the same time-scale; contrary to**
133 **catchments containing large glaciers that respond with a larger temporal inertia to**
134 **environmental changes. Hydrological analyses were restricted to the upper catchment**
135 **because the streamflows measured at the snout of the glacier are not influenced by the signals**
136 **of other environmental processes that may occur downstream (e.g., forest clearing or**
137 **increasing grazing). The methodological approach, including the PCA and the hydrological**
138 **indices computed over sub-daily resolution data demonstrated itself as viable for detecting**
139 **changes on the diurnal cycle of the glacier and can be applied to other small glaciers of the**
140 **tropical Andes that respond rapidly (at sub-annual scales) to environmental forcing. The**
141 **necessity for in situ observations on a fine scale in order to improve accuracy on future**
142 **estimations of water availability related to glacier retreat is emphasized.**

143

144 Reviewer:

145 *They also claim confusingly that the glacier in Colombia lacks seasonality of precipitation*
146 *because of its inner tropical location (L123, 124). However, they show in Fig. 2 and on L187*
147 *that there are two contrasting seasons of precip. Clarify this.*

148 Authors:

149 That’s right, we were referring to seasonal patterns in temperature. We added the next
150 sentence to clarify:

151 Seasonal runoff variation dependent on ablation and accumulation periods at latitudes with
152 markedly variable temperature and/or precipitation seasonal patterns. **In the case of**
153 **temperature**, this does not apply to glaciers in the inner tropics.

154

155 Reviewer:

156 *Study site: Why are the ecological zones of the watershed described for the Rio Claro Basin? Is*
157 *this relevant?*

158 Authors:

159 We included such description for the reader to get a general idea of how the environment is.
160 Moreover, we refer later in the discussion section to the páramos ecosystem as a buffer of
161 hydrological dynamics. Considering that this description only takes 2 lines, we think it’s ok to
162 leave it as it is.

163

164 Reviewer:

165 *Data and Methods: This study refers to a 'network' of observations (How many stations? Where*
166 *are they?) in the Río Claro basin initiating in 2009, but also that for reasons of data quality*
167 *concerns only sensor data from 2013-2017 were used. Still, it is unclear exactly what*
168 *instruments were used, and where. The map shows only two stations. Is this the extent of the*
169 *network?*

170 Authors: For this particular study, we just used the stations located at the surroundings of the
171 glacier, but in the Río Claro basin there is a network hydrological and meteorological stations
172 covering the whole altitudinal gradient. We have included them all in the map of Río Claro
173 basin, and make it clear in the text that from the whole network, in this work we just used the
174 ones in the upper catchment: (English has been revised)

175 **The experimental site of the Río Claro basin has been monitored since 2009, with a**
176 **network of meteorological and hydrological stations located at different tributaries of the**
177 **Río Claro River, covering an altitudinal gradient of 2700 – 4900 m.asl. As this research is**
178 **focused on the upper catchment in which the glacier is located for the present study, we**
179 **used data from just the stations located at the Conejeras glacier snout (Figure 1, bottom**
180 **map).This includes one stream gauge (with associated rating curve) measuring 15-minute**
181 **resolution water discharge (m³ s⁻¹); one temperature station measuring hourly**
182 **temperature (°C) (both stations located at 4662 m.asl); and one rain-gauge measuring 10-**
183 **minute precipitation (mm, the station located at 4413 m. asl).**

184 Reviewer:

185 *What does it mean that in 2013 the "sensors stabilized"? The map in Fig. 1 locates a*
186 *precipitation gage below 4400 m, but the text refers to a station at 4413 m. Is this the same?*
187 *Also, the Brisas climate station that is used for longer time series (Fig. 2) is not described in the*
188 *data section, nor is it identified on the maps, and needs to be. it in the basin? The elevation is*
189 *identified as 2721 m, which means it is at the extreme lower end (the watershed is defined as*
190 *spanning to 2700 m). What actually defines the pour point of this watershed?*

191 Authors:

192 Up to 2013, the sensors and the logging systems experienced technical problems, and they were
193 solved with more or less frequency in the field trips. Thus, from 2009 to 2013 we found many
194 inconsistencies in the data, outliers, inhomogeneities, etc.. which prevented us from using the
195 data from those years. The sensors and loggers stopped giving trouble in 2013, and series from
196 then do not contain any more error. We changed the text as following:

197 **Even though these data have been available since 2009, the sensors and loggers**
198 **experienced technical problems and numerous inhomogeneities; thus, out-of-range values**
199 **and empty records were present in the data series. From 2013, the technical problems**
200 **were solved and the data is suitable for analysis**

201 As for the station at 4413 m, our coordinates for locating the station in the map were wrong. We
202 have corrected it,

203 As for Brisas station, we have included it in the map, and describe it in the text. This station is
204 not in the watershed, indeed, is located right in the opposite hillside.

205 The pour point of the watershed is defined by a gauge station at the Río Claro basin, illustrated
206 in the map

207

208 Reviewer: *Glacier evolution data: Impressive mass balance monitoring has been maintained,*
209 *and this is not easy. But the mass balance data presented are only shown as summation bar*
210 *graphs. We don't see stake specific information.*

211 Authors: We have included a new figure, containing more information (mass balance per
212 elevation ranges and time), and explained it in the results section.

213 Reviewer: *What does the "topographic surveys" comprise? Theodolite? GPS? And how is the*
214 *satellite imagery able to reconstruct elevation to "support direct topographic surveys"? This is*
215 *not explained.*

216 Authors: We used Differential GPS. Satellite imagery analyses were only addressed to compute
217 the surface of the glacier, not to reconstruct elevation.

218

219 Reviewer:


220 *Results:*

221 *Climatology and glacier's evolution:*

222 *There is a strong El Nino signal in temperature. This is not surprising. Precipitation seasonality*
223 *is also unsurprising, and reported in the Setting description, so not as appropriate to repeat*
224 *also in results. There is mention of spectral analysis on the time series showing high power at*
225 *48-month frequency that is not surprising, although those data are not displayed. Finally, it is*
226 *awkward to overlay the glacier mass balance data that are not introduced until Fig. 3. Why not*
227 *show how the temperature data at the station closest to the glacier compares to that at the*
228 *distant Brisas station for the period of overlap (2013-17)?*

229

230 Authors: We have removed the precipitation seasonality plot, and the glacier mass balance data
231 from the plots. Instead we have included the series of temperature at the upper location, as is
232 suggested later by the reviewer. As for the high-power at 48-month frequency, we must say that
233 these data was displayed as small text tables embedded in the plots. The new corrected figure
234 looks like this:



235

236

237


238

239 Reviewer: *The glacier change is actually just a surface area evolution, and not a total mass or*
 240 *topographic (surface elevation) change. The polygons mapped in Fig. 3 are not very*
 241 *meaningful, and seems to have curious boundaries shifting (in the higher elevation). By*
 242 *referring to 'global balance', is this the same as the net cumulative mass balance? Likewise,*
 243 *would a cumulative surface area loss and cumulative mass balance curve not suffice to tell the*

244 story, with maybe a table showing area, uncertainty and satellite image ID used per epoch
245 rather than the current Fig. 3 b?. Relative retreat is not convincingly depicted with these data,
246 and images of varying resolution. For example, the authors make an association of sharp
247 retreat post 2014 in the upper glacier. However, Fig. 3 depicts only overlapped outlines of
248 dated polygons in which the later images of 2016, 2017 seem to feature higher resolution, as
249 well as an offset (gain) in the mapped area in the upper glacier for 2016.

250 Authors: we are aware of the limitations detailed by the reviewer. First of all we must say that
251 the analysis of the mass balance data and changes in glacier area is not an objective of this
252 research, *per se*. We included it in the paper as a general framework to help understanding the
253 hydrological dynamics at the outlet of the glacier. There is already a research article (Rabatel et
254 al., 2017) showing the glacier thickness change, thanks to a campaign of measurements with ice
255 penetrating radar undertaken in 2014.

256 This is why we didn't perform a deep analysis of mass balance and surface evolution. With
257 regard to mass balance, we have included a plot of annual change by elevation ranges:




258

259 As for the different resolution of the polygons' boundaries, we also admit this limitation, whose
260 origin is purely methodological: some polygons were derived from differential GPS
261 measurements, and others from images of different satellites (with different resolution). The
262 resulting figure is more informative than analytical, and we thought it was a good way of
263 describing the recession experienced by the glacier.

264 We must say however, that given that this part of the paper is the most problematic one, we
265 have simplified the figure by putting only 4 polygons, all derived from the same method (NDSI,
266 from satellite), for the years 2006, 2010, 2013 and 2017. Thus, the glacier recession during the
267 study period can be easily observed, and the year-to-year evolution can be extrapolated from the
268 yearly mass balance (above figure). The resulting whole figure would be like this:

269

270




271

272

273 Reviewer: *The authors infer temperature increase and less snowfall as key drivers for*
 274 *accelerated recession, yet do not show the time series of temperature from the upper elevation*
 275 *site. It would be more informative to see the time series of temperature loggers located closer to*
 276 *the glacier being correlated to mass balance. This would also provide a test of the idea that*
 277 *some break occurred around 2013-14. The inference of albedo alteration via volcanism is only*
 278 *anecdotal. Any figures or photographs to substantiate the volcanic ash hypothesis?*

279 Authors: This assertion is not totally right. We don't mention temperature increase (in terms of
 280 a trend), but sustained temperatures above 0°; and is not true that we don't show the time series
 281 of temperature from the upper elevation site; it is clearly depicted in the bottom left plot of
 282 Figure 6. As suggested by the reviewer, to such plot we have added the mass balance series, to
 283 see the correlation of the two variables. (see fragment of the figure, below)



284

285 As for the volcanic ash hypothesis, we have included in figure 3 2 photographs (there is not
 286 actual data from measurements) that show the content of volcanic ashes on the surface (see
 287 below)

288



289

290

291

292

293 Reviewer:

294 *Hydrological dynamics:*

295 *Are the temperatures from the same location as discharge? How well constrained is the*
 296 *discharge (presumably, this is a stage recorder with an associated rating curve, but none of this*
 297 *information is provided)? Using the coded names for variables is awkward; why not use full*
 298 *terms? The analytical approach of using PCA on the hourly stream flow statistics is interesting,*
 299 *but it is unclear if any meaningful trends can be extrapolated given that the data span only 4*
 300 *years.*

301 Authors: yes, temperatures are recorded at the same location as discharge (it is shown in the
 302 map of figure 1). Yes, the discharges are measured at a stream gauge with an associated rating
 303 curve, and now we specify it in the methods section.

304 As for the use of the coded names, we believe that it makes the text more dynamic to say:
 305 “correlations between Q_{max} , Q_{range} , Q_{slope} and $totalQ$, with T_{max} and T_{mean} ”, than
 306 “correlations between the maximum streamflow, the range between baseflow and maximum
 307 streamflow, the slope of the rising limb of streamflow, and the total daily streamflow with
 308 maximum and mean temperature”. In Table 1 we specified the meaning of each coded name,
 309 and with a quick look at it is easy to get familiarized with them.

310

311 It is clear that we cannot extrapolate any climatological trend from our 5-years data (5, not 4),
 312 nor it was our intention. It is explicitly admitted in the discussion. However, when considering

313 the monthly aggregation of the variables (Figure 6) and their evolution, it is correct, at least
314 statistically, to speak about trends. In order to avoid confusion, we have changed the term
315 "trend" throughout the text, by more suitable terminology
316

317
318

319 Reviewer:

320 *Changes in the runoff-climate relationship:*

321 *Similarly, it is difficult to find the observed inflection point in mid-2016 to be significant when*
322 *the total time series is short.*

323 Authors:

324 Here we partially disagree. It is true that the inflection point may not be significant
325 climatologically (at least there is no way to know it, as for the short time series available). But
326 what our analyses from figures 4, 6, 7 and 8 reveal is that not only there is a change from
327 increasing to decreasing flows; there is also a change in the timing of most analyzed indices, as
328 well as a change in the frequency of the two main types of hydrographs observed. Even if it's
329 not climatically significant, it must mean something. For us the key aspect of the analysis is
330 that we don't observe the same increasing-decreasing pattern in either temperature or
331 precipitation. And the correlations (which are statistically significant, as are based on 56
332 monthly data) between flows and temperature (precipitation) change drastically before and
333 after such turning point. In simple words: the flows increase and decrease independently of
334 temperature and precipitation (this is evident from the analysis). This is why we speak of a
335 non-climate-driven hydrological change.

336 There is a scientific debate on the existence of such tipping points in areas subjected to glacier
337 recession, and we think that our work contributes positively to such debate, while admitting
338 the limitations imposed by the data. All these things are properly discussed in the "discussion
339 section", and the limitations are also recognized. In fact, we emphasize that the "turning
340 point" observation cannot be taken conclusively
341

342

343 Reviewer: *The statement that: "...the runoff increases because glacier mass becomes more*
344 *sensitive to energy exchange as it gets smaller" is not evident. All that seems fair to say is that*
345 *runoff decreases in direct proportion to less mass loss. Moreover, the mass balance seems most*
346 *closely controlled by MEI. Yet, we are not shown how well the temperature locally at the glacier*
347 *match the longer record.*

348

349 Authors: We have removed such statement to avoid confusion, and have added that
350 **"streamflow increases and decreases in direct proportion to mass balance change indicating**
351 **the strong dependence of runoff to glacier melt"**.

352 As for the local temperatures, this has been corrected in figure 2, and the match with the
353 longer record seems evident.
354

355
356

357 Reviewer: *The presentation of results is difficult to follow given the use of abbreviated names of*
358 *variables. Also in final Fig. 8, the shift in use of red-blue symbols between (a) and (b) is*
359 *confusing to follow.*

360 Authors: We still think that is less confusing to use the coded names (once you are familiarized
361 with them), rather than the long names of variables (i.e. slope of the rising limb of runoff →
362 Qslope). For the more common variable names, such as temperature, or precipitation, we
363 don't use abbreviations. Finally, there was no shift in the use of red-blue symbols between a)
364 and b) in Figure 8. Blue triangles correspond to the months previous to 2016, and red circles to
365 the months after 2016.
366

367 Reviewer:

368 *General terminology: Glaciation can refer to landscapes previously covered with ice, but no*
369 *longer. Instead of deglaciation, the modern or actual glacier loss is better described as glacier*
370 *recession. Water yield is an unfamiliar term; why not use discharge?*

371 Authors: we are thankful for this observation and have change the “deglaciation” term
372 accordingly throughout the paper. Even if we are familiarized with the term “water yield” (it is
373 used in the fields of hydrology/hydrological modelling to name the volume of water produced
374 by the catchment), it's actually the same as “water discharge”, so we have also changed it across
375 the document.

376

377 Authors: all the line specific edits detailed below have been done as suggested by the reviewer

378 *R: L62 change regarding to of, and just use glacier not glaciers' In the lines following, the list*
379 *of research topics can be made neater by removing the redundant 'and regarding' from each*
380 *line.*

381 *L85 and following: sentence is long and should be rephrased.*

382 *L92 territories is awkward word; landscapes?*

383 *L99 glaciers are not a main source of water for Lima! This hyperbole needs to be moderated.*
384 *Look carefully at Vuille et al. (published in 2018), who make it clear that only in La Paz is*
385 *glacier melt contributing significantly to municipal water supply.*

386 *L112 do streams really remain constant flow? This implies no variability, and that is not the*
387 *case.*

388 *117 do not use "unglaciated" here, as glaciated can be used to describe landscapes at one time*
389 *in past having glacier cover. Better to use "non-glacierized" to refer to regions without actual*
390 *glaciers. Edit this throughout the text.*

391 *L151-155: this sentence is dense and needs clarifying, and perhaps split into two*

392 *L188 should say "the dry seasons" and "wet seasons" to emphasize more than one exist per*
393 *year.*

394 *L198 change has been to have been*

395 *L203 unclear what 'consecutively located' means*

396 *L205 reword sentence; not clear to say located at surroundings*

- 397 *L231 losses should be loses*
- 398 *L238 should be cloud-free*
- 399 *L256 should be over time*
- 400 *L265 missing full stop.*
- 401 *L308 double full stop; delete one.*
- 402 *L325 yield is not a familiar term. Suggest discharge is better. Also, gauging station is better*
403 *than gauge station.*
- 404 *L371 missing "-" to demarcate range of precipitation rates (mm per day)*
- 405 *L402 lacks full stop.*
- 406 *L409 lacks full stop.*
- 407 *Fig. 2 caption: two c)'s ; make last one d) This figure 2(d) should have some range of*
408 *variability around only a mean.*

409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430

431 **Recent evolution and associated hydrological dynamics of a vanishing Tropical Andean**
432 **glacier: *Glaciar de Conejeras*, Colombia**

433

434 Enrique Morán-Tejeda¹, Jorge Luis Ceballos², Katherine Peña², Jorge Lorenzo-Lacruz¹ and
435 Juan Ignacio López-Moreno³

Eliminado: q

436

- 437 1. Department of Geography. University of the Balearic Islands. Palma, Spain
438 2. Instituto de Hidrología, Meteorología y Estudios Ambientales (IDEAM). Bogotá, Colombia
439 3. Pyrenean Institute of Ecology. Consejo Superior de Investigaciones Científicas. Zaragoza, Spain.

440

441

442 **Abstract**

443 Glaciers in the inner tropics are rapidly retreating due to atmospheric warming. In Colombia, this
444 retreat is accelerated by volcanic activity, and most glaciers are in their last stages of existence.
445 There is general concern about the hydrological implications of receding glaciers, as they
446 constitute important freshwater reservoirs and, after an initial increase in melting flows due to
447 glacier retreat, a decrease in water resources is expected in the long term as glaciers become
448 smaller. In this paper, we perform a comprehensive study of the evolution of a small Colombian
449 glacier, Conejeras (Parque Nacional Natural de los Nevados) that has been monitored since
450 2006, with special focus on the hydrological response of the glacierized catchment. The glacier
451 shows great sensitivity to changes in temperature and especially to the evolution of the [El Niño](#)
452 [Southern Oscillation \(ENSO\)](#) phenomenon, with great loss of mass and area during El Niño warm
453 events. Since 2006 it has suffered a 37% reduction from 22.45 ha in 2006 to 12 ha in 2017, with
454 an especially abrupt reduction since 2014. During the period of hydrological monitoring (June
455 2013 to December 2017) streamflows at the outlet of the catchment experienced a noticeable
456 cycle of increasing flows up to mid-2016 and decreasing flows afterwards. The same cycle was
457 observed for other hydrological indicators, such as slope of the rising flow limb and the monthly
458 variability of flows. We observed an evident change in the daily hydrograph: from a predominance
459 of days with a pure melt-driven hydrograph up to mid-2016, to an increase in the frequency of
460 days with flows less influenced by melt after 2016. Such a hydrological cycle is not directly related
461 to fluctuations of temperature or precipitation; therefore, it is reasonable to consider that it is the
462 response of the glacierized catchment to retreat of the glacier. Results confirm the necessity for
463 small-scale studies at a high temporal resolution in order to understand the hydrological response
464 of glacier-covered catchments to glacier retreat and imminent glacier extinction.

Eliminado: ,

Eliminado: kind of

Eliminado: or

465

466

467 **Key words:** glacier retreat, melting flows, tropical glaciers; hydrological change; tipping point

468

469

470

471

472

473

474

475

476

477

482 1. Introduction

483 1.2 Andean glaciers and water resources

484 Glacier retreat is one of the most prominent signals of global warming; glaciers from most mountain
485 regions in the world are disappearing or have already disappeared due to atmospheric warming
486 (Vaughan et al., 2013). Of the retreating mountain glaciers worldwide, those located within the
487 tropics are particularly sensitive to atmospheric warming (Chevallier et al., 2011; Kaser and
488 Omaston, 2002). Their locations in the tropical region involve a larger energy forcing in terms of
489 received solar radiation compared to other latitudes. Unlike glaciers in mid and high latitudes,
490 which are subject to freezing temperatures during a sustained season, tropical glaciers may
491 experience above-zero temperatures all year round, especially at the lowest elevations, involving
492 constant ablation and rapid response of the glacier snout to climate variability and climate change
493 (Francou et al., 2004; Rabatel et al., 2013). As a result of atmospheric warming since the mid-
494 20th century, glaciers in the tropics are seriously threatened, and many of them have already
495 disappeared (Vuille et al., 2008). Of the tropical glaciers, 99% are located in the Central Andes
496 and constitute a laboratory for glaciology (see review in Vuille et al., 2017), including studies of
497 glacier response to climate forcing (e.g. Favier et al., 2004; Francou et al., 2004, 2003; López-
498 Moreno et al., 2014), hydrological and geomorphological consequences of glacier retreat (Bradley
499 et al., 2006; Chevallier et al., 2011; Kaser et al., 2010; López-Moreno et al., 2017; Ribstein et al.,
500 1995; Sicart et al., 2011) and the vulnerability of populations to risks associated with glacier retreat
501 (Mark et al., 2017). Perhaps the glaciers in the most critical situation in the Andean mountains
502 are those located in the inner tropics, including the countries of Ecuador, Venezuela and
503 Colombia (Klein et al., 2006; Rekowski, 2016). In the latter country, a constant glacier recession
504 since the 1970s has been reported, with an acceleration since the 2000s (Ceballos et al., 2006;
505 Rabatel et al., 2013), and most glaciers are in danger of disappearing in the coming years
506 (Poveda and Pineda, 2009; Rabatel et al., 2017). In the outer tropics, the variability of glacier
507 mass balance is highly dependent on seasonal precipitation; thus, during the wet season
508 (December-February) freezing temperatures ensure seasonal snow cover that increases the
509 glaciers' surface albedo and compensates mass balance losses of the dry season. In contrast,
510 for glaciers of the inner tropics, ablation rates remain more or less constant throughout the year
511 due to the absence of seasonal fluctuations of temperature and to a freezing level which is
512 constantly oscillating within the glaciers' elevation ranges. Therefore, the mass balance of these
513 glaciers is more sensitive to inter-annual variations of temperature, and they are much more
514 sensitive to climate warming (Ceballos et al., 2006; Favier et al., 2004; Francou et al., 2004;
515 Rabatel et al., 2013, 2017). In Colombia, this situation is further aggravated by the location of
516 glaciers near or on the top of active volcanoes. The hot pyroclastic material emitted during volcanic
517 eruptions and the reduced albedo of glacier surfaces by the deposition of volcanic ash, have
518 notably contributed to rapid glacier recession in these areas (Granados et al., 2015; Huggel et
519 al., 2007; Rabatel et al., 2013; Vuille et al., 2017).

520 Current glacier recession in the Andes involves the loss of natural scientific laboratories (Francou
521 et al., 2003) and of landscape and cultural emblems of mountainous areas (IDEAM, 2012; Rabatel
522 et al., 2017). But in more practical terms, the vanishing of glaciers has a major impact on
523 livelihoods of communities living downstream, including potential reduction of freshwater storage
524 and changes in the seasonal patterns of water supply by downstream rivers (Kaser et al., 2010).
525 Glaciers constitute natural water reservoirs in the form of ice accumulated during cold and wet
526 seasons, and they provide water when ice melts during above-freezing temperature seasons.
527 The hydrological importance of glaciers for downstream areas depends on the availability of other
528 sources of runoff, including snow melt and rainfall. Therefore, water supply by glaciers becomes
529 critical for arid or semi-arid regions downstream of the glacierized areas, buffering the lack of
530 sustained precipitation or water provided by seasonal melt of snow cover (Rabatel et al., 2013;
531 Vuille et al., 2008). Such is the case for the western slopes of the tropical Andes: in countries like
532 Peru or Bolivia, with a high variability in precipitation and a sustained dry season, the contribution
533 of glacier melt is crucial for socioeconomic activities and for water supply, especially since it is
534 one of the main sources of water for the highly populated capital cities such as La Paz (Kaser et
535 al., 2010; López-Moreno et al., 2014; Soruco et al., 2015; Vuille et al., 2017). In more
536 humid/temperate regions (i.e. the Alps or western North America) the melt of seasonal snow
537 cover provides the majority of water during the melt season (Beniston, 2012; Stewart et al., 2004)

- Eliminado: regarding
- Eliminado: s'
- Eliminado: and regarding
- Eliminado: ,
- Eliminado: also regarding
- Eliminado:
- Eliminado:
- Eliminado: deglaciation

- Eliminado: ; hence

- Eliminado: rs
- Eliminado: '
- Eliminado: ,
- Eliminado: deglaciation
- Código de campo cambiado
- Con formato: Español (alfabetización tradicional)
- Eliminado: (Huggel et al., 2007; Rabatel et al., 2013; Vuille et al., 2017).
- Con formato: Español (alfabetización tradicional)
- Eliminado: Regardless of
- Eliminado: ,
- Eliminado:
- Eliminado: territories
- Eliminado: glaciated
- Eliminado: of Lima and

559 and glacier melt is a secondary contributor. However, even in this region, water availability can
 560 be subject to climate variability, and the occurrence of dry and warm periods that comprise thin
 561 and brief snow cover may involve glacier melt as the main source of water during such events
 562 (Kaser et al., 2010). In the inner tropics, glaciers may not constitute the main source of water for
 563 downstream populations, as the seasonal shift of the Intertropical Convergence Zone (Poveda et
 564 al., 2006) assures two humid seasons every year; however, the loss of water from glacier melt
 565 can affect the eco-hydrological functioning of the wetland ecosystems called “*páramos*”, which
 566 are positioned in the altitudinal tier located below that of the periglacial ecosystem (Rabatel et al.,
 567 2017). Agriculture and livestock in Colombian mountain communities are partly dependent on
 568 water from these important water reservoirs that provide water flow to downstream rivers even
 569 during periods of less precipitation.
 570

Eliminado: located

Eliminado: constant

571 1.2. Hypothesis and objectives

572 The present work is focused on the hydrological dynamics of a Colombian glacier near extinction
 573 due to prolonged deglaciation. Hock et al. (2005) presented a summary of the effects of glaciers
 574 on streamflow compared to non-glacierized areas. The main characteristics of streamflow can be
 575 summarized as follows (Hock et al., 2005):

Eliminado: unglaciated

- 576 - Specific runoff dependence on variability of glacier mass balance. In years of mass
 577 balance loss, total streamflow will increase as water is released from glacier storage. The
 578 opposite will happen in years of positive mass balance.
- 579 - Seasonal runoff variation dependent on ablation and accumulation periods at latitudes
 580 with markedly variable temperature and/or precipitation seasonal patterns. In the case of
 581 temperature, this does not apply to glaciers in the inner tropics
- 582 - Large diurnal fluctuation in the absence of precipitation as a result of the daily cycle of
 583 temperature and derived glacier melt.
- 584 - Moderation of year-to-year variability. Moderate percentages (10 to 40%) of ice cover
 585 fraction within the basin reduces variability to a minimum, but it becomes greater at both
 586 higher and lower glacierization levels.
- 587 - Large glacierization involves a high correlation between runoff and temperature, whereas
 588 low levels of glacier cover increase runoff correlation with precipitation.

Eliminado: T

589 However, under warming conditions that lead to glacier retreat, the hydrological contribution of
 590 the glacier may notably change from the aforementioned characteristics. The retreat of a glacier
 591 is a consequence of prolonged periods of negative mass balance, the result of a disequilibrium
 592 in the accumulation/ablation ratio that involves an upward shift of the equilibrium line (the
 593 elevation at which accumulation and ablation volumes are equal), and an increase of the ablation
 594 area with respect to the accumulation area (Chevallier et al., 2011). As a result, the glacierized
 595 area is increasingly smaller compared to the non-glacierized area within the catchment in which
 596 the glacier is settled. Under such conditions of sustained negative mass balance, the hydrological
 597 response of the glacier will be a matter of time-scales (Chevallier et al., 2011; Hock et al.,
 598 2005). The total runoff production of the retreating glacier comprises a tradeoff between two
 599 processes: on one side, an acceleration of glacier melt that will increase the volume of glacier
 600 outflows independent of the volume precipitated as snowfall or rainfall; on the other side, water
 601 discharges from the catchment decrease because the water reservoir that represents the glacier
 602 is progressively emptying (Huss and Hock, 2018). Thus, the contribution of glacier melt to total
 603 water discharge will initially increase, as the first process will dominate over the other; however,
 604 after reaching a discharge peak, the second process dominates, leading to a decrease in water
 605 discharge until the glacier vanishes. In terms of runoff variability, there is also a different signal
 606 between initial and final stages of glacier retreat: on a daily basis, the typical diurnal cycle of
 607 glacier melt will exacerbate at the initial stages (larger difference between peak and base runoff)
 608 and will moderate at the final stages. However, in terms of year-to-year variability, there can be
 609 a reduction or increase at the initial stages, depending on the original glacierized area. And for the
 610 long term, increasing variability should be expected, as the water discharge will correlate with
 611 precipitation instead of temperature because the percentage of runoff from glacier melt decreases
 612 with decreasing glacierization (Hock et al., 2005).

Eliminado: ,

Eliminado: glaciated

Eliminado: a

Eliminado: ted

Eliminado:

Eliminado: ,

Eliminado: glaciated

Eliminado: , a

Eliminado: an increase in

Eliminado: water yield

627 It is expected that changes will be observed in the hydrological dynamics of vanishing glaciers,
628 independently of climate drivers. Such hydrological changes may serve as indicators of glacier
629 shrinkage, complementing others such as mass balance or areal observations. The objective of
630 this work is to provide a comprehensive analysis of the hydrological dynamics of a glacierized
631 basin, with the glacier in its last stages prior to extinction. Considering the abovementioned
632 characteristics of the hydrology of retreating glaciers, the specific aim is to explore changes on
633 time of streamflow dynamics, focusing on the daily cycle, and to discern whether such changes
634 are driven by climate or are a result of the diminishing glacierized area within the basin.

Eliminado: [1]
Eliminado: glaciated

635 The case study is a small glacier (see description in Section 2) in the Central Colombian Andes,
636 and the catchment that drains the water at the snout of the glacier. It is one of the very few
637 monitored glaciers in the tropical Andes (Mölg et al., 2017; Rabatel et al., 2017) and represents
638 an ideal case where the hydrological signal of the glacier can be studied in isolation from any
639 environmental factors that may occur in the downstream areas. For this reason, the approach
640 used (see Section 3.3) can be applied to similar environments, and the obtained results can be
641 representative of expected hydrological dynamics in other glacierized areas in the Andes, with
642 glaciers close to extinction.

Eliminado: Thanks to sub-hourly meteorological and hydrological data, changes in time of streamflow dynamics as a result of changes in atmospheric conditions and/or changes in the glacier area resulting from mass balance loss are also explored.
Eliminado: R
Eliminado: glaciated

644 2. Study site

645 Our study focuses on the Conejeras glacier, a very small ice mass (14 hectares in 2017) that
646 forms part of a larger glacier system called Nevado de Santa Isabel (1.8 km²), one of the six
647 glaciers that still persist in Colombia. It is located in the Cordillera Central (the central range of
648 the three branches of the Andean chain in Colombia) and, together with the glaciers of Nevado
649 del Ruiz and Tolima, comprises the protected area called Parque Nacional Natural de los
650 Nevados (Fig. 1). The summit of the Santa Isabel glacier reaches 5100 m, being the lowest glacier
651 in Colombia. As a result, it is as well the most sensitive to atmospheric warming and why it has
652 been monitored since 2006, part of the world network of glacier monitoring (IDEAM, 2012). The
653 Santa Isabel glacier has been retreating since the 19th century, with an intensification of
654 deglaciation since the middle of the 20th century. As a result, the glacier is now a set of separated
655 ice fragments instead of a continuous ice mass, as it was a decade ago (IDEAM, 2012). One of
656 the fragments, located at the north-east sector of the glacier, is the Conejeras glacier, which is
657 the object of this study, whose elevation ranges between 4700 and 4895 m. In 2006, at the glacier
658 terminus, hydro-meteorological stations were installed in order to measure glacier contribution to
659 runoff, as well as air temperature and precipitation.

Eliminado: "quebradas"
Eliminado: name
Eliminado: (
Eliminado:)
Eliminado: higher
Eliminado: ,

660 The Conejeras water stream is a tributary of one of the 'quebradas' (Spanish for small mountain
661 rivers in South American countries) flowing into the river Rio Claro. Thus, the Conejeras glacier
662 corresponds to the uppermost headwaters of the Rio Claro basin (Fig. 1). The Rio Claro basin
663 comprises an elevation range of 2700 to 4895 m and, from highest to lowest, presents a
664 succession of typical Andean ecosystems: glacial (4700 to 4894), periglacial (4300 – 4700 m),
665 páramo wetland ecosystem (3600 to 4300 m) and high elevation tropical forest *bosque altoandino*
666 (2700 to 3600 m). Mean annual temperature at the glacier base is $1.3 \pm 0.7^\circ\text{C}$, with very little
667 seasonal variation, and precipitation sums reach 1025 ± 50 mm annually, with two contrasted
668 seasons (see Figure 2), resulting from the seasonal migration of the Intertropical Convergence
669 Zone (ITCZ, Poveda et al., 2006). During the dry seasons (December to January and June to
670 August), mean precipitation barely reaches 75 mm per month, whereas during the wet seasons
671 (March to May and September to October), values exceed 150 mm per month.

676 3. Data and Methods

677

678 3.1. Hydrological and meteorological data

695 Meteorological and hydrological data used in the present work have been collected by the Institute
 696 for Hydrological, Meteorological and Environmental Studies of Colombia (IDEAM, *Instituto de*
 697 *Hidrología, Meteorología y Estudios Ambientales*), thanks to the automatic meteorological and
 698 gauge stations network at the Río Claro basin (Figure 1).

Eliminado: has

699 The experimental site of the Río Claro basin has been monitored since 2009, with a network of
 700 meteorological and hydrological stations located at different tributaries of the Río Claro River,
 701 covering an altitudinal gradient of 2700 – 4900 m.asl. As this research is focused in the upper
 702 catchment in which the glacier is located, for the present study, we used data from just the stations
 703 located at the Conejeras glacier snout (Figure 1, bottom map). This includes one stream gauge
 704 (with associated rating curve) measuring, 15-minute resolution water discharge ($m^3 s^{-1}$), one
 705 temperature station measuring hourly temperature ($^{\circ}C$) (both stations located at 4662 m.asl); and
 706 one rain-gauge measuring 10-minute precipitation (mm, the station located at 4413 m. asl). Even
 707 though these data have been available since 2009, the sensors and loggers experienced
 708 technical problems; thus, numerous inhomogeneities, out-of-range values and empty records
 709 were present in the data series. From 2013 the technical problems were solved and the data is
 710 suitable for analysis. The period covered for analysis ranges from June 2013 to December 2017,
 711 a total of 56 months, and data was aggregated hourly, daily and monthly to perform statistical
 712 analyses. However, in order to obtain a wider perspective and to take advantage of the effort
 713 made by the IDEAM glaciologist who conscientiously took mass balance measurements every
 714 month since 2006, also shown are trends and variability in climate (from a nearby meteorological
 715 station of the Colombian national network, (Brisas), that contain data since 1982 and glacier mass
 716 evolution for the longest time period available. The Multivariate ENSO Index, used for
 717 characterizing influence of the ENSO phenomenon on the glacier evolution, has been
 718 downloaded from NOAA <https://www.esrl.noaa.gov/psd/enso/mei/table.html> (December 2017).

Eliminado: consecutively

Eliminado: the

Eliminado: river

Eliminado: For

Eliminado: data was used from

Eliminado: surroundings

Eliminado: ,

Eliminado: including

Eliminado: :

Eliminado: water yield

Eliminado: ,

Eliminado: and

Eliminado: 4413

Eliminado: quality analysis prevented us from using the entire series, as

Eliminado: automatic sensors stabilized

Eliminado: ,

Eliminado: glaciologist of

Eliminado: for

Eliminado: undertaking

Eliminado: s

Eliminado:)

Eliminado: five

Eliminado: 6

Eliminado: about

Eliminado: ,

Eliminado: s

Eliminado: ,

719

720 3.2. Glacier evolution data

721 The evolution of the Conejeras glacier (Fig. 3) has been monitored by the Department of
 722 Ecosystems of IDEAM. Since March 2006, a network of 14 stakes was installed on the Conejeras
 723 glacier to measure ablation and accumulation area. The 6–12 m long stakes are PVC pipes of 2
 724 m in length. These 14 stakes are vertically inserted into the glacier at a depth not less than 5
 725 meters and they are roughly organized in six cross profiles at approximately 4670, 4700, 4750,
 726 4780, 4830 and 4885 m a.s.l. Accumulation and ablation measurements are performed monthly.
 727 Typical measurements of the field surveys include stake readings (monthly), density
 728 measurement in snow and firn pits (once per year), and re-drilling of stakes (if required) to the
 729 former position. The entire methodology can be found in (Mölg et al., 2017; Rabatel et al., 2017).
 730 The mass balance data is calculated using with the classical glaciological method that represents
 731 the water equivalent that a glacier gains or loses in a given time. This data is used to generate
 732 yearly mappings of mass balance and calculate the equilibrium line altitude (ELA), which is the
 733 altitude point where mass balance is equal to zero equivalent meters of water, and separates the
 734 ablation and accumulation area in the glacier (Francou and Pouyaud, 2004).

735 Changes in glacier surface during the study period were computed by means of satellite imagery
 736 (Landsat and Sentinel constellations) for the years 2006, 2010, 2013 and 2017. Cloud-free cover
 737 Landsat TM images were selected for 2006 and 2010 years, and Landsat OLI and Sentinel
 738 images for 2013 and 2017 respectively. TOA (Top Of Atmosphere) Reflectance was obtained
 739 using specific radiometric calibration coefficients for each image and sensor (Chander et al.,
 740 2009; Padró et al., 2017). BOA (Bottom of the Atmosphere) Reflectance was based on the Dark
 741 Object Substraction (DOS) approach (Chavez, 1988). The Normalized Difference Snow Index
 742 (NDSI) was used to discriminate snow and ice-covered areas from snow-free areas. The NDSI is
 743 expressed as the relationship between reflectance in the visible region and reflectance in the
 744 medium-infrared region (the specific bands vary among different sensors; e.g. TM bands 2 and
 745 5). Pixels in the different images were classified as snow- or ice-covered areas when the NDSI
 746 was greater than 0.4 (Dozier, 1989).

Eliminado: Changes in glacier surface have been measured by direct topographic surveys (in the years 2009, 2012, 2014, and 2017) or computed using satellite imagery (Landsat and Sentinel constellations) for the other years within the 2006-2018 period

Eliminado: Free-cloud cover Landsat TM images were selected until 2011 and then Landsat OLI or Sentinel 2 images were selected when data became available. A set of 8 images was processed in order to support direct topographic surveys for ice-covered area measurements.

Eliminado:

Eliminado:

750 3.3. Statistical Analyses

793 A number of indices were extracted from the streamflow, temperature and precipitation hourly
794 series in order to assess changes in time in the hydrological output of the glacier and their relation
795 to climate (Table 1). These daily indices were subject to statistical analyses, including correlation
796 tests, monthly aggregation and assessment of changes over time.

Eliminado: on

797 Since one of the main objectives of the paper is to characterize daily dynamics of streamflow and
798 changes in time, a principal component analysis (PCA) was conducted in order to extract the
799 main patterns of daily streamflow cycles. The data matrix for the PCA was then composed by
800 streamflow hourly values in 1614 columns as variables (number of days) and 24 rows as cases
801 (hours in a day). As the PCA does not allow the number of variables to exceed the number of
802 cases, PCAs were performed on 25 bootstrapped random samples of days (n=23, with
803 replacement); Results showed that three principal components were stable throughout the
804 samples (see Table 3 in Results sections). After the main PCs were extracted, calculation of
805 correlation between each day of the time series and the selected PCs was determined. The PC
806 that best correlated with the correspondent day was assigned to every day, obtaining a time-
807 series of the three PCs. This allowed assessment of changes in time of the main patterns of daily
808 streamflow cycles observed.

Eliminado: the

Eliminado: it was ensured that results with

Eliminado: table

Eliminado: results

809

810 4. Results

811 4.1. Climatology and glacier's evolution

812 The long-term climatic evolution of the study area is depicted in Figure 2. The temperature and
813 precipitation series (Fig. 2 a, c and d) correspond to the Brisas meteorological station, which is
814 located 25 km from the glacier, at 2721 m elevation. It therefore does not accurately represent
815 the climate conditions at the glacier. It is, however, the closest meteorological station with
816 available meteorological data to study long-term climate. The temperature record measured at
817 the glacier snout is included (blue line). It can be observed that despite the different range of
818 values (temperatures at the glacier are 3.2 °C lower than at Brisas), there is a match in variability
819 for the common period.

Eliminado: , especially due to the lapse rate of temperature with elevation (which makes temperatures at the glacier 3.2 °C lower: annual mean temperature of 1.4 °C at the glacier's base, compared to 4.6 °C at Brisas)

Eliminado:

820 Long-term evolution of temperature does not show any significant trend or pattern from 1982 to
821 2015; however, a spectral analysis shows that the frequency with higher spectral density
822 corresponds with a seasonality of 48 months, indicating a recurrent cycle every four years. By
823 comparing Fig. 2a with Fig. 2b, there is a close match between temperature and evolution of the
824 Multivariate ENSO Index (R = 0.49), which also shows, as well, a high value of power spectra in
825 the 48-month frequency cycle. Notwithstanding other factors whose analysis is far beyond the
826 scope of this paper, it is evident that the evolution of temperature in the study area is highly driven
827 by the ENSO phenomenon. Regarding precipitation (Fig. 2c), no long-term trend is observed, and
828 the most evident pattern is the bi-modal seasonal regime which is confirmed by the frequency
829 analysis showing the highest power spectra in the 6-month cycle.

Eliminado: 4

Eliminado: we observe

Eliminado: ,

Eliminado: (Fig. 2d) with two "humid" seasons and two "dry" seasons every year, typical of the whole country that is driven by a seasonal shifting of the Intertropical Convergence Zone (ITZC).

Eliminado: .

830 The evolution of the glacier since 2006 is shown in Figure 3. Almost every month since
831 measurements began in 2006, the glacier has lost mass (113 months), and very few months (20)
832 recorded a positive mass balance. The global balance in this period is a loss of 34.4 meters of
833 water equivalent. For the sake of visual comparison, we have included the time series of MEI and
834 a close correspondence between the variables is observed (Figure 3 a). During the warm phases
835 of ENSO (Niño events, values of MEI above 0.5), the glacier loses up to 600 mm w.e. per month,
836 as in the Niño event of 2009-2010, when the glacier lost a total of 7000 mm w.e. One could
837 surmise that during La Niña (cold phases of ENSO, MEI values < -0.5) the glacier could
838 recuperate mass. In fact, when MEI values are negative, the glacier experiences much less
839 decrease; however, even during the strongest La Niña events, the balance is negative, with just
840 a few months having a positive balance (e.g. in the 2010-2011 La Niña, the glacier lost 1000 mm
841 w.e.) This occurs because even during La Niña mean temperatures at the glacier are above zero
842 (0.8 ± 0.3 °C). The aforementioned agreement between ENSO and mass balance appears to
843 break from 2012 onwards. There were two events of large mass balance loss around 2013-2014
844 that do not match with El Niño events. A local factor that can affect the glacier's mass balance
845 independent of climatology is reduced albedo of the surface caused by the quantity of deposited
846 ash that comes from the nearby Santa Isabel volcano. This variable has not been considered in

Eliminado: saw

Eliminado: was recorded

Eliminado: the mass balance evolution in the temperature and MEI plot's (Fig. 2a and 2b) are included,

Eliminado: the

Eliminado: , but they do coincide with temperature peaks. This might be due to other factors that affect temperature and mass balance such as increasing radiation due to less cloudiness. But some peaks of mass balance loss that do not correspond to temperature peaks were also observed. .

Eliminado: ly

Eliminado: the

Eliminado: on the ice surface

Eliminado:

882 the present study but there are two pictures of the glacier for visual evidence (Figure 3.d). This
 883 fact, together with prevalence of above-zero degrees (see Figure 2, top plot) at the elevation in
 884 which the glacier is located has induced the large glacier recession observed between 2006 and
 885 2017 (Figure 3.c). During this period, there has been a 37% reduction, from 22.45 ha in 2006 to
 886 12 ha in 2017. However, this reduction has been far from linear. As shown in Figure 3.b, mass
 887 balance losses during the first years of the monitoring period were in general less pronounced
 888 than in the latest years. In 2012, the ice mass retreated up the 4700 m elevation curve, and from
 889 then on the years with larger mass loss were 2015, 2016 and 2014.

890

891 **4.2. Hydrological dynamics**

892 The water discharge of the Conejeras glacier is measured at a gauging station located 300 m
 893 from the glacier snout (when the station was installed in 2009, it was only 10 meters away from
 894 the glacier snout). The water volume measured at this station is a combination of water from
 895 glacier melt and water from precipitation into the watershed area, although the former exerts a
 896 larger control in water discharge variability. Table 2 shows the correlation between hydrological
 897 and temperature indices for samples of days with precipitation, independent of the amount of
 898 fallen precipitation (left), and for samples of days without precipitation (right). On days without
 899 precipitation, most hydrological indices show significant correlation with temperature, except for
 900 the baseflow and hQmax. The highest correlation values are found between Qmax, Qrange,
 901 Qslope and totalQ, with Tmax and Tmean (correlation values in the range of 0.5 – 0.65), indicating
 902 that the higher the temperatures, the more prominent the melting pulse of runoff. Tmin shows
 903 smaller and less significant correlation values. The hpulse also shows high correlation with
 904 temperature, but in this case in a negative fashion, indicating a later occurrence of the daily
 905 melting pulse when minimum temperatures and maximum temperatures are lower. On days with
 906 precipitation, correlation values are generally smaller but, in some cases, they are still significant
 907 as for Qmax, Qrange and Qslope.

908 A Principal Component Analysis (PCA) performed on hourly streamflow data (in a recursive
 909 fashion, see Section 3.3 for explanation of the method) allowed procurement of the main patterns
 910 of daily flow, as well as changes in time during the study period. Three principal components were
 911 obtained, whose values of explained variance were stable throughout the 25 bootstrapped
 912 samples (Table 3). The first PC explained an average of $48 \pm 6\%$ of the variance throughout the
 913 25 samples, and the second PC an average of $35 \pm 5.7\%$. Together they account for 83% of
 914 variance and they both showed a neat pattern of daily streamflows (Fig. 4a). The main difference
 915 between PC1 and PC2 is the time of the day when peak flows are reached and hence the time
 916 range when most daily flows occur. Thus, PC1 corresponds to days with an earlier melt pulse
 917 (towards 10h) and earlier peak flows (towards 14h), compared to PC2, with days of melt pulse at
 918 13h and peak flows at 18h. The remaining PC explains a residual percentage of the variance and,
 919 unlike PC1 and PC2, does not show a stable streamflow pattern across the samples. However, it
 920 was decided to keep it, as it can help explain some peculiarities in the results. In Figure 4b the
 921 evolution of the frequency (days per month) of days corresponding to each PC is shown. Although
 922 there is some degree of variability, the frequency of days with PC1 streamflow pattern increases
 923 over time, and dominates over the frequency of PC2 and PC3 days. This is especially significant
 924 between 2015 and 2016, coinciding with an El Niño event. However, by mid-2016 the frequency
 925 of PC1 days drops considerably and the frequency of PC2 days increases in the same ratio. Thus,
 926 from mid-2016 to the end of the study period, they both maintain similar levels of frequency.

927 In order to understand the underlying factors of each PC, the frequency distribution of the climatic
 928 and hydrological indices for the days corresponding to each PC was computed, in the form of
 929 boxplots (Figure 5). From a hydrological point of view, PC1 better corresponds to days with higher
 930 total runoff and maximum runoff and with a more pronounced slope in both the rising and
 931 decreasing limbs of the peak flow volume than PC2 and PC3. The variability (expressed by the
 932 amplitude of boxes in the boxplots) of such hydrological indicators is, as well, higher amongst
 933 days of PC1, compared to days of PC2 and PC3. Base runoff is higher in PC1 but not significantly.
 934 The contrasted weight of climate may explain such hydrological differences between PCs: days
 935 of PC1 present significantly higher mean temperature (median = 1.7°C) and maximum
 936 temperature (median = 3.8°C) than days of PC2 (0.9°C and 2.4°C respectively) and PC3 (0.5°C

- Eliminado:
- Eliminado: could be the subject of further research, since a meteorological station that includes albedo measurements has been recently installed on the top of the glacier.
- Eliminado:
- Eliminado: In terms of the glacier's area,
- Eliminado:
- Eliminado: i
- Eliminado: 2014
- Eliminado: receded
- Eliminado: Between 2006 and 2014 the area reduction was 9%, occurring in a gradual fashion, and one year (2012) saw a slight recuperation in glaciated area. Reduction during these years was limited to a slight receding of the glacier's snout, and no apparent changes were observed in the upper parts of the ice body (see map in Figure 3). From 2014 to 2017, in contrast, there has been a sharp decrease in the glacier's area, being especially drastic between 2014 and 2015, with substantial receding of the snout combined with a retreat of ice in the upper parts of the glacier.
- Eliminado: water yield
- Eliminado: gauge
- Eliminado: water yield
- Eliminado: Qbase
- Eliminado: d
- Eliminado: section
- Eliminado: the

- Eliminado:
- Eliminado: am

- Eliminado: ,

- Eliminado: s

971 and 1.6°C respectively). In contrast, precipitation is notably higher (and shows greater variability)
972 in days grouped within PC3 (median = 1.9 mm day⁻¹) and PC2 (2.2 mm day⁻¹) compared to days
973 of PC1 (0.3 mm day⁻¹). To summarize, PC1 corresponds to a daily regimen of pure glacier melting,
974 whereas PC2 and PC3 correspond to days with a lower glacier melting pulse with more (PC3) or
975 less (PC2) influence of precipitation.

Eliminado: 2.2
Eliminado: 1.9
Eliminado: 2.2

976
977
978

979 In Figure 4 a notable change occurs in the frequency of the two main patterns of hourly
980 streamflow, PC1 and PC2, by mid-2016. Further details regarding changes in the hydrological
981 yield of the glacier are shown in Figure 6, which presents the evolution of the main hydrological
982 indices computed, along with temperature, precipitation and glacier mass balance, during the
983 study period and averaged monthly. Total and maximum daily streamflow (*totalQ* and *Qmax*)
984 depict an increase up to mid-2016, where they begin to decrease. During the last 18 months, they
985 remain at low levels compared to previous months. This turning point seems to coincide in time
986 with the 2015-16 El Niño event, with higher-than-average temperatures and low levels of
987 precipitation that led to an increasing mass balance loss and therefore increased flows. It is
988 remarkable that streamflow increases and decreases in direct proportion to mass balance
989 change, indicating the strong dependence of runoff to glacier melt. Similar evolution is observed
990 in the difference between base flows and maximum flows (*Qrange*), as well as the slope of the
991 rising limb of diurnal flows (*Qslope*) which are indicators of diurnal variability: they increase up to
992 2016 and decrease afterwards, which coincides with the change in the frequency of daily
993 streamflow patterns in Fig. 5. The mean hour of the day at which maximum flows are reached
994 (*hQmax*) shows a steady evolution until mid-2016 when it begins to rise. This seems surprising
995 when comparing it to the evolution of *hTmax* (i.e. the hour of the day when maximum temperature
996 is reached) which does not show any particular temporal pattern. Regarding the monthly
997 variability of flows (third panel on the right, Fig. 7) the same turning point is observed, with a clear
998 decrease in the coefficient of variation until 2016 and an increase afterwards. It is clear that a
999 hydrological change has occurred at the outlet of the glacier, but when we look at the two most
1000 plausible drivers of change (temperature and precipitation, bottom plots Fig. 7) do not seem to be
1001 responsible for it. They both are affected by the El Niño event, when temperatures increased and
1002 precipitation decreased; however, they do not show an increasing-decreasing temporal pattern
1003 before and after such an event. This leads to the hypothesis that the hydrological change
1004 observed at these last stages of the glacier's life is independent of climate.

Eliminado: ,
Eliminado: ing trend

1005
1006
1007

4.3. Changes in the runoff-climate relationship

1008 In this section, the runoff is isolated from temperature and precipitation in order to determine if
1009 observed hydrological dynamics are driven by climate or are related to shrinkage of the glacier.
1010 Figure 7 shows the mean monthly runoff for days with temperatures lower and higher than 2°C,
1011 i.e. water discharge series independent of temperature. Precipitation has also been added to the
1012 plot. It was noted that water discharge for days warmer than 2°C is significantly higher than water
1013 discharge on days cooler than 2°C. The characteristic evolution of runoff, with increasing amounts
1014 during most of the study period up to mid-2016 and decreasing runoff from that point onwards
1015 was also observed. The same evolution occurs for both days below and days above 2°C, and it
1016 occurs for very similar amounts of precipitation. This indicates that flows from the melting glacier
1017 are becoming less dependent on temperature, or climate in general, and more dependent on the
1018 size of the glacier. The boxplots of Figure 8 (bottom) confirm this observation by showing water
1019 volumes significantly higher before than after the breaking point, but also because the differences
1020 between water discharge at < 2°C and water discharge at > 2°C are also smaller (and not
1021 significant) after the breaking point, indicating the decreasing importance of temperature in the
1022 process of runoff production in the shrinking glacier.

Eliminado: trend
Eliminado: we observe
Eliminado:) they
Eliminado: contrasted sign on trends
Eliminado: a
Eliminado: climatology
Eliminado: The most plausible causes for this will be explained
Eliminado: from temperature and precipitation
Eliminado: the
Eliminado: by
Eliminado: water yield
Eliminado:
Eliminado: water yield
Eliminado: water yield
Eliminado: It seems evident, therefore,
Eliminado: Following the hypothesis of Section 1.2, regarding hydrological changes of shrinking glaciers from 2013 to mid-2016, the runoff increases because glacier mass becomes more sensitive to energy exchange as it gets smaller. From mid-2016 onwards, runoff decreases because the water reservoir present in the ice has reached a threshold where its contribution cannot be offset by incoming precipitation or potentially higher temperatures.
Eliminado: water yield
Eliminado: water yield

1055 Finally, Figure 8 shows correlations between temperature/precipitation and monthly flows for
1056 different time periods. In Figure 8a two years are compared that can be considered analogues in
1057 terms of total flow (similar amounts of monthly flow, see Figure 6), but one year (2013-14) belongs
1058 to the period of increasing flows, before the 2016 breakpoint, and the other year (2017) belongs
1059 to the period of decreasing flows after the breakpoint. Correlation between temperature and flow
1060 is much higher ($R = 0.65$) for 2013-14 than for 2017 ($R = 0.35$), which would corroborate the
1061 previous observation. However, precipitation also shows higher correlation with flow for 2013-14
1062 ($R = 0.67$) than for 2017 ($R = 0.42$), which would contradict the hypothesis. One year, however,
1063 may not be representative of general trends, and so the same analysis is repeated, not for
1064 individual years but for the whole periods pre- and post-2016 breakpoint (Fig. 8b). The pattern
1065 seems more clear and corroborates the aforementioned hypothesis: correlation between
1066 temperature and flow is significant for the pre-2016 period ($R = 0.55$) but is non-existent for the
1067 post-2016 period ($R = -0.1$). Correlation between precipitation and flow is insignificant ($R = -0.23$)
1068 for the pre-2016 period, and it is positive and significant for the post-2016 period ($R = 0.32$). These
1069 previous observations lead to reasoning that during the years of hydrological monitoring (2013-
1070 2017), the observed hydrological dynamic, with a marked breakpoint in 2016, is a result of the
1071 vanishing glacier process and not a response to climate variability.

Eliminado: ... [2]

Eliminado: due to deglaciation

Eliminado:

1072

1073 5. Discussion and conclusions

1074 The present paper shows a comprehensive analysis of the dynamics of an Andean glacier that is
1075 close to extinction, with special focus on its hydrological yield. This research has benefited from
1076 a hydro-climatic monitoring network located in the surroundings of the glacier terminus that has
1077 been fully operative since 2013 and from monthly and annual estimations of mass balance and
1078 glacier extent respectively, derived from ice depth measurements and topographical surveys
1079 since 2006. Everything has been managed by the Institute of Hydrology Meteorology and
1080 Environmental Studies (IDEAM) of Colombia. The Conejeras glacier is currently an isolated small
1081 glacier that used to be part of a larger ice body called Nevado de Santa Isabel. Since
1082 measurements have been available, the glacier has constantly lost mass and, consequently, a
1083 reduction in its area is evident. The extinction of Colombian glaciers has been documented since
1084 1850, with an average loss of 90% of their area, considering current values (IDEAM, 2012). This
1085 reduction, of about 3% per year, has been much larger during the last three decades (57%)
1086 compared to previous decades (23%), which is directly related to the general increase in
1087 temperatures in the region and to re-activation of volcanic activity (IDEAM, 2012; Rabatel et al.,
1088 2017). Since direct measurements began in 2006, the studied glacier has constantly lost area;
1089 however, until 2014 the area loss was gradual and restricted to the glacier front, from 2014 the
1090 sharp retreat also involved higher parts of the glacier. The main reason for this strong shrinkage
1091 is the existence of above-zero temperatures during most of the year and less precipitation fallen
1092 as snow. This involves a constant migration of the equilibrium line to higher positions, and a
1093 decreasing albedo of the ice surface that involves greater energy absorption, the latter
1094 accelerated by intense activity of Nevado de el Ruiz in the last years. In terms of mass balance,
1095 very few months exhibit a gain of ice during the studied period and these tend to coincide with la
1096 Niña events (negative MEI episodes). These episodes cannot compensate for the great losses
1097 that occurred during the majority of months, which are especially large during El Niño events
1098 (positive MEI episodes), when above-normal temperatures are recorded. The ENSO
1099 phenomenon exerts great influence on the evolution of the glacier, similar to that reported for
1100 most inner tropical glaciers (Francou et al., 2004; Rabatel et al., 2013; Vuille et al., 2008);
1101 however, some episodes of great mass balance loss, such as that of 2014, cannot be explained
1102 by the ENSO. Observations of glacier surface during field surveys showed that during some
1103 periods of mass loss, surface ice retreat left ancient layers of volcanic ash exposed. The reduced
1104 energy reflectance caused by such ash layers might have triggered positive feedback that led to
1105 increasing melting and large ice retreat.

Eliminado: We have

Eliminado: our

Eliminado: , and

Eliminado: ,

Eliminado: occurring

Eliminado: then

Eliminado: (which coincides with the sharp decrease in glacier extent)

1106 Glacier retreat is a worldwide phenomenon currently linked to global warming (IPCC, 2013).
1107 Amongst the environmental issues related to glacier retreat, the issue concerning water resources
1108 has produced a vast amount of research. This is because glaciers constitute water reservoirs in
1109 the form of accumulated ice over thousands of years, and they provide water supply to
1110 downstream areas for the benefit of life, ecosystems and human societies. The rapid decrease in

1123 glacier extent during the last decades involves a change in water availability in glacier-dominated
 1124 regions, and, thus, changes in water policies and water management are advisable (Huss, 2011;
 1125 Kundzewicz et al., 2008). In the short term, glacier retreat involves increasing runoff in
 1126 downstream areas but, after reaching a peak, runoff will eventually decrease until the contribution
 1127 of the glacier melt is zero when the glacier completely disappears. From a global perspective,
 1128 such a tipping point is referred to as *peak water* and has given rise to concern from the scientific
 1129 community (Gleick and Palaniappan, 2010; Huss and Hock, 2018; Kundzewicz et al., 2008; Mark
 1130 et al., 2017; Sorg et al., 2014). Research regarding the occurrence of such a runoff peak related
 1131 to glacier retreat demonstrates that it will not occur concurrently worldwide. In some mountain
 1132 areas it has already occurred, i.e. the Peruvian Andes (Baraer et al., 2012), the Western U.S
 1133 mountains (Frans et al., 2016), or Central Asia (Sorg et al., 2012). At the majority of studied glacier
 1134 basins it is expected to occur in the course of the present century (Immerzeel et al., 2013; Ragetti
 1135 et al., 2016; Sorg et al., 2014; Soruco et al., 2015). In recent global-scale research, Huss and
 1136 Hock (2018) state that in nearly half of the 56 large-scale glacierized drainage basins studied, the
 1137 peak water has already occurred. In the other half, it will occur in the next decades, depending
 1138 on extension of the ice cover fraction.

Eliminado: ,

1139 It was not the aim of this study to allocate such a tipping point in our studied glacier; however,
 1140 observations on the characteristics of streamflow along the studied period suggest that it may
 1141 have occurred during our study period. Our observations corroborate glacier melt being the main
 1142 contributor to runoff in the catchment. However, even when correlations between runoff and
 1143 temperature are mostly significant, the values are not as high as could be expected for a
 1144 glacierized catchment. This is due to decreasing dependence of runoff on temperature, and
 1145 therefore to glacier melt, as at a specific point during the study period. We observed a changing
 1146 dynamic in most hydrological indicators, with a turning point in mid-2016, whereas climate
 1147 variables, i.e. temperature and precipitation, do not show such evident variation (besides the
 1148 exceptional conditions during an El Niño event). Both the PCA analysis and the monthly
 1149 aggregation of hydrological indices point to a less glacier-induced hydrological yield once the
 1150 runoff peak of 2016 was reached. According to literature (see Section 1.2.) this change from
 1151 increasing to decreasing runoff and to lesser importance of glacier contribution to total water
 1152 discharge must be expected in glacierized catchments with glaciers close to extinction. The short
 1153 length of our hydrological series (five years) does not allow long-term analysis to determine water
 1154 discharge in years of less glacier loss (i.e. from 2006 to 2012, see Fig. 3), which could verify or
 1155 refute such a hypothesis. However, when we isolated total runoff from climate variables before
 1156 and after the 2016 breakpoint (Figures 8 and 9), we observed that the increase and later decrease
 1157 of flows was mostly independent of temperature and precipitation, which would involve a glacier-
 1158 driven hydrological change. Summarizing, streamflow measured at the glacier's snout showed the
 1159 following characteristics: increasing trend in flow volume until mid-2016 and decreasing trend
 1160 thereafter; increasing diurnal variability (given by the range between high flows and low flows and
 1161 by the slope of the rising flow limb) up to mid-2016 and decreasing thereafter; decreasing and
 1162 increasing monthly variability (given by the coefficient of variation of flows within a given month)
 1163 before and after such date; and high dependence of flow on temperatures before 2016 and low
 1164 or null dependence after 2016, with increasing dependence on precipitation. As well, this is
 1165 supported by an evident change in the type of hydrograph, from a prevalence of days with melt-
 1166 driven hydrographs (low baseflows, sharp melting pulse and great difference between high flows
 1167 and low flows) before 2016, to an increase in the occurrence of days with less influence of melt
 1168 and more influence by precipitation. All these characteristics support the idea of a hydrological
 1169 change driven by the glacier recession in the catchment, as summarized by Hock et al. (2005,
 1170 see Section 1.2). This observation cannot be taken conclusively, because the time period of
 1171 hydrological observation is not long enough to deduce long-term trends and to explore
 1172 hydrological dynamics before the great decline in glacier extent in 2014. However, given the
 1173 current reduced size of the glacier (14 hectares, which represents 35% of the catchment that
 1174 drains into the gauge station), it is likely that water discharge will continue to decrease in the
 1175 upcoming years, until glacier contribution ends and runoff depends only on the precipitation that
 1176 falls within the catchment. Like this glacier, other small glaciers in Colombia are expected to
 1177 disappear in the coming decades (Rabatel et al., 2017); thus, a similar hydrological response can
 1178 be expected.

Eliminado: This is due to a decreasing dependence of runoff on temperature, and therefore to glacier melt, as at a certain point during the study period

Eliminado: water yield

Eliminado: water yield

Eliminado: deglaciation

Eliminado: .

Eliminado: :

Eliminado: deglaciation

Eliminado: of

Eliminado: section

Eliminado: Data on glacierized area fraction supports this idea: the area covered by ice changed from 56-51% of the catchment in the 2006-2014 timespan to 39-35% of the catchment in 2015-2017; therefore, the contribution of ice melt to total flow compared to that of precipitation must necessarily decrease.

Eliminado: water yield

1198 Unlike glaciers in the western semi-arid slopes of the Andes (i.e. Peru, Bolivia), Colombian
1199 glaciers do not constitute the main source of freshwater for downstream populations (IDEAM,
1200 2012). The succession of humid periods provides enough water in mountain areas, most of which
1201 is stored in the deep soils of *Páramos*. These wetland ecosystems are mainly fed by rainfall (the
1202 contribution of glacier melt is mostly unknown, IDEAM, 2012) and act as water buffers, ensuring
1203 water availability during not-so-humid periods. Therefore, the role of glaciers in Colombia
1204 regarding water resources, including the studied ice body, is more marginal, and the occurrence
1205 of the peak water from glacier melt is not a current concern, as it is in Peru or Bolivia (Francou et
1206 al., 2014). Yet this does not diminish the relevance of the results of this work, because they may
1207 be taken as an example of what can happen to the hydrology of glacierized basins in the tropics
1208 whose glaciers are in the process of disappearing. The studied glacier has a very small size
1209 compared to other ice bodies in the region. This makes it respond rapidly to variations in climate,
1210 as well as involving a rapid hydrological response of the catchment to the loss of ice, as was
1211 observed in this work. The increasing/decreasing flow dynamic observed as the glacier retreated
1212 occurred in roughly five years, and this is most likely related to the reduced size of the studied
1213 glacier. Most studies on the hydrological response to glacier retreat consider large river basins
1214 with large glacier coverage, usually by modeling approaches (i.e. Huss and Hock, 2018;
1215 Immerzeel et al., 2013; Ragettli et al., 2016; Sorg et al., 2014, 2012; Stahl et al., 2008), and the
1216 response times reported on either increasing flow at the initial stages or decreasing flow at the
1217 final stages are always on the scale of decades.

Eliminado: our

Eliminado: to our

Eliminado: our observations

Eliminado: Our

Eliminado: our

Eliminado:

1218 The added value of studying the hydrology related to this small-sized and near-extinct glacier is
1219 that the changes observed in the hydrology of the catchment could be directly attributed to the
1220 dynamics of the glacier and the climate that occurs at the same time-scale; contrary to catchments
1221 containing large glaciers that respond with a larger temporal inertia to environmental changes.
1222 Hydrological analyses were restricted to the upper catchment because the streamflows measured
1223 at the snout of the glacier are not influenced by the signals of other environmental processes that
1224 may occur downstream (e.g., forest clearing or increasing grazing). The methodological
1225 approach, including the PCA and the hydrological indices computed over sub-daily resolution
1226 data demonstrated itself as viable for detecting changes on the diurnal cycle of the glacier, and
1227 can be applied to other small glaciers of the tropical Andes that respond rapidly (at sub-annual
1228 scales) to environmental forcing. The necessity for in situ observations on a fine scale in order to
1229 improve accuracy on future estimations of water availability related to glacier retreat is
1230 emphasized.

Eliminado: Our work demonstrates

Eliminado: t

Eliminado: r

1231

1232 Acknowledgments

1233 This work has been possible thanks to the monitoring network installed by the Department of
1234 Ecosystems of the Colombian Institute for Hydrology, Meteorology and Environmental Studies
1235 (*Instituto de Hidrología, Meteorología y Estudios Ambientales*, IDEAM) and to the monthly field
1236 surveys on the Conejeras glacier and Río Claro river basin done by employed staff. Our sincere
1237 gratitude to them, with special thanks to Yina Paola Nocua. The following projects gave economic
1238 support to this paper: "*Estudio hidrológico de la montaña altoandina (Colombia) y su respuesta a*
1239 *procesos de cambio global*" financed by Banco Santander, through the program of exchange
1240 scholarships for young researchers in Ibero-America "*Becas Iberoamérica Jóvenes Profesores e*
1241 *Investigadores*" (2015); and CGL2017- 82216-R (HIDROIBERNIEVE) funded by the Spanish
1242 Ministry of Economy and Competitiveness. We are thankful to the anonymous referees for their
1243 valuable comments and suggestions that helped improve the final version of this manuscript.

Eliminado:

1244

1245

1246

1247

1248

1249 6. References

- 1260 Baraer, M., Mark, B. G., McKenzie, J. M., Condom, T., Bury, J., Huh, K.-I., Portocarrero, C.,
 1261 Gómez, J. and Rathay, S.: Glacier recession and water resources in Peru's Cordillera Blanca, J.
 1262 Glaciol., 58(207), 134–150, doi:10.3189/2012JoG11J186, 2012.
- 1263 Beniston, M.: Impacts of climatic change on water and associated economic activities in the
 1264 Swiss Alps, J. Hydrol., 412–413, 291–296, doi:10.1016/J.JHYDROL.2010.06.046, 2012.
- 1265 Bradley, R. S., Vuille, M., Diaz, H. F. and Vergara, W.: Climate change. Threats to water
 1266 supplies in the tropical Andes., Science, 312(5781), 1755–6, doi:10.1126/science.1128087,
 1267 2006.
- 1268 Cayan, D. R., Dettinger, M. D., Kammerdiener, S. A., Caprio, J. M., Peterson, D. H., Cayan, D.
 1269 R., Dettinger, M. D., Kammerdiener, S. A., Caprio, J. M. and Peterson, D. H.: Changes in the
 1270 Onset of Spring in the Western United States, Bull. Am. Meteorol. Soc., 82(3), 399–415,
 1271 doi:10.1175/1520-0477(2001)082<0399:CITOOS>2.3.CO;2, 2001.
- 1272 Ceballos, J. L., Euscátegui, C., Ramírez, J., Cañon, M., Huggel, C., Haeblerli, W. and Machguth,
 1273 H.: Fast shrinkage of tropical glaciers in Colombia, Ann. Glaciol., 43, 194–201,
 1274 doi:10.3189/172756406781812429, 2006.
- 1275 Chander, G., Markham, B. L. and Helder, D. L.: Summary of current radiometric calibration
 1276 coefficients for Landsat MSS, TM, ETM+, and EO-1 ALI sensors, Remote Sens. Environ.,
 1277 113(5), 893–903, doi:10.1016/J.RSE.2009.01.007, 2009.
- 1278 Chavez, P. S.: An improved dark-object subtraction technique for atmospheric scattering
 1279 correction of multispectral data, Remote Sens. Environ., 24(3), 459–479, doi:10.1016/0034-
 1280 4257(88)90019-3, 1988.
- 1281 Chevallier, P., Pouyaud, B., Suarez, W. and Condom, T.: Climate change threats to
 1282 environment in the tropical Andes: glaciers and water resources, Reg. Environ. Chang., 11(S1),
 1283 179–187, doi:10.1007/s10113-010-0177-6, 2011.
- 1284 Dozier, J.: Spectral signature of alpine snow cover from the landsat thematic mapper, Remote
 1285 Sens. Environ., 28, 9–22, doi:10.1016/0034-4257(89)90101-6, 1989.
- 1286 Favier, V., Wagnon, P. and Ribstein, P.: Glaciers of the outer and inner tropics: A different
 1287 behaviour but a common response to climatic forcing, Geophys. Res. Lett., 31(16), L16403,
 1288 doi:10.1029/2004GL020654, 2004.
- 1289 Francou, B. and Pouyaud, B.: Metodos de observacion de glaciares en los Andes tropicales :
 1290 mediciones de terreno y procesamiento de datos : version-1 : 2004, [online] Available from:
 1291 https://www.researchgate.net/profile/Bernard_Pouyaud/publication/282171220_Metodos_de_ob_servacion_de_glaciares_en_los_Andes_tropicales_mediciones_de_terreno_y_procesamiento_de_datos_version-1_2004/links/561ba9b808ae78721fa0f8ad.pdf (Accessed 12 March 2018),
 1292 2004.
- 1295 Francou, B., Vuille, M., Wagnon, P., Mendoza, J. and Sicart, J.: Tropical climate change
 1296 recorded by a glacier in the central Andes during the last decades of the twentieth century:
 1297 Chacaltaya, Bolivia, 16°S, J. Geophys. Res., 108(D5), 4154, doi:10.1029/2002JD002959, 2003.
- 1298 Francou, B., Vuille, M., Favier, V. and Cáceres, B.: New evidence for an ENSO impact on low-
 1299 latitude glaciers: Antizana 15, Andes of Ecuador, 0°28'S, J. Geophys. Res., 109(D18), D18106,
 1300 doi:10.1029/2003JD004484, 2004.
- 1301 Francou, B., Rabatel, A., Soruco, A., Sicart, J. E., Silvestre, E. E., Ginot, P., Cáceres, B.,
 1302 Condom, T., Villacís, M., Ceballos, J. L., Lehmann, B., Anthelme, F., Dangles, O., Gomez, J.,
 1303 Favier, V., Maisincho, L., Jomelli, V., Vuille, M., Wagnon, P., Lejeune, Y., Ramallo, C. and
 1304 Mendoza, J.: Glaciares de los Andes tropicales: víctimas del cambio climático, Comunidad
 1305 Andina, PRAA, IRD. [online] Available from:
 1306 <http://bibliotecavirtual.minam.gob.pe/biam/handle/minam/1686> (Accessed 22 February 2018),
 1307 2014.
- 1308 Frans, C., Istanbuluoglu, E., Lettenmaier, D. P., Clarke, G., Bohn, T. J. and Stumbaugh, M.:
 1309 Implications of decadal to century scale glacio-hydrological change for water resources of the
 1310 Hood River basin, OR, USA, Hydrol. Process., 30(23), 4314–4329, doi:10.1002/hyp.10872,

Con formato: Inglés (americano)

Con formato: Inglés (americano)

Con formato: Inglés (americano)

- 1311 2016.
- 1312 Gleick, P. H. and Palaniappan, M.: Peak water limits to freshwater withdrawal and use, *Proc.*
 1313 *Natl. Acad. Sci.*, 107(25), 11155–11162, doi:10.1073/pnas.1004812107, 2010.
- 1314 Granados, H. D., Miranda, P. J., Núñez, G. C., Alzate, B. P., Mothes, P., Roa, H. M., Cáceres
 1315 Correa, B. E. and Ramos, J. C.: Hazards at Ice-Clad Volcanoes: Phenomena, Processes,
 1316 and Examples From Mexico, Colombia, Ecuador, and Chile, *Snow Ice-Related Hazards, Risks*
 1317 *Disasters*, 607–646, doi:10.1016/B978-0-12-394849-6.00017-2, 2015.
- 1318 Hock, R., Jansson, P. and Braun, L. N.: Modelling the Response of Mountain Glacier Discharge
 1319 to Climate Warming, in *Global Change and Mountain Regions (A State of Knowledge*
 1320 *Overview)*, pp. 243–252, Springer, Dordrecht., 2005.
- 1321 Huggel, C., Ceballos, J. L., Pulgarín, B., Ramírez, J. and Thouret, J.-C.: Review and
 1322 reassessment of hazards owing to volcano–glacier interactions in Colombia, *Ann. Glaciol.*, 45,
 1323 128–136, doi:10.3189/172756407782282408, 2007.
- 1324 Huss, M.: Present and future contribution of glacier storage change to runoff from macroscale
 1325 drainage basins in Europe, *Water Resour. Res.*, 47(7), doi:10.1029/2010WR010299, 2011.
- 1326 Huss, M. and Hock, R.: Global-scale hydrological response to future glacier mass loss, *Nat.*
 1327 *Clim. Chang.*, 8(2), 135–140, doi:10.1038/s41558-017-0049-x, 2018.
- 1328 IDEAM: Glaciares de Colombia, más que montañas con hielo, edited by Comité de
 1329 Comunicaciones y Publicaciones del IDEAM, Bogotá., 2012.
- 1330 Immerzeel, W. W., Pellicciotti, F. and Bierkens, M. F. P.: Rising river flows throughout the
 1331 twenty-first century in two Himalayan glacierized watersheds, *Nat. Geosci.*, 6(9), 742–745,
 1332 doi:10.1038/ngeo1896, 2013.
- 1333 IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to
 1334 the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, edited by T. F.
 1335 Stocker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex,
 1336 and P. M. Midgley, Cambridge University Press, Cambridge, United Kingdom and New York,
 1337 NY, USA., 2013.
- 1338 Kaser, G. and Omasston, H.: Tropical glaciers, Cambridge University Press. [online] Available
 1339 from: https://books.google.es/books?hl=es&lr=&id=ZEB-I3twN_gC&oi=fnd&pg=PR11&dq=tropical+glaciers&ots=WLwn1fdjig&sig=897EG6q4Pyc113vo9Qb2bnyUo7g#v=onepage&q=tropical+glaciers&f=false (Accessed 21 November 2017), 2002.
- 1340
 1341
- 1342 Kaser, G., Grosshauser, M. and Marzeion, B.: Contribution potential of glaciers to water
 1343 availability in different climate regimes., *Proc. Natl. Acad. Sci. U. S. A.*, 107(47), 20223–7,
 1344 doi:10.1073/pnas.1008162107, 2010.
- 1345 Klein, A. G., Morris, J. N. and Poole, A. J.: Retreat of Tropical Glaciers in Colombia and
 1346 Venezuela from 1984 to 2004 as Measured from ASTER and Landsat Images, in 63 rd
 1347 EASTERN SNOW CONFERENCE, Newark, Delaware USA. [online] Available from:
 1348 <https://www.researchgate.net/publication/228492383> (Accessed 5 July 2018), 2006.
- 1349 Kundzewicz, Z. W., Mata, L. J., W., A. N., Döll, P., Jimenez, B., Miller, K., Oki, T., Şed, Z. and
 1350 Shiklomanov, I.: The implications of projected climate change for freshwater resources and their
 1351 management, *Hydrol. Sci. J.*, 53(1), 3–10, doi:10.1623/hysj.53.1.3, 2008.
- 1352 López-Moreno, J. I., Fontaneda, S., Bazo, J., Revuelto, J., Azorín-Molina, C., Valero-Garcés, B.,
 1353 Morán-Tejeda, E., Vicente-Serrano, S. M., Zubieta, R. and Alejo-Cochachín, J.: Recent glacier
 1354 retreat and climate trends in Cordillera Huaytapallana, Peru, *Glob. Planet. Change*, 112, 1–11,
 1355 doi:10.1016/j.gloplacha.2013.10.010, 2014.
- 1356 López-Moreno, J. I., Valero-Garcés, B., Mark, B., Condom, T., Revuelto, J., Azorín-Molina, C.,
 1357 Bazo, J., Frugone, M., Vicente-Serrano, S. M. and Alejo-Cochachin, J.: Hydrological and
 1358 depositional processes associated with recent glacier recession in Yanamarey catchment,
 1359 Cordillera Blanca (Peru), *Sci. Total Environ.*, 579, 272–282,
 1360 doi:10.1016/J.SCITOTENV.2016.11.107, 2017.

Con formato: Inglés (americano)

Con formato: Inglés (americano)

- 1361 Mark, B. G., French, A., Baraer, M., Carey, M., Bury, J., Young, K. R., Polk, M. H., Wigmore, O.,
 1362 Lagos, P., Crumley, R., McKenzie, J. M. and Lautz, L.: Glacier loss and hydro-social risks in the
 1363 Peruvian Andes, *Glob. Planet. Change*, 159, 61–76, doi:10.1016/J.GLOPLACHA.2017.10.003,
 1364 2017.
- 1365 Mölg, N., Ceballos, J. L., Huggel, C., Micheletti, N., Rabatel, A. and Zemp, M.: Ten years of
 1366 monthly mass balance of Conejeras glacier, Colombia, and their evaluation using different
 1367 interpolation methods, *Geogr. Ann. Ser. A, Phys. Geogr.*, 99(2), 155–176,
 1368 doi:10.1080/04353676.2017.1297678, 2017.
- 1369 Padró, J.-C., Pons, X., Aragonés, D., Díaz-Delgado, R., García, D., Bustamante, J., Pesquer,
 1370 L., Domingo-Marimon, C., González-Guerrero, Ó., Cristóbal, J., Doktor, D. and Lange, M.:
 1371 Radiometric Correction of Simultaneously Acquired Landsat-7/Landsat-8 and Sentinel-2A
 1372 Imagery Using Pseudoinvariant Areas (PIA): Contributing to the Landsat Time Series Legacy,
 1373 *Remote Sens.*, 9(12), 1319, doi:10.3390/rs9121319, 2017.
- 1374 Poveda, G. and Pineda, K.: Reassessment of Colombia's tropical glaciers retreat rates: are they
 1375 bound to disappear during the 2010–2020 decade?, *Adv. Geosci.*, 22, 107–116,
 1376 doi:10.5194/adgeo-22-107-2009, 2009.
- 1377 Poveda, G., Waylen, P. R. and Pulwarty, R. S.: Annual and inter-annual variability of the
 1378 present climate in northern South America and southern Mesoamerica, *Palaeogeogr.*
 1379 *Palaeoclimatol. Palaeoecol.*, 234(1), 3–27, doi:10.1016/j.palaeo.2005.10.031, 2006.
- 1380 Rabatel, A., Francou, B., Soruco, A., Gomez, J., Cáceres, B., Ceballos, J. L., Basantes, R.,
 1381 Vuille, M., Sicart, J.-E., Huggel, C., Scheel, M., Lejeune, Y., Arnaud, Y., Collet, M., Condom, T.,
 1382 Consoli, G., Favier, V., Jomelli, V., Galarraga, R., Ginot, P., Maisincho, L., Mendoza, J.,
 1383 Ménégoz, M., Ramirez, E., Ribstein, P., Suarez, W., Villacis, M. and Wagnon, P.: Current state
 1384 of glaciers in the tropical Andes: a multi-century perspective on glacier evolution and climate
 1385 change, *Cryosph.*, 7(1), 81–102, doi:10.5194/tc-7-81-2013, 2013.
- 1386 Rabatel, A., Ceballos, J. L., Micheletti, N., Jordan, E., Braitmeier, M., González, J., Mölg, N.,
 1387 Ménégoz, M., Huggel, C. and Zemp, M.: Toward an imminent extinction of Colombian glaciers?,
 1388 *Geogr. Ann. Ser. A, Phys. Geogr.*, 1–21, doi:10.1080/04353676.2017.1383015, 2017.
- 1389 Ragettli, S., Immerzeel, W. W. and Pellicciotti, F.: Contrasting climate change impact on river
 1390 flows from high-altitude catchments in the Himalayan and Andes Mountains., *Proc. Natl. Acad.*
 1391 *Sci. U. S. A.*, 113(33), 9222–7, doi:10.1073/pnas.1606526113, 2016.
- 1392 Rekowski, I. C.: Variações de área das geleiras da Colômbia e da Venezuela entre 1985 e
 1393 2015, com dados de sensoriamento remoto. [online] Available from:
 1394 <https://www.lume.ufrgs.br/handle/10183/149546> (Accessed 5 July 2018), 2016.
- 1395 Ribstein, P., Tiriau, E., Francou, B. and Saravia, R.: Tropical climate and glacier hydrology: a
 1396 case study in Bolivia, *J. Hydrol.*, 165(1–4), 221–234, doi:10.1016/0022-1694(94)02572-S, 1995.
- 1397 Sicart, J. E., Hock, R., Ribstein, P., Litt, M. and Ramirez, E.: Analysis of seasonal variations in
 1398 mass balance and meltwater discharge of the tropical Zongo Glacier by application of a
 1399 distributed energy balance model, *J. Geophys. Res.*, 116(D13), D13105,
 1400 doi:10.1029/2010JD015105, 2011.
- 1401 Sorg, A., Bolch, T., Stoffel, M., Solomina, O. and Beniston, M.: Climate change impacts on
 1402 glaciers and runoff in Tien Shan (Central Asia), *Nat. Clim. Chang.*, 2(10), 725–731,
 1403 doi:10.1038/nclimate1592, 2012.
- 1404 Sorg, A., Huss, M., Rohrer, M. and Stoffel, M.: The days of plenty might soon be over in
 1405 glacierized Central Asian catchments, *Environ. Res. Lett.*, 9(10), 104018, doi:10.1088/1748-
 1406 9326/9/10/104018, 2014.
- 1407 Soruco, A., Vincent, C., Rabatel, A., Francou, B., Thibert, E., Sicart, J. E. and Condom, T.:
 1408 Contribution of glacier runoff to water resources of La Paz city, Bolivia (16° S), *Ann. Glaciol.*,
 1409 56(70), 147–154, doi:10.3189/2015AoG70A001, 2015.
- 1410 Stahl, K., Moore, R. D., Shea, J. M., Hutchinson, D. and Cannon, A. J.: Coupled modelling of
 1411 glacier and streamflow response to future climate scenarios, *Water Resour. Res.*, 44(2),

Con formato: Inglés (americano)

1412 doi:10.1029/2007WR005956, 2008.

1413 Stewart, I. T., Cayan, D. R. and Dettinger, M. D.: Changes in Snowmelt Runoff Timing in
1414 Western North America under a 'Business as Usual' Climate Change Scenario, *Clim. Change*,
1415 62(1–3), 217–232, doi:10.1023/B:CLIM.0000013702.22656.e8, 2004.

1416 Vaughan, D. G., Comiso, J. C., Allison, I., Carrasco, J., Kaser, G., Kwok, R., Mote, P., Murray,
1417 T., Paul, F., Ren, J., Rignot, E., Solomina, O., Steffen, K. and Zhang, T.: Observations:
1418 Cryosphere, in *Climate Change 2013: The Physical Science Basis. Contribution of Working*
1419 *Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*,
1420 edited by T. F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels,
1421 Y. Xia, V. Bex, and P. M. Midgley, pp. 317–382, Cambridge University Press, Cambridge,
1422 United Kingdom and New York, NY, USA., 2013.

1423 Vuille, M., Francou, B., Wagon, P., Juen, I., Kaser, G., Mark, B. G. and Bradley, R. S.: Climate
1424 change and tropical Andean glaciers: Past, present and future, *Earth-Science Rev.*, 89(3–4),
1425 79–96, doi:10.1016/J.EARSCIREV.2008.04.002, 2008.

1426 Vuille, M., Carey, M., Huggel, C., Buytaert, W., Rabatel, A., Jacobsen, D., Soruco, A., Villacis,
1427 M., Yarleque, C., Timm, O. E., Condom, T., Salzmann, N. and Sicart, J.-E.: Rapid decline of
1428 snow and ice in the tropical Andes – Impacts, uncertainties and challenges ahead, *Earth-*
1429 *Science Rev.*, 176, 195–213, doi:10.1016/j.earscirev.2017.09.019, 2017.

1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454

1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469

1470 **Table 1.** Hydrologic and climatic indices computed from the hourly streamflow, temperature and
1471 precipitation series. * *hpulse* is computed as the hourly equivalent of the melting-runoff spring
1472 pulse proposed by Cayan et al. (2001) for daily data, i.e.: the time of the day when the minimum
1473 cumulative streamflow anomaly occurs, which is equivalent to finding the hour after which most
1474 flows are greater than the daily average.

Index	Explanation	unit
<i>totalQ</i>	total daily <u>water discharge</u>	m ³ day ⁻¹
<i>Qmax</i>	value of maximum hourly <u>water discharge per day</u>	m ³ hour ⁻¹
<i>hpulse</i> *	hour of the day when the melting streamflow pulse starts	hour of the day
<i>Qbase</i>	mean <u>water discharge</u> value between the start of the day (00:00 h) and the hour when <i>hpulse</i> occurs	m ³ hour ⁻¹
<i>hQmax</i>	hour of the day when	hour of the day
<i>Qrange</i>	difference between <i>Qbase</i> and <i>Qmax</i>	m ³ hour ⁻¹
<i>Qslope</i>	slope of the streamflow rising limb between <i>hpulse</i> and <i>hQmax</i>	slope in %
<i>decayslope</i>	slope of the streamflow decaying limb between <i>hQmax</i> and 23:00h	slope in %
<i>Tmax</i>	value of maximum hourly temperature per day	°C hour ⁻¹
<i>Tmin</i>	value of minimum hourly temperature per day	°C hour ⁻¹
<i>Tmean</i>	mean daily temperature	°C day ⁻¹
<i>Trange</i>	difference between <i>Tmin</i> and <i>Tmax</i>	°C hour ⁻¹
<i>hTmax</i>	hour of the day when the <i>Tmax</i> occurs	hour of the day
<i>Diffh</i>	time difference between <i>hTmax</i> and <i>hQmax</i>	Hours
<i>Pmax</i>	value of maximum hourly precipitation per day	mm hour ⁻¹
<i>hPmax</i>	hour of the day when the <i>Pmax</i> occurs	hour of the day
<i>pp</i>	daily precipitation sum	mm day ⁻¹

Eliminado: water yield

Eliminado: water yield

Eliminado: water yield

Eliminado: d

1475
1476
1477

1478 **Table 2.** Pearson correlation coefficient between daily hydrological indices and temperature for
1479 days with and without precipitation (left) and for days only without precipitation (right) between
1480 July 2013 and June 2017. The correlation values correspond to the average obtained by 100
1481 resampling iterations (n = 99) of the correlation test. * and ** indicate that correlations are
1482 significant at 95% and 99% confidence respectively (two-tailed test).

1487

Index	days with and without precipitation (n = 99)				days without precipitation (n = 99)			
	Tmin	Tmax	Tmean	Trange	Tmin	Tmax	Tmean	Trange
total	0.25**	0.12	0.19	0.02	0.31**	0.54**	0.53**	-0.39**
Qmax	0.25**	0.30**	0.33**	-0.18	0.24*	0.64**	0.57**	-0.54**
Qbase	0.13	-0.13	-0.05	0.22*	0.18	0.05	0.11	0.06
Qrange	0.25**	0.36**	0.37**	-0.25**	0.22*	0.65**	0.58**	-0.57**
Qslope	0.18	0.40**	0.38**	-0.34**	0.12	0.58**	0.48**	-0.55**
hQmax	0.06	-0.03	0.00	0.06	0.04	0.00	0.02	0.02
Hpulse	-0.18	-0.17	-0.21*	0.08	-0.36**	-0.50**	-0.52**	0.31**

Eliminado: d
Eliminado: d

1488

1489

1490 **Table 3.** Mean and standard deviation of variance explained (%) by each PC throughout the 25
1491 bootstrapped samples

PC	Mean	standard deviation
PC1	47.78	5.91
PC2	34.99	5.66
PC3	11.82	6.77

1492

1493

1494

1495 **Figure 1.** Study area, showing the glaciers of the Parque Nacional Natural de los Nevados, and
1496 the Río Claro river basin (top map) and the Conejeras glacier with hydro-meteorological stations
1497 (bottom map).
1498

1499 **Figure 2.** Long-term evolution of climate variables in the study area. a) monthly air temperature
1500 at the Brisas meteorological station (2721 m. asl), 1982-2015 (black line), and the temperature at
1501 the glacier snout (note the difference in the range of values), 2013-2017 (blue line); b) Multivariate
1502 ENSO Index; c) monthly precipitation at the Brisas station, 1982-2015; The frequency and its
1503 equivalent in months (1/frequency) of the two top spectral densities from spectral analysis is
1504 shown for temperature, MEI and precipitation monthly series.

Eliminado: Conejeras glacier's
Eliminado: mass balance 2006-2018
Eliminado: inverted axis in the latter
Eliminado: , and the Conejeras glacier's mass balance (note the inverted axis in the latter)
Eliminado: c) monthly long-term average of precipitation at the Brisas station.
Eliminado: Red dashed lines indicate peak mass balance loss coincident with temperature peaks but not with MEI peaks.
Con formato: Resaltar
Eliminado: b
Eliminado: .

1506 **Figure 3.** Evolution of the Conejeras glacier. a) monthly mass balance in mm w.e. and Multivariate
1507 ENSO Index (not the inverted axis). b) annual mass balance per altitudinal range. c) extension of
1508 the glacier in hectares at the beginning (2006) and end (2017) of monitoring period. d)
1509 Photographs of the glacier surface covered by volcanic ashes, from 2015 and 2016.

1510

1511

1512

1513

1514

1515

1516 **Figure 4.** Principal Component Analysis on hourly streamflow. a) scores of the three main
1517 principal components (patterns of daily streamflow), with gray lines indicating the scores for each
1518 one of the 25 bootstrapped samples in the recursive PCA, and colored lines indicating the
1519 average. b) Evolution of the number of days per month that show maximum correlation with each
1520 PC. Red corresponds to PC1, blue corresponds to PC2 and green corresponds to PC3

Eliminado:

1521

1537 **Figure 5.** Summary of the frequency distributions (boxplots) of the hydrological and
1538 meteorological indicators for days grouped within PC1, PC2 and PC3.

1539

1540

1541 **Figure 6.** Evolution of monthly averaged hydrological indices, temperature, precipitation and
1542 glacier mass balance (in blue bars), for the study period. Dashed lines indicate the 2015-2016
1543 strong El Niño event. 12-months window moving average (black smooth lines) are shown to
1544 represent trends.

1545

1546

1547 **Figure 7.** Mean monthly water discharge (Q), for days with temperature lower than 2°C (blue)
1548 and days with temperature higher than 2°C (red) Top: Inter-annual evolution with indication of El
1549 Niño 2015-16 event (grey shading), breakpoint in water discharge evolution (dashed line), and
1550 monthly precipitation (blue bars); bottom: comparative boxplots for water discharge before and
1551 after breakpoint in May 2016.

1552

1553 **Figure 8.** Correlations between monthly flow and monthly temperature (top plots) and
1554 precipitation (bottom plots) for: a) 2013-14 (blue triangles) and 2017 (red circles) years, which are
1555 considered as analogues in terms of amounts of flow; and b) months before May 2016 breakpoint
1556 (blue triangles) and months after May 2016 breakpoint (red circles).

1557

Eliminado: on the top two plots

Eliminado: water yield

Eliminado: water yield

Eliminado: water yield

The objective of the present work is to