

Dear Editor,

This document provides a point-by-point summary of the feedback on the manuscript ‘Global Phosphorus Recovery for Agricultural Reuse’

Please also find attached with this document: 1) the revised manuscript, and 2) a marked-up document showing all changes made.

Sincerely,

On behalf of all authors,
Dirk-Jan Kok

Editor Feedback:

1. Mathematical notations: Please adhere to the HESS requirements. Remove for example the “*” symbol for multiplication in all equations and brackets around units if they are a normal part of the text.

We have replace asterix multiplication symbol with the recommended ‘.’ symbol.
We have ensured that all units in text are in brackets

2. p. 1: Move “through struvite crystallization” (p. 1, L. 14) to p. 1, L. 8 (after “wastewater”).

Moved “through struvite crystallization” from p. 1, L. 14 to p. 1, L. 9.

3. p. 4, L. 4: Remove comma after “objective”

Removed comma after “objective”

4. p. 4, L. 11: “organisms” instead of “organism”

Corrected spelling of “organisms”

5. p. 4, L. 13 (and elsewhere): What does the symbol “h” stand for in the bracket?

Replaced symbol ‘h’ with ‘head’ (per individual).

6. p. 4, L. 15: The expression “person individual” sounds weird.

Corrected “person individual” to “[...] each individual person”

7. p. 13, L. 23 – 25: “relative accuracy”: Please provide a quantitative description of how well the model performs.

Removed “relative accuracy”. The paragraph now presents the numerical differences between predicted and observed prices.

8. p. 13, L. 33: Table S7 is actually a figure.

Moved to figures section and renamed in text.

9. P. 13, L. 33 – 37: This paragraph is very superficial. Screening through the results that you show in the SI, I think some more findings should be reported. As it appears now it is more an exercise to demonstrate that a sensitivity analysis has been carried out but you do not really provide much insight for the reader. However, your figures reveal quite diverse patterns ranging from a complete absence of sensitivity for certain variable pairs, over linear relationships to very non-linear patterns. Please provide some essential findings and refer for explicitly to specific data that you show in the Supplementary Material.

We now present the figure data also in a table (Figure S3 → Table S7).

We elaborate on the implications of the sensitivity analysis in the Sup. Mats. and mention some essential findings in order to provide insight for the reader.

10. Fig. 3: Please increase the font size for better readability.’

We increased the fontsize.

11. Fig. 4: Please change the colour set you use. It is hardly possible to distinguish the different categories.

We changed the colour set so that it is easier to distinguish the different scenarios.

Referee Feedback:

1. “If this is the case – that prices are not included in the analysis before a certain point (please specify which point), then this has to be mentioned explicitly in the paper.”

Explicitly mentioned in section titles under the results to avoid confusion:

- section 3.1. ‘*Phosphorus recovery potentials **excluding economic dynamics.***’
- section 3.2. ‘*Phosphorus recovery potentials **including economic dynamics***’

6. “I have not found any explanation or sources of data for labour wage estimates in the Supplementary Materials.”

We now explain the origin of the labour wage estimate. This explanation is presented under the transport parameter table (Table S3).

“* 15 [\$ /h] value is an informed estimate of trucking labour wage. In reality, trucking labour wages will vary globally. The model could not account for spatial diversity in wages provided the absence of a global data set on trucking wages per country or even per region. Instead, the 15 [\$ /h] was determined taking into consideration the wage indicators from the U.S. (c.a 20 \$/h) (United States Department of Labor, 2017), the Netherlands (c.a 30\$/h) (MijnZZP, 2018), and China (c.a 5 \$/h) (Salary Expert, 2018).”

It seems unnecessary to reference these sources. We leave the decision up to the editor on whether or not to include these since the information they provide are only used by the authors as inspiration and are not directly used by the model or in the text.

12. “My point here was to make a clear link between spatial variation in P, its concentration in wastewater, and population and livestock density - ideally in a single sentence (e.g. in Section 1.1). While this connection may be very obvious to someone working on phosphorus, it would be useful to spell it out for non-experts.”

Section 2.1.1 reiterates the importance of spatial variation in P (p. 5 L. 0 to p. 5 L. 3):

“The spatial variation in recovered P-production potential is therefore marked by global variation in population and livestock densities. [\\] The locations for the P-mining industry are, instead, geographically concentrated.”

Other changes:

1. Renamed the title ‘Global Phosphorus Recovery for Agricultural Reuse’ to ‘Global Phosphorus Recovery **from Wastewater** for Agricultural Reuse’
2. Minor spelling and grammar fixes throughout the text

Global Phosphorus Recovery from Wastewater for Agricultural Reuse

Dirk-Jan D. Kok^{1*}, Saket Pande¹, Jules B. van Lier¹, Angela R.C. Ortigara², Hubert Savenije¹, Stefan Uhlenbrook^{1,2}

¹ Department of Water Management, Delft University of Technology, Delft, Netherlands.

² World Water Assessment Programme, UNESCO, Perugia, Italy.

5 Correspondence to: Dirk-Jan Kok (D.D.Kok@cml.leidenuniv.nl)

Abstract. Phosphorus is a nutrient necessary for the development of crops and is thus commonly applied as fertilizer to sustain agricultural production. It occurs naturally, at indefinite quantities of uncertain quality in phosphate rock formations, but also accumulates in urban and livestock wastewater wherefrom it is often lost as a pollutant. Recovering phosphorus from wastewater, however, is ~~technologically~~ feasible through struvite crystallization technologies and has the potential to reduce phosphorus pollution of the environment as well as lower the agricultural demand for artificial P-fertilizers. In this study, we developed a model to assess the global potential of P-fertilizer recovery from wastewater ~~and~~ to visualize its trade at sub-national resolution. Results show that humans discharge a maximum of 3.7 [Mt] P into wastewater, thereby potentially satisfying 20% of the global fertilizer demand. Provided 2015 market dynamics, however, the model determines that only 4% of this discharge is technologically and economically recoverable in a market that offers cheap rock phosphate products also. The results of this study demonstrate that in the current economic context, phosphorus recovery from wastewater ~~through struvite crystallization~~ offers only a small contribution to resolving global phosphorus issues ~~provided current economic contexts~~. Nevertheless, this recovery offers many wastewater treatment facilities the opportunity to contribute to creating sustainable communities and protecting the environment locally, while reducing their own operational costs.

Key words: *Phosphorus, circular economy, wastewater, struvite, fertiliser*

20 1 Importance of Phosphorus

Phosphorus (P) is an element necessary for the development of all living beings as it forms an unsubstitutable, key structural component of DNA and RNA. It is also a limiting nutrient, and therefore growth is often restricted by the ~~natural~~ lack of P naturally available. While P-related malnutrition in humans is uncommon, inhibited plant growth due to soil phosphorus deficiency is a much more prevalent issue (MacDonald, Bennett, Potter, & Ramankutty, 2011). For this reason, phosphorus is often applied to croplands in the form of organic fertilizers or synthetically as single or triple superphosphate, or mono-Ammonium or di-Ammonium Phosphate (DAP). These fertilizers are easy to transport and distribute over fields, while also readily absorbed by plants. The most essential resource for the production of artificial phosphorus fertilizers is phosphate rock.

The rates at which we exploit phosphate rock formations is out of proportion to the rates at which they form - essentially classifying phosphate rock as a non-renewable resource. Peak production of phosphate ore could occur as early as 2030 (Cordell et al., 2009) with economically extractable P resources becoming scarce or exhausted within the next 50 to 100 years (Smil, 2000; Steen, 1998; Van Vuuren et al., 2010). Other authors predict more optimistically that reserves will last another 300-400 years (van Kauwenbergh, 2010). After this depletion, it will likely ~~not~~ be economically infeasible to exploit the

remaining ore ~~because as~~ it will be too costly to process due to its poor quality (i.e. low phosphorus and high heavy metal content). Gradual depletion of economically extractable reserves will result in further reduction in accessibility to fertilizer by small-holder and subsistence farmers that comprise areas already struggling to cope with food shortages (Pande and Savenije, 2016). Sub-Saharan Africa is one such region, as nearly 75% of its agricultural soils are nutrient deficient, contributing significantly to the crop yield gaps (Sanchez et al., 1997). The prospect of phosphorus depletion ultimately threatens global food security where especially regions of poor soil nutrition levels are vulnerable to its effects.

1.1 The Environment, Humanity, and Phosphorus

The introduction of intensified (P) fertilization during the Green Revolution of the 1960's, demonstrated P's significant potential to improve crop yields but also the dangers it poses to the environment. Through seepage and runoff processes (Bouwman et al., 2009), as well as the discharge of improperly treated wastewater (Van Drecht et al., 2009; Morée et al., 2013), phosphorus and other nutrient excesses come into contact with open surface water. As a limiting nutrient, even the smallest quantity of P in water can spark the growth of large algal blooms. These algal blooms have a detrimental effect on aquatic ecosystems, in causing the suffocation of aquatic life through eutrophication, resulting in a loss of habitat and biodiversity (EPA, 2010). Such deoxygenated "dead zones" can be found in both lakes and seas, and affect an estimated 245,000 km² of marine ecosystems (Corcoran et al., 2010). If excess fertilization and water pollution is a major threat to water quality around the world, then why not extract this excess from the water system and put it back in the food chain? Proper nutrient management practice in tandem with nutrient recovery from rural and urban water systems may potentially be an important strategy to reduce phosphorus pollution by reducing phosphorus discharge to the environment while simultaneously increasing phosphorus supply for food production.

There exist a broad range of sources that contribute the phosphorus loads in ~~urban~~-wastewater, ranging from excrements to detergents, ~~to-from~~ toothpaste ~~to-~~ dishwashing liquids, ~~and from~~ medicines ~~to-~~ food preparation wastes ~~and-food~~ leftovers. There also exist numerous technologies that may be employed to recover this phosphorus from wastewater (Egle et al., 2016). Unfortunately, ~~they-these technologies~~ are often deemed too costly to implement and operate while their recovery efficiencies vary, among others, with local wastewater composition and existing wastewater treatment infrastructure. Adoption of these technologies is therefore often challenged by (perceived) economic infeasibility or lacking economic incentives and social stigma. Yet, ~~it must be recognized that the~~ economic feasibility of recovery is not globally homogeneous, but varies in space and time. Spatially, the global accretion of phosphorus in wastewater provides recovered products with a (diffused) location-defined competitive advantage over the geographically concentrated rock phosphate mines (fig. 1). While temporally, the appeal for recover will improve over time with the increasing price trends for rock-based fertilizers (e.g. DAP) (fig. 2).¹

1.2 Turning brown waste to green gold

To our knowledge, there exist no studies that evaluate the *spatially dependant* feasibility for phosphorus recovery from wastewater, at sub-national resolution, in a global and economically dynamic context. Insights provided by such a study, however, could accelerate the efficient transitioning to a more sustainable phosphorus fertilizer market by illustrating where

¹ Over the past 15 years the phosphorus price of DAP has increased from 665 [\$ t⁻¹] to 1,552 [\$ t⁻¹]. In that same period, the price has been as high as 5,217 [\$ t⁻¹] (2008) and as low as 656 [\$ t⁻¹] (2002) (IndexMundi, 2017).

recovery may economically be most feasible. We therefore aim to determine the total phosphorus recovery potential from wastewater, as well as the economic feasibility for this recovery, in a global assessment. This is achieved by integrating geospatial data, statistics, and findings from other studies into a model that identifies and connects phosphorus recovery and demand sites based on location, quantities, and prices.

5

Because of the wide array of pathways to phosphorus recovery, the subject of this investigation is constrained to the recovery of phosphorus from urban and livestock wastewater as a struvite and compost pellet fertilizer product, only. Phosphorus lost via other fluxes (e.g. municipal solid waste) are therefore excluded from this assessment. The sole reason for focusing on recovery through ~~the struvite crystallization precipitation technology~~ is because of the fertilizer potential of struvite ~~and as well~~ as the current industrial-scale implementation of the technology itself (Cornel and Schaum, 2009).

10

2 Materials and Methods

The phosphorus cycle is delineated by a combination of both social and physical attributes and as such demands a coupled human water systems perspective. A socio-hydrological approach is endeavored as both these attributes are here expressively and emphatically accounted for (Sivapalan et al., 2012). The social component, however, of this coupled human-phosphorus system is confined to the characteristics of a distinct economic nature. The materials and methods employed in assessing the recovery potential of P from wastewater at global scale is therefore interdisciplinary and extensive, covering largely economics within a sciences context.

15

In general, the methodology can be summarized to consist of three phases which are also used to define the order of the materials and methods section:

20

1. **Identification of sites and quantities:** Identifying the locations of wastewater accumulation sites and agricultural croplands and assessing the potential associated phosphorus production and demand quantities.
2. **Determination of node prices:** Approximating the minimum production costs of recovering P-fertilizers at the wastewater accumulation (recovery) sites, and the maximum paying prices for P-fertilizers at the agricultural (consumption) sites.
3. **Modelling international trade** in phosphorus, involving:
 - a. *Determination of global market price:* Determining an international, free-market price for phosphorus as a function of phosphorus quantities and prices, as well as the distances between the different sites.
 - b. *Visualizing trade flows:* Creating a realistic network of P-trade fluxes at subnational resolution.

25

The main tools employed in this investigation are Geographic Information System (GIS) tools, used to prepare the spatial data, (Q-GIS 2.14) (Quantum GIS Development Team, 2017); and Python 3.6, used to build the trade network model.

30

2.1 Identification of Sites and Quantities

Phosphorus trade occurs between production and demand sites, and therefore two groups of actors are identified in the model: P-producers and P-consumers. The first group, P-producers, consists of three types of actors:

35

1. **Urban wastewater treatment facilities** – Nodes recovering P from domestic wastewater (i.e. phosphorus excreted, used, and discharged by humans);
2. **Livestock keepers** – Nodes recovering P from animal manure (i.e. manure and liquid waste produced by farm cows, chicken, and swine, while stabled);
3. **Phosphate mines** – Nodes extracting P from rock phosphate reserves.

The three actor types recover phosphorus in different forms, yet all act on the same phosphorus market. The only objective characteristic that the model will use to distinguish one type product from another, is the absolute elemental P value of the product (i.e. U.S. dollars per mass P; [$\$ \text{mass}^{-1}$] P). Since both mines and recovery sites have the same purpose in the model (i.e. supplying phosphatic fertilizers), both node types ~~will be~~ are grouped as ‘production nodes’.

2.1.1 Phosphorus Production

The location and phosphorus production quantities of P-production nodes are determined by integrating geospatial datasets with statistics and findings from other studies. As phosphorus is not ‘produced’ by organisms, but only consumed and excreted, or used (i.e. detergents), we will refer to the annual amount of phosphorus discharged in wastewater per individual, as the ‘phosphorus throughput rate’ [$\text{kg head}^{-1} \text{a}^{-1}$] P. Wastewater phosphorus has variable sources ranging from human excrements to detergents, toothpaste, dishwashing liquids, medicines, food preparation wastes, food leftovers, etc., ~~but~~ These sources are not considered individually. Instead we assume that each ~~person~~ individual ~~person~~, globally, contributes an equal contribution to the phosphorus load in wastewater. We combine approximations for these rates from different studies with population density maps for: humans (CIESIN, 2016), cattle, swine, and poultry (Robinson et al., 2014), to determine the spatial distribution of phosphorus excretion rates globally. This spatial distribution of the mass phosphorus produced per unit area, per annum, [$\text{kg km}^{-2} \text{a}^{-1}$], represents the phosphorus production density (a map). Its determination per unit area on this map is summarized for as follows (eq. 1):

$$S = (D * P) * E \quad (1)$$

where S is the maximum organic phosphorus production density [$\text{kg km}^{-2} \text{a}^{-1}$]; D is the population density [heads km^{-2}]; P is the phosphorus throughput rate [$\text{kg head}^{-1} \text{a}^{-1}$]; and E is the estimated base recovery efficiency [-]. For humans, the phosphorus throughput rate (P) is assumed globally homogeneous, fixed at 0.77 [$\text{kg head}^{-1} \text{a}^{-1}$]. This is in close relation to other published findings (0.77, Gilmour et al. (2008); 0.78, CRC (2005); 0.2-0.7, Mihelcic et al. (2011); 0.7, Smil (2000)). For livestock, the throughput rate (P) is taken to be a function of slaughter weight (FAOSTAT, 2018) following the methodology of Sheldrick et al., (2003). Manure recovery efficiencies (E) are taken as: cattle (31%), swine (80%), poultry (77%) (Sheldrick et al., 2003). For humans, it is assumed that 100% of the phosphorus discharged as domestic wastewater reaches the treatment facility. The actual recovery efficiencies will vary per recovery technology implemented at the wastewater treatment plants (WWTP) (section 2.2.). The production density maps for livestock and humans are presented in fig S1a and S1b of the Supplementary Materials. As a final step, the phosphorus production densities per land area are converted into point nodes using GIS tools (see section 2.1.3).

The location and P-production values for the mining industry are acquired from a USGS (2002) dataset. This dataset is adjusted to match the USGS reported phosphate production estimates for different simulation years.

2.1.2 Phosphorus Consumption

Similar to the phosphorus production density, the phosphorus demand density map represents per unit area the yearly amount of phosphorus required by agriculture [kg km⁻² a⁻¹]. It is determined following a comparable methodology to that of phosphorus production, where crop densities, as approximated through crop harvested area maps (Monfreda et al., 2008), are related to phosphorus-requirement rates (UNIDO and IFDC, 1998). The crop phosphorus requirement rates [kg km⁻² harvest⁻¹] as reported by UNIDO and IFDC (1998), however, are determined for optimal yield. Provided that no farmer is going to fertilize for optimal yield when he or she knows that these yields are unachievable provided persistent regional water limitations, the actual P demand is proportionally reduced to the potential *water-constrained* yield. This assessment is made for six major crops: maize, wheat, rice, sorghum, soy bean, and potato. The water-constrained yield is determined by adapting the evaporation-transpiration deficit equation (Steduto et al., 2012) to the following (eq. 2):

$$1 - \frac{Y_a}{Y_m} = K_y \left(1 - \frac{E_a * \frac{T_g}{365} * A_H * C}{E_m} \right) \quad (2)$$

where Y_a is the actual yield [kg km⁻² a⁻¹]; Y_m is the optimal yield [kg km⁻² a⁻¹]; K_y is the crop coefficient [-]; E_a is the cumulative actual evaporation-transpiration per year [mm a⁻¹ Area⁻¹]; T_g is the duration of the crop growing period [d]; A_H is the fractional area harvested [-]; E_m is the evaporation-transpiration for optimal yield per harvest [mm a⁻¹ km²]; and C [-] is a correction factor. In this investigation, E_m is assumed equal to the crop water requirement for optimal yield. Global approximations of K_y and E_m values were retrieved for different crops from FAO sources (2015), and E_a was approximated from MODIS evapotranspiration products (NASA, 2005). A summary of the data used and its sources is presented in Table S2 of the Supplementary Materials.

Eq. 2 is an adaptation of the original evaporation-transpiration deficit equation. The original equation is designed for a single growing period, yet much of ~~our~~^{the other} input data is yearly. As the beginning and ending of a crop's growing season will vary globally, it is not possible to combine global *yearly* harvest area maps with *monthly* evaporation data. Reformulation was therefore necessary to account for a difference in temporal scales of the input data. For the adaptation, it is assumed that of the yearly evaporation, an amount proportional to the duration of a crops growing period is evaporated during the crop growing season ($E_a * \frac{T_g}{365}$). This value is then further reduced by multiplication with the fractional crop harvested area (A_H) to account for evaporation from other landcover types in the area as well. Finally, as these simple manipulations introduce a significant error, a correction factor (C) was added to globally scale potential yields greater than optimum (*i.e.* >1), back down to optimum (*≡*1) and to achieve a total, global phosphorus demand that approaches observed values for these crops.

In this investigation, we assume no change in soil stored phosphorus. The yearly phosphorus demand per area therefore reflects only the crops yearly phosphorus uptake as a function of a crops water-constrained yield and harvested area. This phosphorus demand is described as a linear regression between yield and P-fertilizer requirement through eq. 3:

$$D_{PT} = \frac{Y_a}{Y_m} * A_H * P_{opt}^n \quad (3)$$

where D_{PT} is the calculated phosphorus demand density [$\text{kg ha}^{-1}\text{a}^{-1}$]; P_{opt}^n is the crop specific (n) P-requirement for optimal yield [kg ha^{-1}]; and A_H is the crop harvested area [ha^{-1}]. The parameters per crop are again summarized in Table S2 of the Supplementary Materials. The six crops evaluated for make up roughly 56% of the global demand (Heffer, 2009). Although spatially inaccurate ~~to~~of the actual global phosphorus demand distribution, the total, global phosphorus demand quantity is approached by dividing each area's value for the six major crops by 0.56 (56%). The demand density maps for agriculture are presented in fig S1c of the Supplementary Materials.

10 2.1.3 From raster to nodes

The areas of major production and consumption densities, determined in 2.1.1 and 2.1.2, are aggregated into nodes that are described by a coordinate position, a class (group: urban or livestock, or crop type), and a quantity of yearly phosphorus supply or demand. Each node is systematically positioned in the centre of a larger area of uninterrupted high phosphorus production or demand density as determined by the raster map calculations preformed in steps 2.11 and 2.12. To avoid the aggregation of administratively separate regions into a single node, the areas of continuous high demand/production density are separated by national boundaries for smaller countries, and first level administrative borders (e.g. states and provinces) for large countries (e.g. U.S.A., India, Russia, Canada, etc.). Nodes with a production values of less than three kilo tonnes per year are considered insignificant in the global context and are therefore excluded from further consideration in the economic analysis. This constrains the total number of actors, reduces the complexity of the network, decreases the processing time, and improves visualization of the results. This preselection reduced the global P-recovery quantity by 15%, while reducing the number of actors by 76%. The trade model thus only accounts for trade from ~~25~~24% of all potential recovery sites, which, nevertheless, represent 85% of the global recovery potential.

2.2 Determination of Node Prices

Throughout this investigation, prices ~~will be~~are presented as the price per tonne phosphorus in each fertilizer product. [$\text{\$ t}^{-1}$]. This is because struvite fertilizers (14% P), conventional artificial fertilizers (DAP: 22% P), and compost pellets (1% P) are, although different in form and P contents, acting on the same phosphorus market. In the model market, they are discriminated only based on their total phosphorus content and, as such, their values are determined based solely on the mass of P that they contain.

Whether trade is possible between a demand and a production node depends on the transportation and production costs of the production node and the maximum bid price of the demand node. Although having identified three different producers (section 2.1), each creating their own unique fertilizer products, they are all ~~are~~ subjected to the same economic constrictions. The production cost of each producer is defined by: i) an investment cost for infrastructure, and ii) a variable cost per mass P recovered. Additionally, they are imposed with iii) a transportation cost for selling. The sum of these costs determines how attractive a producer is to the individual, agricultural demand nodes.

2.2.1 Cost for Recovery

The cost for recovery (i.e. production cost) varies depending on the recovery technology whose feasibility for implementation, in turn, depends on wastewater composition and existing infrastructure. In this study, municipal wastewater composition is assumed roughly, globally homogenous. The Sustainable Development Goals (SDG) dataset on *percent of urban population with access to sanitation* is then used to approximate how well developed the existing sanitary infrastructure is in different parts of the world (WHO/UNICEF JMP, 2015). This estimation is then used as an indicator for the feasibility to implement certain recovery technologies. How the SDG dataset is used to interpret the feasibility for implementation of specific recovery technologies, is presented below:

A high national percentage of urban population with access to sanitation (>90%) is likely to be indicative of highly developed countries that observe stricter effluent standards and that therefore have, or are working towards upgrading, conventional wastewater treatment plants (WWTP) for biological nutrient recovery (BNR) or chemical phosphorus removal. The phosphorus recovery costs for these highly developed nodes are those associated with the investment in a large Ostara® Pearl Reactor for struvite precipitation. The production cost for this highly developed group is reduced by the savings in uncontrolled struvite scaling maintenance and sludge handling costs that are associated with controlled struvite precipitation (Shu et al., 2006). Although other struvite recovery technologies are available, Ostara® Pearl Reactors were chosen given their commercially effective implementation in various different countries. There exist other recovery technologies that allow for absolute greater recovery amounts (see Egle et al., (2016)), but few of these are potentially economically competitive producers of pure P-fertilizer products.

Most of the influent phosphorus at a WWTP accumulates at the centrifuge (80-90%), where it is separated in centrifuge cake (~85%) and liquor (~15%) (Jaffer et al., 2002). Struvite crystallization from centrifuge liquor achieve efficiencies higher than 90% (Jaffer et al., 2002; Münch and Barr, 2001). Therefore, assuming some variability amongst WWTP, we make the optimistic estimate that approximately 20% of influent wastewater phosphorus may be recovered at BNR WWTP through struvite precipitation.

Areas with intermediate urban access to sanitation (40-90%) are assumed to be serviced by simple, centralised wastewater treatment facilities. The technology investment cost for these nodes are the same as for the highly developed infrastructure group but excluding this time the sludge handling cost savings. The recovery efficiency is again assumed to equal 20% of the influent P.

Low urban access to sanitation (<40%) is taken to be indicative of low sanitary development and thus offers the flexibility to adopt more novel, less water dependant forms of sanitation. The technology applied for these areas are source separating-, dry composting toilets, where urine and faecal compost are collected separately. The urine is collected by 40,000 litre tank trucks and processed at a centralized struvite precipitation facility. Faecal compost is collected, dried and processed into compost pellets also at a central facility. A 90% efficiency is easily achieved when struvite is precipitated from source-separated urine (Wilsenach et al., 2007). It is furthermore assumed that all of the faecal phosphorus is retained in the compostation, drying and pelletization processes.

For livestock nodes the collected manure is composted, dried and pelletized, also. As opposed to the source-separated faeces of dry composting toilets, the pelletization of manure from livestock farms occurs not at a centralized facility, but on-site. This is assumed feasible provided due to the high volumes of manure produced by livestock in comparison compared to humans.

For all production nodes, a generalized phosphorus recovery cost can be described as follows (eq. 4):

$$f_{min}^i = \frac{R * S_{PT} + B}{S_{PT}} - S + \frac{T^i}{\rho} \quad (4)$$

5 where f_{min}^i is the minimum price for phosphorus produced at node i [\$ t⁻¹]; R is the variable cost per tonne P recovered (i.e. the magnesium cost for struvite precipitation; the pelletization cost for compost pelletization) [\$ t⁻¹] P; B fixed operational cost [\$]; S_{PT} is the total phosphorus recovery potential [t]. Furthermore, where relevant, S is the struvite scaling and sludge handling cost savings per tonne P recovered [\$ t⁻¹] P (i.e. for BNR plants); T^i is the intracity transport cost of collection [\$ t⁻¹] (i.e. for urine and faeces from source-separating toilets); and ρ is the proportion of phosphorus by weight of the transported material
10 [kg kg⁻¹], which are 0.066% and 0.46% for urine and dried, faecal, toilet compost respectively (Vinnerås, 2001). The annual fixed operational costs are taken as the annual costs minus the resource costs as reported in the dissertation of Egle (n.d.), TU Wien. For BNR plants, a struvite and sludge handling cost savings is included of 0.89 [\$ kg⁻¹] struvite removed (Shu et al., 2006). This essentially allows BNR plants to supply struvite for free (excl. transportation costs), which common occurrence for struvite precipitating BNR WWTPs in The Netherlands. The additional variable cost for the dry-toilet solution is attributed
15 to the collection of waste and a pelletization cost of \$30 [\$ t⁻¹] compost (Masayuki Hara, 2001). A summary of these and other data values are presented in Tables S3, S4 and S5 of Supplementary Materials.

The intracity transport cost T^i is described as followed (eq. 5):

$$T^i = D_{ta} \frac{P_d * E_{Lt} + \frac{L_{ct}}{\bar{V}_{Lt}}}{W_{Lt}} \quad (5)$$

20 Where D_{ta} is the average distance of a return journey for a tanker truck in servicing the city per full load [km]; P_d is the price of diesel [\$ L⁻¹]; E_{Lt} is the fuel efficiency of the tanker truck [L km⁻¹]; L_{ct} is the labour wage [\$ head⁻¹]; \bar{V}_{Lt} is the average velocity of the truck in the city [km head⁻¹]; W_{Lt} is the truck load weight [t]. The transportation cost from the processing facility to the consumer is included in the final sale price [\$ t⁻¹] of the producer and is therefore not accounted for yet at this stage.

25

2.2.2 Maximum Buyer Bid Price

The maximum price that demand nodes are willing to purchase phosphorus at depends on the marginal value of phosphorus. This varies per crop type and can be described as followed (eq. 6):

$$30 \quad f_{max}^n = \frac{Y_{opt}^n * C_a^n}{P_{opt}^n} * R^n \quad (6)$$

where f_{max}^n is the maximum price for phosphorus [\$ t⁻¹]; Y_{opt}^n is the optimal yield [t ha⁻¹]; C_a^n is the crop price in year a [\$ t⁻¹]; P_{opt}^n is the optimum fertilizer dosage rate (equal to total P-requirement for optimal, water constrained yield) [t ha⁻¹]; and R^n is the ratio of fertilizer cost to total production costs [-], for crop n .

35

2.2.3 Transportation cost

Lastly, the transportation costs between the production and demand sites are determined with *as-the-crow-flies* distances and the parameters given in Table S3 (Supplementary Materials) substituted into the following transport cost equation (eq. 7):

$$T_c^{i,n} = D^{i,n} \left[F_W * \frac{P_b * E_W + C_F}{\bar{V}_W * W_W} + F_L * \frac{P_d * E_L + L_c}{W_L} \right] \quad (7)$$

5

where $T_c^{i,n}$ is the transportation cost from node i to node n [\$ t^{-1}\$]; $D^{i,n}$ is the distance between node i and node n [km]; P_b is a bunker fuel price (shipping fuel) [\$ t^{-1}\$]; P_d is the price of diesel [\$ L^{-1}\$]; E_W is the fuel efficiency of a container mainliner ship [t d⁻¹]; E_L is the fuel efficiency of a 2x30 tonne truck combination [L km⁻¹]; C_F is the fixed costs per ship [\$ d⁻¹]; L_c is the labour wage [\$ head⁻¹]; \bar{V}_W is the average velocity of the ship over water [km d⁻¹]; \bar{V}_L is the average velocity of the truck over land [km head⁻¹]; W_W is the full ship weight [t]; W_L is the full truck weight [t]. F_W and F_L are the fractions of the total distance that is travelled over land and over sea. The model at present does not distinguish between the transportation over land and over sea based on observed geography. Instead the model employs a cumulative probability curve that approximates the proportion of the total distance likely to have been traversed over water, F_W (eq. 8); and over land, F_L (eq. 9). It is assumed that at least 15% of the total distance is always transversed over land.

$$F_W = \frac{85}{1 + e^{\frac{D - \mu}{S}}} \quad (8)$$

$$F_L = \frac{85}{1 + e^{\frac{D - \mu}{S}}} + 15 \quad (9)$$

where μ and S are function shape constants of 500 and 100 [-], respectively.

2.3 Trade model

A model is constructed to determine i) the market price for international phosphorus trade, and ii) what amounts of phosphorus are being traded between which nodes. Market prices emerge as a function of the individual production/consumption prices and associated supply/demand quantities, which will vary depending on which actors are included in the market scenario. The phosphorus recovery potential is therefore assessed for three different combinations of actors that represent, respectively, the current, potential near-future, and potential far-future markets:

- 1) **Current market - mine supplied products only:** The current phosphorus market is strongly rock-phosphate oriented. When the model runs the data for a ‘current market’ scenario, only rock phosphate products are available on the market. This scenario serves mostly a model validation purpose.
- 2) **Future market - both mine and recovered products:** Recovered phosphorus may become a more important product in the future market. When the model runs the data for a ‘future market’ scenario, it is assumed that both rock phosphate as well as recovered phosphorus products partake.
- 3) **Far-Future market - only recovered products:** In the far future market, most rock phosphate reserves will have been depleted. When the model runs the data for a ‘far-future’ scenario, it is assumed that rock phosphates no longer take part in the market which is then solely dominated by sustainable, recovered products.

3)4)

Please note that the model does not actually make future prediction. All parameters and input data is maintained for the same 2005 to 2015 period. For the 'future' scenario's, only the sets of actors change as described per above.

5 2.3.1. Determination of global market price

The global market price is determined as the price at which total quantity of phosphorus demanded (sum of agricultural demand quantities) is equal to the quantity supplied (sum of P-production quantities). It is approximated as the point where global demand function (defined as cumulative phosphorus demand vs. maximum buying price), intersects the global supply function (cumulative phosphorus production vs. production price). The demand function is the locus of the maximum prices at which demand nodes are willing and able to purchase phosphorus, and the supply function is the locus of minimum prices at which supply nodes can sell certain amounts of phosphorus without going out of business (i.e. without making a loss). Where the two curves intersect, the market for P is cleared providing a best approximation of the market price (Arrow and Debreu, 1954). The Supplementary Text provides an illustration of how market prices for the three scenarios are determined following this principle, and how transportation costs complicate this method of price determination.

15 2.3.2. Quantification of Trade Flows

The quantity of phosphorus traded for each scenario is determined following a method of reduction and elimination. Firstly, a list of all possible combinations of supply and demand nodes is created. Each combination of supply and demand nodes is passed through two 'filters' that removes some pairs provided simple conditional statements. This reduces the list down to a selection of trading node pairs for each market scenario, for each year.

20

Before anything, the first filter in the model removes the pairs that can never trade with each other based on their combination of the minimum production costs, the maximum bidding price, and transportation costs associated with the data of that year (e.g. fuel cost, wastewater flows, etc). The second filter then removes node pairs which cannot trade with each other at a given 'hypothetical market' price imposed on the network. Either the production cost may be above-, or the maximum bidding price may be below the imposed market price, implying that the nodes cannot trade. If both the production cost is below- and bidding price is above the imposed market price, then the node pair is left in the list for that 'theoretical' market price. After these two filters, the list of potential trade partners is reduced significantly and assessment may be made as to whether the saved pairs actually trade and then with what quantities.

30 In the model, phosphorus consumers will look for the cheapest suppliers. The matter becomes obscure here as, in reality, there are no cheaper or more expensive suppliers for a single market price. Supply nodes that could, however, supply at prices far lower than the set market price (due to lower production costs) have a competitive advantage over those that cannot. At the same time, agricultural demand nodes that are willing to pay much more than the hypothetical market price, have a greater financial capacity to outbid those agricultural nodes whose maximum bid price is much closer to the hypothetical market price.

35 The difference between this market price and i) the production and transportation costs for production nodes and ii) maximum bid price for demand nodes, shows how competitive a node pair is. If a node is *able* to produce and transport at prices much lower than market price, and if a demand node is *able* to pay much more than the market price, then the model assumes that

trade between these most competitive nodes occurs first. Therefore, the list of remaining node pairs is sorted according to the greatest difference between production + transport cost, and maximum bid price, with market price.

For each of node pair lists for the different ‘hypothetical market’ prices, trade is executed and the list updated. After each trade, the list is updated in terms of total phosphorus quantity (Q) demanded by demand node (D) number n , or supplied by supply node (S) number i (D_Q^n , and S_Q^i , for demand and supply respectively). The amount traded ($Q_{(i,n)}$) between each node pair is taken to be equal to the minimum of supply or demand as formulated below (eq. 10):

$$Q_{(i,n)} = \begin{cases} S_Q^i & \text{if } D_Q^n > S_Q^i; \\ D_Q^n & \text{if } S_Q^i > D_Q^n; \end{cases} \quad (10)$$

The supply available and quantity demanded at each supply demand nodes are updated as follows (eq. 11),

$$\begin{aligned} S_Q^i &= 0, & D_{Q(new)}^n &= D_{Q(original)}^n - Q_{(i,n)} & \text{if } D_Q^n > S_Q^i; \\ D_Q^n &= 0, & S_{Q(new)}^i &= S_{Q(original)}^i - Q_{(i,n)} & \text{if } S_Q^i > D_Q^n; \end{aligned} \quad (11)$$

By eq. 11, one of the nodes will have 0 production capacity or demand after each trade, and so all possible trade combinations with that node are removed from the list of possible trade partners **remaining** for that hypothetical market price. The trade is recorded in a separate list of ‘successfully executed trades’. This process is continued until the list is empty and thus all feasible trade for that price has been conducted. Plotting the cumulative phosphorus trade for each hypothetical market price simulated, results in a combined version of the supply and demand curves of 2.3.1, where the apex coincides with the determined market price. The trade pairs saved in the list of ‘successfully executed trades’ are then connected through a series of coloured vertices on maps to visualize the trade network.

2.4. Simulation years

The model is setup using human population density, livestock population density, and croplands data for 2005. This data is adjusted using constant growth rates when simulating other years. For example, in extrapolating the human population density (CIESIN, 2016), and therefore phosphorus production potential, from 2005 to another year, the models employs the following equation:

$$P_y = P_{2005}(1 + 0.0122)^{y-2005} \quad (12)$$

Where P_y is the population density for an area in year y ; P_{2005} is the population density for an area in 2005; and 0.0122 represents the 2005 to 2015 averaged, global human population growth rate of 1.22% per year (World Bank, n.d.). The social phenomena of migration and urbanization are not considered in the process. Also, contrasting growth rates of the different areas of the world are lost, as by this definition the population in each node, everywhere, grows equally. A summary of this yearly data and its sources is presented in Table S4 of the Supplementary Materials. This is followed up by Table S5 which presents an overview of most fixed parameters.

3. Results

3.1. Phosphorus recovery potentials excluding economic dynamics

The phosphorus recovery potential can be assessed in three different ways depending on the constraints imposed. Firstly, and most simply, the sum of all global production densities (2.1.1) provide an indication of the *total potential* of recovering all excreted phosphorus, everywhere, without regard for any economic dynamics. For 2015, these are determined to amount to 3.7 [Mt a⁻¹] P for humans, and 17.39 [Mt a⁻¹] P for livestock. Recovering all urban phosphorus discharged as wastewater can thus potentially satisfy 20% of the 19.1 [Mt a⁻¹] calculated agricultural demand. Recovering all phosphorus in animal manure can potentially satisfy 90% of the total agricultural demand. The recycling of all animal manure and human excreta confirms the large potential of recovery to substitute phosphorus fertilizers (Bouwman et al., 2009).

Unfortunately, however, it is not feasible to recover every ounce of phosphorus excreted or to fertilize every crop patch everywhere. More realistically, recovery will be efficient in areas of high population or livestock density, while fertilisation will benefit mainly areas of intensive agriculture. By imposing a size constraint on the nodes determined from the phosphorus density maps (section 2.1.3), a more realistic assessment of the contribution of recovered products to the global P demand can be made. For 2015, the phosphorus recovery potential from high density urban sites is estimated at 1.73 [Mt a⁻¹] P, which is 48% of the original 3.7 [Mt a⁻¹]. This percentage approaches the percentage of global population urbanized, 54% (World Bank, 2018), partially confirming that this constraint indeed excludes smaller rural areas, thereby limiting recovery to the high potential, urban sites only. Recovery from livestock is reduced from 17.39 to 8.8 [Mt a⁻¹] P, now accounting only for the most intensive animal husbandry sites in the world. With a demand of 16.81 [Mt a⁻¹] P from areas of intense agriculture, approximately 10% and 52% of the demand may be satisfied by recovery from urban wastewater and livestock, respectively (62% total), while still excluding any economic dimension.

The results of these optimistic recovery potentials are summarized, per continent, in Table 1. Some continents (i.e. South and North America) show significant disproportionalities in recoverable P from waste vs. phosphorus demand for crop production. (Virtual) phosphorus trade (e.g. soy bean products) can play an important role in determining these continental budget surpluses and deficits. In the end, however, the total phosphorus budget is only slightly off balance at 109% of the total production potential to demand for the density maps. This global 9% surplus suggests that there is an inherent *overestimation* of the phosphorus production (excretion) or *underestimation* of the demand (agriculture), or that some degree of soil nutrient mining by the crops is considered in the phosphorus requirement values presented in 'Fertilizers and Their Use' (FAO & IFA, 2000). Another explanation for this disproportionality is that non-agricultural consumers of phosphorus (e.g. medicine and detergents industries) are not considered as actors even though their consumed products are included in the wastewater discharge figures. The 9% overestimation could suggest a 9% share of these actors in the global phosphorus market. Including these actors would raise the demand and likely close the deficit.

3.3. Phosphorus recovery potentials including economic dynamics

Phosphorus quantities, prices, and the distances between the nodes are used to determine market prices at which phosphorus trade may occur globally. This is used to assess the more realistic, spatially dependant and economically constrained, phosphorus recovery potential. The production costs for phosphorus for the different actors are summarized in fig 3.

In a 2015 market where sustainable products compete with rock-based fertilizers (scenario 2), the model determines that approximately 0.15 [Mt] can be economically recovered, thereby satisfying 0.8% of total agricultural demand. In a market without rock-based fertilizer products, approximately 7.92 [Mt] can be economically recovered, satisfying 41% of the total agricultural demand. Due to differences in total supply and demand amongst the scenarios, both the market prices and total quantities traded will vary. Optimal trade in a near-future scenario of recovered and mined phosphorus products (scenario 2) occurs at a market price of 2,039 [\$ t⁻¹]. For a market of only recovered products (scenario 3), where there exists a strong deficit in phosphorus supply for agriculture, optimal trade occurs at much higher prices of 5,700 [\$ t⁻¹]. The model price and trade determinations for 2015 and other years are summarized in Table 2.

Associated with these quantities and prices are network trade maps. Although easily created for every year and for every price, only those relate to scenario two and three, for 2015, are presented in figures 5 and 6, respectively. The high recovery potential and close proximity of recovery nodes to agricultural demand nodes makes phosphorus recovery in Asia particularly competitive in both scenarios. The struvite scaling maintenance and sludge handling cost savings in developed areas (with BNR equipped WWTPs) results in ~~potential~~ competitive trade in Europe and the United States also. Because compost pellets, which due to their low P-density are far more expensive per tonne phosphorus than other products, the market prices for phosphorus are driven upwards in the 3rd scenario. This is a result of the model economics, where the different commodities (struvite vs. compost pellets) are treated as products acting on the same market - discriminated only based on their phosphorus content. When struvite producers observe consumers buying the relatively expensive (dollars per amount P) compost pellets due to the depletion of struvite suppliers, the initial struvite sellers will adjust their prices upwards forcing consumers to pay more for the same amount ~~for~~ of P simply because they can (they are profit driven). The inverse would be true also if agricultural consumers observed cheaper trades occurring among other actors. They would then demand lower prices from their producer or switch producer all together, resulting in lower market prices.

3.3.3. Model Validation and Sensitivity Assessment

Due to the hypothetical nature of global phosphorus recovery from wastewater, model performance can only be assessed by comparing the quantities and prices produced by the model for the 'mines only scenario' (scenario 1) with observed and recorded DAP price statistics. These prices, along with the results for the other two (more hypothetical) modelling scenarios, are presented in fig. 4. The graph of figure 4 shows that the model is able to reproduce DAP price trends ~~with relative accuracy~~. Only the price estimates of 2009, 2012 and 2013 are slightly higher than those recorded, both of which may be a delayed response to price stabilization after the price inflations of 2008 and 2011, respectively.

Model sensitivity is determined through the *one-at-a-time* (OAT) method. Following this technique, the value of one parameter is adjusted and the model is re-run to evaluate how significant the change in model output is as a result of the parameter change. For each of the individual 31 parameters or input data, we assess how sensitive the model is to a -50% to +50% change in original value. We assess the impact of this change for the model determined i) total phosphorus trade, ii) sustainable phosphorus trade, and iii) optimal market price, for a 2015 market of both recovered and mined products.

The sensitivity analysis shows variable sensitivity to changes in parameter (Table S7 and Figure S3 of Supplementary Materials). Provided that each different year has different input data, these sensitivities will likely vary also depending on the simulation year. For example, in a year where rock phosphate exploitation costs are high (e.g. 2008), the market may be much more dependent on recovered products than in other years. The total phosphorus trade will then likely fluctuate much more with the price of, e.g., magnesium chloride, than determined in the sensitivity analysis conducted for 2005.

What is remarkable is the sensitivity of the optimal price and quantity of sustainable trade with variations in transport parameters. An increase in ship velocity or carry load, for example, results in significantly lower market prices. Also, it increases the competitive position of geographically concentrated, artificial fertilizers, and therefore reduces the quantity of recovered products traded. Oppositely, the model results show to be insensitive to changes in recovery parameters, suggesting that at global scale the transportation cost of products have a much greater weight in determining the feasibility for trade than the production costs do. Also the relatively small contribution of urban recovered P (maximum 10% of global P-demand, section 3.1) may be a reason why the global P market prices are so insensitive to small changes in the parameters that determine the P-recovery cost.

4 Discussion

Many generalizations, assumptions, and simplifications have been made in this study. The lack of consideration for immediate on-site recycling, trans-Atlantic movement, assumptions of free trade, and other economic simplifications, are among those possibly contributing to errors in market price determination and patterns of trade flow. Furthermore, external costs such as those associated to environmental impacts (e.g. CO₂ emissions and energy requirements) for the various fertilizer production/recovery chains has been neglected to keep the economic analysis simple. Including these costs would likely significantly improve the favourability of phosphorus recovery from wastewater considering the reduced transportation distances and relatively CO₂ neutral recovery technologies (Molinos-Senante et al., 2011). Many of the assumptions of this study are partially justified by the global and explorative nature of this investigation on potentials. An overview of all significant assumptions and their possible implications are presented in Table 3.

The lack of studies on the global *economic potentials* for recovery and phosphorus trade patterns at subnational resolution, inhibits comparison of this study's primary results. The model results on *total potentials* and struvite pricing, however, are well aligned to those of other studies. Above we determined that in 2015, 3.7 [Mt a⁻¹] of phosphorus is discharged into wastewater, satisfying 20% of the reported 18.52 [Mt a⁻¹] agricultural demand (Heffer and Prud'homme, 2016), and 19% of determined 19.51 [Mt a⁻¹] agricultural demand.

Smil (2000) found 3 [Mt a⁻¹] potentially recoverable, which would account for 20-25% of the global agricultural demand. Extrapolating Smil's (2000) figure proportionally with a population growth of 1.22% per year would result in a potential urban production of 3.6 [Mt a⁻¹] in 2015, a less than 3% difference from the model determined potential.

Mihelcic et al. (2011), through a study on diets and phosphorus excretion, concludes that the phosphorus excretion rates per individual can vary as much as from 0.18 P [kg a⁻¹] in the Democratic People's Republic of Congo, to 0.73 P [kg a⁻¹] in Israel. This confirms that our 'Western' approximation for phosphorus excretion of 0.77 P [kg a⁻¹] is on the

global high end. For 2009, nonetheless, Mihelcic et al. (2011) determines that 3.4 P [Mt a⁻¹] of human waste produced could account for 22% of the 15 [Mt a⁻¹] of global phosphorus demand.

Van Drecht et al., (2009) considers variability in access to sanitation, livings standards, and other population relevant variables to determine a phosphorus discharge of 1.3-3.1 [Mt a⁻¹] to wastewater systems in the period 2000 to 2050.

Similarly, **Morée et al., (2013)** determines a P discharge of 0.2 to 1.0 [Mt a⁻¹] from urban wastewater over the period 1950 to 2000. These are lower estimates than the ones produced in this study, suggesting that our assumptions of i) everyone being connected to some form of sanitary infrastructure and ii) everyone discharging phosphorus according to western throughput figures, is unrealistically optimistic even for near future scenarios. However, Morée et al., (2013) also determines that, over that same period, 0.08 [Mt a⁻¹] of P was recycled back to agriculture, which is lower, but in range to our estimated economic recovery potential of 0.13 [Mt a⁻¹] for 2015.

Bouwman et al., (2013), also using slaughter weights, determined that 17 [Mt a⁻¹] of P is produced by livestock in 2000, matching closely the model determined 17.11 [Mt a⁻¹]. **Bouwman et al. (2011)** notes the potential for recovery in industrialised countries, which is in line with the general trade patterns presented in fig. 5.

Koppelaar and Weikard (2013) estimate 4.2 [Mt a⁻¹] P domestic discharge, which is 25% of their 16.7 P [Mt a⁻¹] agricultural demand. They also estimated total domestic animal manure production of 28.3 [Mt a⁻¹] P. This is distinctly higher than this study's 17.11 [Mt/year] P from livestock. This is likely in part due to the fact that they account for a much larger variety of livestock types than the cattle, poultry, and swine considered in this study.

The model estimates struvite *production costs* ranging from 0 to 670 [\$ t⁻¹] variable with the nature and location of the recovery site (fig. 3; converted prices²). Phosphorus *market prices* for a market that also offers rock phosphate products range from 273 to 391 [\$ t⁻¹] over the different years (fig. 4; converted prices). The supply deficit resulting from a scenario without rock phosphate products, drives these sale prices upwards to a range of 570 to 955 [\$ t⁻¹]. These costs and prices, like the recovery potentials, are difficult to compare provided no found study has evaluated the prices for economic recovery potential at global scale. Instead, there exist case-studies on the feasibility for phosphorus recovery at specific sites.

Ueno and Fujii (2001) observed that struvite obtained from wastewater in Japan is sold to fertilizer companies at rates of 300 [\$ t⁻¹]. A market study by **Münch and Barr (2001)** revealed that struvite can be sold in Australia for between 220 and 370 [\$ t⁻¹]. **Shu et al. (2006)**, however, estimated that the market price of struvite is around 550 [\$ t⁻¹]. Based on fertilizer market estimation, **Dockhorn (2009)** estimated far higher prices than those mentioned before, and values recovered struvite at 900 [\$ t⁻¹]. Dockhorn's high recovered product prices are approached in the model by the 2015 struvite market price, 955 [\$ t⁻¹], for a market scenario with no rock phosphate competition and a severe P supply deficit. It appears that the model determined price range for global struvite production covers the spectrum of different production costs as determined in various different other studies.

In this investigation we consider struvite precipitation as the primary means to recover phosphorus from wastewater. In reality, there are many other recovery technologies that also offer high recovery rates (Cordell et al., 2011; Egle et al., 2016). It would

² Prices in this study are mostly represented as U.S. dollars per tonne phosphorus. For comparison purposes, we can convert these to DAP prices through multiplication with DAP's P density (24%), or to struvite prices through multiplication with struvite's P density (14%).

be an interesting follow-up study to adapt and run the model for different phosphorus recovery technologies and visualize changes in trade patterns for different phosphorus products.

5 Conclusion

5

Despite the simplifying assumptions, the model developed in this study generates realistic trade networks for different phosphorus supply scenarios, for different prices, at a subnational resolution. However, the credibility of model outputs is only supported by an accurate simulation of DAP prices because data and/or other studies on the purely hypothetical nature of global trade in sustainably recovered P, are lacking. Nevertheless, the model sets a basis that provides some general indication of the spatially dependant recovery feasibility of phosphorus from wastewater. It is furthermore able to provide this indication for potentially any recovery technology for which there exists adequate economic data.

10

Model results reveal a relatively minor potential of economically profitable, struvite fertilizer production from wastewater. This recovery thus appears to offer a limited contribution to resolving the global phosphorus issues of the 21st century. Nevertheless, at a more local scale (namely in the highly populated cities of developed countries), this recovery offers wastewater treatment plants the opportunity to contribute to creating sustainable communities and protecting the environment, while reducing their own operational cost.

15

Although recognizing that there is no single solution to solving phosphorus pollution and insecurity issues (Cordell and White, 2011), recovering phosphorus from all waste sources may come to provide a greater contribution as populations grow and urbanize, technologies develop, and the economically extractable phosphorus reserves deplete. For this reason, it is essential to determine how the widespread implementation of recovery technologies impacts phosphorus market dynamics. Only then can we stimulate and regulate its recovery in such a way that maximum benefits are achieved for both the environment and the urban community, as well as the livestock and agricultural sectors.

20

References

- Arrow, K. J. and Debreu, G.: Existence of an Equilibrium for a Competitive Economy, *Econometrica*, 22(3), 265–290 [online] Available from: <http://links.jstor.org/sici?sici=0012-9682%28195407%2922%3A3%3C265%3AE0AEFA%3E2.0.CO%3B2-B> (Accessed 11 October 2017), 1954.
- 30 Bouwman, A. F., Beusen, A. H. W. and Billen, G.: Human alteration of the global nitrogen and phosphorus soil balances for the period 1970–2050, *Global Biogeochem. Cycles*, 23(4), n/a–n/a, doi:10.1029/2009GB003576, 2009.
- Bouwman, L., Goldewijk, K. K., Van, K. W., Hoek, D., Beusen, A. H. W., Van Vuuren, D. P., Willems, J., Rufino, M. C. and Stehfest, E.: Exploring global changes in nitrogen and phosphorus cycles in agriculture induced by livestock production over the 1900–2050 period, *Proc Natl Acad Sci*, 110(10), 20882–20887, doi:10.1073/pnas.1206191109, 2011.
- 35 Bouwman, L., Goldewijk, K. K., Van Der Hoek, K. W., Beusen, A. H. W., Van Vuuren, D. P., Willems, J., Rufino, M. C. and Stehfest, E.: Exploring global changes in nitrogen and phosphorus cycles in agriculture induced by livestock production over the 1900–2050 period., *Proc. Natl. Acad. Sci. U. S. A.*, 110(52), 20882–7, doi:10.1073/pnas.1012878108, 2013.
- Center for International Earth Science Information Network - CIESIN - Columbia University: Gridded Population of the World, Version 4 (GPWv4): Population Density, [online] Available from: <http://dx.doi.org/10.7927/H4NP22DQ> (Accessed 1 May 2016), 2016.
- 40 Cordell, D. and White, S.: Peak phosphorus: Clarifying the key issues of a vigorous debate about long-term phosphorus security, *Sustainability*, 3(10), 2027–2049, doi:10.3390/su3102027, 2011.
- Cordell, D., Schmid-Neset, T., White, S. and Jan-Olof, D.: Preferred Future Phosphorus Scenarios: A Framework For Meeting Long-Term Phosphorus Needs for Global Food Demand, *Int. Conf. Nutr. Recover. from Wastewater Streams*, 2009.
- 45 Cordell, D., Rosemarin, A., Schröder, J. J. and Smit, A. L.: Towards global phosphorus security: A systems framework for phosphorus recovery and reuse options, *Chemosphere*, 84(6), 747–758, doi:10.1016/J.CHEMOSPHERE.2011.02.032, 2011.

- Cornel, P. and Schaum, C.: Phosphorus recovery from wastewater: needs, technologies and costs, *Water Sci. Technol.*, 59(6), 1069, doi:10.2166/wst.2009.045, 2009.
- CRC: Handbook of Chemistry and Physics, 86th ed., Taylor and Francis., 2005.
- Dockhorn, T.: About the Economy of Phosphorus Recovery, *Int. Conf. Nutr. Recover. from Wastewater Streams*, 2009.
- 5 Van Drecht, G., Bouwman, A. F., Harrison, J. and Knoop, J. M.: Global nitrogen and phosphate in urban wastewater for the period 1970 to 2050, *Global Biogeochem. Cycles*, 23(4), n/a-n/a, doi:10.1029/2009GB003458, 2009.
- Egle, L., Rechberger, H., Krampe, J. and Zessner, M.: Science of the Total Environment Phosphorus recovery from municipal wastewater: An integrated comparative technological, environmental and economic assessment of P recovery technologies, *Sci. Total Environ.*, 571, 522–542, doi:10.1016/j.scitotenv.2016.07.019, 2016.
- 10 Egle, L. A.: Doctoral Thesis Phosphorus recovery from municipal wastewater Dissertation Phosphorrückgewinnung aus kommunalem Abwasser., n.d.
- EPA (United States Environmental Protection Agency): Nutrient Control Design Manual, The Cadmus Group, Inc, Washington, D.C., 2010.
- FAO: Crop Water Information, [online] Available from: www.FAO.org, 2015.
- 15 FAO (Food and Agriculture Organisation) and IFA (International Fertilizer Industry Association): Fertilizers and Their Use: A pocket guide for extension officers, Rome, Italy., 2000.
- FAOSTAT: Animal Live Weights, [online] Available from: <http://www.fao.org/home/en/> (Accessed 1 May 2018), 2018.
- Gilmour, D., Blackwood, D., Comber, S. and Thornell, A.: Identifying human waste contribution of phosphorus loads to domestic wastewater, *Proceedings 11th Int. Conf. Urban Drainage*, Held Edinburgh, Scotland, UK, 1–10, 2008.
- 20 Heffer, P.: Assessment of Fertilizer Use by Crop at the Global Level., 2009.
- Jaffer, Y., Clark, T. ., Pearce, P. and Parsons, S. .: Potential phosphorus recovery by struvite formation, *Water Res.*, 36(7), 1834–1842, doi:10.1016/S0043-1354(01)00391-8, 2002.
- Koppelaar, R. H. E. M. and Weikard, H. P.: Assessing phosphate rock depletion and phosphorus recycling options, *Glob. Environ. Chang.*, 23(6), 1454–1466, doi:10.1016/j.gloenvcha.2013.09.002, 2013.
- 25 Masayuki Hara: Fertilizer Pellets Made from Composted Livestock Manure, *Mie Prefect. Sci. Technol. Promot. Cent.* [online] Available from: <http://www.ftc.agnet.org/library.php?func=view&id=20110801154610> (Accessed 12 January 2018), 2001.
- Mihelcic, J. R., Fry, L. M. and Shaw, R.: Global potential of phosphorus recovery from human urine and feces., *Chemosphere*, 84(6), 832–9, doi:10.1016/j.chemosphere.2011.02.046, 2011.
- Molinos-Senante, M., Hernández-Sancho, F., Sala-Garrido, R. and Garrido-Baserba, M.: Economic Feasibility Study for Phosphorus Recovery Processes, *Ambio*, 40(4), 408–416, doi:10.1007/s13280-010-0101-9, 2011.
- 30 Monfreda, C., Ramankutty, N. and Foley, J. A.: Farming the planet. Part 2: Geographic distribution of crop areas, yields, physiological types, and net primary production in the year 2000, *Global Biogeochem. Cycles*, 22, doi:10.1029/2007GB002947, 2008.
- Morée, A. L., Beusen, A. H. W., Bouwman, A. F. and Willems, W. J.: Exploring global nitrogen and phosphorus flows in urban wastes during the twentieth century, *Global Biogeochem. Cycles*, 27(3), 836–846, doi:10.1002/gbc.20072, 2013.
- 35 Münch, E. V. and Barr, K.: Controlled struvite crystallization for removing phosphorus from anaerobic digester side stream, *Water Res.*, 35(1), 151–159, 2001.
- NASA: MODIS Evapotranspiration, [online] Available from: <https://modis.gsfc.nasa.gov/data/dataproduct/mod16.php> (Accessed 1 May 2018), 2005.
- 40 Pande, S. and Savenije, H. H. G.: A sociohydrological model for smallholder farmers in Maharashtra, India, , 7503–7512, doi:10.1002/2016WR018977, Received, 2016.
- Quantum GIS Development Team: Quantum GIS Geographic Information System 2.14, Open Source Geospatial Found. Proj. [online] Available from: <http://qgis.osgeo.org>, 2017.
- Robinson, T. P., William Wint, G. R., Conchedda, G., van Boeckel, T. P., Ercoli, V., Palamara, E., Cinardi, G., D’Aielli, L.,
- 45 Hay, S. I. and Gilbert, M.: Mapping the global distribution of livestock, *PLoS One*, 9(5), doi:10.1371/journal.pone.0096084, 2014.
- Sheldrick, W., Syers, J. K. and Lingard, J.: Contribution of livestock excreta to nutrient balances, *Nutr. Cycl. Agroecosystems*, 66, 119–131 [online] Available from: <https://link.springer.com/content/pdf/10.1023%2FA%3A1023944131188.pdf> (Accessed 1 May 2018), 2003.
- 50 Shu, L., Schneider, P., Jegatheesan, V. and Johnson, J.: An economic evaluation of phosphorus recovery as struvite from digester supernatant, *Bioresour. Technol.*, 97(17), 2211–2216, doi:10.1016/j.biortech.2005.11.005, 2006.
- Sivapalan, M., Savenije, H. H. G. and Blöschl, G.: Socio-hydrology: A new science of people and water, *Hydrol. Process.*, 26(8), 1270–1276, doi:10.1002/hyp.8426, 2012.
- Smil, V.: Phosphorus in the environment: Natural flows and human interferences, *Annu. Rev. Energy Environ.*, 25, 53–88, doi:10.1146/annurev.energy.25.1.53, 2000.
- 55 Steduto, P., Hsiao C., T., Fereres, E. and Raes, D.: FAO Irrigation and Drainage Paper: Crop Yield Response to Water, Food and Agriculture Organization of the United Nations (FAO), Rome., 2012.
- Steen, I.: Management of a non-renewable resource, Phosphorus and potassium, (217), 25–31 [online] Available from: <http://cat.inist.fr/?aMode=afficheN&cpsid=2380873> (Accessed 24 March 2016), 1998.

- Ueno, Y. and Fujii, M.: Three Years Experience of Operating and Selling Recovered Struvite from Full-Scale Plant, *Environ. Technol.*, 22(11), 1373–1381, doi:10.1080/09593332208618196, 2001.
- UNIDO (United Nations Industrial Development Organisation) and IFDC (International Fertiliser Development Centre): *Fertilizer Manual*, Springer Netherlands. [online] Available from: <https://books.google.nl/books?id=qPkoOU4BvEsC>, 1998.
- 5 USGS (United States Geological Survey): *World phosphate mines, deposits, and occurrences*, [online] Available from: <https://mrdata.usgs.gov/phosphate/>, 2002.
- USGS (United States Geological Survey): *Mineral Commodity Summary Phosphate Rock*, Washington, D.C. [online] Available from: https://minerals.usgs.gov/minerals/pubs/commodity/phosphate_rock/phospmcs06.pdf, 2007.
- USGS (United States Geological Survey): *Mineral Commodity Summary Phosphate Rock*, Washington, D.C. [online] Available from: https://minerals.usgs.gov/minerals/pubs/commodity/phosphate_rock/phospmcs06.pdf, 2010.
- 10 Vinnerås, B.: Faecal separation and urine diversion for nutrient management of household biodegradable waste and wastewater, *J. Biosci.*, 5(2), 29–50, 2001.
- Van Vuuren, D. P., Bouwman, A. F. and Beusen, A. H. W.: Phosphorus demand for the 1970–2100 period: A scenario analysis of resource depletion, *Glob. Environ. Chang.*, 20(3), 428–439, doi:10.1016/j.gloenvcha.2010.04.004, 2010.
- 15 WHO/UNICEF JMP (Joint Monitoring Programme): *Percent of Urban Population with Access to Improved Sanitation Facilities*, 2015.
- Wilsenach, J. A. A., Schuurbiers, C. A. H. A. H. and van Loosdrecht, M. C. M. C. M.: Phosphate and potassium recovery from source separated urine through struvite precipitation., *Water Res.*, 41(2), 458–66, doi:10.1016/j.watres.2006.10.014, 2007.
- World Bank: *Urban population (% of total)*, 2018 [online] Available from: <https://data.worldbank.org/indicator/SP.URB.TOTL.IN.ZS> (Accessed 2 May 2018), 2018.
- 20 World Bank: *Population growth (annual %)* | Data, [online] Available from: <https://data.worldbank.org/indicator/SP.POP.GROW> (Accessed 3 January 2018), n.d.




Figure 1. Estimated global phosphorus reserve distribution. The vast majority (73%) of estimated natural reserves lie in Moroccan and West-Saharan territories (USGS, 2017).

5




Figure 2. Nominal Phosphate rock and DAP price trends (IndexMundi, 2017). The rock phosphate and DAP price trend can be characterized as gradually increasing and vulnerable to market dynamics (i.e. the 2007 global economic recession).




Figure 3. Minimum production costs per ton of recycled P for supply nodes. Urb1-Str shows the prices for struvite from WWTP in countries with highly developed sanitation systems (BNR), where the struvite is offered for free (excl. transport costs) because of the maintenance and sludge handling savings by controlled struvite precipitation; Urb2-Str, from nations with slightly less developed sanitation systems; Urb3-Str, of source-separated urine collection and struvite precipitation in nations with underdeveloped sanitation systems; Urb3-Pellets, the price per tonne P (0.1%) in urban compost pellets; and Livestock shows the price per tonne P in livestock pellets (0.1% P).




Figure 4. Plausible phosphorus price ranges at which trade can occur at global scale for different years: observed prices (grey) versus modelled scenarios (Sc. 1, pink; Sc. 2, blue; Sc. 3, green). The ends of the whiskers on the grey boxes for the observed prices represent the maximum and minimum prices for DAP that year, the box itself shows the upper and lower quartiles for that data and the orange line indicates the median. For the modelled scenario the whisker-ends show the maximum and minimum prices as approximated by the model, and the box indicates the most likely price range, and the orange line marks the most probable price given supply-demand ratio's. Noticeable is that the model price determinations for the current mines only scenario are in close proximity with observed price ranges. Scenario 2 and 3 are hypothetical scenarios and therefore do not have a realistic counterpart dataset that can be used for comparison, cannot be compared. Nevertheless, they show predictable and realistic behaviour.




Figure 5. Phosphorus trade network for trade in both conventional and recycled P (scenario 2) but showing only sustainable trade flows. Optimal trade occurs at a phosphorus market price of 2,039 [\$ t⁻¹] with 16.81 [Mt] being traded in total, of which 0.15 [Mt] is traded sustainably (0.8% of total demand).




Figure 6. Phosphorus trade network for trade in recycled P only (scenario 3) at determined phosphorus market prices of 5,700 [\$ t⁻¹] with 7.92 [Mt] being traded. All of this trade is sustainable but is only able to accommodate 41% of demand.

Continental Phosphorus Budgets

	Livestock Production		Human Production		Agricultural Demand	
	Total [Mt P]	Major Nodes [Mt P]	Total [Mt P]	Major Nodes [Mt P]	Total [Mt P]	Major Nodes [Mt P]
Asia	5.85	-	2.29	-	9.49	-
North America	3.10	-	0.31	-	3.30	-
Europe	2.39	-	0.40	-	1.79	-
Africa	1.72	-	0.53	-	2.51	-
South America	3.89	-	0.22	-	2.17	-
Oceania	0.05	-	0.00	-	0.00	-
Australia	0.28	-	0.01	-	0.24	-
World	17.11	10.47	3.75	1.73	19.52	16.81

Table 1. Approximate continental phosphorus budgets associated to maps. The ‘Total’ values represent an estimation of recovering all phosphorus on the continent, while the ‘Major Nodes’ values represents that which can be recovered at sites of high potential only.

Model Trade and Price Determinations

		Scenario 1	Scenario 2	Scenario 3	Units
2005	Maximum Traded	15.01	15.01	5.59	[mt]
	Sustainably Traded	0	0.12	5.59	[mt]
	% of total demand met	100	100	37	[-]
	% of total demand met sustainably	0	0.80	37	[-]
	Optimal Price	1276	1189	4000	[\$ t ⁻¹] P
	Min price for 90% of max. trade	1000	1000	3700	[\$ t ⁻¹] P
	Max price for 90% of maximum trade	21000	2100	4000	[\$ t ⁻¹] P
2006	Maximum Traded	15.46	15.46	5.89	[mt]
	Sustainably Traded	0	0.12	5.89	[mt]
	% of total demand met	100	100	38	[-]
	% of total demand met sustainably	0	0.78	38	[-]
	Optimal Price	1559	1460	4300	[\$ t ⁻¹] P
	Min price for 90% of max. trade	1100	1100	3800	[\$ t ⁻¹] P
	Max price for 90% of maximum trade	2300	2300	4300	[\$ t ⁻¹] P
2007	Maximum Traded	15.91	15.91	6.59	[mt]
	Sustainably Traded	0	0.12	6.59	[mt]
	% of total demand met	100	100	41	[-]
	% of total demand met sustainably	0	0.75	41	[-]
	Optimal Price	1780	1674	5500	[\$ t ⁻¹] P
	Min price for 90% of max. trade	1500	1500	4000	[\$ t ⁻¹] P
	Max price for 90% of maximum trade	2900	2900	5500	[\$ t ⁻¹] P
2008	Maximum Traded	10.03	10.03	7.12	[mt]
	Sustainably Traded	0	6.91	7.12	[mt]
	% of total demand met	62.69	62.69	45	[-]
	% of total demand met sustainably	0	43.19	45	[-]
	Optimal Price	4551	4445	6900	[\$ t ⁻¹] P
	Min price for 90% of max. trade	4700	3500	3500	[\$ t ⁻¹] P
	Max price for 90% of maximum trade	6900	6900	6900	[\$ t ⁻¹] P
2009	Maximum Traded	16.81	16.81	6.82	[mt]
	Sustainably Traded	0	0.16	6.82	[mt]
	% of total demand met	100.00	100.00	41	[-]
	% of total demand met sustainably	0	0.95	41	[-]
	Optimal Price	2355	2299	5500	[\$ t ⁻¹] P
	Min price for 90% of max. trade	2100	2100	4000	[\$ t ⁻¹] P
	Max price for 90% of maximum trade	2900	2900	5500	[\$ t ⁻¹] P
2010	Maximum Traded	17.26	17.26	7.13	[mt]
	Sustainably Traded	0	0.16	7.12	[mt]
	% of total demand met	100.00	100.00	41	[-]
	% of total demand met sustainably	0	0.93	41	[-]
	Optimal Price	2534	2444	6618	[\$ t ⁻¹] P
	Min price for 90% of max. trade	2200	2200	3400	[\$ t ⁻¹] P

	Max price for 90% of maximum trade	3400	3400	6600	[\$ t ⁻¹] P
2011	Maximum Traded	17.71	17.71	8.16	[mt]
	Sustainably Traded	0	0.6	8.16	[mt]
	% of total demand met	100.00	100.00	46	[-]
	% of total demand met sustainably	0	3.39	46	[-]
	Optimal Price	3481	3374	8000	[\$ t ⁻¹] P
	Min price for 90% of max. trade	3100	3100	3600	[\$ t ⁻¹] P
	Max price for 90% of maximum trade	4200	4200	8000	[\$ t ⁻¹] P
2012	Maximum Traded	18.14	18.16	7.9	[mt]
	Sustainably Traded	0	0.59	7.12	[mt]
	% of total demand met	99.89	100.00	44	[-]
	% of total demand met sustainably	0	3.25	44	[-]
	Optimal Price	3484	3348	7596	[\$ t ⁻¹] P
	Min price for 90% of max. trade	3100	3000	3500	[\$ t ⁻¹] P
	Max price for 90% of maximum trade	3900	3900	7600	[\$ t ⁻¹] P
2013	Maximum Traded	18.61	18.61	8.01	[mt]
	Sustainably Traded	0	0.2	8.01	[mt]
	% of total demand met	100.00	100.00	43	[-]
	% of total demand met sustainably	0	1.07	43	[-]
	Optimal Price	2943	2840	7389	[\$ t ⁻¹] P
	Min price for 90% of max. trade	2500	2500	3500	[\$ t ⁻¹] P
	Max price for 90% of maximum trade	3800	3800	7400	[\$ t ⁻¹] P
2014	Maximum Traded	19.06	19.06	8	[mt]
	Sustainably Traded	0	0.15	8	[mt]
	% of total demand met	100.00	100.00	42	[-]
	% of total demand met sustainably	0	0.79	42	[-]
	Optimal Price	2469	2365	7000	[\$ t ⁻¹] P
	Min price for 90% of max. trade	2000	2000	3500	[\$ t ⁻¹] P
	Max price for 90% of maximum trade	3700	3700	7000	[\$ t ⁻¹] P
2015	Maximum Traded	19.51	19.51	7.92	[mt]
	Sustainably Traded	0	0.15	7.92	[mt]
	% of total demand met	100.00	100.00	41	[-]
	% of total demand met sustainably	0	0.77	41	[-]
	Optimal Price	2155	2039	5700	[\$ t ⁻¹] P
	Min price for 90% of max. trade	1900	1900	4000	[\$ t ⁻¹] P
	Max price for 90% of maximum trade	3000	3000	5700	[\$ t ⁻¹] P

Table 2. Price ranges and amounts traded per scenario for all years.

Overview of Assumptions

Methods	Assumption	In Reality	Implication
Production and demand estimates.	Crop phosphorus requirement only varies with crop type and water-constrained yield.	<ul style="list-style-type: none"> - A particular crop yield may be constrained by other factors than water availability (e.g. soil acidity levels, micronutrient levels, management practices, etc.), also influencing the actual phosphorus demand. - Crop phosphorus requirements depend also on soil dynamics processes determining leaching rates and sorption. 	The phosphorus demand quantities will be spatially more variable and are likely overestimated.
	Domestic wastewater composition is globally homogeneous, and its quantity only varies with population density	<ul style="list-style-type: none"> - Wastewater composition and amount vary amongst different populations. Both quantity and quality factors of wastewater depend on the amount and nature of connected industries and lifestyle characteristics of the people connected (e.g. diets and detergents use), the regional climate, as well as whether or not the sewerage network is a combined or separate system, etc. 	Recovery efficiencies will likely be lower, provided the current assumption of western lifestyles, globally.
	We can approximate the type of wastewater treatment practiced in a country based on data that shows the percent of urban population that have access to sanitary facilities.	<ul style="list-style-type: none"> - The type of wastewater treatment at a node varies with many different socio-economic (and natural) parameters that it can hardly be approximated for using a single dataset. 	Even in the near future, few people in developing countries will have access to sanitation resulting in an even lower recovery efficiency of phosphorus from developing regions than is currently predicted.
	Phosphorus recovery efficiency is determined solely by the recovery technology.	<ul style="list-style-type: none"> - Phosphorus recovery efficiency varies not just per technology, but also with the wastewater composition. 	Node specific phosphorus production potentials can potentially be higher or lower than is currently determined.
	Phosphorus throughput per individual is globally homogeneous	<ul style="list-style-type: none"> - Phosphorus excretion rates vary enormously depending on age, diet, and gender of the individual. - Phosphorus discharge relates also to population lifestyles which determine the diet (as described above) but also detergent use. 	Node specific phosphorus production potential is lower in developing countries than is currently determined.
	Assume container ship as sea transport mode	<ul style="list-style-type: none"> - Bulk trade occurs in bulk carriers. Depending on how finished the product is (e.g. bagged and sealed), it may be transported by container also, 	Sea transport may be significantly cheaper than is current the case.
	Fertilizer maximum bid price depends on crop	<ul style="list-style-type: none"> - Farmers growing the same crop, requiring the same amount of fertilizer, may have different maximum prices for P fertilizers depending on other factors (e.g. experience, subsidies, crop quality and client for which the crop is grown, hence different crop values) 	Recovery feasibility will be higher or lower depending on the profitability and experience in agriculture in an area.
Modelling trade	All phosphorus goes to the international fertilizer market.	<ul style="list-style-type: none"> - Most phosphorus is recycled locally, applied on nearby agricultural soils 	Phosphorus recovery potentials are underestimated given the disregard for immediate local use.

	<p>directly as manure or as treated wastewater sludge (ash).</p> <ul style="list-style-type: none"> - Some is distributed amongst pharmaceutical and detergent industries 	Phosphorus production from mines is overestimated, as not all is used for fertilizer production.
Free trade	<ul style="list-style-type: none"> - Trade does not stand separate from politics. Embargo's, trade sanctions, or trade tariffs can influence the pattern and amount of global phosphorus trade. 	Trade patterns would look different provided international politics.
Non-preferential trade	<ul style="list-style-type: none"> - Trade is not purely rational. Some countries/actors may be more or less likely to trade with each other depending on historical and current relations. 	Trade patterns would look different provided international relations.
Two actor trades	<ul style="list-style-type: none"> - Phosphorus extraction from phosphate rock and its processing into artificial fertilizers may not occur at the same site, nor by the same actor. Often it involves many more parties. One for exploitation, for manufactory, for logistics, etc. 	Trade patterns are simplified, showing only the path from site of initial production to site of final demand.
Transportation distances are calculated as straight lines as opposed to following existing infrastructure.	<ul style="list-style-type: none"> - Phosphorus transport in reality follows existing shipping routes and road infrastructure. 	The model likely underestimates the transportation cost of moving phosphorus from supply to demand areas. However, it does so consistently for both recovered as well as mined products. This nevertheless reduces recovery potential estimates.
No transatlantic trade due planar projection of earth.	<ul style="list-style-type: none"> - International trade is not restricted by cartographic boundaries. 	Few implications for the simulated years as trade in phosphorus between the America's and Asia is unlikely to occur due to the greater distances and relatively balanced continental phosphorus budget of North America.
Near-Future and Far Future scenarios do not consider development of technology	<ul style="list-style-type: none"> - Existing technologies are likely to become cheaper and more efficient in the future, while new technologies may also be developed 	Recovery will rates will likely be higher in the future, thereby also reducing the price of phosphorus. However, provided the uncertainty around making technological development predictions, we decided to exclude this factor from our analysis.

Table 3. An overview of the assumptions made in this study and their possible implications on the results.

Supplementary Material

Supplementary Text


5 The global market price is determined as the price at which total quantity of phosphorus demanded is equal to the quantity supplied (i.e. the market for P is cleared; (Arrow and Debreu, 1954)). This is approximated as the point where global demand function for phosphorus intersects the global supply function (fig. A).



10 **Figure A. Conceptualization of supply-demand curves. The interception point of both curves marks an approximation for the price equilibrium.**

However, good nutrient management leads farmers not to purchase more phosphorus than the optimal amount that is required for their crops. As such, it is assumed that farmers do not over-fertilize (buy more fertilizers than they need for optimal yield) when fertilizer prices are extremely low. Therefore the phosphorus trade will be limited by maximum global demand (Q_m) for price, P. This assumption would flatten the demand curve beyond the price at which the market is saturated. The created supply-demand curves for the network will deviate further from the traditional curves when including transport cost. Transportat cost is determined based on the distance between two trading nodes and therefore cannot be determined for nodes without a trade partner (i.e. not partaking in trade). The curves created for the network are therefore limited to using data only of those nodes partaking in trade.

20 To exemplify this, consider figure A where three production nodes and three demand nodes populate a hypothetical network. The production nodes have a minimum per unit production costs ($P_1 < P_2 < P_3$) and specific quantities (Q_1 , Q_2 , and Q_3) to sell. The demand nodes have maximum prices they are willing to pay ($P_4 < P_5 = P_6$) for their quantities (Q_4 , Q_5 , and Q_6) to buy. The transportation costs ($T_{(i,n)}$) for all production nodes are lowest to demand node D4, highest to D6 and intermediate to D5 ($T_{(i,4)} < T_{(i,5)} < T_{(i,6)}$). This results in the cheapest, and therefore first, trade occurring between S1 and D4, as: $P_1 + T_{(1,4)} < P_i + T_{(i,n)}$ (line 1, Figure B).




5 **Figure B.** Network of demand and supply nodes. Arrows with numbers indicate trade order, including transportation costs.

For this first trade, quantity Q_4 is smaller than Q_1 and therefore the amount traded is equal to the node's entire demand, Q_4 , etc. The next cheapest trade follows until the entire demand is satisfied or supply is depleted. In this hypothetical network the entire demand can be satisfied by the first two production nodes. The third supply node is therefore not involved in trade.

10 Since the transport component of the price cannot be determined for this node, it is disregarded in the creation of the supply-demand curve. Executing this procedure successively for all nodes, in the example network, and plotting the prices inclusive of transportation cost with the networks cumulative quantity traded yields figure C. The cumulative quantity traded for each individually bargained price is summarized in table A.

15



20

25

Figure C. Adapted supply (green) and demand (red) curve, for six node network illustrated in Figure 12, including transportation cost component in price.

TRADE ORDER, PRICES AND QUANTITIES FOR EXAMPLE NETWORK

Trade Order	Node Pair (i,n)	Best 'hypothetical' Prices	Quantity Traded	Cumulative Quantity Traded
1	(1,4)	$P_1 + T_{(1,4)}$	Q_4	Q_4
2	(1,5)	$P_1 + T_{(1,5)}$	$Q_1 - Q_5$	$Q_4 + Q_1 - Q_5$
3	(2,5)	$P_2 + T_{(2,5)}$	$Q_2 - (Q_1 - Q_5)$	$Q_2 + Q_4 + Q_5$
4	(2,6)	$P_2 + T_{(2,6)}$	Q_6	$Q_3 + Q_4 + Q_5$

30 **Table A.** Trade order, prices and quantities for node pairs for supply curve.

Where the supply and demand curves overlap, both consumers and producers are satisfied with the amount of phosphorus traded at those prices. This range is thus indicative of the range wherein the optimal market price will lie. A more precise estimate of the optimal price is made using information on the demand or supply nodes not partaking in trade. Depending on

35 the total proportion of excess supply or demand (i.e. $\frac{Q_1+Q_2+Q_3}{Q_4+Q_5+Q_6}$) the optimal price will lie in the upper or lower half of the range. When the total supply is far greater than the total demand, then demand nodes are in power to bargain for cheaper prices. The opposite is true for when demand is greater than supply. As such, a more precise estimate of optimal market price can be made.

Optimum price determination from range

5 Unless the supply and demand curves intersect each other before the maximum total quantity that can be traded is reached, the optimum price is identified to lie between the prices at which the supply and demand curves reach maximum trade. The optimal price can be further differentiated from this (potentially large) range. Depending on the proportion of global phosphorus supply to demand, it is possible to determine whether the optimum price will lie in the upper or lower half of this range. When the total supply is far greater than the demand, then the demand nodes are in a stronger bargaining position, and
10 are able to shift the prices downward in their favor. The opposite is true when the demand is far greater than the supply, then the supply nodes can shift the prices to the upper end of the spectrum, to their favor. By this premise, we can reduce the large range to either the top half or lower half. Depending on the slope of the supply curve as it meets the satisfiable demand, a large reduction in price may lead to only a minor reduction trade. Extending therefore the upper and lower boundaries to the price that allow for 95% of maximum trade, some small possible errors in accuracy or market flexibility are more
15 appropriately accounted for. The models best-guess price lies in the middle of the final range.

Results: Production prices

The minimum production costs for the production nodes has been determined (fig. 3). The low density of phosphorus (1% phosphorus by weight) results in relatively high production and transportation costs of compost pellets per tonne of phosphorus content. The added value that is not considered in the model is that compost pellets also upgrade the soil in providing substantial amounts of nitrogen and organic matter as well. The significant, general difference in the production cost of P from recycled sources with that of mines means that transportation costs, i.e. distance between trading nodes and fuel costs, will have to play a critical role if sustainable trade in recycled P is to be feasible.

25 Most scrutiny of the results may be directed to assumptions on technological potential of recovery. The technological potential is assessed through the dataset *on urban population with access to improved sanitary facilities* – where improved facilities are defined as those designed to hygienically separate excreta from human contact. These may include anything ranging from pit latrines to flushed piped systems. A country's scoring in this dataset is then used as an indicator for the state
30 of sanitary development in that country. Struvite precipitation from digester liquor is assumed to be the technology appropriate for the highest scoring countries (>90% access). However, there is no guarantee that struvite precipitation from anaerobic digester liquor is possible for nodes in these countries, seeing as by definition *improved sanitary facilities* 100% access may indicate that 100% of the urban population has access to a pit latrine. Even if information on the wastewater treatment infrastructure was known at all locations, then the phosphorus recovery efficiency from these rich sidestreams is
35 still likely vary due to differences in phosphorus concentration of the influent. In summary, large uncertainties in the feasibility to recover arise from the generalised technological assessment carried out. Furthermore, the production and transportation cost per tonne P for compost pellets from source-separated faecal matter and livestock manure will be high due to the relatively low phosphorus content. These products are unlikely able to compete in the international phosphorus market and drive prices upwards when the demand for P is severe, e.g. when rock phosphate products are absent.

Supplementary Figures





Figure S1a. Potential phosphorus recovery from bovine, swine and poultry livestock in tonnes per square kilometre.



5 Figure S1b. Potential phosphorus recovery from human population in tonnes per square kilometre.




Figure S1c. Approximate, combined phosphorus demand for six major crops in tonnes per square kilometre.






Figure S2. Phosphorus trade network for trade in conventional P (Scenario 1) at a calculated phosphorus market price of 2,155 [\$ t⁻¹]. This results in 19.51 [Mt] P being traded in total, which meets 100% of the agricultural demand.


S3. Parameter Sensitivity Figures

5 Crop Parameters





Transport Parameters







Recovery Parameters





Market Parameters (yearly input data)





5

Figure S3. Figures reveal how -50% to +50% change in parameter influences the model determined total quantity of phosphorus traded (1st column), the total quantity of recovered products traded (2nd column), and the optimal market price (3rd column). Green marker shows original parameter value with original output. Yellow marker shows relative change in output (y-axis) vs. relative change in parameter (x-axis).

10

Supplementary Tables

S1. Phosphorus Production Estimate Data

Population	Throughput (kg P)		Site	Author
Bovine (Dairy)	25		US	Barker et al., 2001
	17.16		NL	Blokland, Luesink, & Jongeneel, 2015
	17.9		NL	CBS, 2014
	20.8		US	Weiss & Wyatt, 2004
	9.6	(stabled period)	NL	CBS, 2014
Bovine (Beef)	11.7		NL	CBS, 2014
	13.3		US	Barker et al., 2001
	5.4	(stabled period)	NL	CBS, 2014
Bovine (Unsp.)	10		Global	Sheldrick et al., 2003
Poultry (layer)	1.2		US	Barker et al., 2001
	0.17		NL	CBS, 2014
Poultry (broiler)	0.6		US	Barker et al., 2001
	0.08		NL	CBS, 2014
Poultry (Unsp.)	0.19		Global	Sheldrick et al., 2003
Swine (sow)	6.4		NL	CBS, 2014
Swine (Unsp.)	4		Global	Sheldrick et al., 2003
	4.1		US	Barker et al., 2001
	2.1		NL	CBS, 2014
Human	0.77		UK	Gilmour et al., 2008
	0.2-0.7		Global	Mihelcic et al., 2011
	0.78		-	CRC, 2005
	0.7		US	Smil, 2000

Table S1. Annual phosphorus excretion rate by species, per head.

S2. Crop Phosphorus Requirement Data

Crop	Ky [-]	Water req. [mm/harvest]	Growing Period [days]	P ₂ O ₅ Range [kg/ha]	P ₂ O ₅ Choice [kg/ha]	P [kg/ha]
Maize	1.30	500-800	80`180	36-50	50	22
Wheat	0.55	450-650	120`150	27-60	40	15
Rice ¹	1.00	450-700	90`150	26-50	35	15
Soybean	0.90	450-700	135`150	35	35	15
Sorghum	0.90	450-650	120`130	20-40, 40-60	40	15
Potato	0.90	500-700	105`145	39-80	80	35

Table S2. Crop Data (FAO, n.d.; IFDC & UNIDO, 1998)

¹ Fageria, N.K. The Use of Nutrients in Crop Plants. Google books

S3. Transportation Cost Data

<u>Sea Transport Component Cost</u>				<u>Land Transport Component Cost</u>			
Constant		Value	Source	Constant		Value	Source
E_W	[t d ⁻¹]	93.1	(Počuča, 2006)	E_L	[L km ⁻¹]	0.53	(Nylund and Erkkilä,
W_W	[t]	2,777	(Počuča, 2006)	W_L	[t]	60	2005)
\bar{V}_W	[km d ⁻¹]	924	(Počuča, 2006)	\bar{V}_L	[km h ⁻¹]	80	-
C_F	[\$ d ⁻¹]	9,989	(Počuča, 2006)	L_c	[\$ h ⁻¹]	15	<u>-*</u>
				C_d	[\$ km ⁻¹]	0.5	

Table S3. Constants for transport cost determination equation for (eq. 6)

* 15 [\$ /h] value is an informed estimate of trucking labour wage. In reality, trucking labour wages will vary globally. The model could not account for spatial diversity in wages provided the absence of a global data set on trucking wages per country or even per region. Instead, the 15 [\$ /h] was determined taking into consideration the wage indicators from the U.S. (c.a 20 \$/h) (United States Department of Labor, 2017), the Netherlands (c.a 30\$/h) (MijnZZP, 2018), and China (c.a 5 \$/h) (Salary Expert, 2018).

Formatted Table

Formatted: Font: Not Bold, Italic

Formatted: Font: Italic

Formatted: Font: Not Bold, Italic

Formatted: Font: Not Bold, Italic

Formatted: Font: Not Bold, Italic

Formatted: Font: Not Bold, Italic

Formatted: Font: Not Bold, Italic

Field Code Changed

Formatted: Font: Italic

Formatted: Font: Not Bold, Italic

Field Code Changed

Formatted: Font: Italic

Field Code Changed

Formatted: Font: Italic

Formatted: Font: Not Bold, English (U.K.)

S4. Yearly variable input data

	Global Phosphate Ore extraction cost (World Bank, 2018a) ² [\$ t ⁻¹]	Global Phosphate Ore production (USGS, 2016) ³ [kt]	Food Price Index (FAO, 2018a) ⁴ [-]	Diesel Fuel Price (U.S. Energy Information Administration, 2018) ⁵ [\$ gal ⁻¹]	Bunker Fuel Price (Institut National de la Statistique et des Etudes Economiques, 2017) ⁶ [\$ t ⁻¹]
2005	42.00	147,000	118	2.402	248
2006	44.21	142,000	127	2.705	290
2007	70.93	156,000	161	2.885	341
2008	345.59	161,000	201	3.803	522
2009	121.66	166,000	160	2.467	355
2010	123.02	181,000	188	2.992	464
2011	184.90	198,000	230	3.840	642
2012	185.89	217,000	213	3.968	672
2013	148.11	225,000	210	3.922	613
2014	110.22	218,000	202	3.825	546
2015	117.46	241,000	164	2.707	291

Table S4. Model yearly variable input data

² <http://pubdocs.worldbank.org/en/226371486076391711/CMO-Historical-Data-Annual.xlsx>

³ https://minerals.usgs.gov/minerals/pubs/commodity/phosphate_rock/

⁴ <http://www.fao.org/worldfoodsituation/foodpricesindex/en/>

⁵ https://www.eia.gov/dnav/pet/pet_pri_gnd_dcus_nus_a.htm

⁶ <https://www.insee.fr/en/statistiques/serie/001642883>

S5. Fixed Parameters

Yearly population growth rate	1.22	[%]	(World Bank, n.d.)
Yearly livestock growth rate	0.8	[%]	(FAO, 2018b)
Yearly agricultural expansion rate	3	[%]	[-]
Struvite recovery efficiency WWTP	0.2	[-]	Derived from (Jaffer et al., 2002)
Phosphorus density urine (for intracity transport)	0.00066	[-]	(Vinnerås, 2001)
Phosphorus density faeces (for intracity transport)	0.00457	[-]	(Vinnerås, 2001)
Phosphorus density rock phosphate	0.08	[-]	18% P ₂ O ₅ rock phosphate
Phosphorus density Struvite	0.14	[-]	-
Phosphorus density compost pellets	0.01	[-]	(Cofie and Nikiema, 2012)
Phosphorus density DAP	0.2	[-]	[-]
Price Magnesium Chloride	250	[\$ t ⁻¹]	(Seymour, 2009)
Scaling maint. savings per mass P recovered	0.89	[\$ kg ⁻¹]	(Shu et al., 2006)
Intracitiy collection distance	20	[km]	[-]
People serviced per WWTP	500,000	[p]	(Egle et al., 2016)
Annual costs Struvite precipitation	180,000	[\$ a ⁻¹]	(Egle et al., 2016)
Annual costs pelletizing facilities	20,000	[\$ a ⁻¹]	[-]
Pelletizing cost per mass influent	30	[\$ t ⁻¹]	(Masayuki Hara, 2001)
Annual costs mines (Inv. cost spread over 10 years)	3,100,000	[\$ a ⁻¹]	(World Bank, 2018b)

Table S5. Model yearly variable input data

SSS6. Livestock Phosphorus in Excrement

Global P	Year	Cattle	Swine	Poultry	Livestock Total
	[-]	[Mt]	[Mt]	[Mt]	[Mt]
This study	2006	11.22	3.66	2.51	17.39 (8.8)
Bouwman et al., 2013	2000	-	-	-	17
Sheldrick et al., 2003	1996	10.43	3.55	2.74	16.72

Table SSS6. Comparison of estimates of global phosphorus produced in the form of excrement by different livestock types, for different years. Estimates made following slaughter weight methodology proposed by (Sheldrick et al., 2003). In brackets for this study, shows contribution of major sites (production greater than 0.5 [t km⁻²] P and greater than 3 [kt] P total)

Formatted: Left: 1,25 cm, Top: 1,65 cm, Bottom: 1,65 cm, Width: 27,7 cm, Height: 3,3 cm

Formatted: Line spacing: 1,2

S7. Table of Sensitivity Analysis

Change in P Market Price with Following Changes in Parameter Value:											
Parameter Change	-50%	-40%	-30%	-20%	-10%	+10%	+20%	+30%	+40%	+50%	Sensitivity
Crop Parameter											
P Req. for optimal yield	6	4	3	2	1	-1	-1	-1	-2	-2	0.39
Optimal yield	-7	13	-2	-1	-1	1	1	2	3	3	0.26
Crop sale value	-7	13	-2	-1	-1	1	1	2	3	3	0.26
Proportion of fertilizer of production cost	-7	13	-2	-1	-1	1	1	2	3	3	0.26
Proportion P-fert. cost of total fert. cost	-7	13	-2	-1	-1	1	1	2	3	3	0.25
Recovery Parameters											
Recovery Efficiency Struvite Precipitation	0	0	0	0	0	0	0	0	0	0	0.00
P density of raw wastes	0	0	0	0	0	0	0	0	0	0	0.00
P density of recovered products	0	0	0	0	0	0	0	0	0	0	0.00
Magnesium cost for struvite	0	0	0	0	0	0	0	0	0	0	0.00
Scaling maintenance savings	0	0	0	0	0	0	0	0	0	0	0.00
Intracity waste collection distance	0	0	0	0	0	0	0	0	0	0	0.00
People serviced per recovery installation	0	0	0	0	0	0	0	0	0	0	0.00
Annual fixed cost per recovery installation	0	0	0	0	0	0	0	0	0	0	0.00
Annual cost mines operation	0	0	0	0	0	0	0	0	0	0	0.00
Transport parameters											
Truck fuel efficiency	0	0	0	0	0	0	0	0	0	0	0.00
Ship fuel efficiency	-12	-10	-7	-5	-3	2	5	7	10	12	0.24
Truck maximum carry load	9	6	4	2	1	-1	-1	-2	-3	-3	0.17
Ship maximum carry load	85	33	21	12	5	-5	-8	-11	-14	-16	1.00
Truck velocity	2	2	1	1	0	0	0	-1	-1	-1	0.05
Ship velocity	43	29	18	11	4	-4	-7	-10	-12	-14	0.87
Truck labour cost	-1	-1	-1	-1	0	0	0	1	1	1	0.02
Truck fixed costs	-3	-3	-2	-1	-1	1	1	2	2	3	0.06
Ship fixed cost (incl. Labour cost)	-11	-9	-7	-5	-2	2	4	6	9	11	0.22
Yearly input data											
Ore extraction cost	-21	-17	-13	-9	-4	4	8	13	17	21	0.43
Global ore production	0	0	0	0	0	0	0	0	0	0	0.00

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted Table

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Food commodity price index	0	0	0	0	0	0	0	0	0	0	0.00
Diesel prices	0	0	0	0	0	0	0	0	0	0	0.00
Bunker prices (ship fuel)	-12	-10	-7	-5	-3	2	5	7	10	12	0.24
Population growth rate	0	0	0	0	0	0	0	0	0	0	0.00
Livestock expansion rate	0	0	0	0	0	0	0	0	0	0	0.00
Agricultural expansion rate	0	0	0	0	0	0	0	0	0	0	0.00

Table S7. This table presents the results of a sensitivity analysis. It reveals how sensitive the model price predictions are to changes in parameter values (other parameters kept constant). A single ‘sensitivity’ value is determined for each parameter. A value of ‘1’ indicates that the model is highly sensitive to that value, where a (up to 50%) change in parameter value may result in an equal or greater percent change in model output. A value of ‘0’ indicates that the model price output is insensitive to (up to 50%) changes in parameter value. The table shows that the model is most sensitive to changes in transport parameters. Remarkably, the model market prices are relatively insensitive to changes in recovery costs. This is likely because of the insignificant share of recovered P on the total P market.

Formatted: Font: (Defa
New Roman, 10 pt

Formatted: Font: (Defa
New Roman, 10 pt

Formatted: Font: (Defa
New Roman, 10 pt

Formatted: Font: (Defa
New Roman, 10 pt


Formatted: Font: (Defa
New Roman, 10 pt

Formatted: Font: (Defa
New Roman, 10 pt

Formatted: Font: (Defa
New Roman, 10 pt

|

5



REFERENCES

- Arrow, K. J. and Debreu, G.: Existence of an Equilibrium for a Competitive Economy, *Econometrica*, 22(3), 265–290 [online] Available from: <http://links.jstor.org/sici?sici=0012-9682%28195407%2922%3A3%3C265%3AE0AEFA%3E2.0.CO%3B2-B> (Accessed 11 October 2017), 1954.
- 5 Bouwman, L., Goldewijk, K. K., Van Der Hoek, K. W., Beusen, A. H. W., Van Vuuren, D. P., Willems, J., Rufino, M. C. and Stehfest, E.: Exploring global changes in nitrogen and phosphorus cycles in agriculture induced by livestock production over the 1900-2050 period., *Proc. Natl. Acad. Sci. U. S. A.*, 110(52), 20882–7, doi:10.1073/pnas.1012878108, 2013.
- Cofie, O. and Nikiema, J.: Converting Faecal Sludge to Fertilizer Pellets: the case of Fortifer, [online] Available from: http://www.susana.org/_resources/documents/default/3-1781-22-1425990893.pdf (Accessed 21 July 2018), 2012.
- 10 Egle, L., Rechberger, H., Krampe, J. and Zessner, M.: Science of the Total Environment Phosphorus recovery from municipal wastewater: An integrated comparative technological, environmental and economic assessment of P recovery technologies, *Sci. Total Environ.*, 571, 522–542, doi:10.1016/j.scitotenv.2016.07.019, 2016.
- FAO: FAO Food Price Index | World Food Situation | Food and Agriculture Organization of the United Nations, [online] Available from: <http://www.fao.org/worldfoodsituation/foodpricesindex/en/> (Accessed 21 July 2018a), 2018.
- FAO: Livestock Patterns Data, [online] Available from: <http://www.fao.org/faostat/en/#data/EK/metadata> (Accessed 21 July 2018b), 2018.
- FAO: World Agriculture: Towards 2015/2030 - An FAO perspective, [online] Available from: <http://www.fao.org/docrep/005/y4252e/y4252e07.htm> (Accessed 15 August 2017), n.d.
- 15 IFDC (International Fertilizer Development Center) and UNIDO (United Nations Industrial Development Organisation): Fertiliser Manual, 3rd ed., Springer Verlag. [online] Available from: <https://books.google.nl/books?id=qPkoOU4BvEsC>, 1998.
- Institut National de la Statistique et des Etudes Economiques: International prices of Heavy fuel oil (Rotterdam) - Prices in US dollars per tonne - FOB - 1% of sulfur, [online] Available from: <https://www.insee.fr/en/statistiques/serie/001642883> (Accessed 21 July 2018), 2017.
- Jaffer, Y., Clark, T. ., Pearce, P. and Parsons, S. .: Potential phosphorus recovery by struvite formation, *Water Res.*, 36(7), 1834–1842, doi:10.1016/S0043-1354(01)00391-8, 2002.
- 20 Masayuki Hara: Fertilizer Pellets Made from Composted Livestock Manure, Mie Prefect. Sci. Technol. Promot. Cent. [online] Available from: <http://www.ffc.agnet.org/library.php?func=view&id=20110801154610> (Accessed 12 January 2018), 2001.
- MijnZZP: Beroep Vrachtwagenchauffeur, [online] Available from: <https://www.mijnzzp.nl/Beroep/608-Vrachtwagenchauffeur> (Accessed 5 October 2018), 2018.
- Nylund, N.-O. and Erkkilä, K.: Heavy-duty truck emission and fuel consumption: Simulating real-world driving in laboratory conditions, *Proceeding 2005 DEER Conf.*, 1–23 [online] Available from: http://www1.eere.energy.gov/vehiclesandfuels/pdfs/deer_2005/session5/2005_deer_erkkila.pdf, 2005.
- 25 Počuča, M.: Methodology of day-to-day ship costs assessment, *Promet - Traffic - Traffico*, 18(5), 337–345, 2006.
- Salary Expert: Truck Driver Salary in China, [online] Available from: <https://www.salaryexpert.com/salary/job/truck-driver/china> (Accessed 5 October 2018), 2018.
- Seymour, D.: Can Nutrient Recovery Be a Financially Sustainable Development Objective? P-Recovery Feasibility Study at HCTP, 2009.
- Sheldrick, W., Syers, J. K. and Lingard, J.: Contribution of livestock excreta to nutrient balances, *Nutr. Cycl. Agroecosystems*, 66, 119–131 [online] Available from: <https://link.springer.com/content/pdf/10.1023/A:1023944131188.pdf> (Accessed 1 May 2018), 2003.
- 30 Shu, L., Schneider, P., Jegatheesan, V. and Johnson, J.: An economic evaluation of phosphorus recovery as struvite from digester supernatant, *Bioresour. Technol.*, 97(17), 2211–2216, doi:10.1016/j.biortech.2005.11.005, 2006.
- United States Department of Labor: Occupational Employment and Wages, May 2017 Heavy and Tractor-Trailer Truck Drivers, *Occup. Employ. Stat.* [online] Available from: <https://www.bls.gov/oes/current/oes533032.htm> (Accessed 5 October 2018), 2017.
- United States Energy Information Administration: U.S. Gasoline and Diesel Retail Prices, [online] Available from: https://www.eia.gov/dnav/pet/pet_pri_gnd_dcus_nus_a.htm (Accessed 21 July 2018), 2018.
- 35 USGS (United States Geological Survey): Mineral Commodities Summaries Phosphate Rock, Washington, D.C., 2016.
- Vinnerås, B.: Faecal separation and urine diversion for nutrient management of household biodegradable waste and wastewater, *J. Biosci.*, 5(2), 29–50, 2001.
- World Bank: Commodity Markets: Annual prices, [online] Available from: <http://www.worldbank.org/en/research/commodity-markets> (Accessed 21 July 2018a), 2018.
- World Bank: Projects: Shidiya Phosphate Rock Mining Project | The World Bank, [online] Available from: <http://projects.worldbank.org/P005278/shidiya-phosphate-rock-mining-project?lang=en&tab=financial> (Accessed 21 July 2018b), 2018.
- 40 World Bank: Population growth (annual %) | Data, [online] Available from: <https://data.worldbank.org/indicator/SP.POP.GROW> (Accessed 3 January 2018), n.d.